

Math Assignment

MTH1098_01E
Zhang Yancheng, 11537668

October 28, 2024

Solutions:

1.

Proof:

$\forall \epsilon > 0$, assume $0 < |x - 2| < \delta$, set $f(x) = 4x - 3$ then:

$$|f(x) - 5| = |4x - 8| = 4|x - 2| < 4\delta$$

Choose $\delta = \frac{\epsilon}{4}$. This implies:

$$4|x - 2| < \epsilon$$

That is:

$$|x - 2| < \frac{\epsilon}{4}$$

Which is equivalent to:

$$|x - 2| < \epsilon$$

Therefore, $\forall \epsilon > 0$, we choose $\delta = \frac{\epsilon}{4}$, which satisfies the condition that if $0 < |x - 2| < \delta$, then $|f(x) - 5| < \epsilon$. This proves that $\lim_{x \rightarrow 2} (4x - 3) = 5$.

2.

- (a) The domain of $g(x)$ is $\{x : x \neq 1\}$.
- (b) $\lim_{x \rightarrow 1} g(x) = 3 \cdot x^2 + 2 \cdot x = 5$.
- (c) $f(x)$ is continuous at $x = 1$ because the limit exists and equals the function value.

3.

- (a) We know that the sine function is bounded between -1 and 1 for all real numbers. Therefore:

$$-1 \leq \sin\left(\frac{2}{x^2}\right) \leq 1$$

Multiply by x^2 :

$$-x^2 \leq x^2 \sin\left(\frac{2}{x^2}\right) \leq x^2$$

Apply the Squeeze Theorem:

$$\begin{aligned}\lim_{x \rightarrow 0} -x^2 &= 0 \\ \lim_{x \rightarrow 0} x^2 &= 0\end{aligned}$$

Therefore, we can conclude:

$$\lim_{x \rightarrow 0} x^2 \sin\left(\frac{2}{x^2}\right) = 0$$

- (b) Transfer the question to:

$$\lim_{x \rightarrow \infty} \frac{(\sqrt{x^2 + 2x} - \sqrt{x^2 - 2x})(\sqrt{x^2 + 2x} + \sqrt{x^2 - 2x})}{\sqrt{x^2 + 2x} + \sqrt{x^2 - 2x}}$$

This can be simplified to:

$$\begin{aligned} & \lim_{x \rightarrow \infty} \frac{(x^2 + 2x) - (x^2 - 2x)}{\sqrt{x^2 + 2x} + \sqrt{x^2 - 2x}} \\ &= \lim_{x \rightarrow \infty} \frac{4x}{\sqrt{x^2 + 2x} + \sqrt{x^2 - 2x}} \\ &= \lim_{x \rightarrow \infty} \frac{4x}{x \left(\sqrt{1 + \frac{2}{x}} + \sqrt{1 - \frac{2}{x}} \right)} \\ &= \lim_{x \rightarrow \infty} \frac{4}{\sqrt{1 + \frac{2}{x}} + \sqrt{1 - \frac{2}{x}}} \end{aligned}$$

As x approaches infinity, $\frac{2}{x}$ approaches 0. That is :

$$\begin{aligned} & \frac{4}{\sqrt{1+0} + \sqrt{1-0}} \\ &= \frac{4}{1+1} \\ &= 2 \end{aligned}$$

So, the limit is:

$$\lim_{x \rightarrow \infty} (\sqrt{x^2 + 2x} - \sqrt{x^2 - 2x}) = 2$$

- (c) As t approaches infinity, $\frac{1}{t}$ approaches 0, so:

The Nominator:

$$1 - \frac{t}{t+1} = 1 - \frac{t}{t(1 + \frac{1}{t})} = 1 - \frac{1}{1 + \frac{1}{t}} = 1 - \frac{1}{1+0} = 1 - 1 = 0$$

The Denominator:

$$1 - \sqrt{\frac{t}{t+1}} = 1 - \sqrt{\frac{t}{t(1 + \frac{1}{t})}} = 1 - \sqrt{\frac{1}{1 + \frac{1}{t}}} = 1 - \sqrt{\frac{1}{1+0}} = 1 - \sqrt{1} = 1 - 1 = 0$$

Giving the indeterminate form:

$$\lim_{t \rightarrow \infty} \frac{\left(1 - \frac{t}{t+1}\right)}{\left(1 - \sqrt{\frac{t}{t+1}}\right)} = \frac{0}{0}$$

Clearly, applying L'Hôpital's rule:

$$\lim_{t \rightarrow \infty} \frac{\frac{d}{dt} \left(1 - \frac{t}{t+1}\right)}{\frac{d}{dt} \left(1 - \sqrt{\frac{t}{t+1}}\right)}$$

Differentiate the Numerator:

$$\frac{d}{dt} \left(1 - \frac{t}{t+1}\right) = \frac{d}{dt} \left(\frac{t+1-t}{t+1}\right) = \frac{d}{dt} \left(\frac{1}{t+1}\right) = -\frac{1}{(t+1)^2}$$

Differentiate the Denominator:

$$\frac{d}{dt} \left(1 - \sqrt{\frac{t}{t+1}}\right) = \frac{d}{dt} \left(\sqrt{\frac{t}{t+1}}\right)$$

Using the chain rule:

$$\frac{d}{dt} \left(\sqrt{\frac{t}{t+1}}\right) = \frac{1}{2\sqrt{\frac{t}{t+1}}} \cdot \frac{d}{dt} \left(\frac{t}{t+1}\right)$$

$$\frac{d}{dt} \left(\frac{t}{t+1} \right) = \frac{(t+1) \cdot 1 - t \cdot 1}{(t+1)^2} = \frac{1}{(t+1)^2}$$

So:

$$\frac{d}{dt} \left(\sqrt{\frac{t}{t+1}} \right) = \frac{1}{2\sqrt{\frac{t}{t+1}}} \cdot \frac{1}{(t+1)^2}$$

Combine the Results:

$$\lim_{t \rightarrow \infty} \frac{1 - \frac{1}{t+1}}{1 - \sqrt{\frac{1}{t+1}}} = \lim_{t \rightarrow \infty} \frac{-\frac{1}{(t+1)^2}}{\frac{1}{2\sqrt{\frac{t}{t+1}}} \cdot \frac{1}{(t+1)^2}} = \lim_{t \rightarrow \infty} -2\sqrt{\frac{t}{t+1}} = \lim_{t \rightarrow \infty} -2\sqrt{\frac{1}{1 + \frac{1}{t}}} = -2$$

Therefore, the limit is

$$-2$$

- (d)

Calculate $(\sin(\frac{3\pi}{4}))$ and $(\cos(\frac{3\pi}{4}))$:

$$\sin\left(\frac{3\pi}{4}\right) = \frac{\sqrt{2}}{2}, \quad \cos\left(\frac{3\pi}{4}\right) = -\frac{\sqrt{2}}{2}$$

Thus,

$$\sin\left(\frac{3\pi}{4}\right) + \cos\left(\frac{3\pi}{4}\right) = \frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2} = 0.$$

Calculate $(\cos(2x))$ at $(x = \frac{3\pi}{4})$:

$$\cos(2x) = \cos\left(2 \cdot \frac{3\pi}{4}\right) = \cos\left(\frac{3\pi}{2}\right) = 0.$$

We have an indeterminate form $\frac{0}{0}$.

Thus, apply L'Hôpital's Rule:

$$\lim_{x \rightarrow \frac{3\pi}{4}} \frac{\cos x - \sin x}{-2 \sin(2x)}.$$

Evaluate at $x = \frac{3\pi}{4}$:

$$\begin{aligned} \cos\left(\frac{3\pi}{4}\right) - \sin\left(\frac{3\pi}{4}\right) &= -\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2} = -\sqrt{2} \\ \sin(2x) &= \sin\left(\frac{3\pi}{2}\right) = -1 \end{aligned}$$

Thus:

$$\lim_{x \rightarrow \frac{3\pi}{4}} \frac{-\sqrt{2}}{-2(-1)} = \frac{-\sqrt{2}}{2} = \frac{\sqrt{2}}{2}.$$

Therefore, the limit is

$$\frac{\sqrt{2}}{2}.$$

- (e) As $x \rightarrow 1$:

The numerator: $x \ln x \rightarrow 1 \cdot \ln(1) = 0$.

The denominator: $x^3 - 1 \rightarrow 1^3 - 1 = 0$.

This gives us the indeterminate form

$$\frac{0}{0}.$$

Thus, apply L'Hôpital's Rule

$$\lim_{x \rightarrow 1} \frac{x \ln x}{x^3 - 1} = \frac{\frac{d}{dx}(x \ln x) = \ln x + 1}{\frac{d}{dx}(x^3 - 1) = 3x^2}, = \lim_{x \rightarrow 1} \frac{\ln x + 1}{3x^2}.$$

Now, as $x \rightarrow 1$:

The numerator

$$\ln(1) + 1 \rightarrow 0 + 1 = 1.$$

The denominator

$$3(1^2) \rightarrow 3.$$

Thus, we have:

$$\lim_{x \rightarrow 1} \frac{\ln x + 1}{3x^2} = \frac{1}{3}.$$

Therefore, the limit is

$$\frac{1}{3}$$

4.

- (a) $\forall x \in (0, 4)$,

$$\therefore \frac{d}{dx}(\sqrt[3]{2x} + 3x - 4) > 0 \implies f(x) \nearrow,$$

and

$$\begin{cases} f(0) = -4 < 0 \\ f(4) = 10 > 0 \end{cases}$$

\therefore

$$\exists x \in (0, 4), \text{ that } f(x) = 0$$

- (b) The function $f(x)$ is continuous on $[0, 4]$ as all its terms are differentiable. Thus we calculate $f(0)$ and $f(4)$:

$$\begin{cases} f(0) = -4 \\ f(4) = 10 \end{cases}$$

According to MVT, $\exists x \in (0, 4)$, such that

$$f'(x) = \frac{f(4) - f(0)}{4 - 0} = \frac{7}{2}$$

\therefore There exist $c \in (0, 4)$ such that $f'(c) = 3\frac{1}{2}$

5.

Let

$$f(x) = \begin{cases} 11 + c^2x & \text{if } x < 2 \\ 1 - 6cx & \text{if } x \geq 2 \end{cases}$$

- (a) Find $\lim_{x \rightarrow 2^-} f(x)$:

$$\lim_{x \rightarrow 2^-} f(x) = 11 + 2c^2$$

- (b) Find $\lim_{x \rightarrow 2^+} f(x)$:

$$\lim_{x \rightarrow 2^+} f(x) = 1 - 12c$$

To ensure $f(x)$ is continuous at $x = 2$, set the limits equal:

$$11 + 2c^2 = 1 - 12c$$

Thus:

$$c = -1 \quad \text{or} \quad c = -5$$

6.

- (a) Rewrite the function:

$$f(x) = (4x + 3)^{1/2}$$

Using the Chain Rule:

$$f'(x) = \frac{1}{2}(4x + 3)^{-1/2} \cdot \frac{d}{dx}(4x + 3) = \frac{1}{2}(4x + 3)^{-1/2} \cdot 4 = \frac{2}{\sqrt{4x + 3}}$$

- (b)

The definition of the derivative is:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}$$

Calculating $f(x + h)$:

$$f(x + h) = \sqrt{4(x + h) + 3} = \sqrt{4x + 4h + 3}$$

Setting up the limit:

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{\sqrt{4x + 4h + 3} - \sqrt{4x + 3}}{h} = \lim_{h \rightarrow 0} \frac{(4x + 4h + 3) - (4x + 3)}{h(\sqrt{4x + 4h + 3} + \sqrt{4x + 3})} \\ &= \lim_{h \rightarrow 0} \frac{4h}{h(\sqrt{4x + 4h + 3} + \sqrt{4x + 3})} = \lim_{h \rightarrow 0} \frac{4}{\sqrt{4x + 4h + 3} + \sqrt{4x + 3}} = \frac{4}{2\sqrt{4x + 3}} = \frac{2}{\sqrt{4x + 3}} \end{aligned}$$

Therefore:

$$f'(x) = \frac{2}{\sqrt{4x + 3}}$$

7.

- (a)

$$\left(\frac{fg}{h}\right)' = \frac{f'g' \cdot h - fg \cdot h'}{h^2}$$

- (b) According to (a), we can calculate:

$$\left(\frac{x^2 \sin x}{e^x}\right)' = \frac{(2x \sin x + x^2 \cos x)e^x - (x^2 \sin x \cdot e^x)}{e^{2x}} = \frac{2x \sin x + x^2 \cos x - x^2 \sin x}{e^x}$$

8.

- (a)

$$\frac{d}{dx} \sqrt{5^x} = \frac{d}{dx} 5^{\frac{x}{2}} = \frac{1}{2} 5^{\frac{x}{2}} \ln(5) = \frac{\ln(5)}{2} \cdot 5^{\frac{x}{2}}$$

- (b)

$$\begin{aligned} \frac{d}{dx} e^{\tan(2x)} \ln(\sin x) &= (e^{\tan(2x)} \sec^2 2x \cdot 2)(\ln \sin x) + (e^{\tan(2x)} \cdot \frac{1}{\sin x} \cdot \cos x) \\ &= 2e^{\tan(2x)} \sec^2(2x) + e^{\tan(2x)} \cot x \end{aligned}$$

- (c) Differentiate the equation:

$$\frac{d}{dx}(xy^2) + \frac{d}{dx}(y \ln x) + \frac{d}{dx}(e^x) = 0$$

Simplify each element:

$$\frac{d}{dx}(xy^2) = x \cdot \frac{d}{dx}(y^2) + y^2 \cdot \frac{d}{dx}(x) = x(2y \frac{dy}{dx}) + y^2$$

$$\frac{d}{dx}(y \ln x) = \frac{dy}{dx} \ln x + y \cdot \frac{1}{x}$$

$$\frac{d}{dx}(e^x) = e^x$$

Thus

$$x(2y \frac{dy}{dx}) + y^2 + \left(\frac{dy}{dx} \ln x + \frac{y}{x} \right) + e^x = 0$$

This equals to:

$$\frac{dy}{dx}(2xy + \ln x) = - \left(y^2 + \frac{y}{x} + e^x \right)$$

Therefore:

$$\frac{dy}{dx} = \frac{-(y^2 + \frac{y}{x} + e^x)}{2xy + \ln x}$$

- (d) The derivative of $\sin^{-1}(u)$ is given by:

$$\frac{d}{du}[\sin^{-1}(u)] = \frac{1}{\sqrt{1-u^2}}$$

Here, $u = 4x$, so we apply the chain rule:

$$\frac{dy}{dx} = \frac{d}{du}[\sin^{-1}(u)] \cdot \frac{du}{dx}$$

Next, we find $\frac{du}{dx}$:

$$u = 4x \implies \frac{du}{dx} = 4$$

Substituting back into the derivative, we have:

$$\frac{dy}{dx} = \frac{1}{\sqrt{1-(4x)^2}} \cdot 4$$

Thus, the derivative is:

$$\frac{dy}{dx} = \frac{4}{\sqrt{1-16x^2}}$$

9.

Given:

- Ladder length (L) = 17 m
- Rate at which the foot is pulled away ($\frac{dx}{dt}$) = 0.8 m/s
- Distance from the wall (x) = 8 m

Using the Pythagorean theorem:

$$x^2 + y^2 = L^2 \implies y^2 = L^2 - x^2$$

Substituting $x = 8$:

$$y^2 = 17^2 - 8^2 \implies y = 15 \text{ m}$$

Differentiating:

$$x \frac{dx}{dt} + y \frac{dy}{dt} = 0 \implies \frac{dy}{dt} = -\frac{x}{y} \frac{dx}{dt}$$

Substituting values:


$$\frac{dy}{dt} = -\frac{8}{15} \cdot 0.8 = -\frac{6.4}{15} \approx -0.427 \text{ m/s}$$

Conclusion: The top of the ladder slides down at approximately 0.427 m/s.

10.

- **Domain:** $x \in (-\infty, -4) \cup (-4, 4) \cup (4, \infty)$
- **Intercepts:**
 - x -intercept: $(0, 0)$
 - y -intercept: $(0, 0)$
- **Asymptotes:**
 - Vertical: $x = -4, 4$
 - Horizontal: $y = 0$
- **Local Extrema:** None in the domain.

Graph:



11.

A piece of cardboard measures 2 m by 3 m. A square with side length x is cut from each corner. The volume V of the resulting box is given by:

$$V = x(2 - 2x)(3 - 2x) = 6x - 6x^2$$

To find the maximum volume, take the derivative:

$$\frac{dV}{dx} = 6 - 12x$$

Setting the derivative to zero gives:

$$12x = 6 \implies x = \frac{1}{2}$$

The dimensions of the box become:

- Length: $3 - 2\left(\frac{1}{2}\right) = 2$ m - Width: $2 - 2\left(\frac{1}{2}\right) = 1$ m - Height: $x = \frac{1}{2}$ m

The maximum volume is calculated as:

$$V = \left(\frac{1}{2}\right)(2)(1) = 1 \text{ m}^3$$

Thus, the dimensions of the box are $2 \text{ m} \times 1 \text{ m} \times 0.5 \text{ m}$ and the maximum possible volume is 1 m^3 .

After all, I would like to express my gratitude for your guidance throughout this course. I used LaTeX to format my assignment, which greatly helped me in presenting the mathematical concepts clearly.