

# Partial Differential Equations - Class Notes

Steven Blythe

January 21, 2022

# 1 Chapter 1

## Sidenotes

### 1.1 January 19, 2022

#### What is a PDE?

A PDE is an equation which contains partial derivatives of an unknown function and we want to find that unknown function.

Example:  $F(t, x, y, z, u, \frac{\partial u}{\partial t}, \frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial u}{\partial z}, \frac{\partial^2 u}{\partial t^2}, \frac{\partial^2 u}{\partial x \partial y}, \dots) = 0$ .

Note, the first partial derivatives are considered 1<sup>st</sup> ordered partials whereas the second ordered partials are considered 2<sup>nd</sup> ordered partials.

The variables that are not  $u$  are considered independent variables and  $u$  is considered a dependent variable.

What PDEs do we study?

Generally, we restrict our attention to equations that model some phenomenon from physics, engineering, economics, geology, ... etc. We can use physical intuition to help guide the math.

#### Classification of PDEs

1. Order of PDE: Highest derivative.

Example:  $\frac{\partial^3 u}{\partial x^3} - \sin(y)u^7 = 3$  is a third order PDE.

Example:  $(\frac{\partial y}{\partial t})^5 - \frac{\partial^2 y}{\partial x \partial t} = e^x$  is a second order PDE.

2. Number of independent variables.

Example:  $\frac{du}{dt} = \frac{\partial^2 u}{\partial x^2}$  has two independent variables:  $t, x$ .

This is the 1 -  $D$  heat equation.

Example:  $\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = \Delta u$  has 4 independent variables.

This is the 3 -  $D$  heat equation.  $\Delta u$  is Laplacian of  $u$ .

Notation

$\Delta u = \nabla^2 u = \nabla \cdot \nabla u = (\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z}) \cdot (\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial u}{\partial z}) = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2}$

$\Delta u = 0$  is considered Laplace's equation.

3. Linear vs non-linear

A linear PDE is any equation of the form  $L[u(x)] = f(x)$  where  $f(x)$  is a known function is a linear partial differential operator.

Definition: A differential operator is any rule that takes a function as its input and returns an expression that involves the derivatives of that function.

Example:

$$u(x, t) \quad v(x, t) \tag{1}$$

$$O[u] = \frac{\partial^2 u}{\partial x^2} + \sin x + \pi - 7e^{tu} \tag{2}$$

$$O[u + 3v] = \frac{\partial^2}{\partial x^2}(u + 3v) + \sin x + \pi - 7e^{tu+3tv} \tag{3}$$

$$= \frac{\partial^2 u}{\partial x^2} + 3\frac{\partial^2 v}{\partial x^2} + \sin x + \pi - 7e^{tu+3tv} \tag{4}$$

Definition: A linear operator,  $L$ , is an operator that has the property:

$$L[au + bv] = aL[u] + bL[v] \tag{5}$$

Where  $a$  and  $b$  are constants.

Theorem: If  $u$  and  $v$  are vectors and  $L$  is linear, then  $L$  can be represented by a matrix.

Theorem: If  $L$  is linear ordinary operator, it must take the form:

$$L[u] = f_0(x)u + f_1(x)u' + f_2(x)u'' + \dots + f_n(x)u^{(n)} \tag{6}$$

Where the  $f_i$ 's are known functions.

Definition: A linear ODE is any ODE of the form where  $f(x)$  is known is the following:

$$L[u] = f(x) \tag{7}$$

If  $f(x) = 0$ , then the equation is homogeneous. Otherwise, the equation is non-homogeneous.

Ex:  $(u')^2 = 0 \Rightarrow u' = 0 \rightarrow$  linear, homogeneous.

Theorem: If  $L$  is a linear partial differential operator, it must take the form ( $x$  is a vector with  $n$  unknowns)

$$L[u(x)] = f_0(x)u + \sum_{i=1}^n f_i(x)\frac{\partial u}{\partial x_i} + \sum_{i=1}^n \sum_{j=1}^n f_{ij}(x)\frac{\partial^2 u}{\partial x_i \partial x_j} + \dots \tag{8}$$

Definition: A linear PDE is any PDE of the form

$$L[u(x)] = f(x) \tag{9}$$

If  $f(x) = 0$ , the equation is homogeneous, else it is non-homogeneous.

Ex:  $u_t = 4u_x$  - Linear, homogeneous.

## 1.2 January 21, 2022

Example:

$$u_{tt} = u_{xx} + uyy \quad \text{Linear, homogeneous} \quad (10)$$

$$\cos(xt) = u + u_t + u_{xyz} \quad \text{Linear, non-homogeneous} \quad (11)$$

$$u_t u_{xt} = 0 \quad \text{non-linear} \quad (12)$$

$$u_{xt} + e^x \cos t \, u_t = 0 \quad \text{linear, homogeneous} \quad (13)$$

$$u_t + u_{xx} + ue^u = 0 \quad \text{non-linear} \quad (14)$$

Note: You can add linear combinations of solutions to linear homogeneous equations and still get a solution.

Example:  $u_x = u_t$ .

Some solutions to this are:

1.  $u_1(x, t) = 3$
  2.  $u_2(x, t) = x + t$
  3.  $u_3(x, t) = e^{x+t} \cos(x + t)$
  4.  $\vdots$
- $Au_1 + Bu_2 + Cu_3$  is also a solution.

### How do we solve an ODE?

1. Use some technique to find an explicit solution.
2. Use power series and determine the coefficients

$$y(x) = \sum_{n=0}^{\infty} a_n x^n \quad (15)$$

3. Laplace Transforms

### How do we solve PDEs?

1. Try to locate an explicit solution
2. We don't use power series, instead, we use a trigonometric series  $\Rightarrow$  Fourier Series.

$$y(x) = \sum_{n=0}^{\infty} a_n \sin(nx) + b_n \cos(nx) \quad (16)$$

3. Laplace Transforms are good if the domain is  $[0, \infty)$ .  
Fourier Transforms are good if the domain is  $(-\infty, \infty)$ .
4. Reduce the PDE to a system of ODEs.

### Initial Condition

1. For ODEs, to solve a 1<sup>st</sup> order equation, you need  $y(0)$ .  
2<sup>nd</sup> order  $\rightarrow y(0), y'(0)$   
3<sup>rd</sup> order  $\rightarrow y(0), y'(0), y''(0)$   
 $\vdots$   
 $n^{\text{th}}$  order  $\rightarrow y(0), y'(0), y''(0), \dots, y^{(n-1)}(0)$
2. For PDEs, it's more complicated  $\Rightarrow$  it depends on the PDE.  
Example:  $u(x, t), x \in [a, b], t \in [0, \infty)$   
If  $u_t = u_{xx}$
3. Boundary conditions:

$$u(a, t) = g_1(t) \quad (17)$$

$$u(b, t) = g_2(t) \quad (18)$$

If  $u_{tt} = u_{xx}$ , we must specify:

(a) Initial Conditions

$$u(x, 0) = f_1(x) \quad (19)$$

$$u_t(x, 0) = f_2(x) \quad (20)$$

(b) Boundary Conditions

$$u(a, t) = g_1(t) \quad (21)$$

$$u(b, t) = g_2(t) \quad (22)$$

### 1-D Heat Equation

Assume cross sections are uniform Imagine a cross section:

$$O \text{ } 0 \text{ } \text{=====} \text{ } 0 \text{ } L$$

Assume cross sections are uniform and the lateral sides are well insulated  $\Rightarrow$  heat only flows in the x-direction.

We need the following:

- $u(x, t)$  : Temperature of rod at position  $x$  and time  $t$ .
- $u(x, 0)$  : Initial temperature
- $u(0, t)$  and  $u(L, t)$  : Boundary Conditions

Definition:

- $g(x, t)$  : heat flux (energy / area time)
- $Q(x, t)$  : heat energy density (energy / volume)
- $A$  : Cross sectional area
- $C_P$  : Heat capacity or specific heat
- $\rho$  : Density
- $K$  : Thermal conductivity

We want to find an equation for the temperature evolution. We will use conservation of energy : Look at a little  $\Delta x$  section of the rod starting at  $x_0$ .

$$\begin{array}{c} \Delta x \\ 0 \text{=====} |0| \text{=====} 0 \\ x_0 \quad x_0 \Delta x \end{array}$$

Conservation of energy : heat in - heat out = heat accumulated

$$\text{Heat in} = \int_{t_0}^{t_0 + \Delta t} q A \Delta t' = A \int_{t_0}^{t_0 + \Delta t} q(x_0, t) \, dt$$

$$\text{Heat out} = A \int_{t_0}^{t_0 + \Delta t} q(x_0 + \Delta x, t) \, dt$$

$$\text{Heat Accumulated} = Q A \Delta x|_{t_0 + \Delta t} - Q A \Delta x|_{t_0}$$

$$= A \int_{x_0}^{x_0 + \Delta x} Q(x, t_0 + \Delta t) \, dx - A \int_{x_0}^{x_0 + \Delta x} Q(x, t_0) \, dx \quad (23)$$