

Fourier Series

$$f(x) = a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$$

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \, dx$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx$$

Laplace's Equation

Recall the Fourier Sine Series:

$$A_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) \, dx$$

$$A_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) \, dx$$

Recall general solutions: sine, exp, sinh

Fill in the blanks at last step.

Steady State Solution

Here, T is our u at the start or end, creates a line.

$$v(x) = \frac{T_2 - T_1}{L}x + T_1$$

$$u(x, t) = w(x, t) + v(x)$$

Continue with $w(x, t)$ using the interval.

$$w(x, t) = u(x, t) - v(x)$$

Use the appropriate condition (*e.g.* $u(x, 0)$)

Continue to solve as usual with w . Once finished,

$$u(x, t) = w(x, t) + v(x)$$