

**Example 1.** Let  $X$  be a zero-dimensional, compact  $L$ -space (hereditarily Lindeloff and non-separable). It is a fact that there exists a point-countable collection  $\mathcal{U} = \{U_\alpha : \alpha < \omega_1\}$  of clopen sets in  $X$ , and it is also true that any point-finite subcollection of  $\mathcal{U}$  is countable.

Let  $C = \{c_\alpha : \alpha < \omega_1\}$  be any uncountable subset of the Cantor space  $2^\omega$ . Let  $X_s = X \times \{s\}$  for each  $s \in 2^{<\omega}$ , and  $U_{\alpha,s} = U_\alpha \times \{s\}$ .

Finally, let

$$\mathbb{X} = C \cup \bigcup_{s \in 2^{<\omega}} X_s$$

be a tree of  $2^{<\omega}$  copies of  $X$ , and where

$$c_\alpha \cup \bigcup_{n < \omega} U_{\alpha, x_\alpha \upharpoonright n}$$

is an open set about each  $c_\alpha$ .

**Definition 2.** Let  $S \in [\omega_1]^{<\omega}$  and  $m < \omega$ . Define

$$\begin{aligned} K_S &= \bigcup_{\alpha \in S} \left( c_\alpha \cup \left( \bigcup_{s < c_\alpha} U_{\alpha,s} \right) \right) \\ A &= \{z \smallfrown \langle 1 \rangle : z \in 1^{<\omega}\} \\ K_S^* &= K_S \setminus \bigcup_{s \in A} X_s \end{aligned}$$

and

$$L_m = \bigcup_{s \in 2^{<m}} X_s$$

and observe that every compact set is dominated by  $K_S^* \cup L_m$  for some  $S, m$ . Intuitively,  $K_S^*$  collects the branches of  $U_\alpha$  converging up to  $c_\alpha$  for each  $\alpha \in S$  while avoiding copies  $X_s$  of  $X$  for each  $s$  in an antichain  $A$ , and  $L_m$  collects the copies  $X_s$  of  $X$  with  $|s| < m$  at the base of the tree.

**Proposition 3.** Without loss of generality,  $P$  always plays points in  $\bigcup_{s \in 2^{<\omega}} X_s$ .

**Proposition 4.**  $K \upharpoonright LF_{K,P}(\mathbb{X})$ .

*Proof.* In response to a point  $\langle x, s \rangle$ ,  $K$  observes that there are only countably many  $\alpha$  such that  $U_\alpha \times \{s\}$  contains  $\langle x, s \rangle$  (by point-countability of  $X$ ). Enumerate these as  $\alpha_n$ .  $K$  makes a promise that during round  $m$ ,  $K$  will forbid some superset of  $K_{\{\alpha_n : n \leq m\}}$ . Finally,  $K$  also always forbids a superset of  $L_{|s|+1}$ .

Suppose  $P$ 's moves clustered at some point. Since  $K$  forbade  $L_{|s|+1}$  during each round, that point must be  $c_\alpha$  for some  $\alpha$ .  $P$ 's play then must have included a subsequence of points  $\langle x_0, s_0 \rangle, \langle x_1, s_1 \rangle, \langle x_2, s_2 \rangle \dots$  such that  $x_n \in U_\alpha$  and  $s_n \leq s_{n+1} \leq c_\alpha$ . However, in response to  $\langle x_0, s_0 \rangle$ ,  $K$  made a promise to eventually forbid a superset of  $K_{\{\alpha\}}$ , making every  $\langle x_n, t_n \rangle$  illegal after that round.  $\square$

**Theorem 5.**  $K \not\vdash_{tact} LF_{K,P}(\mathbb{X})$ .

*Proof.* This is actually a corollary of G's theorem in [?]. The following is a direct game-theoretic proof.

Suppose that  $\sigma(\langle x, s \rangle)$  was a winning strategy for  $K$  and assume

$$\sigma(\langle x, s \rangle) = \bigcup_{|t| \leq |s|} \sigma(\langle x, t \rangle) = \sigma'(x, |s|)$$

Thus there exists some  $f : \omega_1 \rightarrow \omega$  such that  $\sigma'(x, f(\alpha))$  covers every neighborhood of  $c_\alpha$  for all  $x \in U_\alpha$ . (If not,  $P$  wins by taking the  $\alpha$  for which  $f$  is not defined, and may always play  $\langle x, s \rangle$  in a neighborhood of  $c_\alpha$  for which  $\sigma'(x, |s|)$  doesn't cover a neighborhood of  $c_\alpha$ .) Fix  $n$  for which  $f(\alpha) = n$  for  $\alpha$  in an uncountable set  $A$ .

Since the collection  $\{U_\alpha : \alpha \in A\}$  is uncountable, it is not point-finite. Fix  $x$  so that  $x$  belongs to  $U_\alpha$  for all  $\alpha$  in an infinite  $B \subseteq A$ . Finally, consider  $\sigma'(x, n)$ . For each  $\alpha \in B$ ,  $\sigma'(x, f(\alpha)) = \sigma'(x, n)$  covers  $c_\alpha$ . Since  $\{c_\alpha : \alpha \in B\}$  is a closed infinite discrete set, we have a contradiction to the compactness of  $\sigma'(x, n)$ .  $\square$

**Theorem 6.**  $K \not\vdash_{2-tact} LF_{K,P}(\mathbb{X})$ .

*Proof.* Suppose  $\sigma(\langle x, s \rangle, \langle y, t \rangle)$  was a winning 2-tactical strategy. We may define  $S(x, y, n) \in [\omega_1]^{<\omega}$  (increasing on  $n$ ) and  $n < m(x, y, n) < \omega$  such that for each  $(x, y)$ ,

$$\bigcup_{s, t \in 2^{\leq n}} \sigma(\langle x, s \rangle, \langle y, t \rangle) \subseteq K_{S(x, y, n)}^* \cup L_{m(x, y, n)}$$

and so we assume

$$\sigma(\langle x, s \rangle, \langle y, t \rangle) = K_{S(x, y, \max(|s|, |t|))}^* \cup L_{m(x, y, \max(|s|, |t|))}$$

Select an arbitrary point  $x' \in X$ . We define a tactical strategy

$$\tau(x, s) = K_{S(x, x', m(x, x', |s|)+1)}^* \cup L_{m(x, x', m(x, x', |s|)+1)}$$

We complete the proof by showing  $\tau$  is a winning tactical strategy (a contradiction).

Suppose

$$\langle x_0, s_0 \rangle, \langle x_1, s_1 \rangle, \langle x_2, s_2 \rangle, \dots$$

successfully countered  $\tau$  by clustering at  $c \in C$  (the strategy trivially prevents clustering elsewhere). Let  $z_n = \langle 0, \dots, 0 \rangle$  with  $n$  zeros. We claim

$$\langle x_0, s_0 \rangle, \langle x', z_{m(x_0, x', |s_0|)} \cap \langle 1 \rangle \rangle, \langle x_1, s_1 \rangle, \langle x', z_{m(x_1, x', |s_1|)} \cap \langle 1 \rangle \rangle, \langle x_2, s_2 \rangle, \langle x', z_{m(x_2, x', |s_2|)} \cap \langle 1 \rangle \rangle, \dots$$

is a successful counter to  $\sigma$ .

We will need the fact that, as  $\langle x_{i+1}, s_{i+1} \rangle$  was legal against  $\tau$ :

$$\begin{aligned} |s_i| &< m(x_i, x', |s_i|) + 1 = |z_{m(x_i, x', |s_i|)} \frown \langle 1 \rangle| \\ &< m(x_i, x', m(x_i, x', |s_i|) + 1) = m(x_i, x', |z_{m(x_i, x', |s_i|)} \frown \langle 1 \rangle|) \leq |s_{i+1}| \end{aligned}$$

Note that  $m(x, y, \max(|s|, |t|))$  is increasing throughout this play of the game versus  $\sigma$ :

$$\begin{aligned} &m(x_i, x', \max(|s_i|, |z_{m(x_i, x', |s_i|)} \frown \langle 1 \rangle|)) \\ &= m(x_i, x', |z_{m(x_i, x', |s_i|)} \frown \langle 1 \rangle|) \\ &\leq |s_{i+1}| \\ &< m(x_{i+1}, x', |s_{i+1}|) \\ &= m(x_{i+1}, x', \max(|s_{i+1}|, |z_{m(x_i, x', |s_i|)} \frown \langle 1 \rangle|)) \\ &= |z_{m(x_{i+1}, x', |s_{i+1}|)}| \\ &< |z_{m(x_{i+1}, x', |s_{i+1}|)} \frown \langle 1 \rangle| \\ &< m(x_{i+1}, x', |z_{m(x_{i+1}, x', |s_{i+1}|)} \frown \langle 1 \rangle|) \\ &= m(x_{i+1}, x', \max(|s_{i+1}|, |z_{m(x_{i+1}, x', |s_{i+1}|)} \frown \langle 1 \rangle|)) \end{aligned}$$

We turn to showing that  $\langle x', z_{m(x_{i+1}, x', |s_{i+1}|)} \frown \langle 1 \rangle \rangle$  is always a legal move. Since  $z_{m(x_{i+1}, x', |s_{i+1}|)} \frown \langle 1 \rangle$  is on the antichain avoided by any  $K^*$ , the problem is reduced to showing that this move isn't forbidden by

$$L_{m(x_{i+1}, x', \max(|s_{i+1}|, |z_{m(x_i, x', |s_i|)} \frown \langle 1 \rangle|))}$$

which we can see here:

$$m(x_{i+1}, x', \max(|s_{i+1}|, |z_{m(x_i, x', |s_i|)} \frown \langle 1 \rangle|)) = m(x_{i+1}, x', |s_{i+1}|) < |z_{m(x_{i+1}, x', |s_{i+1}|)} \frown \langle 1 \rangle|$$

We can conclude by showing that  $\langle x_{i+1}, s_{i+1} \rangle$  is always a legal move. We can see it avoids

$$L_{m(x_i, x', \max(|s_i|, |z_{m(x_i, x', |s_i|)} \frown \langle 1 \rangle|))}$$

since

$$m(x_i, x', \max(|s_i|, |z_{m(x_i, x', |s_i|)} \frown \langle 1 \rangle|)) = m(x_i, x', |z_{m(x_i, x', |s_i|)} \frown \langle 1 \rangle|) \leq |s_{i+1}|$$

Since  $\langle x_{i+1}, s_{i+1} \rangle$  was legal against  $\tau$ , it avoided

$$K_{S(x_h, x', m(x_h, x', |s_h|)+1)}^* = K_{S(x_h, x', \max(|s_h|, |z_{m(x_h, x', |s_h|)} \frown \langle 1 \rangle|))}^*$$

for  $h \leq i$ . And when  $h < i$ , we see it avoids:

$$\begin{aligned} K_{S(x_{h+1}, x', \max(|s_{h+1}|, |z_{m(x_h, x', |s_h|)} \frown \langle 1 \rangle))}^* &= K_{S(x_{h+1}, x', |s_{h+1}|)}^* \\ &\subseteq K_{S(x_{h+1}, x', m(x_{h+1}, x', |s_{h+1}|)+1)}^* \end{aligned}$$

This concludes the proof.  $\square$

**Theorem 7.**  $K \nVdash_{k\text{-tact}} LF_{K,P}(\mathbb{X})$ .

*Proof.* The proof proceeds in parallel to the proof of  $K \nVdash_{2\text{-tact}} LF_{K,P}(\mathbb{X})$ .

Suppose  $\sigma(\langle x_0, s_0 \rangle, \dots, \langle x_k, s_k \rangle)$  was a winning  $(k+1)$ -tactical strategy. We may define  $S(x_0, \dots, x_k, n) \in [\omega_1]^{<\omega}$  (increasing on  $n$ ) and  $n < m(x_0, \dots, x_k, n) < \omega$  such that for each  $(x_0, \dots, x_k)$ ,

$$\bigcup_{s_0, \dots, s_k \in 2^{\leq n}} \sigma(\langle x_0, s_0 \rangle, \dots, \langle x_k, s_k \rangle) \subseteq K_{S(x_0, \dots, x_k, n)}^* \cup L_{m(x_0, \dots, x_k, n)}$$

and so we assume

$$\sigma(\langle x_0, s_0 \rangle, \dots, \langle x_k, s_k \rangle) = K_{S(x_0, \dots, x_k, \max(|s_0|, \dots, |s_k|))}^* \cup L_{m(x_0, \dots, x_k, \max(|s_0|, \dots, |s_k|))}$$

Select an arbitrary point  $x' \in X$ . Let  $M^0(x, n) = m(x, x', \dots, x', n)$  and  $M^{i+1}(x, n) = M^0(x, M^i(x, n) + 1)$ . We define a tactical strategy

$$\tau(x, s) = K_{S(x, x', \dots, x', M^{k-1}(x, |s|)+1)}^* \cup L_{m(x, x', \dots, x', M^{k-1}(x, |s|)+1)}$$

We complete the proof by showing  $\tau$  is a winning tactical strategy (a contradiction).

Suppose

$$\langle x_0, s_0 \rangle, \langle x_1, s_1 \rangle, \langle x_2, s_2 \rangle, \dots$$

successfully countered  $\tau$  by clustering at  $c \in C$  (the strategy trivially prevents clustering elsewhere). Let  $z_n = \langle 0, \dots, 0 \rangle$  with  $n$  zeros. We claim

$$\begin{aligned} &\langle x_0, s_0 \rangle, \langle x', z_{M^0(x_0, |s_0|)} \frown \langle 1 \rangle \rangle, \langle x', z_{M^1(x_0, |s_0|)} \frown \langle 1 \rangle \rangle, \dots, \langle x', z_{M^{k-1}(x_0, |s_0|)} \frown \langle 1 \rangle \rangle, \\ &\langle x_1, s_1 \rangle, \langle x', z_{M^0(x_1, |s_1|)} \frown \langle 1 \rangle \rangle, \langle x', z_{M^1(x_1, |s_1|)} \frown \langle 1 \rangle \rangle, \dots, \langle x', z_{M^{k-1}(x_1, |s_1|)} \frown \langle 1 \rangle \rangle, \dots \end{aligned}$$

is a successful counter to  $\sigma$ .

We will need the fact that, as  $\langle x_{i+1}, s_{i+1} \rangle$  was legal against  $\tau$ :

$$\begin{aligned} |s_i| &< M^0(x_i, |s_i|) + 1 = |z_{M^0(x_i, |s_i|)} \frown \langle 1 \rangle| < M^0(x_i, M^0(x_i, |s_i|) + 1) + 1 \\ &= M^1(x_i, |s_i|) + 1 = |z_{M^1(x_i, |s_i|)} \frown \langle 1 \rangle| < \dots < |z_{M^{k-1}(x_i, |s_i|)} \frown \langle 1 \rangle| \end{aligned}$$

$$= M^{k-1}(x_i, |s_i|) + 1 < m(x_i, x', \dots, x', M^{k-1}(x_i, |s_i|) + 1) \leq |s_{i+1}|$$

Note that  $m(x_0, \dots, x_k, \max(|s_0|, \dots, |s_k|))$  is increasing throughout this play of the game versus  $\sigma$ :

$$\begin{aligned}
& m(x_i, x', \dots, x', \max(|s_i|, |z_{M^0(x_i, |s_i|)} \frown \langle 1 \rangle|, \dots, |z_{M^{k-1}(x_i, |s_i|)} \frown \langle 1 \rangle|)) \\
&= m(x_i, x', \dots, x', |z_{M^{k-1}(x_i, |s_i|)} \frown \langle 1 \rangle|) \\
&= m(x_i, x', \dots, x', M^{k-1}(x_i, |s_i|) + 1) \\
&\leq |s_{i+1}| \\
&< M^0(x_{i+1}, |s_{i+1}|) \\
&= m(x_{i+1}, x', \dots, x', |s_{i+1}|) \\
&= m(x_{i+1}, x', \dots, x', \max(|s_{i+1}|, |z_{M^0(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle|, \dots, |z_{M^{k-1}(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle|)) \\
&= |z_{m(x_{i+1}, x', \dots, x', |s_{i+1}|)}| \\
&= |z_{M^0(x_{i+1}, |s_{i+1}|)}| \\
&< |z_{M^0(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle| \\
&< m(x_{i+1}, x', \dots, x', |z_{M^0(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle|) \\
&= m(x_{i+1}, x', \dots, x', \max(|s_{i+1}|, |z_{M^0(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle|, |z_{M^1(x_i, |s_i|)} \frown \langle 1 \rangle|, \dots, |z_{M^{k-1}(x_i, |s_i|)} \frown \langle 1 \rangle|)) \\
&\quad \vdots \\
&< m(x_{i+1}, x', \dots, x', \max(|s_{i+1}|, |z_{M^0(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle|, \dots, |z_{M^{k-1}(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle|))
\end{aligned}$$

We turn to showing that  $\langle x', z_{M^j(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle \rangle$  is always a legal move. Since  $z_{M^j(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle$  is on the antichain avoided by any  $K^*$ , the problem is reduced to showing that this move isn't forbidden by

$$\begin{aligned}
& Lm(x_{i+1}, x', \dots, x', \max(|s_{i+1}|, |z_{M^0(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle|, \dots, |z_{M^{j-1}(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle|, |z_{M^j(x_i, |s_i|)} \frown \langle 1 \rangle|, \dots, |z_{M^k(x_i, |s_i|)} \frown \langle 1 \rangle|)) \\
&= Lm(x_{i+1}, x', \dots, x', |z_{M^{j-1}(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle|)
\end{aligned}$$

which we can see here:

$$\begin{aligned}
& m(x_{i+1}, x', \dots, x', |z_{M^{j-1}(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle|) \\
&= m(x_{i+1}, x', \dots, x', M^{j-1}(x_{i+1}, |s_{i+1}|) + 1) \\
&= M^0(x_{i+1}, M^{j-1}(x_{i+1}, |s_{i+1}|) + 1) \\
&= M^j(x_{i+1}, s_{i+1})
\end{aligned}$$

$$< |z_{M^j(x_{i+1}, |s_{i+1}|)} \frown \langle 1 \rangle|$$

We can conclude by showing that  $\langle x_{i+1}, s_{i+1} \rangle$  is always a legal move. We can see it avoids

$$L_{m(x_i, x', \dots, x', \max(|s_i|, |z_{M^0(x_i, |s_i|)} \frown \langle 1 \rangle|, \dots, |z_{M^{k-1}(x_i, |s_i|)} \frown \langle 1 \rangle|))}$$

since

$$\begin{aligned} m(x_i, x', \dots, x', \max(|s_i|, |z_{M^0(x_i, |s_i|)} \frown \langle 1 \rangle|, \dots, |z_{M^{k-1}(x_i, |s_i|)} \frown \langle 1 \rangle|)) \\ = m(x_i, x', \dots, x', |z_{M^{k-1}(x_i, |s_i|)} \frown \langle 1 \rangle|) \\ = m(x_i, x', \dots, x', M^{k-1}(x_i, |s_i|) + 1) \\ \leq |s_{i+1}| \end{aligned}$$

Since  $\langle x_{i+1}, s_{i+1} \rangle$  was legal against  $\tau$ , it avoided

$$\begin{aligned} K_{S(x_h, x', \dots, x', M^{k-1}(x_h, |s_h|) + 1)}^* \\ = K_{S(x_h, x', \dots, x', \max(|s_h|, |z_{M^0(x_h, |s_h|)} \frown \langle 1 \rangle|, \dots, |z_{M^{k-1}(x_h, |s_h|)} \frown \langle 1 \rangle|))}^* \end{aligned}$$

for  $h \leq i$ . And when  $h < i$ , we see it avoids both:

$$\begin{aligned} K_{S(x_{h+1}, x', \dots, x', \max(|s_{h+1}|, |z_{M^0(x_{h+1}, |s_{h+1}|)} \frown \langle 1 \rangle|, \dots, |z_{M^{j-1}(x_{h+1}, |s_{h+1}|)} \frown \langle 1 \rangle|, |z_{M^j(x_h, |s_h|)} \frown \langle 1 \rangle|, \dots, |z_{M^k(x_h, |s_h|)} \frown \langle 1 \rangle|))}^* \\ = K_{S(x_{h+1}, x', \dots, x', |z_{M^{j-1}(x_{h+1}, |s_{h+1}|)} \frown \langle 1 \rangle|)}^* \\ = K_{S(x_{h+1}, x', \dots, x', M^{j-1}(x_{h+1}, |s_{h+1}|) + 1)}^* \\ \subseteq K_{S(x_{h+1}, x', \dots, x', M^{k-1}(x_{h+1}, |s_{h+1}|) + 1)}^* \end{aligned}$$

and:

$$\begin{aligned} K_{S(x_{h+1}, x', \dots, x', \max(|s_{h+1}|, |z_{M^0(x_h, |s_h|)} \frown \langle 1 \rangle|, \dots, |z_{M^k(x_h, |s_h|)} \frown \langle 1 \rangle|))}^* \\ = K_{S(x_{h+1}, x', \dots, x', |s_{k+1}|)}^* \\ \subseteq K_{S(x_{h+1}, x', \dots, x', M^{k-1}(x_{h+1}, |s_{h+1}|) + 1)}^* \end{aligned}$$

This concludes the proof. □