

# Limited information strategies for a topological proximal game

Spring Topology Conference 2015


Steven Clontz <http://stevenclontz.com>

Auburn, AL

May 13, 2015

## Game

Bell's absolutely proximal game  $Bell_{D,P}^{\rightarrow}(X)$  [1] (2014)



$\mathcal{D}$  wins the game if the points chosen by  $\mathcal{P}$  converge. Otherwise,  $\mathcal{P}$  wins.

If  $\mathcal{D} \uparrow Bell_{D,P}^{\rightarrow}(X)$ , then  $X$  is called an *absolutely proximal space*. “Absolutely proximal” is a strengthening of “proximal”, but these concepts coincide for (uniformly locally) compact spaces.

This game connects to a game of Gary Gruenhage: [1]

### Theorem

*Every proximal space is a  $W$ -space. So*  
 $\mathcal{D} \uparrow Bell_{D,P}^{\rightarrow}(X) \Rightarrow \mathcal{O} \uparrow Gru_{O,P}^{\rightarrow}(X, x) \text{ for all } x \in X.$

If  $\mathcal{D} \uparrow \text{Bell}_{D,P}^{\rightarrow}(X)$ , then  $X$  is called an *absolutely proximal space*. “Absolutely proximal” is a strengthening of “proximal”, but these concepts coincide for (uniformly locally) compact spaces.

This game connects to a game of Gary Gruenhage: [1]

### Theorem

*Every proximal space is a  $W$ -space. So*  
 $\mathcal{D} \uparrow \text{Bell}_{D,P}^{\rightarrow}(X) \Rightarrow \mathcal{O} \uparrow \text{Gru}_{O,P}^{\rightarrow}(X, x) \text{ for all } x \in X.$

Proximal spaces have strong preservation properties, as any closed subset or  $\Sigma$ -product of proximal spaces is proximal.

Since any metrizable space is proximal, and any proximal space is collectionwise normal, Bell's game gives an elegant proof of the classic result of Rudin and Gulko:

### Theorem

*A  $\Sigma$ -product of metrizable spaces is collection-wise normal.*

Proximal spaces have strong preservation properties, as any closed subset or  $\Sigma$ -product of proximal spaces is proximal.

Since any metrizable space is proximal, and any proximal space is collectionwise normal, Bell's game gives an elegant proof of the classic result of Rudin and Gulko:

### Theorem

*A  $\Sigma$ -product of metrizable spaces is collection-wise normal.*

Nyikos [4] observed that

### Proposition

*Corson compact spaces are proximal.*

and asked if the converse holds as well.

With Gruenhage, I showed that the answer is yes: [2]

### Theorem

*A compact space is Corson compact if and only if it is proximal.*

Nyikos [4] observed that

### Proposition

*Corson compact spaces are proximal.*

and asked if the converse holds as well.

With Gruenhage, I showed that the answer is yes: [2]

### Theorem

*A compact space is Corson compact if and only if it is proximal.*



Player  $\mathcal{D}$  chooses *entourages* of the diagonal: elements of a *uniformity* inducing the topology of the space.

A uniformity  $\mathbb{D}$  on  $X$  is a filter of subsets of  $X^2$  satisfying:

- $\bigcap \mathbb{D} = \Delta = \{\langle x, x \rangle : x \in X\}$
- $D \in \mathbb{D}$  implies  $D^{-1} = \{\langle y, x \rangle : \langle x, y \rangle \in D\} \in \mathbb{D}$
- for each  $D \in \mathbb{D}$  there is  $\frac{1}{2}D \in \mathbb{D}$  such that  $\frac{1}{2}D \circ \frac{1}{2}D \subseteq D$

The topology induced by a uniformity is the smallest topology such that  $D[x] = \{y : \langle x, y \rangle \in D\}$  is a neighborhood of  $x$  for each  $D \in \mathbb{D}$ .

Player  $\mathcal{D}$  chooses *entourages* of the diagonal: elements of a *uniformity* inducing the topology of the space.

A uniformity  $\mathbb{D}$  on  $X$  is a filter of subsets of  $X^2$  satisfying:

- $\bigcap \mathbb{D} = \Delta = \{\langle x, x \rangle : x \in X\}$
- $D \in \mathbb{D}$  implies  $D^{-1} = \{\langle y, x \rangle : \langle x, y \rangle \in D\} \in \mathbb{D}$
- for each  $D \in \mathbb{D}$  there is  $\frac{1}{2}D \in \mathbb{D}$  such that  $\frac{1}{2}D \circ \frac{1}{2}D \subseteq D$

The topology induced by a uniformity is the smallest topology such that  $D[x] = \{y : \langle x, y \rangle \in D\}$  is a neighborhood of  $x$  for each  $D \in \mathbb{D}$ .

Player  $\mathcal{D}$  chooses *entourages* of the diagonal: elements of a *uniformity* inducing the topology of the space.

A uniformity  $\mathbb{D}$  on  $X$  is a filter of subsets of  $X^2$  satisfying:

- $\bigcap \mathbb{D} = \Delta = \{\langle x, x \rangle : x \in X\}$
- $D \in \mathbb{D}$  implies  $D^{-1} = \{\langle y, x \rangle : \langle x, y \rangle \in D\} \in \mathbb{D}$
- for each  $D \in \mathbb{D}$  there is  $\frac{1}{2}D \in \mathbb{D}$  such that  $\frac{1}{2}D \circ \frac{1}{2}D \subseteq D$

The topology induced by a uniformity is the smallest topology such that  $D[x] = \{y : \langle x, y \rangle \in D\}$  is a neighborhood of  $x$  for each  $D \in \mathbb{D}$ .

Our goal is to obtain a purely topological characterization of the proximal properties.

As it turns out, the union of all uniformities inducing a topology is itself a uniformity inducing that topology, called the *universal uniformity*. Furthermore,  $\mathcal{D} \uparrow \text{Bell}_{\vec{D}, P}^{\rightarrow}(X)$  if and only if  $\mathcal{D}$  is required to choose from the universal uniformity.

This reduces our goal to characterizing the universal entourages.

Our goal is to obtain a purely topological characterization of the proximal properties.

As it turns out, the union of all uniformities inducing a topology is itself a uniformity inducing that topology, called the *universal uniformity*. Furthermore,  $\mathcal{D} \uparrow Bell_{\vec{D}, P}(X)$  if and only if  $\mathcal{D}$  is required to choose from the universal uniformity.

This reduces our goal to characterizing the universal entourages.

Our goal is to obtain a purely topological characterization of the proximal properties.

As it turns out, the union of all uniformities inducing a topology is itself a uniformity inducing that topology, called the *universal uniformity*. Furthermore,  $\mathcal{D} \uparrow Bell_{\vec{D}, P}(X)$  if and only if  $\mathcal{D}$  is required to choose from the universal uniformity.

This reduces our goal to characterizing the universal entourages.

From Willard's topology text [5], we have such a characterization:

### Theorem

*A neighborhood  $U$  of the diagonal is a universal uniformity if and only if there exist neighborhoods  $U_n$  for  $n < \omega$  where  $U = U_0$  and  $U_{n+1} \circ U_{n+1} \subseteq U_n$ .*

As a bonus, for strongly collectionwise normal ( $\Leftarrow$  paracompact) spaces, *all* neighborhoods of the diagonal have this property.

So we topologize Bell's game by saying an “entourage” is any (open symmetric) neighborhood of the diagonal with this property, and discard the need to consider a specific uniform structure.

From Willard's topology text [5], we have such a characterization:

### Theorem

*A neighborhood  $U$  of the diagonal is a universal uniformity if and only if there exist neighborhoods  $U_n$  for  $n < \omega$  where  $U = U_0$  and  $U_{n+1} \circ U_{n+1} \subseteq U_n$ .*

As a bonus, for strongly collectionwise normal ( $\Leftarrow$  paracompact) spaces, *all* neighborhoods of the diagonal have this property.

So we topologize Bell's game by saying an “entourage” is any (open symmetric) neighborhood of the diagonal with this property, and discard the need to consider a specific uniform structure.



From Willard's topology text [5], we have such a characterization:

### Theorem

*A neighborhood  $U$  of the diagonal is a universal uniformity if and only if there exist neighborhoods  $U_n$  for  $n < \omega$  where  $U = U_0$  and  $U_{n+1} \circ U_{n+1} \subseteq U_n$ .*

As a bonus, for strongly collectionwise normal ( $\Leftarrow$  paracompact) spaces, *all* neighborhoods of the diagonal have this property.

So we topologize Bell's game by saying an “entourage” is any (open symmetric) neighborhood of the diagonal with this property, and discard the need to consider a specific uniform structure.

A *perfect information strategy* uses full information of the previous moves of the opponent.  $(\mathcal{A} \uparrow G)$

A *k-tactical strategy* only uses the last  $k$  previous moves of the opponent.  $(\mathcal{A} \underset{k\text{-tact}}{\uparrow} G)$

A *k-Markov strategy* only uses the round number and the last  $k$  previous moves of the opponent.  $(\mathcal{A} \underset{k\text{-mark}}{\uparrow} G)$

If omitted, assume  $k = 1$ .

A *perfect information strategy* uses full information of the previous moves of the opponent.  $(\mathcal{A} \uparrow G)$

A *k-tactical strategy* only uses the last  $k$  previous moves of the opponent.  $(\mathcal{A} \underset{k\text{-tact}}{\uparrow} G)$

A *k-Markov strategy* only uses the round number and the last  $k$  previous moves of the opponent.  $(\mathcal{A} \underset{k\text{-mark}}{\uparrow} G)$

If omitted, assume  $k = 1$ .

A *perfect information strategy* uses full information of the previous moves of the opponent.  $(\mathcal{A} \uparrow G)$

A *k-tactical strategy* only uses the last  $k$  previous moves of the opponent.  $(\mathcal{A} \underset{k\text{-tact}}{\uparrow} G)$

A *k-Markov strategy* only uses the round number and the last  $k$  previous moves of the opponent.  $(\mathcal{A} \underset{k\text{-mark}}{\uparrow} G)$

If omitted, assume  $k = 1$ .

A *perfect information strategy* uses full information of the previous moves of the opponent.  $(\mathcal{A} \uparrow G)$

A *k-tactical strategy* only uses the last  $k$  previous moves of the opponent.  $(\mathcal{A} \underset{k\text{-tact}}{\uparrow} G)$

A *k-Markov strategy* only uses the round number and the last  $k$  previous moves of the opponent.  $(\mathcal{A} \underset{k\text{-mark}}{\uparrow} G)$

If omitted, assume  $k = 1$ .

Bell showed that a winning perfect information strategy may always be passed down to win in a closed subspace.

### Proposition

*If  $\mathcal{D} \uparrow Bell_{D,P}^{\rightarrow}(X)$ , then  $\mathcal{D} \uparrow Bell_{D,P}^{\rightarrow}(H)$  for every closed subspace  $H$  of  $X$ .*

This actually holds in general:

### Proposition

*If  $\mathcal{D} \underset{k\text{-tact}}{\uparrow} Bell_{D,P}^{\rightarrow}(X)$ , then  $\mathcal{D} \underset{k\text{-tact}}{\uparrow} Bell_{D,P}^{\rightarrow}(H)$  for every closed subspace  $H$  of  $X$ .*

### Proposition

*If  $\mathcal{D} \underset{k\text{-mark}}{\uparrow} Bell_{D,P}^{\rightarrow}(X)$ , then  $\mathcal{D} \underset{k\text{-mark}}{\uparrow} Bell_{D,P}^{\rightarrow}(H)$  for every closed subspace  $H$  of  $X$ .*

Bell showed that a winning perfect information strategy may always be passed down to win in a closed subspace.

### Proposition

*If  $\mathcal{D} \uparrow Bell_{D,P}^{\rightarrow}(X)$ , then  $\mathcal{D} \uparrow Bell_{D,P}^{\rightarrow}(H)$  for every closed subspace  $H$  of  $X$ .*

This actually holds in general:

### Proposition

*If  $\mathcal{D} \overset{k\text{-tact}}{\uparrow} Bell_{D,P}^{\rightarrow}(X)$ , then  $\mathcal{D} \overset{k\text{-tact}}{\uparrow} Bell_{D,P}^{\rightarrow}(H)$  for every closed subspace  $H$  of  $X$ .*

### Proposition

*If  $\mathcal{D} \overset{k\text{-mark}}{\uparrow} Bell_{D,P}^{\rightarrow}(X)$ , then  $\mathcal{D} \overset{k\text{-mark}}{\uparrow} Bell_{D,P}^{\rightarrow}(H)$  for every closed subspace  $H$  of  $X$ .*

Bell's also showed that winning strategies are preserved for  $\Sigma$ -products.

### Theorem

*If  $\mathcal{D} \uparrow \text{Bell}_{D,P}^{\rightarrow}(X_{\alpha})$  for  $\alpha < \kappa$ , then  $\mathcal{D} \uparrow \text{Bell}_{D,P}^{\rightarrow}(\sum_{\alpha < \kappa} X_{\alpha})$ .*

Idea of proof: during round  $n$ , consider the first  $n$  non-zero coordinates of the previous  $n$  moves by  $\mathcal{P}$  and use the winning strategies for those finite coordinates. Note that this uses the round number and perfect information of all previous moves.



Bell's also showed that winning strategies are preserved for  $\Sigma$ -products.

### Theorem

*If  $\mathcal{D} \uparrow \text{Bell}_{D,P}^{\rightarrow}(X_{\alpha})$  for  $\alpha < \kappa$ , then  $\mathcal{D} \uparrow \text{Bell}_{D,P}^{\rightarrow}(\sum_{\alpha < \kappa} X_{\alpha})$ .*

Idea of proof: during round  $n$ , consider the first  $n$  non-zero coordinates of the previous  $n$  moves by  $\mathcal{P}$  and use the winning strategies for those finite coordinates. Note that this uses the round number and perfect information of all previous moves.

We settle for considering a countable product instead (but still require knowledge of the round number).

## Theorem

If  $\mathcal{D} \xrightarrow[k\text{-mark}]{\uparrow} \text{Bell}_{D,P}^{\rightarrow}(X_i)$  for  $i < \omega$ , then  $\mathcal{D} \xrightarrow[k\text{-mark}]{\uparrow} \text{Bell}_{D,P}^{\rightarrow}(\prod_{i < \omega} X_i)$ .

Probable counter-example for generalization:

$\mathcal{D} \not\xrightarrow[\text{mark}]{\uparrow} \text{Bell}_{D,P}^{\rightarrow}(\sum_{\alpha < \omega_1} 2)$ ?

Existence of a winning limited information strategy characterizes a stronger topological property than the existence of a winning perfect information strategy.

As it turns out:

### Theorem

*A compact space  $X$  is strongly Eberlein compact if and only if*  
 $\mathcal{D} \uparrow_{tact} Bell_{D,P}^{\rightarrow}(X).$

Existence of a winning limited information strategy characterizes a stronger topological property than the existence of a winning perfect information strategy.

As it turns out:

### Theorem

*A compact space  $X$  is strongly Eberlein compact if and only if*

$$\mathcal{D} \underset{\text{tact}}{\uparrow} \text{Bell}_{D,P}^{\rightarrow}(X).$$

# Sketch of Proof

Easy direction:

## Definition

Strong Eberlein compacts embed in  $\sigma 2^\kappa$  for some  $\kappa$ .

## Lemma

$\mathcal{D} \uparrow_{tact} Bell_{D,P}^{\rightarrow}(\sigma 2^\kappa)$ .

## Sketch of Proof (cont.)

Lemmas which give the other direction:

### Lemma (Gruenhage [3])

*Scattered proximal compacts are strong Eberlein compact.*

### Lemma

*Non-scattered proximal compacts contain copies of the Cantor space  $2^\omega$ .*

### Lemma

$\mathcal{D} \not\downarrow_{\text{tact}} \text{Bell}_{D,P}^{\rightarrow}(2^\omega).$

Any questions?



Jocelyn R. Bell.

An infinite game with topological consequences.  
*Topology Appl.*, 175:1–14, 2014.



Steven Clontz and Gary Gruenhage.

Proximal compact spaces are Corson compact.  
*Topology Appl.*, 173:1–8, 2014.



Gary Gruenhage.

Covering properties on  $X^2 \setminus \Delta$ ,  $W$ -sets, and compact subsets of  $\Sigma$ -products.  
*Topology Appl.*, 17(3):287–304, 1984.



Peter Nyikos.

Proximal and semi-proximal spaces.  
*Questions Answers Gen. Topology*, 32(2):79–91, 2014.



Stephen Willard.

*General topology*.  
Dover Publications, Inc., Mineola, NY, 2004.  
Reprint of the 1970 original [Addison-Wesley, Reading, MA; MR0264581].