

50TH SPRING TOPOLOGY AND DYNAMICAL SYSTEMS
CONFERENCE
CONTRIBUTED PROBLEMS IN SET-THEORETIC TOPOLOGY

1. PROBLEMS ON MONOTONE COVERING PROPERTIES

John E. Porter

Murray State University, Murray, Kentucky U.S.A.
jporter@murraystate.edu

Given a covering property \mathcal{P} , one can define a monotone version of the covering property by requiring an operator r that assigns the right kind of refinement to each acceptable open covering \mathcal{U} in such a way that $r(\mathcal{U})$ refines $r(\mathcal{V})$ whenever \mathcal{U} refines \mathcal{V} . For example, Popvassilev [4] called a space is *monotonically (countably) metacompact* if one can assign to every (countable) open cover \mathcal{U} a point-finite open cover $r(\mathcal{U})$ that refines \mathcal{U} so that $r(\mathcal{U})$ refines $r(\mathcal{V})$ whenever \mathcal{U} refines \mathcal{V} .

Chase and Gruenhage [2] showed that compact monotonically countably metacompact spaces are metrizable, and Gruenhage announced at the conference that he and Chase [3] have proven that separable monotonically countable metacompact spaces are also metrizable. Chase and Gruenhage's results simultaneously generalize similar results for proto-metrizable spaces and Moore spaces. Can Chase's and Gruenhage's results be furthered generalized? Recall that a space is orthocompact if every open cover has an interior preserving open refinement. That is, every open cover has an open refinement, with the further property that at any point, the intersection of all open sets in the refinement containing that point, is also open.

Problem 1.1. *Are compact monotonically orthocompact spaces metrizable?*

Remark 1.2. *The separable version of Problem 1.1 has a negative answer. Popvassilev [5] has shown the Sorgenfrey line is monotonically orthocompact.*

Recall that a space X is proto-metrizable if X is paracompact and has an ortho-base. Gartside and Moody [1] showed that a space is proto-metrizable if and only if X has a monotone operator r such that $r(\mathcal{U})$ star-refines \mathcal{U} . Popvassilev and Porter [6] showed that proto-metrizable spaces possess a monotone locally finite operator.

Problem 1.3. *Are metacompact spaces with an ortho-base monotonically (countably) metacompact?*

In general, it seems that little is known about the class of metacompact spaces with an ortho-base.

REFERENCES

- [1] P. M. Gartside, P. J. Moody, *A note on protometrizable spaces*, Topology Appl., **52** (1993), no.1, 1-9.

- [2] Chase, T., Gruenhage, G., *Compact monotonically metacompact spaces are metrizable*, *Compact monotonically metacompact spaces are metrizable*, Topology Appl. **160** (2013), no.1, 45–49.
- [3] Chase, T., Gruenhage, G., *Monotone covering properties and properties they imply, to appear*.
- [4] Popvassilev, S. G., $\omega_1 + 1$ is not monotonically countably metacompact, Questions and Answers in General Topology, **27** (2009), 133–135.
- [5] Popvassilev, S. G., *Versions of monotone covering properties*, 29th Summer Conference on Topology and its Applications July 2014, College of Staten Island, CUNY. Abstract retrieved from <http://at.yorku.ca/c/b/j/p/06.htm>.
- [6] Popvassilev, S. G., Porter, J. E., *On monotone paracompactness*, Topology Appl. **167** (2014), no. 19.

2. 2-MARKOV STRATEGIES IN SELECTION GAMES

Steven Clontz

The University of North Carolina at Charlotte, Charlotte, NC U.S.A.
 steven.clontz@gmail.com

Let $S_{fin}^\omega(\mathcal{A}, \mathcal{B})$ be the statement that whenever $A_n \in \mathcal{A}$ for $n < \omega$, there exist $B_n \in [A_n]^{<\omega}$ such that $\bigcup\{B_n : n < \omega\} \in \mathcal{B}$. This *selection principle* characterizes a property of a topological space X when \mathcal{A}, \mathcal{B} are defined in terms of X . For example, if \mathcal{O}_X is the collection of open covers of X , then $S_{fin}^\omega(\mathcal{O}_X, \mathcal{O}_X)$ is the well-known Menger covering property.

This property may be made stronger by considering the following two-player game of length ω : $G_{fin}^\omega(\mathcal{A}, \mathcal{B})$. During each round $n < \omega$, the first player \mathcal{A} chooses $A_n \in \mathcal{A}$, followed by \mathcal{B} choosing $B_n \in [A_n]^{<\omega}$. \mathcal{B} wins the game if $\bigcup\{B_n : n < \omega\} \in \mathcal{B}$; otherwise \mathcal{A} wins. If \mathcal{B} has a *winning strategy* for the game (a function which defines a move for each finite sequence of previous moves by \mathcal{A} , and beats every possible response by \mathcal{A}), then we write $\mathcal{B} \uparrow G_{fin}^\omega(\mathcal{A}, \mathcal{B})$.

These concepts were first introduced by Scheepers in [1]. Of course, $\mathcal{B} \uparrow G_{fin}^\omega(\mathcal{A}, \mathcal{B}) \Rightarrow S_{fin}^\omega(\mathcal{A}, \mathcal{B})$, but the converse need not hold, since each B_n may be defined in $S_{fin}^\omega(\mathcal{A}, \mathcal{B})$ using knowledge of all A_n , not just those “previously played”. Thus for each topological property P characterized by $S_{fin}^\omega(\mathcal{A}, \mathcal{B})$, we denote the (possibly) stronger property $\mathcal{B} \uparrow G_{fin}^\omega(\mathcal{A}, \mathcal{B})$ as *strategic P*.

Such notions may be made even stronger using *limited information strategies*. A *k-Markov strategy* for \mathcal{B} uses only the last k moves of \mathcal{A} and the round number. When \mathcal{B} has a winning k -Markov strategy for $G_{fin}^\omega(\mathcal{A}, \mathcal{B})$, we write $\mathcal{B} \uparrow G_{fin}^\omega(\mathcal{A}, \mathcal{B})$. Similarly, for each topological property P characterized by $S_{fin}^\omega(\mathcal{A}, \mathcal{B})$, we denote property $\mathcal{B} \uparrow G_{fin}^\omega(\mathcal{A}, \mathcal{B})$ as *k-Markov P*.

In the case of the selection game $G_{fin}^\omega(\mathcal{A}, \mathcal{B})$, it may be shown that a $(k+2)$ -Markov strategy may always be improved to a 2-Markov strategy, as shown in [3] with regards to $G_{fin}^\omega(\mathcal{O}_X, \mathcal{O}_X)$.

The following natural question is open:

Question 2.1. *Do there exist (interesting/topological) \mathcal{A}, \mathcal{B} such that $\mathcal{B} \uparrow G_{fin}^\omega(\mathcal{A}, \mathcal{B})$ but $\mathcal{B} \not\uparrow G_{fin}^\omega(\mathcal{A}, \mathcal{B})$?*
2-mark

Consider the case that $\mathcal{A} = \mathcal{B} = \mathcal{O}_X$, i.e. the Menger game. The following summarize results from [2] [3] and [4].

Definition 2.2. For any cardinal κ , let $\kappa^\dagger = \kappa \cup \{\infty\}$ denote the one-point Lindelöfication of discrete κ , where points in κ are isolated, and the neighborhoods of ∞ are co-countable.

Proposition 2.3. $\mathcal{B} \uparrow G_{fin}^\omega(\mathcal{O}_{\kappa^\dagger}, \mathcal{O}_{\kappa^\dagger})$.

Definition 2.4. For two functions f, g we say f is almost compatible with g if $|\{x \in \text{dom}(f) \cap \text{dom}(g) : f(x) \neq g(x)\}| < \omega$.

Definition 2.5. $\mathcal{A}'(\kappa)$ states that there exists a collection of pairwise almost compatible finite-to-one functions $\{f_A \in \omega^A : A \in [\kappa]^{\leq \omega}\}$.

Theorem 2.6. $\mathcal{A}'(\omega_n)$ holds for all $n < \omega$.

Theorem 2.7. $\mathcal{A}'(\kappa)$ implies $\mathcal{B} \uparrow_{2\text{-mark}} G_{fin}^\omega(\mathcal{O}_{\kappa^\dagger}, \mathcal{O}_{\kappa^\dagger})$.

Theorem 2.8. For any cardinal κ , κ Cohen reals may be added to a model of $ZFC + CH$ while preserving $\mathcal{A}'(\mathfrak{c})$.

Theorem 2.9. There exists a model of ZFC where $\mathcal{A}'(\omega_\omega)$ fails.

Theorem 2.10. $\mathcal{B} \uparrow_{2\text{-mark}} G_{fin}^\omega(\mathcal{O}_{\omega_\omega^\dagger}, \mathcal{O}_{\omega_\omega^\dagger})$.

It remains open whether $\mathcal{B} \uparrow_{2\text{-mark}} G_{fin}^\omega(\mathcal{O}_{\omega_{\omega+1}^\dagger}, \mathcal{O}_{\omega_{\omega+1}^\dagger})$ might fail when $\mathcal{A}'(\omega_\omega)$ fails. Due to the above, any attempt to show $\mathcal{B} \not\uparrow_{2\text{-mark}} G_{fin}^\omega(\mathcal{O}_{\kappa^\dagger}, \mathcal{O}_{\kappa^\dagger})$ cannot happen solely within ZFC .

REFERENCES

- [1] Scheepers, M., *Combinatorics of open covers. I. Ramsey theory*. Topology Appl. 69 (1996), no. 1, 3162. MR1378387
- [2] Scheepers, M., *Concerning n -tactics in the countable-finite game*. J. Symbolic Logic 56 (1991), no. 3, 786794. MR1129143
- [3] Clontz, S., *Game-theoretic strengthenings of Menger's property*. Preprint.
- [4] Clontz, S., Dow, A., *Almost compatible functions and topological games*. Preprint.