

Example 1. $\mathcal{D} \uparrow_{\text{tact}} \text{Bell}_{D,P}^{\rightarrow}(\kappa^*)$, so $\mathcal{D} \uparrow_{\text{mark}} \text{Bell}_{D,P}^{\rightarrow}((\kappa^*)^\omega)$.

Definition 2. A space X is *strong Eberlein compact* if it embeds in $\sigma 2^\kappa = \{x \in 2^\kappa : |\{\alpha : x(\alpha) = 1\}| < \omega\}$.

Theorem 3 (Gruenhage). *For compact spaces X , X is strong Eberlein compact if and only if X is scattered and X is a W -space ($\mathcal{O} \uparrow \text{Gru}_{O,P}^{\rightarrow}(X, x)$ for all $x \in X$).*

Theorem 4. *If X is strong Eberlein compact, then $\mathcal{D} \uparrow_{\text{tact}} \text{Bell}_{D,P}^{\rightarrow}(X)$.*

Proof. Consider $\text{Bell}_{D,P}^{\rightarrow}(\sigma 2^\kappa)$. Let $\text{supp}(x) = \{\alpha : x(\alpha) = 1\} \in [\kappa]^{<\kappa}$.

Define the tactic σ for \mathcal{D} such that

$$\sigma(\langle x \rangle) = \bigcap \{P_\alpha(\Delta) : \alpha \in \text{supp}(x)\}$$

Fix a legal attack $p : \omega \rightarrow \sigma 2^\kappa$, and let $\alpha < \kappa$. If $p_\alpha : \omega \rightarrow \sigma 2^\kappa$ defined by $p_\alpha(n) = p(n)(\alpha)$ converges, then σ is a winning tactic. So assume $p_\alpha(n) \neq 0$ for all $n < \omega$. Then $p_\alpha(n) = 1$ for some n , and as $\alpha \in \text{supp}(p(n))$, $\sigma(p(n)) \subseteq P_\alpha(\Delta)$. As p is a legal attack, it follows that $p_\alpha(m) = p_\alpha(m+1)$ for all $m > n$, so p_α converges.

Since every strong Eberlein compact X embeds as a closed subset of $\sigma 2^\kappa$, it follows that $\mathcal{D} \uparrow_{\text{tact}} \text{Bell}_{D,P}^{\rightarrow}(X)$. \square

Theorem 5. *If X contains a copy of the Cantor set, then $\mathcal{D} \not\uparrow_{\text{tact}} \text{Bell}_{D,P}^{\rightarrow}(X)$.*

Proof. The result follows from showing that $\mathcal{D} \not\uparrow_{\text{tact}} \text{Bell}_{D,P}^{\rightarrow}(2^\omega)$ (any copy of the Cantor set within a Hausdorff space is a compact and thus closed subspace). Let σ be a tactic for \mathcal{D} in $\text{Bell}_{D,P}^{\rightarrow}(2^\omega)$ and let $D_k = \{\langle f, g \rangle : f \upharpoonright k = g \upharpoonright k\}$. Since $\{D_k : k < \omega\}$ is a base for the uniformity on 2^ω , we may fix $k(f) < \omega$ for each $f \in 2^\omega$ such that $D_{k(f)} \subseteq \sigma(\langle f \rangle)$.

Then there exists $k < \omega$ such that $\{f : k = k(f)\}$ is uncountable, and therefore there exist distinct f, g such that $k = k(f) = k(g)$ and $f \upharpoonright k = g \upharpoonright k$. Then $p : \omega \rightarrow 2^\omega$ defined by $p(2n) = f$ and $p(2n+1) = g$ is an attack against σ which obviously doesn't converge. This attack is legal since $f \in D_k[g] \subseteq \sigma(\langle g \rangle)[g]$ and $g \in D_k[f] \subseteq \sigma(\langle f \rangle)[f]$. \square

Lemma 6. *Every non-scattered Corson compact space contains a homeomorphic copy of the Cantor set.*

Proof. Every non-scattered space contains a closed subspace without isolated points. Let X be such a subspace, and assume that this Corson compact is embedded in $\Sigma \mathbb{R}^\kappa$. Let $B_{\alpha,\epsilon}(x) = \{y : d(x(\alpha), y(\alpha)) < \epsilon\}$. For each $x \in X$ and $n < \omega$, let $\beta(x, n) < \kappa$ be defined such that $\{\alpha : x(\alpha) \neq 0\} = \{\beta(x, n) : n < \omega\}$.

Choose an arbitrary $x_\emptyset \in X$ and $\epsilon_0 > 0$, and let $A_0 = \emptyset$.

Suppose then that for some $n < \omega$, $x_s \in X$ is defined for all $s \in 2^n$, and $\epsilon_n > 0$ and $A_n \in [\kappa]^{<\omega}$ are defined. Since each x_s is not isolated in X , let U_s be the open set

$$U_s = X \cap \bigcap_{\alpha \in A_{|s|}} B_{\alpha, \epsilon_{|s|}}(x_s)$$

and choose $x_{s \smallfrown \langle 0 \rangle}, x_{s \smallfrown \langle 1 \rangle} \in U_s$ distinct. Then let $\alpha_s < \kappa$ such that $x_{s \smallfrown \langle 0 \rangle}(\alpha_s) \neq x_{s \smallfrown \langle 1 \rangle}(\alpha_s)$. Let

$$A_{n+1} = \{\alpha_s : s \in 2^{\leq n}\} \cup \{\beta(x_s, i) : s \in 2^{\leq n}, i \leq n\}$$

Then choose $0 < \epsilon_{n+1} < \frac{1}{2}\epsilon_n$ such that

$$B_{\alpha_s, \epsilon_{n+1}}(x_{s \smallfrown \langle 0 \rangle}) \cap B_{\alpha_s, \epsilon_{n+1}}(x_{s \smallfrown \langle 1 \rangle}) = \emptyset$$

and

$$\overline{\bigcap_{\alpha \in A_{n+1}} B_{\alpha, \epsilon_{n+1}}(x_{s \smallfrown \langle 0 \rangle})} \cup \overline{\bigcap_{\alpha \in A_{n+1}} B_{\alpha, \epsilon_{n+1}}(x_{s \smallfrown \langle 1 \rangle})} \subseteq \bigcap_{\alpha \in A_n} B_{\alpha, \epsilon_n}(x_s)$$

for all $s \in 2^n$.

Let $x_f = \lim_{n < \omega} x_{f|n} \in X$ for each $f \in 2^\omega$. We claim $C = \{x_f : f \in 2^\omega\}$ is a copy of the Cantor set. This will follow if we can show that $\{U_s : s \in 2^{<\omega}\}$ is a base for C , since it has the structure of the Cantor tree.

Consider x_f for some $f \in 2^\omega$, and a subbasic open ball $B_{\alpha, \epsilon}(x_f)$. Observe that $x_f \in \bigcap_{n < \omega} U_{f|n}$ since $x_{f|n} \in U_{f|n}$ for all $n < \omega$.

If $\alpha \in \{\beta(x_s, n) : s \in 2^{<\omega}, n < \omega\}$, choose $k < \omega$ with $\alpha \in A_k$. Then choose $l < \omega$ such that $\epsilon_l < \epsilon$. Then $U_{f|(l+k)} \subseteq B_{\alpha, \epsilon}(x_f)$.

Otherwise, $x_g(\alpha) = 0$ for all $g \in 2^\omega$, and $C \subseteq B_{\alpha, \epsilon}(x_f)$. □

Corollary 7. *For compact spaces X , X is strong Eberlein compact if and only if $\mathcal{D} \uparrow_{tact} Bell_{D,P}^{\rightarrow}(X)$.*

Proof. Suppose X is not strong Eberlein compact; then X is either not a W -space or not scattered. If $\mathcal{D} \not\uparrow Bell_{D,P}^{\rightarrow}(X)$, then the result follows immediately, which only leaves non-scattered proximal compact to be considered. But non-scattered proximal compacts are non-scattered Corson compacts, and thus contain a copy of the Cantor set, so the result follows from Theorem 5. □