

# Game-theoretic strengthenings of Menger's property

AMS Fall Sectional Meeting at UNCG

Steven Clontz  
<http://stevenclontz.com>

Department of Mathematics and Statistics  
Auburn University

November 9, 2014

# The Menger property

## Definition

A space  $X$  is Menger if for every sequence  $\langle \mathcal{U}_0, \mathcal{U}_1, \dots \rangle$  of open covers of  $X$  there exists a sequence  $\langle \mathcal{F}_0, \mathcal{F}_1, \dots \rangle$  such that  $\mathcal{F}_n \in [\mathcal{U}_n]^{<\omega}$  and  $\bigcup_{n < \omega} \mathcal{F}_n$  is a cover of  $X$ .

## Proposition

$X$  is  $\sigma$ -compact  $\Rightarrow X$  is Menger  $\Rightarrow X$  is Lindelöf.

# The Menger property

## Definition

A space  $X$  is Menger if for every sequence  $\langle \mathcal{U}_0, \mathcal{U}_1, \dots \rangle$  of open covers of  $X$  there exists a sequence  $\langle \mathcal{F}_0, \mathcal{F}_1, \dots \rangle$  such that  $\mathcal{F}_n \in [\mathcal{U}_n]^{<\omega}$  and  $\bigcup_{n < \omega} \mathcal{F}_n$  is a cover of  $X$ .

## Proposition

$X$  is  $\sigma$ -compact  $\Rightarrow X$  is Menger  $\Rightarrow X$  is Lindelöf.

# The Menger game

## Game

Let  $Men_{\mathcal{C}, \mathcal{F}}(X)$  denote the *Menger game* with players  $\mathcal{C}$ ,  $\mathcal{F}$ . In round  $n$ ,  $\mathcal{C}$  chooses an open cover  $\mathcal{C}_n$ , followed by  $\mathcal{F}$  choosing  $\mathcal{F}_n \in [\mathcal{C}_n]^{<\omega}$ .

$\mathcal{F}$  wins the game ( $\mathcal{F} \uparrow Men_{\mathcal{C}, \mathcal{F}}(X)$ ) if  $\bigcup_{n < \omega} \mathcal{F}_n$  is a cover for the space  $X$ , and  $\mathcal{C}$  wins otherwise.

Theorem (Hurewicz 1926, effectively)

$X$  is Menger if and only if  $\mathcal{C} \nuparrow Men_{\mathcal{C}, \mathcal{F}}(X)$ .

# The Menger game

## Game

Let  $Men_{\mathcal{C},F}(X)$  denote the *Menger game* with players  $\mathcal{C}$ ,  $\mathcal{F}$ . In round  $n$ ,  $\mathcal{C}$  chooses an open cover  $\mathcal{C}_n$ , followed by  $\mathcal{F}$  choosing  $\mathcal{F}_n \in [\mathcal{C}_n]^{<\omega}$ .

$\mathcal{F}$  wins the game ( $\mathcal{F} \uparrow Men_{\mathcal{C},F}(X)$ ) if  $\bigcup_{n < \omega} \mathcal{F}_n$  is a cover for the space  $X$ , and  $\mathcal{C}$  wins otherwise.

## Theorem (Hurewicz 1926, effectively)

$X$  is Menger if and only if  $\mathcal{C} \nuparrow Men_{\mathcal{C},F}(X)$ .

Menger suspected that the subsets of the real line with his property were exactly the  $\sigma$ -compact spaces; however:

Theorem (Fremlin, Miller 1988)

*There are ZFC examples of non- $\sigma$ -compact subsets of the real line which are Menger.*

But metrizable non- $\sigma$ -compact Menger spaces will be *undetermined* for the Menger game.

Theorem (Telgarsky 1984, Scheepers 1995)

*Let  $X$  be metrizable.  $\mathcal{F} \uparrow \text{Men}_{C,F}(X)$  if and only if  $X$  is  $\sigma$ -compact.*

Menger suspected that the subsets of the real line with his property were exactly the  $\sigma$ -compact spaces; however:

### Theorem (Fremlin, Miller 1988)

*There are ZFC examples of non- $\sigma$ -compact subsets of the real line which are Menger.*

But metrizable non- $\sigma$ -compact Menger spaces will be *undetermined* for the Menger game.

### Theorem (Telgarsky 1984, Scheepers 1995)

*Let  $X$  be metrizable.  $\mathcal{F} \uparrow \text{Men}_{C,F}(X)$  if and only if  $X$  is  $\sigma$ -compact.*

Menger suspected that the subsets of the real line with his property were exactly the  $\sigma$ -compact spaces; however:

### Theorem (Fremlin, Miller 1988)

*There are ZFC examples of non- $\sigma$ -compact subsets of the real line which are Menger.*

But metrizable non- $\sigma$ -compact Menger spaces will be *undetermined* for the Menger game.

### Theorem (Telgarsky 1984, Scheepers 1995)

*Let  $X$  be metrizable.  $\mathcal{F} \uparrow \text{Men}_{C,F}(X)$  if and only if  $X$  is  $\sigma$ -compact.*



Note that for Lindelöf spaces, metrizability is characterized by regularity and second countability.

By considering winning *limited-information strategies*, it turns out Scheeper's proof essentially factors into two lemmas: one for regularity and one for second-countability.

Note that for Lindelöf spaces, metrizability is characterized by regularity and second countability.

By considering winning *limited-information strategies*, it turns out Scheeper's proof essentially factors into two lemmas: one for regularity and one for second-countability.

# Limited information strategies

## Definition

A *(perfect information) strategy* has knowledge of all the past moves of the opponent.

## Definition

A  *$k$ -Markov strategy* has knowledge of only the past  $k$  moves of the opponent and the round number.

# Limited information strategies

## Definition

A *(perfect information) strategy* has knowledge of all the past moves of the opponent.

## Definition

A  *$k$ -Markov strategy* has knowledge of only the past  $k$  moves of the opponent and the round number.

## Proposition

If  $X$  is  $\sigma$ -compact, then  $\mathcal{F} \underset{1\text{-mark}}{\uparrow} \text{Men}_{C,F}(X)$ .

## Proof.

Let  $X = \bigcup_{n < \omega} K_n$ . During round  $n$ ,  $\mathcal{F}$  picks a finite subcollection of the last open cover played by  $\mathcal{C}$  (the only one  $\mathcal{F}$  remembers) which covers  $K_n$ . □

In general that implication cannot be reversed.

### Definition

A subset  $Y$  of  $X$  is *relatively compact* if for every open cover for  $X$ , there exists a finite subcollection which covers  $Y$ .

### Proposition

If  $X$  is  $\sigma$ -relatively-compact, then  $\mathcal{F} \xrightarrow[1\text{-mark}]{\uparrow} \text{Men}_{C,F}(X)$ .

In general that implication cannot be reversed.

### Definition

A subset  $Y$  of  $X$  is *relatively compact* if for every open cover for  $X$ , there exists a finite subcollection which covers  $Y$ .

### Proposition

If  $X$  is  $\sigma$ -relatively-compact, then  $\mathcal{F} \xrightarrow[1\text{-mark}]{\uparrow} \text{Men}_{C,F}(X)$ .

## Theorem

$\mathcal{F} \uparrow$  1-mark  $\text{Men}_{C,F}(X)$  if and only if  $X$  is  $\sigma$ -relatively-compact.

## Proof.

Let  $\sigma(\mathcal{U}, n)$  represent a 1-Markov strategy. For every open cover  $\mathcal{U} \in \mathfrak{C}$ ,  $\sigma(\mathcal{U}, n)$  witnesses relative compactness for the set

$$R_n = \bigcap_{\mathcal{U} \in \mathfrak{C}} \bigcup \sigma(\mathcal{U}, n)$$

If  $X$  is not  $\sigma$ -relatively compact, fix  $x \notin R_n$  for any  $n < \omega$ . Then  $\mathcal{C}$  can beat  $\sigma$  by choosing  $\mathcal{U}_n \in \mathfrak{C}$  during each round such that  $x \notin \bigcup \sigma(\mathcal{U}_n, n)$ . □



## Theorem

$\mathcal{F} \uparrow$   
1-mark  $\text{Men}_{C,F}(X)$  if and only if  $X$  is  $\sigma$ -relatively-compact.

## Proof.

Let  $\sigma(\mathcal{U}, n)$  represent a 1-Markov strategy. For every open cover  $\mathcal{U} \in \mathfrak{C}$ ,  $\sigma(\mathcal{U}, n)$  witnesses relative compactness for the set

$$R_n = \bigcap_{\mathcal{U} \in \mathfrak{C}} \bigcup \sigma(\mathcal{U}, n)$$

If  $X$  is not  $\sigma$ -relatively compact, fix  $x \notin R_n$  for any  $n < \omega$ . Then  $\mathcal{C}$  can beat  $\sigma$  by choosing  $\mathcal{U}_n \in \mathfrak{C}$  during each round such that  $x \notin \bigcup \sigma(\mathcal{U}_n, n)$ . □

## Theorem

$\mathcal{F} \uparrow_{1\text{-mark}} Men_{C,F}(X)$  if and only if  $X$  is  $\sigma$ -relatively-compact.

## Proof.

Let  $\sigma(\mathcal{U}, n)$  represent a 1-Markov strategy. For every open cover  $\mathcal{U} \in \mathfrak{C}$ ,  $\sigma(\mathcal{U}, n)$  witnesses relative compactness for the set

$$R_n = \bigcap_{\mathcal{U} \in \mathfrak{C}} \bigcup \sigma(\mathcal{U}, n)$$

If  $X$  is not  $\sigma$ -relatively compact, fix  $x \notin R_n$  for any  $n < \omega$ . Then  $\mathcal{C}$  can beat  $\sigma$  by choosing  $\mathcal{U}_n \in \mathfrak{C}$  during each round such that  $x \notin \bigcup \sigma(\mathcal{U}_n, n)$ . □

## Corollary

For regular spaces  $X$ ,  $\mathcal{F} \uparrow \text{Men}_{C,F}(X)$  if and only if  $X$  is  $\sigma$ -compact.

1-mark

## Theorem

For second countable spaces  $X$ ,  $\mathcal{F} \uparrow \text{Men}_{C,F}(X)$  if and only if  $\mathcal{F} \uparrow \text{Men}_{C,F}(X)$ .

1-mark

## Corollary

For regular spaces  $X$ ,  $\mathcal{F} \overset{1\text{-mark}}{\uparrow} \text{Men}_{C,F}(X)$  if and only if  $X$  is  $\sigma$ -compact.

## Theorem

For second countable spaces  $X$ ,  $\mathcal{F} \overset{1\text{-mark}}{\uparrow} \text{Men}_{C,F}(X)$  if and only if  $\mathcal{F} \overset{1\text{-mark}}{\uparrow} \text{Men}_{C,F}(X)$ .

# Proof

It's sufficient to consider only basic open sets, and since  $X$  is a second-countable space, there are only countably many finite collections of basic open sets.

Let  $\sigma$  be a perfect information strategy, and suppose we've defined open covers  $\mathcal{U}_{s'}$  for  $s' \leq s \in \omega^{<\omega}$ . If  $\mathcal{U}$  is an arbitrary open cover, then there are only countably many choices for the finite subcollection

$$\sigma(\mathcal{U}_{s|1}, \dots, \mathcal{U}_s, \mathcal{U}) \subseteq \mathcal{U}$$

# Proof

It's sufficient to consider only basic open sets, and since  $X$  is a second-countable space, there are only countably many finite collections of basic open sets.

Let  $\sigma$  be a perfect information strategy, and suppose we've defined open covers  $\mathcal{U}_{s'}$  for  $s' \leq s \in \omega^{<\omega}$ . If  $\mathcal{U}$  is an arbitrary open cover, then there are only countably many choices for the finite subcollection

$$\sigma(\mathcal{U}_{s \upharpoonright 1}, \dots, \mathcal{U}_s, \mathcal{U}) \subseteq \mathcal{U}$$

## Proof (cont.)

Thus we may define open covers  $\mathcal{U}_{s \smallfrown \langle n \rangle}$  for each  $n < \omega$  such that for an arbitrary open cover  $\mathcal{U}$ ,

$$\sigma(\mathcal{U}_{s \upharpoonright 1}, \dots, \mathcal{U}_s, \mathcal{U}) = \sigma(\mathcal{U}_{s \upharpoonright 1}, \dots, \mathcal{U}_s, \mathcal{U}_{s \smallfrown \langle n \rangle})$$

for some  $n < \omega$ .

Let  $t : \omega \rightarrow \omega^{<\omega}$  be a bijection. We define a 1-Markov strategy  $\tau$  as follows:

$$\tau(\mathcal{U}, n) = \sigma(\mathcal{U}_{t(n) \upharpoonright 1}, \dots, \mathcal{U}_{t(n)}, \mathcal{U})$$

## Proof (cont.)

Thus we may define open covers  $\mathcal{U}_{s \smallfrown \langle n \rangle}$  for each  $n < \omega$  such that for an arbitrary open cover  $\mathcal{U}$ ,

$$\sigma(\mathcal{U}_{s \upharpoonright 1}, \dots, \mathcal{U}_s, \mathcal{U}) = \sigma(\mathcal{U}_{s \upharpoonright 1}, \dots, \mathcal{U}_s, \mathcal{U}_{s \smallfrown \langle n \rangle})$$

for some  $n < \omega$ .

Let  $t : \omega \rightarrow \omega^{<\omega}$  be a bijection. We define a 1-Markov strategy  $\tau$  as follows:

$$\tau(\mathcal{U}, n) = \sigma(\mathcal{U}_{t(n) \upharpoonright 1}, \dots, \mathcal{U}_{t(n)}, \mathcal{U})$$



## Proof (cont.)

Suppose there exists a counter-attack  $\langle \mathcal{V}_0, \mathcal{V}_1, \dots \rangle$  which defeats the 1-Markov strategy  $\tau$ . Then there exists  $f : \omega \rightarrow \omega$  such that, if  $\mathcal{V}^n = \mathcal{V}_{t^{-1}(f \upharpoonright n)}$

$$\begin{aligned} x &\notin \bigcup \tau(\mathcal{V}^n, t^{-1}(f \upharpoonright n)) \\ &= \bigcup \sigma(\mathcal{U}_{f \upharpoonright 1}, \dots, \mathcal{U}_{f \upharpoonright n}, \mathcal{V}^n) \\ &= \bigcup \sigma(\mathcal{U}_{f \upharpoonright 1}, \dots, \mathcal{U}_{f \upharpoonright n}, \mathcal{U}_{f \upharpoonright (n+1)}) \end{aligned}$$

Thus  $\langle \mathcal{U}_{f \upharpoonright 1}, \mathcal{U}_{f \upharpoonright 2}, \dots \rangle$  is a successful counter-attack by  $\mathcal{C}$  against the perfect information strategy  $\sigma$ , showing

$$\mathcal{O} \uparrow \text{Men}_{\mathcal{C}, F}(X) \Rightarrow \mathcal{O} \uparrow \underset{\text{1-mark}}{\text{Men}_{\mathcal{C}, F}(X)}.$$



## Proof (cont.)

Suppose there exists a counter-attack  $\langle \mathcal{V}_0, \mathcal{V}_1, \dots \rangle$  which defeats the 1-Markov strategy  $\tau$ . Then there exists  $f : \omega \rightarrow \omega$  such that, if  $\mathcal{V}^n = \mathcal{V}_{t^{-1}(f \upharpoonright n)}$

$$\begin{aligned} x &\notin \bigcup \tau(\mathcal{V}^n, t^{-1}(f \upharpoonright n)) \\ &= \bigcup \sigma(\mathcal{U}_{f \upharpoonright 1}, \dots, \mathcal{U}_{f \upharpoonright n}, \mathcal{V}^n) \\ &= \bigcup \sigma(\mathcal{U}_{f \upharpoonright 1}, \dots, \mathcal{U}_{f \upharpoonright n}, \mathcal{U}_{f \upharpoonright (n+1)}) \end{aligned}$$

Thus  $\langle \mathcal{U}_{f \upharpoonright 1}, \mathcal{U}_{f \upharpoonright 2}, \dots \rangle$  is a successful counter-attack by  $\mathcal{C}$  against the perfect information strategy  $\sigma$ , showing

$$\mathcal{O} \uparrow \text{Men}_{C,F}(X) \Rightarrow \mathcal{O} \uparrow_{\text{1-mark}} \text{Men}_{C,F}(X).$$



It's speculated that there are spaces  $X_k$  such that for the Banach-Mazur game,  $\mathcal{N} \uparrow_{k+1\text{-tact}} BM_{E,N}(X_k)$  but

$\mathcal{N} \not\uparrow_{k\text{-tact}} BM_{E,N}(X_k)$ . (This is true for  $k = 1$ .)

## Theorem

$\mathcal{F} \uparrow_{k+2\text{-mark}} Men_{C,F}(X)$  if and only if  $\mathcal{F} \uparrow_{2\text{-mark}} Men_{C,F}(X)$ .

## Proof.

$$\tau(\langle \mathcal{U}, \mathcal{V} \rangle, n+1) = \bigcup_{m < k+2} \sigma(\underbrace{\langle \mathcal{U}, \dots, \mathcal{U} \rangle}_{k+1-m}, \underbrace{\langle \mathcal{V}, \dots, \mathcal{V} \rangle}_{m+1}, (n+1)(k+2)+m)$$



It's speculated that there are spaces  $X_k$  such that for the Banach-Mazur game,  $\mathcal{N} \uparrow_{k+1\text{-tact}} BM_{E,N}(X_k)$  but

$\mathcal{N} \not\uparrow_{k\text{-tact}} BM_{E,N}(X_k)$ . (This is true for  $k = 1$ .)

## Theorem

$\mathcal{F} \uparrow_{k+2\text{-mark}} Men_{C,F}(X)$  if and only if  $\mathcal{F} \uparrow_{2\text{-mark}} Men_{C,F}(X)$ .

## Proof.

$$\tau(\langle \mathcal{U}, \mathcal{V} \rangle, n+1) = \bigcup_{m < k+2} \sigma(\underbrace{\langle \mathcal{U}, \dots, \mathcal{U} \rangle}_{k+1-m}, \underbrace{\langle \mathcal{V}, \dots, \mathcal{V} \rangle}_{m+1}, (n+1)(k+2)+m)$$



Having knowledge of *two* of an opponent's moves allows a player to react when the opponent changes her moves, something impossible to do using a 1-tactical or 1-Markov strategy.

### Definition

Let  $\kappa^\dagger = \kappa \cup \{\infty\}$  be the *one point Lindelöf-ication* of discrete  $\kappa$ : neighborhoods of  $\infty$  are exactly the co-countable sets containing it.

$\kappa^\dagger$  is a simple space which is a regular and Lindelöf, but not second-countable or  $\sigma$ -compact. Thus  $\mathcal{F} \not\uparrow_{1\text{-mark}} \text{Men}_{C,F}(\kappa^\dagger)$ , but

it's easy to see that  $\mathcal{F} \uparrow \text{Men}_{C,F}(\kappa^\dagger)$ . What about 2-Markov strategies?

Having knowledge of *two* of an opponent's moves allows a player to react when the opponent changes her moves, something impossible to do using a 1-tactical or 1-Markov strategy.

## Definition

Let  $\kappa^\dagger = \kappa \cup \{\infty\}$  be the *one point Lindelöf-ication* of discrete  $\kappa$ : neighborhoods of  $\infty$  are exactly the co-countable sets containing it.

$\kappa^\dagger$  is a simple space which is a regular and Lindelöf, but not second-countable or  $\sigma$ -compact. Thus  $\mathcal{F} \not\uparrow \text{Men}_{C,F}(\kappa^\dagger)$ , but  
 1-mark

it's easy to see that  $\mathcal{F} \uparrow \text{Men}_{C,F}(\kappa^\dagger)$ . What about 2-Markov strategies?

Having knowledge of *two* of an opponent's moves allows a player to react when the opponent changes her moves, something impossible to do using a 1-tactical or 1-Markov strategy.

## Definition

Let  $\kappa^\dagger = \kappa \cup \{\infty\}$  be the *one point Lindelöf-ication* of discrete  $\kappa$ : neighborhoods of  $\infty$  are exactly the co-countable sets containing it.

$\kappa^\dagger$  is a simple space which is a regular and Lindelöf, but not second-countable or  $\sigma$ -compact. Thus  $\mathcal{F} \not\uparrow \underset{1\text{-mark}}{Men_{C,F}(\kappa^\dagger)}$ , but

it's easy to see that  $\mathcal{F} \uparrow Men_{C,F}(\kappa^\dagger)$ . What about 2-Markov strategies?

The game  $Men_{C,F}(\kappa^\dagger)$  essentially involves choosing countable and finite subsets of  $\kappa$ . Conveniently, there already exists an infinite game also involving the countable and finite subsets of  $\kappa$  in the literature (Scheepers 1991). An adaptation follows:

### Game

Let  $Fill_{C,F}^\cap(\kappa)$  denote the *intersection filling game* with two players  $\mathcal{C}$ ,  $\mathcal{F}$ . In round 0,  $\mathcal{C}$  chooses  $C_0 \in [\kappa]^{\leq \omega}$ , followed by  $\mathcal{F}$  choosing  $F_0 \in [\kappa]^{< \omega}$ . In round  $n+1$ ,  $\mathcal{C}$  chooses  $C_{n+1} \in [\kappa]^{\leq \omega}$ , followed by  $\mathcal{F}$  choosing  $F_{n+1} \in [\kappa]^{< \omega}$ .  $\mathcal{F}$  wins the game if  $\bigcup_{n < \omega} F_n \supseteq \bigcap_{n < \omega} C_n$ ; otherwise,  $\mathcal{C}$  wins.



The game  $Men_{C,F}(\kappa^\dagger)$  essentially involves choosing countable and finite subsets of  $\kappa$ . Conveniently, there already exists an infinite game also involving the countable and finite subsets of  $\kappa$  in the literature (Scheepers 1991). An adaptation follows:

## Game

Let  $Fill_{C,F}^\cap(\kappa)$  denote the *intersection filling game* with two players  $\mathcal{C}$ ,  $\mathcal{F}$ . In round 0,  $\mathcal{C}$  chooses  $C_0 \in [\kappa]^{\leq \omega}$ , followed by  $\mathcal{F}$  choosing  $F_0 \in [\kappa]^{< \omega}$ . In round  $n+1$ ,  $\mathcal{C}$  chooses  $C_{n+1} \in [\kappa]^{\leq \omega}$ , followed by  $\mathcal{F}$  choosing  $F_{n+1} \in [\kappa]^{< \omega}$ .  $\mathcal{F}$  wins the game if  $\bigcup_{n < \omega} F_n \supseteq \bigcap_{n < \omega} C_n$ ; otherwise,  $\mathcal{C}$  wins.

## Theorem

$\mathcal{F} \uparrow_{k\text{-mark}} \text{Men}_{C,F}(\kappa^\dagger)$  if and only if  $\mathcal{F} \uparrow_{k\text{-mark}} \text{Fill}_{C,F}^\cap(\kappa)$ .

## Definition

For two functions  $f, g$  we say  $f$  is *almost compatible* with  $g$  if  $|\{x \in \text{dom}(f) \cap \text{dom}(g) : f(x) \neq g(x)\}| < \omega$ .

## Definition

$S(\kappa)$  states that there exist functions  $f_A : A \rightarrow \omega$  for each  $A \in [\kappa]^{\leq \omega}$  such that  $|f_A^{-1}(n)| < \omega$  for all  $n < \omega$  and  $f_A, f_B$  are almost compatible for all  $A, B \in [\kappa]^\omega$ .

## Theorem

$\mathcal{F} \xrightarrow[k\text{-mark}]{} \text{Men}_{C,F}(\kappa^\dagger)$  if and only if  $\mathcal{F} \xrightarrow[k\text{-mark}]{} \text{Fill}_{C,F}^\cap(\kappa)$ .

## Definition

For two functions  $f, g$  we say  $f$  is *almost compatible* with  $g$  if  $|\{x \in \text{dom}(f) \cap \text{dom}(g) : f(x) \neq g(x)\}| < \omega$ .

## Definition

$S(\kappa)$  states that there exist functions  $f_A : A \rightarrow \omega$  for each  $A \in [\kappa]^{\leq \omega}$  such that  $|f_A^{-1}(n)| < \omega$  for all  $n < \omega$  and  $f_A, f_B$  are almost compatible for all  $A, B \in [\kappa]^\omega$ .

## Theorem

$\mathcal{F} \xrightarrow[k\text{-mark}]{} \text{Men}_{C,F}(\kappa^\dagger)$  if and only if  $\mathcal{F} \xrightarrow[k\text{-mark}]{} \text{Fill}_{C,F}^\cap(\kappa)$ .

## Definition

For two functions  $f, g$  we say  $f$  is *almost compatible* with  $g$  if  $|\{x \in \text{dom}(f) \cap \text{dom}(g) : f(x) \neq g(x)\}| < \omega$ .

## Definition

$S(\kappa)$  states that there exist functions  $f_A : A \rightarrow \omega$  for each  $A \in [\kappa]^{\leq \omega}$  such that  $|f_A^{-1}(n)| < \omega$  for all  $n < \omega$  and  $f_A, f_B$  are almost compatible for all  $A, B \in [\kappa]^\omega$ .

## Theorem (Scheepers 1991)

$S(\omega_1); \neg S(\kappa)$  for  $\kappa > 2^\omega$ ;  $\text{Con}(S(2^\omega) + \neg CH)$ .

## Theorem

If  $S(\kappa)$ , then  $\mathcal{F} \underset{2\text{-mark}}{\uparrow} \text{Fill}_{C,F}^\cap(\kappa)$ .

## Theorem (Scheepers 1991)

$S(\omega_1); \neg S(\kappa)$  for  $\kappa > 2^\omega$ ;  $\text{Con}(S(2^\omega) + \neg CH)$ .

## Theorem

If  $S(\kappa)$ , then  $\mathcal{F} \xrightarrow[2\text{-mark}]{\uparrow} \text{Fill}_{C,F}^\cap(\kappa)$ .

## Theorem

If  $S(\kappa)$ , then  $\mathcal{F} \xrightarrow[2\text{-mark}]{} \text{Fill}_{C,F}^{\cap}(\kappa)$ .

## Proof.

Let  $f_A : A \rightarrow \omega$  witness  $S(\kappa)$ . Then we define the winning 2-Markov strategy  $\sigma$  as follows:

$$\sigma(\langle A \rangle, 0) = \{\alpha \in A : f_A(\alpha) = 0\}$$

$$\sigma(\langle A, B \rangle, n+1) = \{\alpha \in A \cap B : f_B(\alpha) \leq n+1 \text{ or } f_A(\alpha) \neq f_B(\alpha)\}$$



## Corollary

$\mathcal{F} \xrightarrow[2\text{-mark}]{} \text{Men}_{C,F}(\omega_1^{\dagger})$ , but  $\mathcal{F} \not\xrightarrow[1\text{-mark}]{} \text{Men}_{C,F}(\omega_1^{\dagger})$ .

## Theorem

If  $S(\kappa)$ , then  $\mathcal{F} \xrightarrow[2\text{-mark}]{} \text{Fill}_{C,F}^{\cap}(\kappa)$ .

## Proof.

Let  $f_A : A \rightarrow \omega$  witness  $S(\kappa)$ . Then we define the winning 2-Markov strategy  $\sigma$  as follows:

$$\sigma(\langle A \rangle, 0) = \{\alpha \in A : f_A(\alpha) = 0\}$$

$$\sigma(\langle A, B \rangle, n+1) = \{\alpha \in A \cap B : f_B(\alpha) \leq n+1 \text{ or } f_A(\alpha) \neq f_B(\alpha)\}$$



## Corollary

$\mathcal{F} \xrightarrow[2\text{-mark}]{} \text{Men}_{C,F}(\omega_1^{\dagger})$ , but  $\mathcal{F} \not\xrightarrow[1\text{-mark}]{} \text{Men}_{C,F}(\omega_1^{\dagger})$ .



# Characterizing 2-Markov strategies topologically

## Game

Let  $Men_{C,F}(X, Y)$  proceed analogously to the Menger game, except that  $\mathcal{F}$  only need cover  $Y \subseteq X$ .

## Definition

A subset  $Y$  of  $X$  is *relatively robustly Menger* if there exist functions  $r_{\mathcal{V}} : Y \rightarrow \omega$  for each open cover  $\mathcal{V}$  of  $X$  such that for all open covers  $\mathcal{U}, \mathcal{V}$  and numbers  $n < \omega$ , the following sets are finitely coverable by  $\mathcal{V}$ :

$$c(\mathcal{V}, n) = \{x \in Y : r_{\mathcal{V}}(x) \leq n\}$$

$$p(\mathcal{U}, \mathcal{V}, n+1) = \{x \in Y : n < r_{\mathcal{U}}(x) < r_{\mathcal{V}}(x)\}$$

# Characterizing 2-Markov strategies topologically

## Game

Let  $Men_{C,F}(X, Y)$  proceed analogously to the Menger game, except that  $\mathcal{F}$  only need cover  $Y \subseteq X$ .

## Definition

A subset  $Y$  of  $X$  is *relatively robustly Menger* if there exist functions  $r_{\mathcal{V}} : Y \rightarrow \omega$  for each open cover  $\mathcal{V}$  of  $X$  such that for all open covers  $\mathcal{U}, \mathcal{V}$  and numbers  $n < \omega$ , the following sets are finitely coverable by  $\mathcal{V}$ :

$$c(\mathcal{V}, n) = \{x \in Y : r_{\mathcal{V}}(x) \leq n\}$$

$$p(\mathcal{U}, \mathcal{V}, n+1) = \{x \in Y : n < r_{\mathcal{U}}(x) < r_{\mathcal{V}}(x)\}$$

## Theorem

*If  $Y$  is relatively robustly Menger to  $X$ , then*

$$\mathcal{F} \xrightarrow[2\text{-mark}]{\uparrow} \text{Men}_{C,F}(X, Y).$$

## Theorem

*If  $\mathcal{F} \xrightarrow[2\text{-mark}]{\uparrow} \text{Men}_{C,F}(X, X_i)$  for  $i < \omega$ , then*

$$\mathcal{F} \xrightarrow[2\text{-mark}]{\uparrow} \text{Men}_{C,F}(X, \bigcup_{i < \omega} X_i).$$

## Theorem

$$\text{Men}_{C,F}(X, X) = \text{Men}_{C,F}(X).$$

## Theorem

$S(\kappa)$  implies  $\omega_1^\dagger$  is relatively robustly Menger to itself.

## Definition

Let  $R_\omega$  be the real line with the topology generated by open intervals with countably many points removed.

## Theorem

$S(2^\omega)$  implies  $\mathcal{F} \underset{2\text{-mark}}{\uparrow} \text{Men}_{C,F}(R_\omega)$

## Theorem

$S(\kappa)$  implies  $\omega_1^\dagger$  is relatively robustly Menger to itself.

## Definition

Let  $R_\omega$  be the real line with the topology generated by open intervals with countably many points removed.

## Theorem

$S(2^\omega)$  implies  $\mathcal{F} \underset{2\text{-mark}}{\uparrow} \text{Men}_{C,F}(R_\omega)$

## Theorem

$S(2^\omega)$  implies  $\mathcal{F} \xrightarrow[2\text{-mark}]{\uparrow} \text{Men}_{C,F}(R_\omega)$

## Proof.

Proceeds by showing that  $[0, 1]$  is relatively robustly Menger (as a subspace of  $R_\omega$ ). Let  $f_C$  witness  $S([0, 1])$ .

Choose a finite subcover  $\mathcal{V}$  for  $[0, 1] \setminus C_\mathcal{V}$ . Let  $r_\mathcal{V}(x) = 0$  for  $x \notin C_\mathcal{V}$ , and  $r_\mathcal{V}(x) = f_{C_\mathcal{V}}(x)$  otherwise. Then  $c(\mathcal{V}, n)$  is finitely coverable by  $\mathcal{V}$  and  $p(\mathcal{U}, \mathcal{V}, n+1)$  is just finite. □

$$S(2^\omega) \text{ implies } \mathcal{F} \underset{2\text{-mark}}{\uparrow} \text{Men}_{C,F}(R_\omega)$$

Proceeds by showing that  $[0, 1]$  is relatively robustly Menger (as a subspace of  $R_\omega$ ). Let  $f_C$  witness  $S([0, 1])$ .

Choose a finite subcover of  $\mathcal{V}$  for  $[0, 1] \setminus C_{\mathcal{V}}$ . Let  $r_{\mathcal{V}}(x) = 0$  for  $x \notin C_{\mathcal{V}}$ , and  $r_{\mathcal{V}}(x) = f_{C_{\mathcal{V}}}(x)$  otherwise. Then  $c(\mathcal{V}, n)$  is finitely coverable by  $\mathcal{V}$  and  $p(\mathcal{U}, \mathcal{V}, n + 1)$  is just finite.

Some questions:

- $S(\kappa)$  implies  $\mathcal{F} \xrightarrow[2\text{-mark}]{\uparrow} \text{Fill}_{C,F}^{\cap}(\kappa)$ ... what about the other direction?
- Is there a space with  $\mathcal{F} \xrightarrow{\uparrow} \text{Men}_{C,F}(X)$  but  $\mathcal{F} \not\xrightarrow[2\text{-mark}]{\uparrow} \text{Men}_{C,F}(X)$ ?
- Is there a non- $\sigma$ -relatively-robustly-Menger space where  $\mathcal{F} \xrightarrow[2\text{-mark}]{\uparrow} \text{Men}_{C,F}(X)$ ?



Questions from any of you? Thanks!