

# Packet 2

## Part 2: Sections 14.1-14.3

### 14.1 Functions of Several Variables

**Definition 1.** A **function  $f$  of two variables** is a rule which assigns a real number  $f(x, y)$  to each pair of real numbers  $(x, y)$  for which that rule is defined. The collection of such well-defined pairs is called the **domain**  $\text{dom}(f)$  of the function, and the set of real numbers which can possibly be produced by the function is called its **range**  $\text{ran}(f)$ .

**Definition 2.** The **level curve** for each  $k \in \text{ran}(f)$  is given by the equation  $f(x, y) = k$ . The **graph** of  $f$  is a surface in 3D space which visualizes the function, given by the equation  $z = f(x, y)$ .

**Definition 3.** A **function  $f$  of three variables** is a rule which assigns a real number  $f(x, y, z)$  to each triple of real numbers  $(x, y, z)$  for which that rule is defined. The collection of such well-defined triples is called the **domain**  $\text{dom}(f)$  of the function, and the set of real numbers which can possibly be produced by the function is called its **range**  $\text{ran}(f)$ .

**Problem 4.** Let  $f(x, y) = x \sin(x + y)$ . Give the value of  $f(\pi, \frac{\pi}{2})$ .

**Solution.**

◇

**Problem 5.** Let  $f(x, y) = -x - y + 2$ . In the  $xy$ -plane, plot the domain of  $f$ , as well as its level curves for  $k = -3, 0, 3$ . Then plot the graph of  $f$  in  $xyz$  space.

**Solution.**

◇

**Problem 6.** Let  $f(x, y) = \sqrt{4 - x^2 - y^2}$ . In the  $xy$ -plane, plot the domain of  $f$ , as well as its level curves for  $k = 0, \frac{1}{\sqrt{2}}, 1$ . Then plot the graph of  $f$  in  $xyz$  space.

**Solution.**

◇

**Definition 7.** The **level surface** for each  $k \in \text{ran}(f)$  is given by the equation  $f(x, y, z) = k$ . (Since the graph of a three variable function would require four variables and therefore is a four-dimensional object, we typically don't consider it.)

**Problem 8.** Let  $f(x, y, z) = \frac{x+3y^2}{z-2x}$ . Give the value of  $f(3, -2, 1)$ .

**Solution.**

◇

**Problem 9.** Let  $f(x, y, z) = -x^2 + y - z^2$ . In  $xyz$  space, plot the level surfaces for  $k = -2, 0, 2$ .

**Solution.**

◇

**Remark 10.** If  $P = (x, y)$ , then we assume that  $f(x, y) = f(P) = f(\vec{P})$ . If  $P = (x, y, z)$ , then we assume that  $f(x, y, z) = f(P) = f(\vec{P})$ .

## 14.2 Limits and Continuity

**Definition 11.** If the value of the function  $f(P)$  becomes arbitrarily close to the number  $L$  as points  $P$  close to  $P_0$  are plugged into the function, then the **limit of  $f(P)$  as  $P$  approaches  $P_0$**  is  $L$ :

$$\lim_{P \rightarrow P_0} f(P) = L$$

**Theorem 12.** Let  $f(x, y)$  be a function of two variables. If there exists a curve  $y = g(x)$  passing through the point  $(x_0, y_0)$  such that  $\lim_{x \rightarrow x_0} f(x, g(x))$  does not exist, then  $\lim_{(x, y) \rightarrow (x_0, y_0)} f(x, y)$  does not exist.

**Problem 13.** Prove that

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{x + y}{|x + y|}$$

does not exist by considering the function  $g(x) = x$ .

**Solution.**

◇

**Theorem 14.** Let  $f(x, y)$  be a function of two variables. If there exist curves  $y = g(x)$  and  $y = h(x)$  passing through the point  $(x_0, y_0)$  such that  $\lim_{x \rightarrow x_0} f(x, g(x)) \neq \lim_{x \rightarrow x_0} f(x, h(x))$ , then  $\lim_{(x, y) \rightarrow (x_0, y_0)} f(x, y)$  does not exist.

**Problem 15.** Prove that

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{x^6 + y^2}{x^3y + x^6}$$

does not exist by considering the functions  $g(x) = x^3$  and  $h(x) = 2x^3$ .

**Solution.**

◇

**Theorem 16.** The “Limit Laws” for single-variable functions also hold for multi-variable functions.

$$\lim_{P \rightarrow P_0} (f(P) \pm g(P)) = \lim_{P \rightarrow P_0} f(P) \pm \lim_{P \rightarrow P_0} g(P)$$

$$\lim_{P \rightarrow P_0} (f(P) \cdot g(P)) = \lim_{P \rightarrow P_0} f(P) \cdot \lim_{P \rightarrow P_0} g(P)$$

$$\lim_{P \rightarrow P_0} (kf(P)) = k \lim_{P \rightarrow P_0} f(P)$$

$$\lim_{P \rightarrow P_0} \frac{f(P)}{g(P)} = \frac{\lim_{P \rightarrow P_0} f(P)}{\lim_{P \rightarrow P_0} g(P)}$$

$$\lim_{P \rightarrow P_0} (f(P))^{r/s} = \left( \lim_{P \rightarrow P_0} f(P) \right)^{r/s}$$

**Theorem 17.** Let  $P_0 = (x_0, y_0, z_0)$ . Multi-variable limits which only use one variable may be reduced to a single-variable limit.

$$\lim_{P \rightarrow P_0} f(x) = \lim_{x \rightarrow x_0} f(x)$$

$$\lim_{P \rightarrow P_0} g(y) = \lim_{y \rightarrow y_0} g(y)$$

$$\lim_{P \rightarrow P_0} h(z) = \lim_{z \rightarrow z_0} h(z)$$

**Problem 18.** Use the above theorems to rigorously prove that

$$\lim_{(x,y) \rightarrow (1,2)} \frac{2x+y}{y^2} = 1$$

**Solution.**

◇

**Remark 19.** Due to the limit laws, the “just plug it in” rule applies when plugging in does not result in an undefined operation.

**Problem 20.** Compute the limit

$$\lim_{(x,y,z) \rightarrow (3,0,-1)} \frac{x \cos y}{z + x}$$

**Remark 21.** There is no L’Hopital rule for multi-variable limits. However, you may still use it once the limit has been reduced to a single-variable limit.

**Problem 22.** Compute the limit

$$\lim_{(x,y) \rightarrow (3,0)} \frac{xy + \sin(2y)}{y}$$

**Solution.**

◇

**Remark 23.** Factoring and canceling (including conjugation tricks) is also effective for computing multi-variable limits.

**Problem 24.** Compute the limit

$$\lim_{(x,y,z) \rightarrow (1,2,4)} \frac{\sqrt{z} - xy}{z - x^2y^2}$$

**Solution.**

◇

**Definition 25.** A function  $f(P)$  is **continuous** if  $\lim_{P \rightarrow P_0} f(P) = f(P_0)$  for all points  $P_0$  in its domain.

**Theorem 26.** If a multi-variable function is composed of continuous single-variable functions, then it is also continuous.

## 14.3 Partial Derivatives

**Definition 27.** The **partial derivative of  $f$  with respect to a variable** is the rate of change of  $f$  as that variable changes and all other variables are held constant. For example:

$$\frac{\partial f}{\partial x} = f_x(x, y) = \lim_{h \rightarrow 0} \frac{f(x + h, y) - f(x, y)}{h}$$

$$\frac{\partial g}{\partial z} = g_z(x, y, z) = \lim_{h \rightarrow 0} \frac{g(x, y, z + h) - g(x, y, z)}{h}$$

**Problem 28.** Let  $f(x, y, z) = xy^2 + 2z$ . Use the definition of a partial derivative to prove that  $\frac{\partial f}{\partial y} = 2xy$ .

**Solution.**

◇

**Theorem 29.** Partial derivatives may be computed in the usual way by treating all other variables as constants.

**Problem 30.** Compute both partial derivatives of  $f(x, y) = 4x^2 - 5y^3 + xy - 1$ .

**Solution.**

◇

**Problem 31.** Compute both partial derivatives of  $f(x, y) = \sin(x + 3y)$ .

**Solution.**

◇

**Problem 32.** Compute both partial derivatives of  $f(x, y) = e^{xy^2}$ .

**Solution.**

◇

**Definition 33.** **Second-order partial derivatives** are the result of taking the partial derivative of a partial derivative.

**Theorem 34.** For sufficiently well-behaved functions, the order in which partial derivatives are taken is irrelevant. (This is sometimes called the **Mixed Derivative Theorem**.)

**Problem 35.** Verify the Mixed Derivative Theorem for  $f(x, y) = 3x^2y^2 - x^3 + y^4 - 7$ .

**Solution.**

◇