

# AN INVESTIGATION IN TOPOLOGICAL PROPERTIES

CODY MARTIN

ABSTRACT. This is the culmination of topics and topological properties that have been studied over the course of the Spring 2018 semester, mentored by Dr. Steven Clontz.

## 1. BASIC TOPOLOGICAL PROPERTIES AND DEFINITIONS

**Proposition 1.** *Find an infinite collection of open intervals in  $\mathbb{R}$  whose intersection is not an open interval.*

*Proof.*  $\bigcap \{(\frac{-1}{n}, \frac{1}{n}) : n \in \mathbb{Z}^+\} = \{0\}$  which is not an open interval or even open.  $\square$

**Proposition 2.** *Any finite union of closed sets is closed, and any arbitrary intersection of closed sets is closed.*

*Proof.* We proceed by showing that any finite union of closed sets is closed. Let  $C, D$  be closed sets. Let  $A, B$  be compliments of  $C, D$  so  $A, B$  are open. Then,  $A \cap B$  is also open. Thus,  $X \setminus (A \cap B)$  is closed. By Demorgan's Law,  $X \setminus (A \cap B) = X \setminus A \cup X \setminus B = C \cup D$  which is closed.

Now that we've shown that  $C \cup D$  is closed for all  $C, D$ ; Assume  $C_1 \cup C_2 \cup \dots \cup C_n$  is closed for closed sets  $C_i$ . Then for  $C_1 \cup \dots \cup C_n \cup C_{n+1}$ , let  $K = C_1 \cup \dots \cup C_n$ . Thus  $K \cup C_{n+1} = C_1 \cup \dots \cup C_n \cup C_{n+1}$  is closed.

Now we show that any arbitrary intersection of closed sets is closed. Let  $\mathcal{C}$  be a collection of closed sets. Let  $\mathcal{U} = \{X \setminus C : C \in \mathcal{C}\}$ , so  $\mathcal{U}$  is a collection of open sets. Then,  $\bigcup \mathcal{U}$  is also open. Thus,  $X \setminus \bigcup \mathcal{U}$  is closed. By Demorgan's Law,  $X \setminus \bigcup \mathcal{U} = \bigcap \mathcal{C}$  Therefore,  $\bigcap \mathcal{C}$  is closed.  $\square$

**Lemma 3.** *A set  $U$  is open if and only if for every point  $x \in U$ , there exists an open set  $U_x$  where  $x \in U_x \subseteq U$*

*Proof.* Suppose  $U$  is open, then for all  $x \in U$  there exists an open  $U_x = U$ , such that  $x \in U_x \subseteq U$ .

To show the converse, suppose that for each  $x \in U$  there is an open set  $U_x$  where  $x \in U_x \subseteq U$ . For  $x \in U, x \in U_x$  so  $x \in \bigcup \{U_x : x \in U\}$ . Thus  $U \subseteq \bigcup \{U_x : x \in U\}$ . Now let  $y \in U_x$  for some  $x \in U$ , then  $U_x \subseteq U$ . Thus  $U \supseteq \bigcup \{U_x : x \in U\}$ . Therefore  $U = \bigcup \{U_x : x \in U\}$ .  $\square$

**Proposition 4.** *A set  $K$  in a topological space  $X$  is closed if and only if  $K$  contains all its limit points.*

*Proof.* Suppose  $K$  contains all its limit points. If  $K = X$ , then  $K$  is closed because  $\emptyset$  is open. Otherwise let  $x \in X \setminus K$ , so  $x$  is not a limit point of  $K$ . Then,  $\exists x \in U_x \in \tau$

such that  $U_x \cap K = \emptyset$  since  $x \notin K$ . So,  $x \in U_x \subseteq X \setminus K$ . By the lemma,  $X \setminus K$  is open so  $K$  is closed.

To show the converse is true as well, let  $x \in X \setminus K$ , which is an open set by the lemma. Since  $(X \setminus K) \cap K$  is the empty set,  $x$  is not a limit point of  $K$ . So, if  $\ell$  is any limit point of  $K$ , then  $\ell \notin X \setminus K$ , so  $\ell \in K$  and  $K$  contains all of its limit points.  $\square$

**Proposition 5.** *Verify the discrete and indiscrete topologies are topologies*

*Proof.* To prove the discrete topology is an actual topology, first we define that the discrete topology on a set  $X$  is  $\tau = \mathcal{P}(X)$ .  $\emptyset, X \in \mathcal{P}(X)$ , So  $\emptyset, X \in \tau$ . Now, let  $\mathcal{U} \subseteq \tau = \mathcal{P}(X)$ , then the  $\bigcup \mathcal{U} \in \mathcal{P}(X)$ . Let  $U, V \in \tau = \mathcal{P}(X)$  then the intersection  $U \cap V \in \mathcal{P}(X) = \tau$ . Now to show that the indiscrete topology is an actual topology, we define that the indiscrete topology on a set  $X$  is given as  $\tau = \{\emptyset, X\}$ . Clearly,  $\emptyset, X \in \tau$ . Let  $\mathcal{U}$  be a collection of open sets in  $X$ , then  $\bigcup \mathcal{U} = X$ , thus  $\bigcup \mathcal{U} \in \tau$ .  $\square$

**Definition 6.** A collection of sets  $\mathcal{B} \subseteq \mathcal{P}(X)$  is called a **basis** if:

- (1) For all  $x \in X$ , there exists  $B \in \mathcal{B}$  such that  $x \in B$ .
- (2) For all  $B_1, B_2 \in \mathcal{B}$  with  $x \in B_1 \cap B_2$ , there exists  $B_3 \in \mathcal{B}$  with  $x \in B_3 \subseteq B_1 \cap B_2$ .

The set  $\{\bigcup \mathcal{B}' : \mathcal{B}' \subseteq \mathcal{B}\}$  is called the topology generated by  $\mathcal{B}$ . [1]

**Proposition 7.** *The “topology generated by  $\mathcal{B}$ ” is actually a topology.*

**Theorem 8.** *Let  $\tau$  be a topology on  $X$ . Then  $\mathcal{B} \subseteq \tau$  generates  $\tau$  if:*

1. *For all  $x \in U \in \tau$ , there exists  $B \in \mathcal{B}$  where  $x \in B \subseteq U \in \tau$*
2. *For all  $B_1, B_2 \in \mathcal{B}$  with  $x \in (B_1 \cap B_2)$ , there exists  $B_3 \in \mathcal{B}$  with  $x \in B_3 \subseteq B_1 \cap B_2$ .*

*Proof.* :

- (1) For  $x \in \mathbb{X} \in \tau$ , select  $x \in B \subseteq \mathbb{X} \in \tau$ , so 1 of definition 6 holds.
- (2) See definition 6-2.
- (3) Let  $U \in \tau$ , then for every  $x \in U$ , there exists  $B_x \in \mathcal{B}$  where  $x \in B_x \subseteq U \in \tau$ . Let  $\mathcal{B}' = \{B_x : B_x \in \mathcal{B}\}$  so  $U = \bigcup \mathcal{B}'$ . Now let  $\mathcal{B}' \subseteq \mathcal{B} \subseteq \tau$ , and since  $\mathcal{B}' \subseteq \tau, \bigcup \mathcal{B}' \in \tau$ .

$\square$

**Theorem 9.**  $\{\mathbb{X}\}$  is a basis for  $\tau = \{\emptyset, \mathbb{X}\}$ . (indiscrete)

*Proof.* :

- (1) Let  $\mathcal{B} = \{\mathbb{X}\}$  and let  $x \in \mathbb{X} = B \in \mathcal{B}$ .
- (2) Now consider  $B_1, B_2 \in \mathcal{B}$ .  $B_1 = \mathbb{X}$  and  $B_2 = \mathbb{X}$  Let  $x \in B_1 \cap B_2$ , then  $x \in B_3 \subseteq B_1 \cap B_2$ .
- (3) Choose  $\mathcal{B}_\emptyset = \emptyset \subseteq \mathcal{B}$  and  $\mathcal{B}_\mathbb{X} = \{\mathbb{X}\} \subseteq \mathcal{B}$  where  $\bigcup \mathcal{B}_\emptyset = \emptyset$  and  $\bigcup \mathcal{B}_\mathbb{X} = \mathbb{X}$ , so since  $\mathcal{P}(\mathcal{B}) = \{\emptyset, \{\mathbb{X}\}\}$ ,  $\{\bigcup \mathcal{B}' : \mathcal{B}' \subseteq \mathcal{B}\} = \{\bigcup \mathcal{B}_\emptyset, \bigcup \mathcal{B}_\mathbb{X}\} = \tau = \{\emptyset, \{\mathbb{X}\}\}$ .

$\square$

**Theorem 10.**  $\mathcal{B} = \{\{x\} : x \in \mathbb{X}\}$  is a basis for  $\tau = \mathcal{P}(\mathbb{X}) = \{U : U \subseteq \mathbb{X}\}$ . (discrete)

*Proof.* :

- (1) Let  $x \in U \in \tau = \mathcal{P}(\mathbb{X})$ . Then for  $B = \{x\} \in \mathcal{B}, x \in B \subseteq U$ .
- (2) Let  $B_1, B_2 \in \mathcal{B}$ . Let  $y \in B_1 \cap B_2$ . Let  $B_3 = \{y\} \in \mathcal{B} \subseteq B_1 \cap B_2$ .
- (3) Let  $U \in \tau = \mathcal{P}(\mathbb{X})$ . Then  $U = \{x : x \in U\} = \bigcup \{x : x \in U\} = \bigcup \mathcal{B}_U$  for  $\mathcal{B}_U = \{\{x\} : x \in U\} \subseteq \mathcal{B}$ .  
Let  $\mathcal{B}' \subseteq \mathcal{B}$ . Then  $\bigcup \mathcal{B}' \in \mathcal{P}(\mathbb{X}) \in \tau$ .

□

**Definition 11.** The **Euclidean topology** on  $\mathbb{R}$  is the topology generated by the **basis**

$$\{(a, b) : a < b \in \mathbb{R}\}.$$

**Theorem 12.**  $\{(a, b) : a < b \in \mathbb{R}\}$  is a basis.

*Proof.* :

- (1) Let  $\mathcal{B} = \{(a, b) : a < b \in \mathbb{R}\}$ . Let  $x \in \mathbb{R}$  and let  $B = (x - 1, x + 1)$ , then  $x \in B \subseteq \mathbb{R}$  and  $B \in \mathcal{B}$ .
- (2) For  $B_1 = (a_1, b_1)$ ,  $B_2 = (a_2, b_2)$ ,  $a_1 < b_1$ ,  $a_2 < b_2 \in \mathbb{R}$  with  $x \in B_1 \cap B_2$ , then there is  $B_3 = (\max(a_1, a_2), \min(b_1, b_2)) \subseteq B_1 \cap B_2$  such that  $a_1, a_2 < x < b_1, b_2$  so  $x \in B_3$ .

□

**Proposition 13.**  $\{(a, b) : a < b \in \mathbb{Q}\}$  is a basis for the Euclidean topology.

## 2. COVERING PROPERTIES

**Definition 14.** A subset  $K$  of a topological space  $X$  is said to be **compact** if for every open cover  $\mathcal{U}$  of  $K$  (every collection  $\mathcal{U}$  of open sets such that  $\bigcup \mathcal{U} \supseteq K$ ) there exists a finite subcollection  $\mathcal{F} \subseteq \mathcal{U}$  that also covers  $K$  (that is,  $\bigcup \mathcal{F} \supseteq K$ ). [1]

**Theorem 15.** Any finite union of compact subsets is compact.

*Proof.*  $K_1$  and  $K_2$  are compact, so let  $\mathcal{U}_3$  be an open cover of  $K_3 = K_1 \cup K_2$ . So let  $\mathcal{U}_1 = \mathcal{U}_3, \bigcup \mathcal{U}_3 \supseteq K_1$  and  $\mathcal{U}_2 = \mathcal{U}_3, \bigcup \mathcal{U}_3 \supseteq K_2$ . Since  $K_1, K_2$  are compact choose finite  $\mathcal{F}_1 \subseteq \mathcal{U}_1 = \mathcal{U}_3$  and  $\mathcal{F}_2 \subseteq \mathcal{U}_2 = \mathcal{U}_3$ , so  $\bigcup \mathcal{F}_1 \supseteq K_1$  and  $\bigcup \mathcal{F}_2 \supseteq K_2$  and  $\bigcup (\mathcal{F}_1 \cup \mathcal{F}_2) \supseteq K_3$ . Since  $\mathcal{F}_1 \cup \mathcal{F}_2$  is a finite subset of  $\mathcal{U}_3$ ,  $K_3$  is compact. □

**Proposition 16.** Any finite subset of a topological space is compact.

*Proof.* For  $F = \{x_i : 0 \leq i < n\} \subseteq \mathbb{X}$ , let  $\mathcal{U}$  be an open covering of  $F$  so  $\bigcup \mathcal{U} \supseteq F$ . For each  $0 \leq i < n$ , let  $U_i \in \mathcal{U}$  satisfy  $x_i \in U_i$ . Let finite  $\mathcal{F} = \{U_i : 0 \leq i < n\} \subseteq \mathcal{U}$ , so  $\bigcup \mathcal{F} \supseteq F$  and  $F$  is compact. □

**Proposition 17.** The subset  $\{0\} \cup \{\frac{1}{n} : n \in \mathbb{Z}^+\}$  of  $\mathbb{R}$  is compact.

*Proof.* Let  $K = \{0\} \cup \{\frac{1}{n} : n \in \mathbb{Z}^+\} \subseteq \mathbb{R}$  and let  $\mathcal{U}$  be an open covering of  $K$ . We may choose  $U_0 \in \mathcal{U}$  where  $0 \in U_0$ . So pick  $a < b \in \mathbb{R}$  where  $0 \in (a, b) \subseteq U_0$ . Let  $N \in \mathbb{Z}^+$  where  $0 < \frac{1}{N} < b$ . Now let finite  $F = \{\frac{1}{m} : 1 \leq m < N\}$  so it is compact, and choose  $\mathcal{F} \subseteq \mathcal{U}$  with  $\bigcup \mathcal{F} \supseteq F$ . So,  $\bigcup \mathcal{F} \cup (a, b) \cup U_0 \supseteq K$ . □

**Proposition 18.** Any unbounded subset of  $\mathbb{R}$  is not compact.

*Proof.* Let  $K$  be an unbounded subset of  $\mathbb{R}$ . Let  $\mathcal{U} = \{(n-2, n) : n \in \mathbb{Z}\}$ . Let  $n_i \in \mathbb{Z}$  be arbitrary for  $0 \leq i < n$ . Then  $\mathcal{F} = \{(n_i-2, n_i) : 0 \leq i < n\}$  is an arbitrary finite sub collection of  $\mathcal{U}$ . However,  $\bigcup \mathcal{F} \not\supseteq K$ , for  $\min(n_i-2 : 0 \leq i < n)$  and  $\max(n_i : 0 \leq i < n)$  we can find a  $k \in K : k < \min(n_i-2 : 0 \leq i < n)$  or  $k > \max(n_i : 0 \leq i < n)$ , so  $k \notin \bigcup \mathcal{F}$  and  $K$  is not compact.  $\square$

**Proposition 19.** Any open interval  $(a, b) \subseteq \mathbb{R}$  is not compact.

*Proof.* Let  $K = (a, b) \subseteq \mathbb{R}$ . Let  $\mathcal{U} = \{(x, y) : x, y \in \mathbb{R}, a < x < y < b\}$ . So,  $\bigcup \mathcal{U} \supseteq K$ , and  $\bigcup \mathcal{U} = K$ . Let  $n \in \mathbb{Z}$ , let  $x_i, y_i \in \mathbb{R}$  be arbitrary with  $0 \leq i < n$ . Then,  $\mathcal{F} = \{(x_i, y_i) : 0 \leq i < n\}$  is an arbitrary finite subset of  $\mathcal{U}$ . However  $\bigcup \mathcal{F} \not\supseteq K$ , for  $\min(x_i : 0 \leq i < n)$  we can find a  $k \in K : k < \min(x_i : 0 \leq i < n)$  which is greater than  $a$  so  $k \notin \bigcup \mathcal{F}$  and  $K$  is not compact.  $\square$

**Definition 20.** A subset  $K$  of a topological space  $X$  is said to be **Lindelöf** if for every open cover  $\mathcal{U}$  of  $K$  (every collection  $\mathcal{U}$  of open sets such that  $\bigcup \mathcal{U} \supseteq K$ ) there exists a countable subcollection  $\mathcal{F} \subseteq \mathcal{U}$  that also covers  $K$  (that is,  $\bigcup \mathcal{F} \supseteq K$ ). [2]

**Definition 21.** A subset  $K$  of a topological space  $X$  is said to be  **$\sigma$ -compact** if there exist compact subspaces  $K_n$  of  $X$  for  $n \in \mathbb{N}$  such that  $K = \bigcup_{n \in \mathbb{N}} K_n$ . [2]

**Theorem 22.** Every  **$\sigma$ -compact** subset of a topological space is **Lindelöf**.

*Proof.* Let  $K$  be  $\sigma$ -compact. Let  $\mathcal{F}_n = \{U_{n0}, U_{n1}, \dots, U_{nm}\}$  be a finite subcover of  $\mathcal{U}$  for  $K_n$ . So,  $\mathcal{C} = \bigcup_{n \in \mathbb{N}} \mathcal{F}_n$  is a countable subcover of  $\mathcal{U}$  for  $K$ . Therefore  $K$  is Lindelöf.  $\square$

**Definition 23.** A subset  $K$  of a topological space  $X$  is said to be **hemicompact** if there exists compact sets  $K_n \subseteq K$  for  $n \in \mathbb{N}$  such that for every compact subset  $H \subseteq K$ ,  $H \subseteq K_n$  for some  $n \in \mathbb{N}$ . [2]

**Theorem 24.** Every **hemicompact** subset of a topological space is  **$\sigma$ -compact**.

*Proof.* Let  $K_n \subseteq K$  witness hemicompactness, of course  $\bigcup_{n \in \mathbb{N}} K_n \subseteq K$ , since  $K_n \subseteq K$ . Let  $k \in K$ , so  $\{k\}$  is compact, thus there is some  $n \in \mathbb{N}$  such that  $\{k\} \subseteq K_n$ . Therefore  $k \in K_n \subseteq K$ , so  $K = \bigcup_{n \in \mathbb{N}} K_n$ .  $\square$

**Definition 25.**  $X$  is **locally compact** if for every  $x \in X$  there exists an open set  $U$  and compact  $K$  with  $x \in U \subseteq K$ . [1]

**Definition 26.**  $x$  is a **cluster point** of  $\langle p_n \rangle_{n < \omega}$  if and only if for every neighborhood  $U$  of  $x$ ,  $U$  hits infinitely many  $p_n$ .

**Lemma 27.** Let  $K$  be a compact subset of  $X$ . If  $\langle p_n \rangle$  is a sequence of points in  $K$ , then  $\langle p_n \rangle$  has a cluster point.

*Proof.* This is obviously true if the sequence only uses finitely many different points (one point must be repeated infinitely often, so it is its own cluster point). So without loss of generality  $\langle p_n \rangle$  uses infinitely many different points. If  $\langle p_n \rangle$  didn't have a cluster point, then for each  $x \in K$ , let  $U_x$  contain only finitely many  $p_n$ .  $\{U_x : x \in K\}$  covers  $K$ , so choose  $x_i$  for  $i \leq n : \{U_{x_i} : i \leq n\}$  covers  $K$ . But each  $U_{x_i}$  contains only finitely many  $p_n$ , so  $\langle p_n \rangle$  cannot have infinitely many different points to choose, contradiction. So  $\langle p_n \rangle$  does have a cluster point.  $\square$

## 3. TOPOLOGICAL GAMES

**Definition 28.**  $Gru_{\mathcal{K}, \mathcal{P}}(X)$  is a game defined as the *Gruenhagecompact/pointgame* with players  $\mathcal{K}$ , *mathcal{P}* played on a topological space  $X$ . During round  $n$ ,  $\mathcal{K}$  chooses a compact subset  $K_n$  of  $X$ , followed by  $\mathcal{P}$  choosing a point  $p_n \in X$  such that  $p_n \notin \bigcup_{m \leq n} K_m$ .  $\mathcal{K}$  wins if and only if  $\langle p_n \rangle_{n < \omega}$  lacks a **cluster point**. [2]

**Definition 29.**  $\mathcal{K} \uparrow_{pre} Gru_{\mathcal{K}, \mathcal{P}}(X)$  is the statement that  $\mathcal{K}$  has a winning strategy  $\sigma : \omega \rightarrow \mathcal{K}(X)$  that depends only on the round number (ignores the moves of  $\mathcal{P}$ ). [2]

**Theorem 30.** If  $X$  is **Lindelöf** and locally compact, then  $\mathcal{K} \uparrow_{pre} Gru_{\mathcal{K}, \mathcal{P}}(X)$

*Proof.* For  $x \in X$ , let  $x \in U_x \subseteq K_x$  for  $U_x$  open and  $K_x$  compact. Then by definition of Lindelöf choose  $U_n \in \mathcal{U} = \{U_x : x \in X\}$  such that  $\bigcup_{n \in \mathbb{N}} U_n = X$ . Then

for each  $U_n$ , there exists compact  $K_n$  where  $U_n \subseteq K_n$ .

For  $Gru_{\mathcal{K}, \mathcal{P}}(X)$ , let  $\sigma(n) = K_n$  denote a strategy for  $\mathcal{K}$ ,  $\mathcal{K}$  chooses compact  $K_n \supseteq U_n \in \mathcal{C}$  and  $\mathcal{P}$  chooses  $p \in X$  such that  $p \notin \bigcup_{m \leq n} K_m$ .  $\{K_i : i < \omega\}$  is countable and

$\bigcup \{K_i : i < \omega\} \supseteq X$ . For  $\mathcal{P}$  to win, there must be an  $x \in X$  such that for every neighborhood  $U$  of  $x$ ,  $U$  hits infinitely many  $p_n$ , but  $x \in U_m \subseteq K_m$  for some  $n < \omega$  and  $p_n \notin K_m$  for  $n \geq m$ . So there is no such  $x$  and  $\mathcal{K} \uparrow_{pre} Gru_{\mathcal{K}, \mathcal{P}}(X)$ .  $\square$

**Theorem 31.**  $\mathcal{K} \uparrow_{pre} Gru_{\mathcal{K}, \mathcal{P}}(X) \Rightarrow X$  is **hemicompact**.

*Proof.* We prove the contrapositive. Let  $\sigma$  be a predetermined strategy, and assume  $X$  isn't hemicompact. So there exists a compact  $K$  that isn't a union of finite  $\sigma(n)$ , so player  $\mathcal{P}$  may always legally choose a point  $p_n \in K$ . Then by the lemma,  $\langle p_n \rangle$  has a cluster point because for each  $x \in K$  there is a  $U_x$  that contains infinitely many  $p_n$ , and  $\{U_x : x \in K\}$  covers  $K$ . Therefore  $\sigma$  isn't a winning strategy.  $\square$

## REFERENCES

- [1] Steven Clontz. Applications of stationary sets in set theoretic topology. 2010.
- [2] Steven Clontz. Zero-markov information in topological games. *Alabama Journal of Mathematics*, 39, 2015.

1313 SCHILLINGER RD. S. APT. 3407 MOBILE, AL 36695

E-mail address: cdm1322@jagmail.southalabama.edu