

1. The Substitution Rule is the opposite of which derivative rule? (2017-06-05, 2.1)
 - A. Chain Rule
 - B. Product Rule
 - C. Quotient Rule
 - D. Power Rule

2. What is incorrect about the following attempt at using the Substitution Rule?

$$\int_0^1 (3 - 2x)^5 dx = \int_0^1 u^5 \left(-\frac{1}{2} du \right)$$

(2017-06-05, 2.1)

- A. dx should have been replaced with $+\frac{1}{2} du$.
 - B. The bounds are incorrect.
 - C. u shouldn't be raised to the 5th power.
 - D. dx should have been replaced with $-2 du$.
3. Which of these formulas would be most useful in finding $\int \sin^4 \theta \cos^2 \theta d\theta$? (2017-06-05, 2.2)
 - A. $\sin^2(\theta) = \frac{1}{2} + \frac{1}{2} \sin(2\theta)$
 - B. $\cos^2(\theta) = 1 - \sin^2(\theta)$
 - C. $\sin^2(\theta) = 1 - \cos^2(\theta)$
 - D. $\cos^2(\theta) = \frac{1}{2} + \frac{1}{2} \cos(2\theta)$
4. Which of these formulas would be most useful in finding $\int \sec^4(\theta) d\theta$? (2017-06-05, 2.2)
 - A. $\sec^2(\theta) = 1 - \tan^2(\theta)$
 - B. $\tan^2(\theta) = 1 + \sec^2(\theta)$
 - C. $\sec^2(\theta) = 1 + \tan^2(\theta)$
 - D. $\tan^2(\theta) = 1 - \sec^2(\theta)$

5. Which of these substitutions would be most useful in finding $\int \frac{1}{25x^2+9} dx$? (2017-06-05, 2.3)
- A. Let $25x^2 + 9 = 25 \sec^2 \theta + 25$.
 - B. Let $25x^2 + 9 = 9 \tan^2 \theta + 9$.
 - C. Let $25x^2 + 9 = 9 \sin^2 \theta + 9$.
 - D. Let $25x^2 + 9 = 25 \cos^2 \theta + 25$.
6. Which of these substitutions would be most useful in finding $\int \frac{1}{x\sqrt{4-16x^2}} dx$? (2017-06-05, 2.3)
- A. Let $4 - 16x^2 = 16 - 16 \cos^2 \theta$.
 - B. Let $4 - 16x^2 = 4 - 4 \sin^2 \theta$.
 - C. Let $4 - 16x^2 = 4 + 4 \tan^2 \theta$.
 - D. Let $4 - 16x^2 = 16 + 16 \sec^2 \theta$.
7. Which of these substitutions would be most useful in finding $\int_3^5 \frac{1}{\sqrt{x^2-9}} dx$? (2017-06-05, 2.3)
- A. Let $x^2 - 9 = 9 \sin^2 \theta + 9$.
 - B. Let $x^2 - 9 = \tan^2 \theta - 1$.
 - C. Let $x^2 - 9 = 9 \sec^2 \theta - 9$.
 - D. Let $x^2 - 9 = \cos^2 \theta + 1$.

8. Which of these sums is the first step in expanding $\frac{4x^2+16x+17}{(x+2)^2(x^2+1)^2}$ into partial fractions?
(2017-06-12, 2.4)
- A. $\frac{A}{x+2} + \frac{Bx}{x+2} + \frac{C}{(x^2+1)^2}$
 - B. $\frac{A}{x+2} + \frac{B}{(x+2)^2} + \frac{Cx+D}{x^2+1} + \frac{Ex+F}{(x^2+1)^2}$
 - C. $\frac{A}{x+2} + \frac{Bx}{(x+2)^2} + \frac{Cx+D}{(x^2+1)^2}$
 - D. $\frac{Ax+B}{(x+2)^2} + \frac{Cx+D}{(x^2+1)^2}$
9. Why must $\frac{3+5x^5}{(x+1)(x+3)^2}$ first be simplified using long polynomial division before using the method of partial fractions? (2017-06-12, 2.4)
- A. It is a rational function of x .
 - B. The degree of its numerator is odd, while the degree of its denominator is even.
 - C. It is an irrational function of x .
 - D. The degree of its numerator is greater than or equal to the degree of its denominator.
10. Integration by Parts is the opposite of which derivative rule? (2017-06-12, 2.5)
- A. Chain Rule
 - B. Quotient Rule
 - C. Power Rule
 - D. Product Rule
11. Which choice is most appropriate for using integration by parts to find $\int 3x \cos(x) dx$?
(2017-06-12, 2.5)
- A. $u = 3, dv = x \cos(x) dx$
 - B. $u = \cos(x), dv = 3x dx$
 - C. $u = 3x, dv = \cos(x) dx$
 - D. $u = x \cos(x), dv = 3 dx$
12. Which of these techniques is most appropriate as the first step to find $\int z^2 \sin(z^3) dz$?
(2017-06-12, 2.6)
- A. Integration by Substitution
 - B. Integration by Parts
 - C. Method of Partial Fractions
 - D. Trigonometric Identities

13. Which of these techniques is most appropriate as the first step to find $\int \frac{t^2+3t+1}{t^3+t} dt$?
(2017-06-12, 2.6)
- A. Method of Partial Fractions
 - B. Trigonometric Substitution
 - C. Trigonometric Identities
 - D. Integration by Substitution
14. Which of these integrals represents the area bounded by the curves $x = y^2$ and $x = 4$?
(2017-06-12, 3.1)
- A. $\int_0^2 2y \, dy$
 - B. $\int_{-2}^2 ((4) - (y^2)) \, dy$
 - C. $\int_0^4 ((\sqrt{y}) - (2)) \, dx$
 - D. $\int_{-2}^0 ((4y^2) - (2\sqrt{y})) \, dx$
15. Which of these integrals also represents the area bounded by the curves $x = y^2$ and $x = 4$? (2017-06-12, 3.1)
- A. $\int_0^2 ((x^2) - (4)) \, dx$
 - B. $\int_2^4 ((x^2) - (-x^2)) \, dx$
 - C. $\int_0^4 ((\sqrt{x}) - (-\sqrt{x})) \, dx$
 - D. $\int_2^0 ((x^2) - (\sqrt{x})) \, dx$

16. Let $A(x)$ be the area of the cross-section at x for a solid defined between $a \leq x \leq b$. Which of these integrals gives its volume? (2017-06-19, 3.2)
- A. $\int_b^a (A(x) - a - b) dx$
 - B. $\int_a^b [A(x)]^2 dx$
 - C. $\int_a^b A(x) dx$
 - D. $\int_b^a \frac{1}{2} A(x) dx$
17. Suppose the cross-sections of a solid defined between $0 \leq x \leq 5$ are triangles with base and height both equal to x . Find $A(x)$. (2017-06-19, 3.2)
- A. $\frac{1}{3}bh$
 - B. $(x^2 + 5x)$
 - C. πx^2
 - D. $\frac{1}{2}x^2$
18. Suppose the cross-sections of a solid are circular washers with outside radius $R(x)$ and inside radius $r(x)$. Which of these gives the area of such a cross-section? (2017-06-19, 3.3)
- A. $A(x) = 2\pi R(x)r(x)$
 - B. $A(x) = \pi R(x)r(x)$
 - C. $A(x) = 2\pi([R(x)]^2 - [r(x)]^2)$
 - D. $A(x) = \pi([R(x)]^2 - [r(x)]^2)$
19. Find the area of a cylindrical shell with radius 4 and height 3. (2017-06-19, 3.4)
- A. 12π
 - B. 30π
 - C. 6π
 - D. 24π

20. In the work integral $\int_a^b F(x) dx$, the function $F(x)$ represents... (2017-06-19, 3.5)
- A. Friction
 - B. Force
 - C. Speed
 - D. Mass
21. To calculate the work done in lifting an unladen swallow, which of the following properties would be most useful to know? (2017-06-19, 3.5)
- A. Airspeed velocity
 - B. Weight
 - C. Volume
 - D. Color
22. Suppose a tank of liquid has its base at height c and its top at height d . If dW describes the work required to pump a infinitesimal cross-section of liquid at height y , which integral gives the work required to completely pump this tank? (2017-06-19, 3.5)
- A. $\int_{y=c}^{y=d} dW$
 - B. $\int_{y=c}^{y=d} w^2 dW$
 - C. $\int_{y=c}^{y=d} -\frac{1}{2}y^2 dW$
 - D. $\int_{y=c}^{y=d} y dW$

23. The equations $x = 3 + 4 \cos(t)$, $y = -2 + 4 \sin(t)$ for $0 \leq t \leq 2\pi$ parametrize which kind of curve? (2017-06-26, 4.1)
- A. A parabola
 - B. A line segment
 - C. A circle oriented counter-clockwise
 - D. A circle oriented clockwise
24. The equations $x = 3 + 4t$, $y = -2 + 4t$ for $0 \leq t \leq 2\pi$ parametrize which kind of curve? (2017-06-26, 4.1)
- A. A parabola
 - B. A circle oriented counter-clockwise
 - C. A line segment
 - D. A circle oriented clockwise
25. The formula $\frac{dy}{dt} = \frac{dy}{dx} \frac{dx}{dt}$ for computing the slope of a tangent line to a curve defined by parametric equations is a result of the... (2017-06-26, 4.2)
- A. Product Rule
 - B. Chain Rule
 - C. Pythagorean Theorem
 - D. Method of Partial Fractions
26. The formula $L = \int_a^b \sqrt{[dx/dt]^2 + [dy/dt]^2} dt$ for computing arclength is a result of the... (2017-06-26, 4.2)
- A. Product Rule
 - B. Method of Partial Fractions
 - C. Chain Rule
 - D. Pythagorean Theorem

27. The polar coordinate $(3, \frac{2\pi}{3})$ equals which of the following Cartesian coordinates? (2017-06-26, 4.3)
- A. $(-3\sqrt{3}, 3)$
 - B. $(-\frac{3}{2}, \frac{3\sqrt{3}}{2})$
 - C. $(3, \frac{3}{2})$
 - D. $(-\frac{3}{2}, -3\sqrt{3})$
28. The polar equation $r = 3 \sec \theta$ parametrizes which kind of curve? (2017-06-26, 4.3)
- A. A vertical line
 - B. A circle
 - C. A horizontal line
 - D. A cardioid
29. In the polar area formula $\frac{1}{2} \int_{\alpha}^{\beta} ((R(\theta))^2 - (r(\theta))^2) d\theta$, the greek letter β represents which of the following? (2017-06-26, 4.4)
- A. The clockwise-most angle of the region.
 - B. The inner-most radius of the region.
 - C. The outer-most radius of the region.
 - D. The counter-clockwise-most angle of the region.