MATH 1121 (Calculus for Engineering Technology) Course Outline

1.3 Rectangular Coordinates

- Illustrate the following concepts:

    - rectangular coordinate system,

    - x-axis,

    - y-axis,

    - origin,

    - quadrants,

    - coordinates

- Examples:

    - (Example 1) Plot $A = (2, 1)$ and $B = (-4, -3)$.

    - (Example 3) Three verticies of a rectangle are $A = (-3, -2)$, $B = (4, -2)$, $C = (4, 1)$. What is the fourth vertex?

- *HW: 1-9, 15-16, 21-24*

2.1 Some Basic Definitions

- Distance Formula

    - $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

    - (Example 2) Find the distance between $(3, -1)$ and $(-2, -5)$.

- Slope Formula

    - $m = \frac{y_2 - y_1}{x_2 - x_1}$

    - (Example 3) Find the slope of the line joining $(3, -5)$, $(-2, -6)$.

    - (Example 4) Find the slope of the line joining $(3, 4)$, $(4, -6)$.

    - $m = \tan\theta$

    - (Example) Find the slope of the line with inclination $120°$.

- Identify parallel/perpendicular lines by slopes.

    - Parallel: $m_1 = m_2$

    - Perpendicular: $m_1 = -\frac{1}{m_2}$

- ■ (Example 7) Prove that the triangle with vertices $A = (-5, 3)$, $B = (6, 0)$, and $C = (5, 5)$ is a right triangle.

- *HW: 1-20, 29-36*

2.2 The Straight Line

- Point-slope form

    - ■ $y - y_1 = m(x - x_1)$
    - ■ (Example 2) Find the equation of the line passing through $(2, -1)$ and $(6, 2)$.

- Slope-intercept form

    - ■ $y = mx + b$
    - ■ (Example 4) Find the slope and y-intercept of the straight line with equation $2y + 4x - 5 = 0$.

- *HW: 1-21, 33-40*

2.3 The Circle

- Definition

    - ■ A circle is a collection of points equidistant from its center.

- Standard form

    - ■ $(x - h)^2 + (y - k)^2 = r^2$
    - ■ (Example 1) Sketch $(x - 1)^2 + (y + 2)^2 = 16$.
    - ■ (Example 2) Find an equation for the circle with center $(2, 1)$ which passes through $(4, 8)$.

- General form

    - ■ $x^2 + y^2 + Dx + Ey + F = 0$
    - ■ (Example 4) Find the center and radius of the circle $x^2 + y^2 - 6x + 8y - 24 = 0$.
    - ■ (Example) Find two functions whose graphs represent the circle with the previous equation.

- *HW: 1-32, 37-38*

2.4 The Parabola

- Definition

  - A parabola is a colleciton of points equidistant from a focus point and a directrix line.
  - The vertex of a parabola is the point closest to the focus and directix.
  - (Example 6) Find an equation for the parabola with focus $(2, 3)$ and directrix $(y = -1)$

- Standard forms with vertex at origin and horizontal/vertical directrix

  - $y^2 = 4px$ with directrix at $x = -p$ and focus at $(p, 0)$
  - $y^2 = -4px$ with directrix at $x = p$ and focus at $(-p, 0)$
  - $x^2 = 4py$ with directrix at $y = -p$ and focus at $(0, p)$
  - $x^2 = -4py$ with directrix at $y = p$ and focus at $(0, -p)$
  - (Example 2) Find an equation for the parabola with focus $(-2, 0)$ and directrix $(x = 2)$.
  - (Example 4) Find the focus and directrix of the parabola with equation $2x^2 = -9y$.

- *HW: 1-22, 25-28*

2.5 The Ellipse

- Definition

  - An ellipse is a collection of points where the sum of distances from two fixed points (called foci) is kept constant.
  - The two points furthest/closest apart from each other on an ellipse are the end-points of the major/minor axis.
  - The sum of distances between each point and the foci is the same as the length of the major axis. The major axis passes through both foci.

- Standard form with center at the origin

  - $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, with foci given by $(c, 0), (-c, 0)$, where $a^2 - b^2 = c^2$
  - $\frac{y^2}{a^2} + \frac{x^2}{b^2} = 1$, with foci given by $(0, c), (0, -c)$, where $a^2 - b^2 = c^2$
  - (Example 3) Sketch the ellipse with equation $4x^2 + 16y^2 = 64$, and compute the locations of its foci.

- (Example 5) Find the equation of the ellipse centered at the origin with an end of its minor axis at $(2,0)$ and containing the point $(-1, \sqrt{6})$.

- *HW: 1-26*

2.6 The Hyperbola

- Definition

  - A hyperbola is a collection of points where the difference of distances from two fixed points (called foci) is kept constant.
  - Hyperbolas are split into two curves. The two closest points on opposite curves are called vertices and give the transverse axis.

- Standard form with center at the origin

  - $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$, with foci given by $(c,0), (-c,0)$ and asymptotes $y = \pm\frac{bx}{a}$, where $a^2 + b^2 = c^2$
  - $\frac{y^2}{a^2} - \frac{x^2}{b^2} = 1$, with foci given by $(0,c), (0,-c)$ and asymptotes $x = \pm\frac{by}{a}$, where $a^2 + b^2 = c^2$
  - (Example 2) Sketch $\frac{y^2}{4} - \frac{x^2}{16} = 1$, labeling its vertices, asymptotes, and foci.
  - (Example 3) Sketch $4x^2 - 9y^2 = 36$, labeling its vertices, asymptotes, and foci.

- Hypberbola with coordinate axis asymptotes

  - $xy = c^2$, with vertices given by (c,c) and $(-c,-c)$
  - $xy = -c^2$, with vertices given by $(c,-c)$ and $(-c,c)$
  - (Example 5) Sketch $xy = 4$.

- *HW: 1-14, 17-24*

2.7 Translation of Axes

- Vertical/horizontal translation:

  - Shift right h: replace x with $x - h$.
  - Shift up k: replace y with $y - k$.
  - (Example 1) Give an equation of the parabola with vertex $(2,4)$ and focus $(4,4)$.
  - (Example 2) Sketch the curve with equation $\frac{(x-3)^2}{25} + \frac{(y+2)^2}{9} = 1$.

- *HW: 1-36*

1.2 Algebraic Functions

- Definition of a function $y = f(x)$.

- Types of functions

  - Polynomials $P(x) = a_0 + a_1 x + \cdots + a_n x^n$
  - Rational functions $R(x) = \frac{P(x)}{Q(x)}$ for polynomials P, Q
  - (Example 1) Voltage equals current multiplied by resistance. If the voltage at time t is given by $E(t) = 2t^2 = y + 5$ and the resistance at time t is given by $R(t) = 3t + 20$, then find a function $I(t)$ which measures the current at time t. Identify it as a polynomial and/or rational function.

- Combinations of functions

  - Addition/Subtraction/Multiplication/Division
  - Compositions $f \circ g$ and $g \circ f$
  - (Example 2) Express $f+g$, $f \circ g$, and $g \circ f$ for the functions given by $f(x) = 2x^2 - 3$ and $g(x) = \sqrt{x+2}$.

- Domain/Range

  - The domain of a function is all real numbers which may be plugged into it without causing division by zero, even roots of negatives, or any other undefined operations.
  - The range of a function is all real numbers which may possibly be attained by the function.
  - (Example 5) Find the domain and range of $f(x) = x^2 + 2$ and $g(t) = \frac{1}{t+2}$.
  - (Example 7) Find the domain of $f(x) = 16\sqrt{x} + \frac{1}{x}$.

- Piecewise functions

  - Piecewise functions are defined differently for different parts of their domains.
  - (Example 9) Find the domain for

$$f(t) = \begin{cases} 8 - 2t & 0 \le t \le 4 \\ 0 & t > 4 \end{cases}$$

  and compute $f(3), f(6), f(-1)$ if possible.

- Exponent laws

  - $a^m a^n = a^{m+n}$

- $\frac{a^m}{a^n} = a^{m-n}$
- $(a^m)^n = a^{mn}$
- $(ab)^m = a^m b^m$
- $(\frac{a}{b})^m = \frac{a^m}{b^m}$
- $a^0 = 1$
- $a^{-n} = \frac{1}{a^n}$
- $a^{1/n} = \sqrt[n]{a}$
- Note $\sqrt{a^2} = |a|$ but $\sqrt[3]{a^3} = a$
- (Example 4) Simplify

$$f(x) = \frac{(3x^2 - 1)^{1/3}(2x) - (2x^3)(3x^2 - 1)^{-2/3}}{(3x^2 - 1)^{2/3}}$$

- *HW: 1-18, 21-34*

1.4 The Graph of a Function

- Definition

  - The graph of a function is the collection of all ordered pairs (x, y) such that $y = f(x)$
  - Graphing Method 1: using Chapter 2
  - Graphing Method 2: using xy chart
  - Vertical line test: the graph of any function hits every vertical line at most once

- Examples

  - (Example 1) Graph $f(x) = 3x - 5$.
  - (Example 3) Graph $f(x) = 1 + \frac{1}{x}$.
  - (Example 4) Graph $f(x) = \sqrt{x + 1}$.
  - (Example 6) Graph

$$f(x) = \begin{cases} 2x + 1 & x \leq 1 \\ 6 - x^2 & x > 1 \end{cases}$$

- *HW: 1-12, 37-40*

Remaining Topics

- 3.1 Limits

- 3.2 The Slope of a Tangent to a Curve

- 3.3 The Derivative

- 3.4 The Derivative as an Instantaneous Rate of Change

- 3.5 Derivatives of Polynomials

- 3.6 Derivatives of Products and Quotients of Functions

- 3.7 The Derivative of a Power of a Function

- 3.8 Differentiation of Implicit Functions

- 3.9 Higher Derivatives

- 4.1 Tangents and Normals

- 4.4 Related Rates

- 4.5 Using Derivatives in Curve Sketching

- 4.6 More on Curve Sketching

- 4.7 Applied Maximum and Minimum Problems

- 4.8 Differentials and Linear Approximations

- 5.1 Antiderivatives

- 5.2 The Indefinite Integral

- 5.3 The Area Under a Curve

- 5.4 The Definite Integral

- 7.1 The Trigonometric Functions

- 7.2 Basic Trigonometric Relations

- 7.3 Derivatives of the Sine and Cosine Functions

- 7.4 Derivatives of the Other Trigonometric Functions

- 8.1 Exponential and Logarithmic Functions

- 8.2 Derivative of the Logarithmic Functions

- 8.3 Derivative of the Exponentials Function

- 9.1 The General Power Formula

- 9.2 Basic Logarithmic Form

- 9.3 Exponential Form

- 9.4 Basic Trigonometric Forms