MATH 1121 (Calculus for Engineering Technology) Course Outline

1.3 Rectangular Coordinates

- Illustrate the following concepts:
 - rectangular coordinate system,
 - \blacksquare x-axis,
 - \blacksquare y-axis,
 - origin,
 - quadrants,
 - coordinates
- Examples:
 - (Example 1) Plot A = (2, 1) and B = (-4, -3).
 - (Example 3) Three vertices of a rectangle are A = (-3, -2), B = (4, -2), C = (4, 1). What is the fourth vertex?
- HW: 1-9, 15-16, 21-24

2.1 Some Basic Definitions

- Distance Formula
 - $d = \sqrt{(x_2 x_1)^2 + (y_2 y_1)^2}$
 - (Example 2) Find the distance between (3, -1) and (-2, -5).
- Slope Formula
 - $\blacksquare m = \frac{y_2 y_1}{x_2 x_1}$
 - (Example 3) Find the slope of the line joining (3, -5), (-2, -6).
 - (Example 4) Find the slope of the line joining (3,4), (4,-6).
 - $m = \tan \theta$
 - (Example) Find the slope of the line with inclination 120°.
- Identify parallel/perpendicular lines by slopes.
 - Parallel: $m_1 = m_2$
 - Perpendicular: $m_1 = -\frac{1}{m_2}$

- (Example 7) Prove that the triangle with vertices A = (-5,3), B = (6,0), and C = (5,5) is a right triangle.
- HW: 1-20, 29-36

2.2 The Straight Line

- Point-slope form
 - $y y_1 = m(x x_1)$
 - (Example 2) Find the equation of the line passing through (2,-1) and (6,2).
- Slope-intercept form
 - y = mx + b
 - (Example 4) Find the slope and y-intercept of the straight line with equation 2y + 4x 5 = 0.
- HW: 1-21, 33-40

2.3 The Circle

- Definition
 - A circle is a collection of points equidistant from its center.
- Standard form
 - $(x-h)^2 + (y-k)^2 = r^2$
 - (Example 1) Sketch $(x-1)^2 + (y+2)^2 = 16$.
 - (Example 2) Find an equation for the circle with center (2,1) which passes through (4,8).
- General form
 - $x^2 + y^2 + Dx + Ey + F = 0$
 - (Example 4) Find the center and radius of the circle $x^2 + y^2 6x + 8y 24 = 0$.
 - (Example) Find two functions whose graphs represent the circle with the previous equation.
- HW: 1-32, 37-38

2.4 The Parabola

- Definition
 - A parabola is a collection of points equidistant from a focus point and a directrix line.
 - The vertex of a parabola is the point closest to the focus and directix.
 - (Example 6) Find an equation for the parabola with focus (2,3) and directrix (y=-1)
- Standard forms with vertex at origin and horizontal/vertical directrix
 - $y^2 = 4px$ with directrix at x = -p and focus at (p, 0)
 - $y^2 = -4px$ with directrix at x = p and focus at (-p, 0)
 - $x^2 = 4py$ with directrix at y = -p and focus at (0, p)
 - $x^2 = -4py$ with directrix at y = p and focus at (0, -p)
 - (Example 2) Find an equation for the parabola with focus (-2,0) and directrix (x=2).
 - (Example 4) Find the focus and directrix of the parabola with equation $2x^2 = -9y$.
- HW: 1-28

2.5 The Ellipse

- Definition
 - An ellipse is a collection of points where the sum of distances from two fixed points (called foci) is kept constant.
 - The two points furthest/closest apart from each other on an ellipse are the endpoints of the major/minor axis.
 - The sum of distances between each point and the foci is the same as the length of the major axis. The major axis passes through both foci.
- Standard form with center at the origin
 - $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, with foci given by (c,0), (-c,0), where $a^2 b^2 = c^2$
 - $\frac{y^2}{a^2} + \frac{x^2}{b^2} = 1$, with foci given by (0,c), (0,-c), where $a^2 b^2 = c^2$
 - (Example 3) Sketch the ellipse with equation $4x^2 + 16y^2 = 64$, and compute the locations of its foci.

- (Example 5) Find the equation of the ellipse centered at the origin with an end of its minor axis at (2,0) and containing the point $(-1,\sqrt{6})$.
- HW: 1-26

2.6 The Hyperbola

- Definition
 - A hyperbola is a collection of points where the difference of distances from two fixed points (called foci) is kept constant.
 - Hyperbolas are split into two curves. The two closest points on opposite curves are called vertices and give the transverse axis.
- Standard form with center at the origin
 - $\frac{x^2}{a^2} \frac{y^2}{b^2} = 1$, with foci given by (c,0), (-c,0) and asymptotes $y = \pm \frac{bx}{a}$, where $a^2 + b^2 = c^2$
 - $\frac{y^2}{a^2} \frac{x^2}{b^2} = 1$, with foci given by (0,c), (0,-c) and asymptotes $x = \pm \frac{by}{a}$, where $a^2 + b^2 = c^2$
 - (Example 2) Sketch $\frac{y^2}{4} \frac{x^2}{16} = 1$, labeling its vertices, asymptotes, and foci.
 - (Example 3) Sketch $4x^2 9y^2 = 36$, labeling its vertices, asymptotes, and foci.
- Hypberbola with coordinate axis asymptotes
 - $xy = c^2$, with vertices given by (c, c) and (-c, -c)
 - $xy = -c^2$, with vertices given by (c, -c) and (-c, c)
 - (Example 5) Sketch xy = 4.
- HW: 1-14, 17-24

2.7 Translation of Axes

- Vertical/horizontal translation:
 - Shift right h: replace x with x h.
 - Shift up k: replace y with y k.
 - \blacksquare (Example 1) Give an equation of the parabola with vertex (2,4) and focus (4,4).
 - (Example 2) Sketch the curve with equation $\frac{(x-3)^2}{25} + \frac{(y+2)^2}{9} = 1$.
- HW: 1-36

1.2 Algebraic Functions

- Definition of a function y = f(x).
- Types of functions
 - Polynomials $P(x) = a_0 + a_1 x + \dots + a_n x^n$
 - Rational functions $R(x) = \frac{P(x)}{Q(x)}$ for polynomials P, Q
 - (Example 1) Voltage equals current multiplied by resistance. If the voltage at time t is given by $E(t) = 2t^2 = y + 5$ and the resistance at time t is given by R(t) = 3t + 20, then find a function I(t) which measures the current at time t. Identify it as a polynomial and/or rational function.
- Combinations of functions
 - Addition/Subtraction/Multiplication/Division
 - Compositions $f \circ g$ and $g \circ f$
 - (Example 2) Express f+g, $f \circ g$, and $g \circ f$ for the functions given by $f(x) = 2x^2 3$ and $g(x) = \sqrt{x+2}$.
- Domain/Range
 - The domain of a function is all real numbers which may be plugged into it without causing division by zero, even roots of negatives, or any other undefined operations.
 - The range of a function is all real numbers which may possibly be attained by the function.
 - (Example 5) Find the domain and range of $f(x) = x^2 + 2$ and $g(t) = \frac{1}{t+2}$.
 - (Example 7) Find the domain of $f(x) = 16\sqrt{x} + \frac{1}{x}$.
- Piecewise functions
 - Piecewise functions are defined differently for different parts of their domains.
 - (Example 9) Find the domain for

$$f(t) = \begin{cases} 8 - 2t & 0 \le t \le 4\\ 0 & t > 4 \end{cases}$$

and compute f(3), f(6), f(-1) if possible.

- Exponent laws
 - $a^m a^n = a^{m+n}$

$$(a^m)^n = a^{mn}$$

$$ab)^m = a^m b^m$$

$$a^0 = 1$$

$$a^{-n} = \frac{1}{a^n}$$

$$a^{1/n} = \sqrt[n]{a}$$

■ Note
$$\sqrt{a^2} = |a|$$
 but $\sqrt[3]{a^3} = a$

■ (Example 4) Simplify

$$f(x) = \frac{(3x^2 - 1)^{1/3}(2x) - (2x^3)(3x^2 - 1)^{-2/3}}{(3x^2 - 1)^{2/3}}$$

• HW: 1-18, 21-34

1.4 The Graph of a Function

- Overview of the table method for plotting function graphs.
- Examples:
 - (Example 1) Graph f(x) = 3x 5.
 - (Example 3) Graph y = 3x 5.
 - (Example 4) Graph $y = \sqrt{x+1}$.
 - (Example 6) Graph

$$f(x) = \begin{cases} 2x+1 & x \le 1\\ 6-x^2 & x > 1 \end{cases}$$

• HW: 1-12, 37-40

Remaining Topics

- 3.1 Limits
- 3.2 The Slope of a Tangent to a Curve
- 3.3 The Derivative
- 3.4 The Derivative as an Instantaneous Rate of Change

- 3.5 Derivatives of Polynomials
- 3.6 Derivatives of Products and Quotients of Functions
- 3.7 The Derivative of a Power of a Function
- 3.8 Differentiation of Implicit Functions
- 3.9 Higher Derivatives
- 4.1 Tangents and Normals
- 4.4 Related Rates
- 4.5 Using Derivatives in Curve Sketching
- 4.6 More on Curve Sketching
- 4.7 Applied Maximum and Minimum Problems
- 4.8 Differentials and Linear Approximations
- 5.1 Antiderivatives
- 5.2 The Indefinite Integral
- 5.3 The Area Under a Curve
- 5.4 The Definite Integral
- 7.1 The Trigonometric Functions
- 7.2 Basic Trigonometric Relations
- 7.3 Derivatives of the Sine and Cosine Functions
- 7.4 Derivatives of the Other Trigonometric Functions
- 8.1 Exponential and Logarithmic Functions
- 8.2 Derivative of the Logarithmic Functions
- 8.3 Derivative of the Exponentials Function
- 9.1 The General Power Formula
- 9.2 Basic Logarithmic Form
- 9.3 Exponential Form
- 9.4 Basic Trigonometric Forms