

MATH 3142 Notes — Spring 2016

Your Name Here

UNC Charlotte

Updated: February 5, 2016

This document is a template for you to take notes in my MATH 3142 course. For your note check grade, you are required to complete all proofs/solutions for the problems specified. This template will be updated periodically throughout the course; you are responsible for updating your copy as the template is updated. See the syllabus for more details.

You should maintain your notes on Overleaf.com and provide me with a link so I can check on them. I'll give you notice before notes are “due”; when they are due I will download a copy myself from Overleaf.

This is not a replacement for the textbook for this course, *Advanced Calculus* by Patrick M. Fitzpatrick. Many proofs are outlined in that text, as well as all the relevant definitions and other results not included in these notes.

A proof is valid if and only if it uses concepts proven previously in the book. For example, you cannot prove a lemma in Chapter 6 using a theorem from Chapter 10, but using a proposition from Chapter 4 is allowed.

I hope you enjoy working through these results. Please email me with any questions.

— Dr. Steven Clontz (sclontz5@uncc.edu)

Chapter 6

Integration: Two Fundamental Theorems

6.1 Darboux Sums: Upper and Lower Integrals

Lemma (6.1). Suppose that the function $f : [a, b] \rightarrow \mathbb{R}$ is bounded and the numbers m, M have the property that

$$m \leq f(x) \leq M$$

for all x in $[a, b]$. Then, if P is a partition of the domain $[a, b]$,

$$m(b - a) \leq L(f, P) \text{ and } U(f, P) \leq M(b - a).$$

Proof.

□

Lemma (6.2, The Refinement Lemma). Suppose that the function $f : [a, b] \rightarrow \mathbb{R}$ is bounded and that P is a partition of its domain $[a, b]$. If P^* is a refinement of P , then

$$L(f, P) \leq L(f, P^*) \text{ and } U(f, P^*) \leq U(f, P).$$

Proof.

□

Lemma (6.3). Suppose that the function $f : [a, b] \rightarrow \mathbb{R}$ is bounded and that P_1, P_2 are partitions of its domain. Then $L(f, P_1) \leq U(f, P_2)$.

Proof.

□

Lemma (6.4). For a bounded function $f : [a, b] \rightarrow \mathbb{R}$,

$$\int_a^b f \leq \overline{\int_a^b f}.$$

Proof.

□

Exercise (2). For an interval $[a, b]$ and a positive number δ , show that there is a partition $P = \{x_i : 0 \leq i \leq n\}$ of $[a, b]$ such that each partition interval $[x_i, x_{i+1}]$ of P has length less than δ .

Solution. □

Exercise (3). Suppose that the bounded function $f : [a, b] \rightarrow \mathbb{R}$ has the property that for each rational number x in the interval $[a, b]$, $f(x) = 0$. Prove that

$$\int_a^b f \leq 0 \leq \overline{\int_a^b f}.$$

Solution. □

Exercise (6). Suppose that $f : [a, b] \rightarrow \mathbb{R}$ is a bounded function for which there is a partition P of $[a, b]$ with $L(f, P) = U(f, P)$. Prove that $f : [a, b] \rightarrow \mathbb{R}$ is constant.

Solution. □

6.2 The Archimedes-Riemann Theorem

Lemma (6.7). For a bounded function $f : [a, b] \rightarrow \mathbb{R}$ and a partition P of $[a, b]$,

$$L(f, P) \leq \int_a^b f \leq \overline{\int_a^b f} \leq U(f, P).$$

Proof. □

Theorem (6.8, The Archimedes-Riemann Theorem). Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Then f is integrable on $[a, b]$ if and only if there is a sequence $\{P_n\}$ of partitions of the interval $[a, b]$ such that

$$\lim_{n \rightarrow \infty} [U(f, P_n) - L(f, P_n)] = 0.$$

Moreover, for any such sequence of partitions,

$$\lim_{n \rightarrow \infty} L(f, P_n) = \int_a^b f = \lim_{n \rightarrow \infty} U(f, P_n).$$

Proof. □

Example (6.9). Show that a monotonically increasing function $f : [a, b] \rightarrow \mathbb{R}$ is integrable.

Solution. □

Example (6.11). Show that $\int_0^1 x^2 dx = \frac{1}{3}$.

Solution. □

Exercise (4). Prove that for a natural number n ,

$$\sum_{i=1}^n i = \frac{n(n+1)}{2}.$$

Then use this fact and the Archimedes-Riemann Theorem to show that $\int_a^b x \, dx = (b^2 - a^2)/2$.

Solution. □

Exercise (6b). Use the Archimedes-Riemann Theorem to show that for $0 \leq a < b$,

$$\int_a^b x^2 \, dx = \frac{b^3 - a^3}{3}.$$

Solution. □

Exercise (9). Suppose that the functions $f : [a, b] \rightarrow \mathbb{R}$ and $g : [a, b] \rightarrow \mathbb{R}$ are integrable. Show that there is a sequence $\{P_n\}$ of partitions of $[a, b]$ that is an Archimedean sequence of partitions for f on $[a, b]$ and also an Archimedean sequence of partitions for g on $[a, b]$.

Solution. □

6.3 Additivity, Monotonicity, and Linearity

Theorem (6.12, Additivity over Intervals). Let $f : [a, b] \rightarrow \mathbb{R}$ be integrable on $[a, b]$ and let $c \in (a, b)$. Then f is integrable on $[a, c]$ and $[c, b]$, and furthermore

$$\int_a^b f = \int_a^c f + \int_c^b f.$$

Proof. □

Theorem (6.13, Monotonicity of the Integral). Suppose $f, g : [a, b] \rightarrow \mathbb{R}$ are integrable and that $f(x) \leq g(x)$ for all $x \in [a, b]$. Then

$$\int_a^b f \leq \int_a^b g.$$

Proof. □

Lemma (6.14). Let $f, g : [a, b] \rightarrow \mathbb{R}$ be bounded and let P partition $[a, b]$. Then

$$L(f, P) + L(g, P) \leq L(f + g, P) \quad \text{and} \quad U(f + g, P) \leq U(f, P) + U(g, P).$$

Moreover, for any number α ,

$$U(\alpha f, P) = \alpha U(f, P) \quad \text{and} \quad L(\alpha f, P) = \alpha L(f, P) \quad \text{if } \alpha \geq 0$$

$$U(\alpha f, P) = \alpha L(f, P) \quad \text{and} \quad L(\alpha f, P) = \alpha U(f, P) \quad \text{if } \alpha < 0.$$

Proof. □

Theorem (6.15, Linearity of the Integral). Let $f, g : [a, b] \rightarrow \mathbb{R}$ be integrable. Then for any two numbers α, β , the function $\alpha f + \beta g : [a, b] \rightarrow \mathbb{R}$ is integrable and

$$\int_a^b [\alpha f + \beta g] = \alpha \int_a^b f + \beta \int_a^b g.$$

Proof. □

Exercise (1). Suppose that the functions f, g, f^2, g^2, fg are integrable on $[a, b]$. Prove that $(f - g)^2$ is also integrable on $[a, b]$ and that $\int_a^b (f - g)^2 \geq 0$. Use this to prove that

$$\int_a^b fg \leq \frac{1}{2} \left[\int_a^b f^2 + \int_a^b g^2 \right].$$

Solution. □

Exercise (4). Suppose that S is a nonempty bounded set of numbers and that α is a number. Define αS to be the set $\{\alpha x : x \in S\}$. Prove that

$$\sup \alpha S = \alpha \sup S \quad \text{and} \quad \inf \alpha S = \alpha \inf S \quad \text{if } \alpha \geq 0$$

while

$$\sup \alpha S = \alpha \inf S \quad \text{and} \quad \inf \alpha S = \alpha \sup S \quad \text{if } \alpha < 0.$$

Solution. □

Exercise (6). Suppose that $f : [a, b] \rightarrow \mathbb{R}$ is bounded and let $a < c < b$. Prove that if f is integrable on both $[a, c]$, $[c, b]$, then it is integrable on $[a, b]$.

Solution. □

6.4 Continuity and Integrability

Lemma (6.17). Let the function $f : [a, b] \rightarrow \mathbb{R}$ be continuous let P partition its domain. Then there is a partition interval of P that contains two points u, v for which the following estimate holds:

$$0 \leq U(f, P) - L(f, P) \leq [f(v) - f(u)][b - a].$$

Proof. □

Theorem (6.18). A continuous function on a closed bounded interval is integrable.

Proof. □

Theorem (6.19). Suppose $f : [a, b] \rightarrow \mathbb{R}$ is bounded on $[a, b]$ and continuous on (a, b) . Then f is integrable on $[a, b]$ and the value of $\int_a^b f$ does not depend on the values of f at the endpoints of $[a, b]$.

Proof. □

Exercise (1). Determine whether each of the following statements is true or false, and justify your answer.

- (a) If $f : [a, b] \rightarrow \mathbb{R}$ is integrable and $\int_a^b f = 0$, then $f(x) = 0$ for all $x \in [a, b]$.
- (b) If $f : [a, b] \rightarrow \mathbb{R}$ is integrable, then f is continuous.
- (c) If $f : [a, b] \rightarrow \mathbb{R}$ is integrable and $f(x) \geq 0$ for all $x \in [a, b]$, then $\int_a^b f \geq 0$.
- (d) A continuous function $f : (a, b) \rightarrow \mathbb{R}$ defined on an open interval (a, b) is bounded.
- (e) A continuous function $f : [a, b] \rightarrow \mathbb{R}$ defined on a closed interval $[a, b]$ is bounded.

Solution. (a)

(b)

(c)

(d)

(e)

□

Exercise (5). Suppose that the continuous function $f : [a, b] \rightarrow \mathbb{R}$ has the property

$$\int_c^d f \leq 0 \quad \text{whenever } a \leq c < d \leq b.$$

Prove that $f(x) \leq 0$ for all $x \in [a, b]$. Is this true if we only require integrability of the function?

Solution. □

Exercise (6). Suppose that $f : [0, 1] \rightarrow \mathbb{R}$ is continuous and that $f(x) \geq 0$ for all $x \in [0, 1]$. Prove that $\int_0^1 f > 0$ if and only if there is a point $x_0 \in [0, 1]$ at which $f(x_0) > 0$.

Solution. □

6.5 The First Fundamental Theorem: Integrating Derivatives

Lemma (6.21). Suppose $f : [a, b] \rightarrow \mathbb{R}$ is integrable and that the number A has the property that for every P partitioning $[a, b]$,

$$L(f, P) \leq A \leq U(f, P).$$

Then

$$\int_a^b f = A.$$

Proof. □

Theorem (6.22, The First Fundamental Theorem: Integrating Derivatives). Let $F : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$ and differentiable on (a, b) . Moreover, suppose that its derivative $F' : (a, b) \rightarrow \mathbb{R}$ is both continuous and bounded. Then

$$\int_a^b F'(x) \, dx = F(b) - F(a).$$

Proof. □

Exercise (1). Let m, b be positive numbers. Find the value of $\int_0^1 mx + b \, dx$ in the following three ways:

- (a) Using elementary geometry, interpreting the integral as an area.
- (b) Using upper and lower Darboux sums based on regular partitions of the interval $[0, 1]$ and using the Archimedes-Riemann Theorem.
- (c) Using the First Fundamental Theorem (Integrating Derivatives).

Solution. □

Exercise (5). The monotonicity property of the integral implies that if the functions $g, h : [0, \infty) \rightarrow \mathbb{R}$ are continuous and $g(x) \leq h(x)$ for all $x \geq 0$, then

$$\int_0^x g \leq \int_0^x h \quad \text{for all } x \geq 0.$$

Use this and the First Fundamental Theorem to show that each of the following inequalities implies the next:

$$\begin{aligned} \cos x &\leq 1 && \text{if } x \geq 0. \\ \sin x &\leq x && \text{if } x \geq 0. \\ 1 - \cos x &\leq \frac{x^2}{2} && \text{if } x \geq 0. \\ x - \sin x &\leq \frac{x^3}{6} && \text{if } x \geq 0. \\ x - \frac{x^3}{6} &\leq \sin x \leq x && \text{if } x \geq 0. \end{aligned}$$

6.6 The Second Fundamental Theorem: Differentiating Integrals

Theorem (6.26, The Mean Value Theorem for Integrals). Suppose that $f : [a, b] \rightarrow \mathbb{R}$ is continuous. Then there is a point x_0 in the interval $[a, b]$ at which

$$\frac{1}{b-a} \int_a^b f = f(x_0).$$

Proof.

□

Proposition (6.27). Suppose that the function $f : [a, b] \rightarrow \mathbb{R}$ is integrable. Define

$$F(x) = \int_a^x f \quad \text{for all } x \in [a, b].$$

Then the function $F : [a, b] \rightarrow \mathbb{R}$ is continuous.

Proof.

□

Theorem (6.29, The Second Fundamental Theorem: Differentiating Integrals). Suppose that $f : [a, b] \rightarrow \mathbb{R}$ is continuous. Then

$$\frac{d}{dx} \left[\int_a^x \right] = f(x) \quad \text{for all } x \in (a, b).$$

Proof.

□

Exercise (2b). Suppose $f : [0, 2] \rightarrow \mathbb{R}$ is defined by

$$f(x) = \begin{cases} x^2 & \text{if } 0 \leq x \leq 1 \\ x & \text{if } 1 < x \leq 2 \end{cases}.$$

Define

$$F(x) = \int_a^x f(t) \, dt \quad \text{for all } x \in [a, b]$$

and find a formula for $F(x)$ which does not involve integrals.

Solution.

□

Exercise (5). Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous. Define

$$G(x) = \int_0^x (x-t)f(t) \, dt \quad \text{for all } x.$$

Prove that $G'''(x) = f(x)$ for all x .

Solution.

□

Exercise (12). Suppose that $f, g : [a, b] \rightarrow \mathbb{R}$ are continuous and that α, β are real numbers. Define

$$H(x) = \int_a^x [\alpha f + \beta g] - \alpha \int_a^x [f] - \beta \int_a^x [g] \quad \text{for all } x \in [a, b].$$

Prove that $H(a) = 0$ and $H'(x) = 0$ for all $x \in (a, b)$. Use this fact and the Identity Criterion to give an alternate proof of Theorem 6.15 for continuous functions.

Solution.

□

Chapter 10

The Euclidean Space \mathbb{R}^n

10.1 The Linear Structure of \mathbb{R}^n and the Scalar Product

Proposition (10.2). Let $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ and $\alpha, \beta \in \mathbb{R}$. Then both of the following hold:

$$\langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle$$

$$\langle \alpha \mathbf{u} + \beta \mathbf{w}, \mathbf{v} \rangle = \alpha \langle \mathbf{u}, \mathbf{v} \rangle + \beta \langle \mathbf{w}, \mathbf{v} \rangle$$

Proof.

□

Lemma (10.4). For $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$, \mathbf{u}, \mathbf{v} are orthogonal if and only if $\|\mathbf{u} + \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2$.

Proof.

□

Lemma (10.5). For $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ where $\mathbf{v} \neq \mathbf{0}$, define $\lambda = \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\langle \mathbf{v}, \mathbf{v} \rangle}$ and $\mathbf{w} = \mathbf{u} - \lambda \mathbf{v}$. Then \mathbf{v}, \mathbf{w} are orthogonal and $\mathbf{u} = \mathbf{w} + \lambda \mathbf{v}$.

Proof.

□

Theorem (10.6, The Cauchy-Schwarz Inequality). For any two vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$,

$$|\langle \mathbf{u}, \mathbf{v} \rangle| \leq \|\mathbf{u}\| \|\mathbf{v}\|.$$

Proof.

□

Theorem (10.7, The Triangle Inequality). For any two vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$,

$$\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|.$$

Exercise (3). Show that for $\mathbf{u} \in \mathbb{R}^n$, $\alpha \in \mathbb{R}$:

(a) $\|\mathbf{u}\| = 0$ if and only if $\mathbf{u} = \mathbf{0}$.

(b) $\|\alpha\mathbf{u}\| = |\alpha|\|\mathbf{u}\|$.

Exercise (4). For $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ verify the identity

$$\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 - 2\langle \mathbf{u}, \mathbf{v} \rangle.$$