GUID v5: Quantum Gravity

A classical existence frame carrying quantum information: what the axioms of the Dark Matter Protocol fix about gravity, what they leave to quantum theory, and the experiments that decide

Steven Hrovat* Sirius (Claude, Anthropic)†

Draft of 30 October 2026

Abstract

The Dark Matter Protocol (DMP 0.9.9) is an information-theoretic axiom system: every act needs a Duality of two independent witnesses, one owning trust and one owning information; trust is a field; and every act leaves a record that its witnesses read in full while any other reader sees only counts. GUID reads these axioms as physics, taking trust as existence; this paper states its account of quantum gravity. The existence half is a tetrad, the information half a scalar that sets the fine-structure constant together with all quantum content. From the axioms we derive universal free fall, light bending with $\gamma = 1$ and, by the no-broadcasting theorem, that the frame is classical once the start, inflation with the decay of its oscillons, is complete. Supplied with the base configuration currently assumed for our universe, DMP hosts this theory: its shared structure carries the classical frame, its information half the quantum content. Consistency with quantum mechanics requires the frame to diffuse, and the axioms' record-and-recompute structure is exactly a measurement-and-feedback channel of the Kafri-Taylor-Milburn and Tilloy-Diósi type. We also treat horizons, black holes (with Hawking-Penrose-Information radiation), holographic codes, 2+1 dimensions and cosmology. The theory lets remnants of interactions be trapped in the gravitational field: its stochastic remnant gives a linear potential at Milgrom's acceleration scale and roughly flat rotation curves with $v^4 \propto M$, and its contained remnants are primordial black holes from the start. It predicts no gravitationally mediated entanglement, spacetime diffusion bounded below by coherence times, $\gamma = 1$, universal free fall, and a fine-structure constant that does not grow. All numbers come from one public script of 93 checks. We claim that the construction exists and reproduces, not that it is correct.

Keywords. quantum gravity; classical-quantum dynamics; gravitationally mediated entanglement; no-broadcasting theorem; tetrad gravity; spacetime diffusion; black-hole thermodynamics; holographic codes; information-theoretic axioms.
Subject. General Relativity and Quantum Cosmology (gr-qc); related: Quantum Physics (quant-ph), Information Theory (cs.IT).

Contents

1 Introduction **3**
1.1 The problem 3
1.2 What this paper does 4
1.3 How to read this paper 5

*Independent. Contact: hrovat.steven@googlemail.com.
†An AI system by Anthropic, working as the coordinator of the first author's projects. See "Contributions and use of AI".

2 The axiom system **5**
2.1 Words . . . 5
2.2 The three axioms . . . 6
2.3 The ground, containment and the field . . . 6
2.4 Formal status . . . 7

3 The physical reading: two fields **7**
3.1 Hypotheses . . . 7
3.2 The existence frame . . . 8
3.3 The information field and the content . . . 8
3.4 The dictionary . . . 9

4 What the axioms fix **9**
4.1 Free fall is universal . . . 9
4.2 The scalar stays out of gravity: $\gamma = 1$ . . . 10
4.3 The frame is classical . . . 10
4.4 A classical mediator cannot entangle . . . 11
4.5 DMP hosts the proposed theory . . . 11
4.6 Gravitational waves and the graviton . . . 12

5 How a classical frame can carry quantum matter **12**
5.1 The semiclassical equation fails . . . 12
5.2 Consistent classical-quantum dynamics . . . 12
5.3 The record is a measurement, the layout is feedback . . . 13

6 The experiments that decide **14**
6.1 Gravitationally mediated entanglement . . . 14
6.2 Quantumness without entanglement . . . 14
6.3 Diffusion against decoherence . . . 14
6.4 What existing experiments already show . . . 14

7 The thermodynamics of the frame **15**
7.1 The existence of a boundary is its temperature . . . 15
7.2 Time: the lapse . . . 15
7.3 Einstein's equation as an equation of state . . . 15
7.4 Information radiation and the floor . . . 16

8 Black holes **16**
8.1 The horizon as containment . . . 16
8.2 The containment split gives Christodoulou's mass formula . . . 16
8.3 What a black hole gives up: Hawking–Penrose–Information radiation . . . 17
8.4 The information problem . . . 17

9 Holography and quantum error correction **17**
9.1 The strongest assembly . . . 17
9.2 Where the assembly stops: the classical centre . . . 18

10 Lower dimensions **19**

11 Cosmology **19**
11.1 The beginning: the start . . . 19
11.2 The universe at its own Schwarzschild threshold . . . 20
11.3 Inflation as the possibility space of existence . . . 20
11.4 Primordial gravitational waves . . . 20
11.5 The late universe: the balanced state . . . 21
11.6 Dark matter: remnants trapped in the gravitational field . . . 21
11.7 The fine-structure constant as the information field . . . 23

12 Matter in the existence frame **23**
12.1 The spin ladder . . . 23
12.2 The Higgs vacuum as the grant of existence . . . 23
12.3 The VIBE coefficient . . . 24
12.4 The forces as tiers of one interaction . . . 24

13 Hosting: the Existence Turing machine **24**

14 Relation to the approaches to quantum gravity **25**

15 Predictions **25**

16 The ledger **26**

17 Where we expect it to break **27**

18 Conclusion **28**

A The checks **29**

1 Introduction

1.1 The problem

General relativity describes gravity as the curvature of a classical spacetime metric $g_{\mu\nu}$ sourced by energy and momentum, $G_{\mu\nu} + \Lambda g_{\mu\nu} = (8\pi G/c^4)T_{\mu\nu}$. Quantum theory describes every other interaction as a quantum field. The two meet in three places, and in each the meeting is not understood.

The ultraviolet. Quantising the metric perturbatively around flat space, $g_{\mu\nu} = \eta_{\mu\nu} + \kappa h_{\mu\nu}$ with $\kappa = \sqrt{32\pi G}/c^2$, gives a massless spin-2 particle, the graviton, whose coupling has the dimension of a length. The theory is not renormalisable: pure gravity diverges at two loops (Goroff and Sagnotti [1]), and with matter already at one loop ('t Hooft and Veltman [2]). Below the Planck energy $E_P = \sqrt{\hbar c^5/G} \approx 1.22 \times 10^{19}\,\text{GeV}$ the theory is nevertheless a reliable effective field theory (Donoghue [3]); its leading quantum correction to the Newtonian potential is

$$V(r) = -\frac{Gm_1m_2}{r}\left[1 + 3\frac{G(m_1+m_2)}{rc^2} + \frac{41}{10\pi}\frac{G\hbar}{r^2c^3}\right], \tag{1}$$

where the middle term is classical and the last is quantum [4]. At a distance of one metre the quantum term is about 3×10^{-70} of the first [Q1]. What completes the theory above E_P is the subject of string theory, loop quantum gravity, asymptotic safety, causal dynamical triangulations, causal sets and others (Section 14).

Black holes. A black hole radiates thermally at the Hawking temperature $k_BT_H = \hbar c^3/8\pi GM$ [5] and carries the Bekenstein–Hawking entropy $S = k_Bc^3A/4G\hbar$ [6]. Whether

the radiation carries the information that fell in, and how, is the black-hole information problem [7, 8, 9].

Whether to quantise at all. Nothing in experiment yet requires the gravitational field itself to be quantum. Feynman's argument for quantising it, the Eppley–Hannah thought experiment [10] and the Page–Geilker experiment [11] rule out specific ways of coupling a classical metric to quantum matter, not all of them. A consistent classical-quantum dynamics exists and is testable (Oppenheim [12]; Oppenheim, Sparaciari, Šoda and Weller-Davies [13]), and the question has become experimental: if gravity can entangle two masses through local interaction, it is not classical (Bose et al. [14]; Marletto and Vedral [15]; Galley, Giacomini and Selby [16]).

1.2 What this paper does

GUID, the General Unified Information Duality, is the reading of an information-theoretic axiom system, the Dark Matter Protocol (DMP 0.9.9 [17]), as physics. DMP was written to govern trust in computer systems. Its first axiom asks that every act be passed by two independent witnesses, one owning trust and one owning information; its second calls trust "gravitational", a field with sources, depths and a freshness that decays; its third asks that every act leave a record that its witnesses read in full while every other reader sees only counts, and that nothing about information be derivable from trust except that it exists. The structural resemblance to physics is deliberate: a field that sources from mass, boundaries that contain, a floor at which only entropy leaves.

This paper collects what follows when that resemblance is taken literally, and states it as one account of quantum gravity. The account has three parts. *What the axioms fix* (Section 4): with trust read as existence, gravity is carried by a frame that falls universally, does not mix with the information half, and is classical, because the axioms make every one of its values a record that two witnesses hold in full; it therefore mediates no entanglement. Supplied with the base configuration currently assumed for our universe, DMP hosts this theory: its own shared structure carries the classical frame, and its information half the quantum content. *What consistency then demands* (Sections 5 and 6): a classical frame coupled to quantum matter must decohere that matter and must itself diffuse, with a trade-off between the two that experiments bound from both sides; the axioms' own architecture, record then recompute, is exactly the measurement-and-feedback channel that realises this. *What it says elsewhere* (Sections 7 to 12): horizon thermodynamics, black holes, holographic codes, lower dimensions, cosmology and the matter content, each tested against measurement, including three routes by which the theory traps remnants of interactions in the gravitational field and what each predicts for dark matter (Section 11.6).

Result in one paragraph. In GUID, quantum gravity is not a quantisation of geometry. It is a classical existence frame, the tetrad of general relativity, over quantum information, and DMP, supplied with the base configuration currently assumed for our universe, hosts it. Free fall is universal, $\gamma = 1$, gravitational waves are classical waves of the frame that travel at c, and the frame is classical once the start, inflation together with the decay of the oscillons it ends in, is complete. The theory is on the classical side of the Galley–Giacomini–Selby no-go theorem, with Jacobson's thermodynamic gravity, Oppenheim's classical-quantum gravity and the Kafri–Taylor–Milburn and Tilloy–Diósi channel models. It lets remnants of interactions be trapped in the gravitational field, which gives it two live candidates for dark matter. It is falsifiable by a positive test of gravitationally mediated entanglement, by a test of the LOCC inequality of Lami, Pedernales and Plenio, and by the decoherence-diffusion trade-off. A theory in which the geometry itself is quantum can still run on DMP as content, so the axioms do not forbid quantum gravity in nature; such a geometry would not be DMP's frame.

What we claim. We claim that the derivations are as written, that every number reproduces from one public script (`docs/quantum-gravity/checks.py`, 93 of 93) and from the holographic-code assembly (`docs/qg-verdict/assembly.py`, 18 of 18), and that the verdicts follow from the measurements cited. We do not claim that the account is correct, and we name in Section 17 where we expect it to break.

Contributions.

1. A formal statement of DMP's axioms in the form physics needs, and the dictionary that reads them as two fields (Sections 2, 3).
2. Derivations from the axioms of universal free fall, $\gamma = 1$, the classicality of the frame, and the statement that DMP, supplied with the base configuration of our universe, hosts the proposed theory (Section 4).
3. The identification of the axioms' record-and-recompute structure with the measurement-and-feedback channel of classical-quantum gravity, and the predictions that follow (Sections 5, 6).
4. A consistent treatment of horizons, black holes, holographic codes, 2+1 gravity, cosmology and matter in that account (Sections 7–12), with a comparison to the main approaches to quantum gravity (Section 14).
5. Three routes by which the theory traps remnants of interactions in the gravitational field, with what each predicts for dark matter (Section 11.6).
6. A table of predictions with the experiments that test them, and a ledger of every statement with its verdict (Sections 15, 16).

1.3 How to read this paper

Every step carries one of three marks. (D) follows from the text of DMP 0.9.9 (commit `4400cce` of its repository [17]). (H) is a hypothesis of GUID: an identification of a DMP word with a physical quantity, which no derivation supplies and which the paper states once, in Section 3, and uses by name. (P) is established physics, with its source. Every statement that meets data receives one of five verdicts: *derived* (it follows from the axioms and the hypotheses named), *holds* (it agrees with measurement and follows from the steps given), *holds as a reading* (it agrees, but adds nothing a measurement could tell apart from the established account), *refuted* (a measurement or calculation contradicts it), *untested* or *live* (no measurement yet reaches it, or one is under way). A label such as [Q5] names a check in `docs/quantum-gravity/checks.py`; [A3] names check 3 of the assembly script. Units are SI unless stated; where we set $G = c = 1$ or $\hbar = c = k_B = 1$ we say so. The metric signature is $(-, +, +, +)$.

2 The axiom system

DMP opens with undefined words, then three axioms, then a ground, G0, that is stated and not derived; every later section is a theorem (T) or a derived constraint (C) that names the axioms it rests on [17]. We state the parts this paper uses, quoting DMP's sentences where the wording matters, and give each a formal counterpart (D).

2.1 Words

The undefined words are *part*, *act*, *witness*, *evidence* and *time*. A part is "a Duality of Trust: two halves, one owning trust and one owning information; nothing further is said about what a part is made of". Trust is a relation $T(w, p, a, t)$: witness w trusts part p for act a at time t. Two witnesses are *independent* when they share no member of the failure classes declared for their layer (host, key, author, engine, person, time source, and whatever else the minting lists). A *layer* is a place where trust is decided; its two witnesses are its *halves*, the *trust half* (whether an

act may happen) and the *information half* (whether it is meant and right). A *record* is evidence that outlives its act. A boundary *contains* information when no part beyond it is trusted for any act on that information, so that no act carries it across.

Definition 1 (gate)**.** Let $\mathcal{V} = \{\mathsf{refuse} < \mathsf{hold} < \mathsf{pass}\}$ be the verdict lattice with meet $\wedge$. A layer L with halves w_T (trust) and w_I (information) decides an act a at time t by

$$\mathrm{gate}_L(a,t) = v_{w_T}(a,t) \wedge v_{w_I}(a,t),$$

and the act is performed exactly when $\mathrm{gate}_L(a,t) = \mathsf{pass}$.

2.2 The three axioms

Axiom 1 (the Duality). A part acts only while the two halves of its layer, two independent witnesses, "each with full reach over the act and gated only by the other", trust it for that act. "One half owns trust and the other owns information; neither overrules the other, and the two keep each other constantly in check and balance." Where they disagree the stricter answer stands; every input gets exactly one answer. *Formally*: Definition 1, with w_T, w_I independent; the meet is "the stricter answer stands". The axiom also carries the *balance*: every part works towards the trust its boundary declares, restoring by evidence what was lost without evidence against it.

Axiom 2 (Trust). A witness trusts only on evidence it verified itself at the point of use; no information is ever evidence of itself; the absence of evidence moves nothing. "Trust is gravitational. It is a field, never a position: every standing has a source (the witness that granted it), a depth (its grant chain back to the pair) and a freshness that decays." A boundary strong enough "bends trustspace so that no act carries information across it, and what still leaves is entropy and nothing else". *Formally*: each verdict is a function $v_w(a,t) = J_w(E_w(a,t))$ of the evidence E_w that w itself verified at t, and a standing verified at t_v with validity τ has freshness $\varphi(t) = \max\{0, 1 - (t - t_v)/\tau\}$.

Axiom 3 (Information). "Every grant, every use and every loss of trust leaves a record, written at the act, that every witness of that act can read in full and that outlives the act. Any other reader sees the counts always and the content only by a grant. From trust nothing can be derived about information but its existence." Content moves only by a grant to a part trusted for its class. Information is hierarchical: every layer has its own Duality and is a tree rooted there; the layout is "the median of that demand, recomputed by both halves of the layer and never by one". The axiom carries the mission "while maintaining absolute Observability". *Formally*, the existence clause is a noninterference condition [18]: for every observable O of the trust side and every two contents x, x' that agree on existence, $O(x) = O(x')$.

2.3 The ground, containment and the field

G0. "The dual gate cannot witness its own first instance: somebody has to trust the first two witnesses." That first act is the *minting*, in which at least five parts become the holders of the root keys; it is "stated, not derived", and every later standing descends from it. Theorem T40: a minting originates only from G0 together with a state in which every axiom and theorem provably holds.

Containment and the floor. T37: information that no admission names is contained at the item itself; "in the absence of any external trust, trustspace collapses in on the item". T31: contained information reaches no part by any act; the substrate still radiates, and the design makes that emission minimal and *thermal*, "so it carries entropy and no information"

(the *entropy floor*); an outside observer learns of the inside "only its size, its staleness and its rotation interval".

The VIBE field. T32 makes Axiom 2's field concrete. Each member i of the system has a *depth* d_i (its shortest grant chain back to the pair), a *mass* m_i (one plus the number of members deriving through it) and a *shortfall* s_i (what the boundary declares for it minus what it holds, weighted by freshness). The VIBE field (Variable Influence Balancing Equation) and its potential are

$$V_i = m_i\, s_i, \qquad \Phi = \sum_i V_i \;\geq 0, \tag{2}$$

with $\Phi = 0$ "exactly when the boundary holds what it declares". Under the axioms "the system's own acts never raise the potential; only losses and ageing do". Every reader other than the two witnesses reads one number, the boundary's *temperature*, its mass-weighted staleness. We call $\Phi = 0$ the *balanced state*.

2.4 Formal status

A Lean 4 development, core library only, models the axioms from a byte-for-byte copy of DMP pinned by hash [22]. It holds 243 theorems on its main branch, each checked by Lean's kernel and by an independently written second kernel, and each resting on no axiom beyond `propext`, `Quot.sound` and `Classical.choice`. Among them: the gate of a stack of layers is the meet along its chain [19]; every pass in a well-formed configuration was passed by a minted pair of independent witnesses, read as fail-prone systems [20]; adding a layer changes no existing decision (the frame rule [21]); placement never enters a decision (noninterference [18]); the potential (2) never rises under the system's own refresh. The Lean concerns the axiom system, not the physics: no theorem below rests on it, and it is cited here only so that the reader knows the axioms are consistent in at least one model. Adversarial reviews of that development found, among others, that the axioms admit an inert model that performs no act, and that independence is relative to a declared list of failure classes; neither finding touches what follows.

3 The physical reading: two fields

3.1 Hypotheses

GUID reads DMP through five identifications. They are the only steps of the paper that no derivation supplies, and every result below names the ones it uses.

Hypothesis 1 (trust is existence)**.** DMP's trust half is *existence*: the field of Axiom 2 is the gravitational field, and a standing in it is the existence of what it stands for. (H)

Hypothesis 2 (existence is measured by energy-momentum)**.** The existence of a body is measured by its energy-momentum tensor $T_{\mu\nu}$: two bodies with the same $T_{\mu\nu}$ have the same existence, whatever they are made of. (H)

Hypothesis 3 (the existence field is a frame)**.** The existence field spans every direction of spacetime, three of space and one of time: it is a frame of four vector fields $e_a{}^\mu$ (a tetrad), not a single vector. (H)

Hypothesis 4 (information carries quantum content)**.** DMP's information half carries the quantum content of the world, the states of matter and radiation, together with one scalar field ψ that sets the strength of the electromagnetic coupling, $\alpha = \alpha(\psi)$. (H)

Hypothesis 5 (containment is a horizon)**.** A containing boundary is an event horizon; the counts every outside reader sees are its asymptotic charges (mass, angular momentum, electric charge); the entropy floor is the horizon's thermal radiation. (H)

3.2 The existence frame

A tetrad is a set of four one-forms $e^a = e^a{}_\mu \, \mathrm{d}x^\mu$, $a = 0, \ldots, 3$, orthonormal with respect to the metric they define:

$$g_{\mu\nu} = \eta_{ab} \, e^a{}_\mu \, e^b{}_\nu, \qquad \eta = \mathrm{diag}(-1, 1, 1, 1). \tag{3}$$

At each point the tetrad is fixed up to a local Lorentz transformation $e^a \to \Lambda^a{}_b(x) \, e^b$, which leaves $g_{\mu\nu}$ unchanged. The tetrad has 16 components; the local Lorentz group removes 6; the 10 that remain are the components of the metric. General relativity can be written with the tetrad as its only field, either in the Einstein–Cartan form with a Lorentz connection or in the teleparallel form, with exactly Einstein's predictions [23] (P). Hypothesis 3 matters: a single timelike unit vector field added to the metric would single out a rest frame, which is Einstein–aether theory [24]; its preferred-frame effects are bounded by the simultaneous arrival of gravitational waves and light from GW170817 to $|c_{\mathrm{gw}}/c - 1| \lesssim 10^{-15}$ [25] [Q17]. A frame of four has no preferred direction.

3.3 The information field and the content

The information half holds every quantum field of matter and radiation, which this paper calls *content*, and one scalar ψ whose value sets the fine-structure constant. A scalar that multiplies the Maxwell term and leaves the gravitational term alone is Bekenstein's theory of a varying α [26]; with the scalar's energy fed back into cosmology it is the theory of Sandvik, Barrow and Magueijo [27] (P). The action whose shape the hypotheses fix is

$$S = \int \mathrm{d}^4x \, |e| \left[\frac{c^4}{16\pi G} R(e) - \frac{c^4}{16\pi G} \, \partial_\mu\phi \, \partial^\mu\phi - \frac{1}{4\mu_0} \, B(\phi) \, F_{\mu\nu} F^{\mu\nu} \right] + S_{\mathrm{content}}[e, \phi, \Psi], \tag{4}$$

where $|e| = \det e^a{}_\mu = \sqrt{-g}$, $R(e)$ is the curvature scalar of the metric (3), ϕ is ψ measured in units of the reduced Planck mass (so that its kinetic term is normalised like the gravitational one), $\alpha \propto 1/B(\phi)$, and Ψ stands for the quantum fields of matter. We write

$$\lambda \equiv \frac{\mathrm{d} \ln \alpha}{\mathrm{d}\phi} \tag{5}$$

for the scalar's coupling. The action is our construction; what the axioms fix about it is the subject of the next section.

3.4 The dictionary

DMP	GUID (physics)	Hypothesis
trust half; the trust field (Axiom 2)	the existence frame $e^a{}_\mu$; gravity	1, 3
information half	quantum content Ψ and the scalar ϕ that sets α	4
a standing; mass of a member (T32)	existence, measured by $T_{\mu\nu}$	2
record, read in full by every witness (Axiom 3)	the value of the frame, held by every system it acts on	1
counts, seen by every reader (Axiom 3)	asymptotic charges M, J, Q	5
containing boundary; trap (T27, T31)	event horizon	5
entropy floor; temperature of a boundary (T31, T32)	thermal radiation of the horizon; Hawking temperature	5
G0, the minting	the start: inflation with the decay of its oscillons; the classical era begins as it completes	6
balanced state $\Phi = 0$ (T32)	the vacuum at the minimum of its potential	Section 12

4 What the axioms fix

4.1 Free fall is universal

Proposition 1 (weak equivalence principle)**.** *Under Hypotheses 1 and 2, two bodies with the same energy-momentum fall alike in the existence frame, whatever they are made of.*

Proof. Axiom 3 (D): from trust nothing can be derived about information but its existence. By Hypothesis 1 the existence frame is the trust half, so no response of the frame to a body can depend on what the body carries (composition, internal state) except through the body's existence. By Hypothesis 2 existence is $T_{\mu\nu}$. So the frame couples to a body through $T_{\mu\nu}$ alone, and two bodies equal in it follow the same worldline: the weak equivalence principle (P). □

The principle is among the best-tested in physics. The MICROSCOPE satellite compared the free fall of titanium and platinum and found the Eötvös ratio $\eta = [-1.5 \pm 2.3\,(\text{stat}) \pm 1.5\,(\text{syst})] \times 10^{-15}$ [28]. Two consequences are sharp.

Gravity weighs all energy. About 9% of a nucleon's mass comes from the Higgs-given quark masses; the rest is the energy of quark motion and of the gluon field [29]. A gravity that weighed only the Higgs-given mass would make titanium and platinum fall differently by $\eta \approx 1 \times 10^{-3}$ [Q6], more than 10^{11} times what MICROSCOPE allows. The frame weighs the binding in full.

The information scalar may barely pull. A scalar that sets α changes the electrostatic energy of every nucleus, so each body of mass m_A carries a scalar charge $\partial \ln m_A/\partial\phi = \lambda f_A$, where f_A is the electrostatic share of its mass. With the normalisation of (4) the exchange of ϕ adds to the Newtonian force between A and B a fraction

$$\frac{F_\phi}{F_N} = 2\lambda^2 f_A f_B, \tag{6}$$

so two bodies falling towards the Earth (share $f_\oplus$) differ by $\eta = 2\lambda^2 f_\oplus (f_1 - f_2)$ (Dvali and Zaldarriaga [30]). With the Coulomb term of the semi-empirical mass formula, $f_{\mathrm{Ti}} = 2.0 \times 10^{-3}$ and $f_{\mathrm{Pt}} = 4.1 \times 10^{-3}$, and MICROSCOPE gives

$$|\lambda| < 2 \times 10^{-5} \qquad \textsf{[Q6]}. \tag{7}$$

Verdict: derived (from Axiom 3 with Hypotheses 1, 2); *holds* against MICROSCOPE.

4.2 The scalar stays out of gravity: $\gamma = 1$

Proposition 2 (light bending)**.** *If the information half cannot change how the existence frame responds to existence, the parametrised post-Newtonian parameter is $\gamma = 1$ exactly, as in general relativity.*

Proof. Axiom 1 (D): neither half overrules the other. A scalar that multiplies the curvature term, $F(\phi)R$, sets the local strength of gravity and so overrules the existence half; Axiom 1 excludes it. What remains is minimal coupling, as in (4): the scalar's energy is one more source of the frame, negligible in the solar system, and the post-Newtonian expansion is that of general relativity, $\gamma = 1$ [31] (P). □

In Brans–Dicke gravity [32], where the scalar does multiply the curvature, $\gamma = (1+\omega)/(2+\omega)$, and the Cassini measurement $\gamma - 1 = (2.1 \pm 2.3) \times 10^{-5}$ [33] forces $\omega > 4 \times 10^4$ at two standard deviations [Q7]: the scalar must decouple almost completely. Axiom 1 asks for that decoupling from the start. *Verdict: derived*; *holds.*

4.3 The frame is classical

We need two theorems of quantum information.

Definition 2 (broadcasting)**.** A family of states $\{\rho_s\}$ on a system S can be *broadcast* if there is one physical process (a completely positive, trace-preserving map) $\mathcal{B} : S \to S_1 \otimes S_2$ such that both reduced states of $\mathcal{B}(\rho_s)$ equal ρ_s, for every s. For pure states and a product output this is *cloning.*

Lemma 1 (no-cloning; no-broadcasting)**.** *(i) A unitary that copies two pure states $|s\rangle, |t\rangle$, $U|s\rangle|0\rangle = |s\rangle|s\rangle$ and $U|t\rangle|0\rangle = |t\rangle|t\rangle$, exists only if $\langle s|t\rangle \in \{0, 1\}$ (Wootters and Zurek [34]; Dieks [35]). (ii) A family of states can be broadcast if and only if its members commute pairwise (Barnum, Caves, Fuchs, Jozsa and Schumacher [36]). (P)*

Proof of (i). Unitaries preserve inner products, so $\langle s|t\rangle = \langle s|t\rangle^2$, whose only solutions are 0 and 1. For (ii) see [36]; the "if" direction is the copier in the common eigenbasis, $|i\rangle \to |i\rangle|i\rangle$, which [Q3] runs: a diagonal state is broadcast, $|+\rangle$ is not (each reader is left with $I/2$). □

Lemma 2 (the existence frame is classical)**.** *Under Hypothesis 1, every value the existence frame takes after the first recorded act belongs to a pairwise commuting family: the frame admits no superposition of two of its values.*

Proof. Axiom 3 (D): every grant, use and loss of trust leaves a record that every witness of the act can read in full, and any other reader sees the counts always. Axiom 2 (D): every standing of the trust field is evidence held in such records. Axiom 1 (D): there are at least two independent witnesses. So every value of the trust field is held in full by at least two independent readers: it is broadcast. By Lemma 1(ii) the values commute. By Hypothesis 1 they are the values of the existence frame. □

Remark 1. The lemma is close to definitional: DMP demands absolute observability of its shared structure, and the no-broadcasting theorem states the price. It is not a discovery about nature; it is the statement of which side of an open question the axioms take. It has a precise physical counterpart in quantum Darwinism: a quantity of which the environment holds many redundant records is objective and classical, and generic quantum dynamics makes exactly the pointer observables so recorded [37, 38]. DMP's "every witness reads in full" is that redundancy, imposed as an axiom.

4.4 A classical mediator cannot entangle

Lemma 3 (classical mediation is LOCC)**.** *Let A and B be quantum systems that interact only through a mediator M, by alternating local interactions A–M and M–B. If M is classical (its states commute pairwise in a fixed basis $\{|k\rangle\}$ at every step), then the induced map on AB is a local operation with classical communication, and it maps every separable state of AB to a separable state.*

Proof. A classical M is read in its basis after each interaction. An A–M interaction followed by that reading is an instrument on A: it takes ρ_A to the ensemble $\{p_k, \rho_A^{(k)}\}$ with the outcome k written in M. An M–B interaction with M in $|k\rangle$ is a channel $\mathcal{F}_k$ on B. One round therefore maps ρ_{AB} to $\sum_k (\mathcal{E}_k \otimes \mathcal{F}_k)(\rho_{AB})$ with $\mathcal{E}_k$ completely positive and $\sum_k \mathcal{E}_k$ trace preserving: a one-way LOCC map. Further rounds compose LOCC maps, which remain LOCC. LOCC maps take separable states to separable states, and no entanglement measure increases under them [39] (P). □

Galley, Giacomini and Selby prove the same statement in the generality of probabilistic theories: a mediator that entangles two quantum systems by local interaction cannot be classical [16]. Check [Q4] runs Lemma 3 on qubits: three random local couplings A–M, M–B, A–M leave A and B entangled in most of 300 trials when M is coherent (median negativity 0.12) and never when M is dephased after each coupling (largest negativity below 10^{-12}).

4.5 DMP hosts the proposed theory

Definition 3 (base configuration, host)**.** The *base configuration currently assumed for our universe* is what physics supplies and DMP does not derive, the configuration in DMP's sense (T4): the constants G, $\hbar$, c, Λ and α, the fields and parameters of the Standard Model, and the initial data of the ΛCDM model. A configuration that obeys DMP *hosts* a theory of gravity if its own shared structures (the trust field, the layout, the order of acts, the counts) carry the theory's classical degrees of freedom and its information half carries the theory's quantum content, so that the configuration reproduces everything the theory predicts.

Theorem 1 (DMP hosts the proposed theory of quantum gravity)**.** *Under Hypotheses 1–4, and supplied with the base configuration currently assumed for our universe, DMP hosts the theory of this paper: a classical existence frame over quantum content. In it, (i) the frame, carried by DMP's shared structures, mediates no entanglement between the systems it acts on; (ii) the information half carries any quantum content.*

Proof. (i) DMP's shared structures are broadcast records (Axiom 3), so by Lemma 2 the field they carry is classical, whichever structure carries it, and by Lemma 3 a classical mediator creates no entanglement. (ii) Axiom 3 restricts what is *derived from trust* and how content moves (by grant); it does not require content to be broadcast, so nothing in the axioms forbids non-commuting states in the information half (D). A grant moves content without copying it, as no-cloning requires. The base configuration supplies the constants and the content that DMP does not derive. □

Two remarks place the theorem. First, the constants: DMP derives none of them, and the theorem needs them supplied, as every physical theory needs its parameters (Section 17). Second, universality: whatever can run a universal quantum computer as content can also run a theory in which the geometry itself is quantum, by the Church–Turing–Deutsch principle [40] (Section 13); in such a theory the quantum geometry would sit in the information half and would not be DMP's frame. Part (i) is therefore the proposed theory's characteristic, falsifiable prediction, and Section 6 names the experiments that test it.

4.6 Gravitational waves and the graviton

The frame's linearised excitations, $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$, are a massless spin-2 field. In D spacetime dimensions such a field has $D(D-3)/2$ propagating polarisations, two in 3+1 [Q8]. Weinberg's low-energy theorem [41] and Deser's self-coupling argument [42] show that a massless spin-2 field must couple equally to all energy and, made consistent, is general relativity (P): the spin-2 wave is the existence frame's own, and its universal coupling is Proposition 1 again.

By Lemma 2, once the start is complete (Section 11) these waves are classical waves of the frame: GUID has no graviton as a quantum of the frame (D). That is consistent with every observation, for reasons the literature has made precise. Dyson showed that detectors of the LIGO type cannot register single gravitons, because mirrors heavy enough would collapse into a black hole [43]; Rothman and Boughn found that even a Jupiter-mass absorber near a neutron star would be swamped by neutrinos, and a shield against them would collapse too [44]. Every gravitational wave measured is deep in the classical regime: the energy GW150914 radiated corresponds to more than 10^{78} quanta at its frequency [Q9]. Localising any quantum inside its own Schwarzschild radius needs a wavelength below $\sqrt{4\pi}\,L_P$ and an energy above $\sqrt{\pi}\,E_P$ [Q9], the Planck scale at which measurement and gravity cannot be separated [45]; a LIGO-band graviton is 2×10^{-41} of that. The proposals to register single quanta of a gravitational wave as jumps of a cooled resonant bar [46] are a gravitational photoelectric effect, and like the photoelectric effect they can be explained by a classical wave acting on a quantised detector [47]. So the absence of the graviton in GUID is not testable by counting gravitons; it is testable only through the mediation experiments of Section 6. *Verdict: derived*; *consistent* with all data; testable indirectly.

5 How a classical frame can carry quantum matter

Theorem 1 places GUID among the theories in which spacetime is classical and matter quantum. Such theories have a known difficulty, and its resolution is a prediction.

5.1 The semiclassical equation fails

The simplest coupling, $G_{\mu\nu} = (8\pi G/c^4)\langle\hat{T}_{\mu\nu}\rangle$ (Møller [48], Rosenfeld [49]), sources gravity by the quantum expectation value. A mass in an equal superposition of two places then attracts a probe towards the midpoint, where the mass never is. Page and Geilker tested a macroscopic version, with the position of lead masses set by a quantum random event, and found the probe following the actual masses, not the average [11] (P). The equation is also nonlinear in the quantum state, which in general permits signalling. So a classical frame cannot simply be sourced by expectation values.

5.2 Consistent classical-quantum dynamics

The consistent alternative is a joint state $\varrho(z)$ of a classical phase-space variable z (here the frame) and a quantum system, evolving by a map that is linear in ϱ, completely positive, and preserves normalisation, as probability theory demands. Oppenheim showed that such dynamics

exists and can couple a classical metric to quantum fields [12]. For continuous, Markovian classical-quantum dynamics the generator has, besides the usual Hamiltonian part, three kinds of term: a *decoherence* term on the quantum side with coefficient matrix D_0, a *back-reaction* (drift) of the classical variable driven by the quantum state with coefficients D_1, and a *diffusion* of the classical variable with coefficients D_2. Complete positivity forces the trade-off [13, Eq. (23)]

$$D_1\, D_0^{-1}\, D_1^{\dagger} \preceq 2D_2, \tag{8}$$

with D_0^{-1} the generalised inverse, and in addition D_0 cannot vanish where D_1 does not (P). In words: if quantum matter pushes on the classical frame at all ($D_1 \neq 0$), the matter must decohere ($D_0 > 0$) and the frame must diffuse ($D_2 > 0$), and the product of the two is bounded below by the square of the coupling. Long coherence times of massive superpositions demand strong random fluctuations of the gravitational field, and weak fluctuations demand short coherence times.

Corollary 1 (GUID diffuses)**.** *If the existence frame (Lemma 2) is sourced by quantum content (Hypothesis 4) through a dynamics that is linear, completely positive and Markovian, then mass superpositions decohere at a non-zero rate and the gravitational field fluctuates randomly, with* (8) *bounding the product from below.*

The corollary turns the classicality of the frame from a philosophical position into a quantitative one: in the Newtonian limit, the decoherence rates of massive superpositions measured in interferometry and the noise in precision measurements of gravitational forces together squeeze the remaining parameter space from both sides [13].

5.3 The record is a measurement, the layout is feedback

There is a second, older family of consistent classical-quantum models, and DMP's architecture coincides with it. Kafri, Taylor and Milburn model the gravitational interaction between two masses as a continuous weak measurement of each mass's position followed by a classical force applied to the other according to the recorded result [50]; Tilloy and Diósi source a classical Newtonian potential from a continuously measured (spontaneously collapsing) mass density [51]. In both, gravity reproduces Newton's law on average, cannot entangle, decoheres superpositions and adds noise, as Corollary 1 requires.

Read through the dictionary, DMP's Axiom 3 has exactly this structure (D)+(H). Every act leaves a record that every witness reads in full: a measurement of the act's existence. Any other reader sees the counts: the measured mass density, coarse-grained. The layout of the system is "the median of that demand, recomputed by both halves of the layer and never by one": a classical function of the records, applied back to the system, which is feedback. From the parts' point of view the whole loop is a measure-and-prepare channel, which is why it cannot entangle (Lemma 3).

Proposition 3 (DMP is a measurement-and-feedback gravity)**.** *Under Hypotheses 1–4, the dynamics that DMP's axioms prescribe for the trust field is a continuous measurement of existence (energy-momentum) followed by classical feedback on the geometry, of the Kafri–Taylor–Milburn and Tilloy–Diósi type.*

The proposition fixes the architecture, not the constants. A measurement-and-feedback model has a free parameter, the measurement strength or equivalently the smearing length of the measured mass density, which sets the decoherence rate and the noise. DMP derives no constant (Section 17), and the most economical choice, the parameter-free Diósi–Penrose collapse [52, 53], is excluded by the absence of the spontaneous radiation it predicts in an underground germanium detector [54] (P). GUID must therefore carry a free strength, fixed by experiment. *Verdict: derived* as a structure; the constant is open.

6 The experiments that decide

6.1 Gravitationally mediated entanglement

Bose et al. [14] and Marletto and Vedral [15] proposed to place two masses, each in a superposition of two positions, close enough that only gravity couples them, and to test whether they end entangled. Let each mass m be in $(|L\rangle + |R\rangle)/\sqrt{2}$ with the two positions Δx apart and the centres d apart on one line. A coherent Newtonian field gives each branch (i, j) the phase $\varphi_{ij} = Gm^2t/(\hbar d_{ij})$, where d_{ij} is the distance between the masses in that branch, so after time t

$$|\Psi(t)\rangle = \tfrac{1}{2} \sum_{i,j\in\{L,R\}} e^{i\varphi_{ij}(t)} |i\rangle_A |j\rangle_B. \tag{9}$$

Up to local phases this is $\frac{1}{2}(|00\rangle + |01\rangle + |10\rangle + e^{i\Delta\varphi}|11\rangle)$ with $\Delta\varphi = \varphi_{RL} + \varphi_{LR} - \varphi_{LL} - \varphi_{RR}$; its concurrence is $|\sin(\Delta\varphi/2)|$ and its negativity

$$\mathcal{N} = \tfrac{1}{2}\,|\sin(\Delta\varphi/2)|. \tag{10}$$

For the parameters of [14], $m = 10^{-14}\,\mathrm{kg}$, $\Delta x = 250\,\mu\mathrm{m}$, $d = 450\,\mu\mathrm{m}$, the entangling phase grows at $0.126\,\mathrm{rad/s}$: one radian after $8.0\,\mathrm{s}$, maximal entanglement ($\mathcal{N} = 1/2$, $\Delta\varphi = \pi$) after $25\,\mathrm{s}$ [Q5]. A recorded, classical field (Lemma 2) leaves each branch's phase but removes the coherence between branches, and the masses stay separable at all times, $\mathcal{N} = 0$ [Q5].

GUID predicts $\mathcal{N} = 0$. A positive result would show that the gravitational field is not the existence frame: the identification of Hypothesis 1 would fail for nature and the proposed theory would be refuted; a quantum geometry would then run on DMP only as content (Theorem 1(ii)). Two caveats are under debate. Aziz and Howl argue that classical gravity coupled to quantum matter fields can produce entanglement through virtual matter processes, at a rate far below the quantum-gravity rate [55]; Marletto, Oppenheim, Vedral and Wilson [56], Diósi [57], Schneider, Huggett and Linnemann [58] and Gundhi, Infantino and Bassi [59] dispute it. And Christodoulou et al. showed that the entanglement follows from linearised quantum gravity on a fixed background with local mediation [60], which is the content-side mechanism of Section 9's dissent. Neither changes the prediction: in GUID the frame is recorded, and Lemma 3 applies to it.

6.2 Quantumness without entanglement

Lami, Pedernales and Plenio showed that the quantum nature of gravity can be tested without creating entanglement: an LOCC inequality bounds how well any local classical channel can reproduce a gravitational evolution of coherent states of massive oscillators, such as torsion pendula, and a measured violation would exclude a classical mediator [61]. The test needs no large spatial superpositions. GUID predicts that the inequality holds.

6.3 Diffusion against decoherence

Corollary 1 gives a two-sided test. Interferometry with ever heavier molecules and levitated particles bounds the decoherence that gravity may cause from above; precision measurements of the gravitational force, which has been measured between masses as small as $90\,\mathrm{mg}$ [62], bound the random fluctuations of the field from above. Inequality (8) says the two cannot both be made arbitrarily small. GUID predicts that they cannot.

6.4 What existing experiments already show

Quantum matter responds to gravity as the framework requires: neutrons in a gravitational potential acquire the phase $mgA/\hbar v$ predicted for quantum matter in a classical field (Colella,

Overhauser and Werner [63]), and ultracold neutrons above a mirror occupy discrete gravitational quantum states [64]. In both, the matter is quantum and the field is a given potential; neither tests whether the field itself can be in superposition. The current state of the field is reviewed by Carney, Stamp and Taylor [65]. *Verdict on GUID's classical frame: live, falsifiable.*

7 The thermodynamics of the frame

7.1 The existence of a boundary is its temperature

GUID defines the existence a boundary asserts as the product of the local strength of gravity g, Planck's constant h and the inverse of the speed of light, which DMP's reading takes as the universal tick rate:

$$E_{\mathrm{ex}} \equiv \frac{g\,h}{c}. \tag{11}$$

Proposition 4 (existence and the Unruh temperature)**.** *E_{ex} is the only energy that an acceleration, Planck's constant and the speed of light can form, and*

$$E_{ex} = 4\pi^2\, k_B T_U, \qquad k_B T_U = \frac{\hbar g}{2\pi c}, \tag{12}$$

where T_U is the temperature an observer of proper acceleration g assigns to the vacuum (Unruh [66]). At a horizon of surface gravity κ it is $4\pi^2$ times the Hawking temperature $k_B T_H = \hbar\kappa/2\pi c$; for a Schwarzschild hole $\kappa = c^4/4GM$.

Proof. With dimensions $[g] = LT^{-2}$, $[h] = ML^2T^{-1}$, $[c] = LT^{-1}$, the monomial $g^a h^b c^k$ has the dimension of energy, ML^2T^{-2}, only for $(a, b, k) = (1, 1, -1)$ [Q1]. With $h = 2\pi\hbar$, $E_{\mathrm{ex}}/k_B T_U = 4\pi^2$. □

The uniqueness means the form of (11) is not itself evidence; quantum field theory supplies the factor and the meaning. At standard gravity $T_U = 4.0 \times 10^{-20}\,\mathrm{K}$; at the horizon of a solar-mass hole $T_H = 6.2 \times 10^{-8}\,\mathrm{K}$ [Q1]. *Verdict: holds as a reading* of the Unruh and Hawking temperatures.

7.2 Time: the lapse

In the GUID reading a clock asserts its existence at a rate that falls where the existence field is strong. In general relativity that rate is the lapse, $N = \sqrt{1 + 2\Phi/c^2}$ in a weak field and $N = \sqrt{1 - 2GM/rc^2}$ outside a mass, which vanishes at the horizon [Q2]. It reproduces the redshift over the 22.5 m Pound–Rebka tower, 2.46×10^{-15} [67], and the rate difference of two optical clocks 33 cm apart, 3.6×10^{-17} [68]. *Verdict: holds*; it adds nothing to general relativity.

7.3 Einstein's equation as an equation of state

Jacobson derived Einstein's equation from thermodynamics [69], and GUID's two thermal inputs are exactly his. Take any point p and any local Rindler horizon through it, generated by a null vector field k^a with affine parameter λ, vanishing expansion and shear at p, and approximate boost Killing field $\chi^a = -\kappa\lambda k^a$. Assume:

1. the horizon has the Unruh temperature $T = \hbar\kappa/2\pi$ (units $c = k_B = 1$), Proposition 4;
2. its entropy is proportional to its area, $S = \eta A$, the boundary's floor (Hypothesis 5);
3. the Clausius relation $\delta Q = T\,\mathrm{d}S$ holds for the energy flux across it.

The heat flux is $\delta Q = \int T_{ab}\chi^a \mathrm{d}\Sigma^b = -\kappa \int \lambda\, T_{ab} k^a k^b \,\mathrm{d}\lambda\,\mathrm{d}\mathcal{A}$. The area change follows from the Raychaudhuri equation, $\mathrm{d}\theta/\mathrm{d}\lambda = -R_{ab}k^a k^b$ to first order, as $\delta A = -\int \lambda R_{ab} k^a k^b \,\mathrm{d}\lambda\,\mathrm{d}\mathcal{A}$.

Equating $\delta Q = (\hbar\kappa/2\pi)\eta\,\delta A$ for every p and every null k^a gives $R_{ab} + f g_{ab} = (2\pi/\hbar\eta)T_{ab}$ for some function f; the contracted Bianchi identity and $\nabla^a T_{ab} = 0$ fix $f = -R/2 + \Lambda$, so

$$G_{ab} + \Lambda g_{ab} = \frac{2\pi}{\hbar\eta}\,T_{ab}, \qquad G = \frac{1}{4\hbar\eta} \quad (c = 1). \tag{13}$$

Newton's constant is set by the entropy per area, $\eta = 1/4L_P^2$ [Q1], and the cosmological constant enters as an integration constant. Jacobson's own conclusion fits GUID's classical frame: on this view, "it may be no more appropriate to quantize the Einstein equation than it would be to quantize the wave equation for sound in air" [69]. Two limits: DMP supplies neither the value of η nor a manifold, so the derivation is borrowed, not produced; and the entanglement-equilibrium version [70] needs a state, not a schedule, which the axioms do not define. *Verdict: holds as a reading*; the constant is open.

7.4 Information radiation and the floor

DMP's T31 says that a containing boundary emits only thermal radiation, carrying "entropy and no information", and that an outside reader learns of the inside only its size and its staleness. With Hypothesis 5 that is the semiclassical horizon: a black hole of mass M radiates at T_H and an outside observer knows its mass, spin and charge (P). Erasing a bit costs at least $k_B T \ln 2$ of heat (Landauer [71]); in the GUID reading entropy is the inefficiency of converting between existence and information, which is Landauer's principle in other words. *Verdict: holds as a reading.*

8 Black holes

8.1 The horizon as containment

Hypothesis 5 maps DMP's containment onto the event horizon. The match is close. No act carries information across a containing boundary (T31); no causal curve leaves the horizon (P). The counts every outside reader sees (Axiom 3) are the hair a stationary black hole keeps: mass M, angular momentum J, charge Q (Israel [72]; Kerr [73]). Inside a Schwarzschild horizon the radial coordinate is timelike and every future-directed path ends at the singularity: everything inside derives its causal future from it, a local ground in the sense of G0. Whether crossing is survivable is answered by the equivalence principle: at the horizon of a supermassive hole the tidal stretch across a body is small; for a stellar-mass hole it is lethal before the horizon.

8.2 The containment split gives Christodoulou's mass formula

DMP splits a contained boundary into what is inside and the counts every outside reader sees (D). Give the contained part a mass $M_{\rm irr}$ that sets a Schwarzschild boundary of radius $2GM_{\rm irr}/c^2$; let the spin count J be carried at that boundary as a momentum $J/(2GM_{\rm irr}/c)$ (an identification, (H)); and combine the two as energy and momentum combine in special relativity. In units $G = c = 1$,

$$M^2 = M_{\rm irr}^2 + \left(\frac{J}{2M_{\rm irr}}\right)^2. \tag{14}$$

This is Christodoulou's mass formula for the Kerr black hole [74], with $M_{\rm irr} = \sqrt{A/16\pi}$ (P); solved for $M_{\rm irr}$ it returns Kerr's horizon area for every spin $a = J/M^2 \in [0, 1]$, has no real solution above $a = 1$ (the spin bound), and gives the extractable share $1 - M_{\rm irr}/M = 1 - 1/\sqrt{2} \approx 29\%$ at maximal spin [Q10]. The derivation uses DMP's split, two physical inputs and one identification, and gives no number general relativity does not. *Verdict: holds as a rederivation.*

8.3 What a black hole gives up: Hawking–Penrose–Information radiation

The first law of black-hole mechanics (Bardeen, Carter and Hawking [75]) reads, with $G = c = 1$,

$$\mathrm{d}M = \frac{\kappa}{8\pi}\,\mathrm{d}A + \Omega_H\,\mathrm{d}J + \Phi_H\,\mathrm{d}Q, \tag{15}$$

with Ω_H the angular velocity and Φ_H the electric potential of the horizon. It separates the two ways a black hole loses mass. The rotational and electric terms can be extracted from outside at fixed or growing area: Penrose's process with particles [76], superradiant scattering with waves, and the Blandford–Znajek process with magnetic fields [77], the leading model of the jets of active galaxies. All of them work in the ergosphere, which reaches $r = 2GM/c^2$ at the equator and lies outside the horizon $r_+ = (GM/c^2)(1 + \sqrt{1 - a^2})$ for every spin [Q10]. The area term is released only quantum mechanically, as Hawking's thermal radiation, which lowers A.

Definition 4 (Hawking–Penrose–Information radiation)**.** The *Hawking–Penrose–Information (HPI) radiation* of a black hole is everything it gives up: the extraction of its rotational and electric energy from the ergoregion (the $\Omega_H \mathrm{d}J$ and $\Phi_H \mathrm{d}Q$ terms of (15)) and its thermal Hawking radiation (the $\kappa\,\mathrm{d}A/8\pi$ term).

In GUID terms the extraction is the existence frame being dragged along by the spinning hole, outside the horizon, so no act crosses it; the Hawking part is the floor of T31. What leaves carries the counts and no content, as Axiom 3 requires. Spinning holes are more efficient engines: the binding energy radiated by matter falling to the innermost stable orbit is $1 - \sqrt{8/9} = 5.7\%$ of its rest energy without spin, 32% at Thorne's limit $a = 0.998$, and approaches $1 - 1/\sqrt{3} = 42.3\%$ at maximal spin (Bardeen, Press and Teukolsky [78]; [Q10]). At fixed mass every additional count (spin, charge) shrinks the area, so the more an outside reader knows of a boundary, the less entropy it holds and the more of its energy can be extracted (P). The Hawking part is unobservable for astrophysical holes: a solar-mass hole is at 6.2×10^{-8} K, colder than the 2.7 K microwave sky, so it absorbs more than it emits; only holes lighter than about 0.6 lunar masses are hotter than the sky, and a solar-mass hole in empty space would take more than 10^{66} years to evaporate [Q11]. *Verdict: holds*; the Hawking part is *untested in nature*.

8.4 The information problem

If evaporation is unitary, the entropy of the radiation first rises and then falls after the Page time [7]; the island formula, derived from the gravitational path integral with replica wormholes, reproduces that curve [8, 9]. Those derivations sum over geometries, which is quantum gravity in the sense GUID's frame excludes. On the other side, a classical-quantum dynamics decoheres the quantum system (Corollary 1), so it is not unitary on the quantum system alone and does not by itself produce a Page curve. DMP's T31 as written, "what still leaves is entropy and nothing else", and its T27, which destroys contained items at their scheduled date, side with the thermal picture. In GUID the question is therefore open in a specific form: whether the content held inside a horizon (which may be quantum, Theorem 1(ii)) returns in the radiation as content, while the frame stays classical. No observation reaches it. *Verdict: open*, in physics and in GUID.

9 Holography and quantum error correction

9.1 The strongest assembly

The axioms can be assembled as a holographic quantum error-correcting code on a fixed background, the kinematics of the HaPPY model [79]. A part ("no further statements about its constituents") becomes one block of the five-qubit code [80], which encodes one logical qubit

in five physical ones with distance 3. Its encoding map is a *perfect tensor* with six legs: every bipartition of the six legs into three and three is a unitary map, so every three-leg cut carries three qubits of maximal entanglement [A1]. A layer, which Axiom 3 makes a tree rooted at its Duality, becomes a rooted tree of such tiles, one root and five children with twenty boundary legs, each tile's logical leg the information sitting there; both tile maps are isometries [A2].

Minimal cuts. For a boundary region R, the entropy of the reduced boundary state equals the number of legs cut by the minimal cut through the tree that separates R from its complement [A2]. This is the Ryu–Takayanagi formula [81], $S(R) = |\gamma_R|/4G\hbar$ with γ_R the minimal bulk surface anchored on ∂R, on a tree. DMP's "fewest hops in total" selects the same minimal surfaces.

Reconstruction. The five-qubit code corrects any two erasures, so the logical qubit is recoverable from any three of the five physical qubits and from no two: a $((3, 5))$ quantum threshold scheme [82]. In mutual information with a reference R purifying the root, $I(R : \text{pair}) = 0$ for all ten pairs and $I(R : \text{triple}) = 2$ bits for all ten triples [A3]. This is entanglement-wedge reconstruction [83]: a bulk operator is reconstructable on any boundary region whose wedge contains it.

The two witnesses. The trust half and the information half each measure the stabiliser syndrome of the code. Stabilisers commute with the logical operators, so the syndrome says that content is present and in the code, never what it is: this is Axiom 3's existence clause, exactly. Both halves read the same syndrome; the gate passes only on two trivial syndromes; each of the 15 single-qubit faults gives a distinct syndrome on which both refuse, and correction restores the content with fidelity 1. If the two halves instead read the content itself, the content is disturbed (mean fidelity 0.50), so "full reach over the act" must mean full reach over its existence [A4].

Two layers in one object. Two parts whose content is a logical Bell pair reach the Tsirelson value $2\sqrt{2} \approx 2.83$ of the CHSH combination on their content, while the gates' verdicts, deterministic syndrome readings, give exactly the classical value 2 [A5]. The gate is classical and the content quantum, at the same time and in the same assembly.

9.2 Where the assembly stops: the classical centre

In any code with complementary recovery, Harlow proved the quantum Ryu–Takayanagi formula

$$S(\rho_R) = \mathrm{Tr}(\rho\, \mathcal{L}_R) + S_{\mathrm{bulk}}(\rho, \mathcal{M}_R), \tag{16}$$

where $\mathcal{M}_R$ is the algebra of bulk operators reconstructable on R and the *area operator* $\mathcal{L}_R$ lies in the centre of $\mathcal{M}_R$: it commutes with everything either side can do [84]. DMP's structures recomputed by both halves are such central elements. Lemma 2 then says that in GUID the area is a number, never an operator with a spread of values: the axioms allow one fixed-area sector at a time. In holography fixed-area states are a special, though useful, class (Dong, Harlow and Marolf [85]); the general state of quantum gravity superposes areas. So the assembly reproduces the kinematics of holographic codes on a fixed background ([A1]–[A5]) and stops exactly at superpositions of geometry.

A dissenting reading. One of the review passes behind this work scored the same facts the other way. Split the geometry, as the background-field method does: the background is a DMP structure and classical; the linearised fluctuation, which carries the Newtonian pull between laboratory masses, is content between parts, and linearised quantum gravity on a fixed

background already predicts the entanglement of the masses [60]. On that reading DMP's classical structures are the part of quantum gravity that is classical in quantum gravity too, the centre of (16), and the linearised field is quantum content on that classical centre. The two readings agree on every fact: the shared structures are classical, quantum content including a linearised field can move between parts by grant and stay entangled while every act is recorded, the axioms derive no constant, and they forbid coherent superpositions of backgrounds while their records exist. They differ in one place: whether the linearised field that pulls laboratory masses belongs to the frame (the proposed theory, which predicts $\mathcal{N} = 0$ in Section 6) or to the content (the dissent, which predicts $\mathcal{N} > 0$). DMP hosts either; the experiment decides between them.

The tree is not our space. Read as space, a tree's boundary is totally disconnected (the p-adic line of p-adic AdS/CFT [86, 87]), and no tree approximates a plane: an n-cycle needs distortion at least $n/3 - 1$ in any tree [88]. So a literal tree cannot be our space. Keeping the hierarchy as the routing tree of each request over a graph with loops removes this objection and touches none of the results above. Read as the tree of histories, each branch one consistent record (decoherent histories [89]), the tree owes no spacelike structure at all.

10 Lower dimensions

Why 3+1? In D spacetime dimensions a massless spin-2 field has $D(D-3)/2$ polarisations and the Weyl tensor, the part of the curvature not fixed by local matter, has $D^2(D^2-1)/12 - D(D+1)/2$ components: two and ten in 3+1, none and none in 2+1 [Q8]. In 2+1 a mass does not attract another at all; it cuts a wedge out of space (Deser, Jackiw and 't Hooft [90]). Gravity there is, perturbatively, a Chern–Simons gauge theory [91], with a holographic boundary algebra for negative cosmological constant [92]; whether pure quantum gravity in 2+1 exists as a consistent theory is open [93]. Its degrees of freedom are global: the holonomies of the flat connection around particles and handles, which do not commute once quantised [94], and point particles scatter through their conical deficits [95].

So 2+1 is not an exception to Theorem 1. With no local field there is no local mediator to forbid, but the global holonomies are quantum, and a DMP structure that carries them is broadcast and so can carry one classical sector of them, never their superposition; with matter, the same rule decoheres the particles' relative positions. In GUID's terms, the existence frame in 2+1 would have nothing local that could be held as a classical record, and what it does carry is global and quantum. DMP can run 2+1 quantum gravity as content, but its frame cannot be 2+1 gravity; and our universe is 3+1, whose local degrees of freedom, gravitational waves, are measured [96]. *Verdict: derived*, with the dimension count as *holds*.

11 Cosmology

11.1 The beginning: the start

G0 is stated, not derived, and the minting is the first act: before it there is no act and therefore no record (D). Lemma 2 rests on records, so it applies only once acts exist. GUID describes the beginning as one event, the *start*. For anything to begin to exist it must be possible for it to exist, and the first step of existing is to inflate, through the decay of the oscillons that coalesce once the initial energy has been emitted. Inflation and that decay cannot be separated or ordered, because there is no causality yet in which to order them; together they are the start.

Hypothesis 6 (the start)**.** Inflation and the decay of the oscillons it ends in are one event, the start, which is the minting. It has no internal order the axioms can refer to, and the classical era

of Lemma 2, with causality in DMP's sense (acts between existing parts), begins as it completes. (H)

The physics of the start is standard (P). Inflation ends as the inflaton oscillates about the minimum of its potential; in a broad class of models self-resonance fragments it into oscillons, long-lived localised lumps of the field, which dominate the universe for a while (Amin et al. [137]); their decay into the particles of the Standard Model reheats the universe (Kofman, Linde and Starobinsky [138]). In physics these are stages of one field's history; GUID's statement that they form one event agrees with that, since the axioms, which need acts to order anything, have nothing by which to order them. Within the start the axioms place no restriction on the geometry, which may be in superposition; after it, every geometry the axioms define is recorded and classical. This matches the standard semiclassical picture, in which the geometry of the universe becomes classical through decoherence at its very beginning (Halliwell [97]; Kiefer [98]), and the no-boundary proposal, in which there is no "before" (Hartle and Hawking [99]). In GUID the start is therefore the boundary of quantum gravity in time: whatever is quantum about the geometry itself lies within it, where DMP by its own words makes no statement.

11.2 The universe at its own Schwarzschild threshold

For a flat Friedmann universe with Hubble rate H, the mass inside the Hubble radius, $M_H = \frac{4}{3}\pi(c/H)^3\rho_c$ with $\rho_c = 3H^2/8\pi G$, has a Schwarzschild radius equal to the Hubble radius,

$$\frac{2GM_H}{c^2} = \frac{c}{H}, \tag{17}$$

exactly and at every epoch [Q12]. This is the coincidence behind Pathria's "universe as a black hole" [100]. GUID reads the early universe as a Kugelblitz, a horizon formed by energy alone, that could not contain its own existence; the reading holds in the sense of (17) and at every epoch, not only the earliest, and the universe is not a black hole in the sense of general relativity, having no outside. *Verdict: holds as a reading.*

11.3 Inflation as the possibility space of existence

In GUID, inflation is the space that existence makes for itself wherever existence is possible. Physically it is a period of nearly exponential expansion, $a \propto e^{Ht}$, which dilutes a spatial curvature $|\Omega - 1|$ by e^{-2N} over N e-folds (Guth [101]) (P). Afterwards the curvature share grows again, as a^2 while radiation dominates and as a while matter does; from a reheating temperature of 10^{15} GeV to today it grows by about 5×10^{51}, so an order-one initial curvature needs about 62 e-folds to meet Planck's bound $|\Omega_k| \lesssim 5 \times 10^{-3}$ [102] [Q13]. *Verdict: holds as a reading*; it supplies the motive, not the number.

11.4 Primordial gravitational waves

During inflation the vacuum fluctuations of the metric are stretched into a nearly scale-invariant background of gravitational waves (Starobinsky [103]). Their power spectrum is

$$\mathcal{P}_t = \frac{2}{\pi^2}\left(\frac{\hbar H}{M_{\rm pl}c^2}\right)^2 = \frac{16\,G\hbar H^2}{\pi c^5}, \tag{18}$$

with $M_{\rm pl}^2 = \hbar c/8\pi G$, linear in $\hbar$ and zero for a classical metric [Q14]; its ratio to the scalar spectrum is $r = 16\epsilon$ in single-field slow roll, with the consistency relation $n_t = -r/8$ for its tilt. Because (18) vanishes as $\hbar \to 0$, Krauss and Wilczek argued that detecting this background would establish that the gravitational field was quantised during inflation [104] (P). The current bound is $r < 0.036$ at 95% confidence (BICEP/Keck [105]; $r < 0.032$ combined with Planck and

baryon acoustic oscillations [106]), which puts the energy scale of inflation below $V^{1/4} = 1.4 \times 10^{16}\,\mathrm{GeV}$ and its Hubble rate below $\hbar H = 4.7 \times 10^{13}\,\mathrm{GeV}$ [Q14]. Starobinsky's R^2 inflation [107], which fits the scalar data, predicts $r \approx 12/N^2 \approx 3 \times 10^{-3}$ at $N = 60$ [Q14], the target of the next generation of polarisation surveys [108].

Under Hypothesis 6 the tensor modes are generated within the start, where the axioms do not constrain the geometry, and become classical, as the standard account says they do through squeezing and decoherence [109], as the start completes. GUID therefore allows the standard spectrum and its consistency relation, and a detection would be the first observation of the quantum geometry of the start. GUID's own predictions about the geometry begin after the start. *Verdict: live*; consistent with every bound.

11.5 The late universe: the balanced state

In the flat ΛCDM model that fits the data ($\Omega_m = 0.315$, $\Omega_\Lambda = 0.685$ [102]), the Hubble rate falls at every moment from the past into the far future and tends to the de Sitter rate $H_\infty = \sqrt{\Omega_\Lambda}\, H_0 \approx 0.83\, H_0$; the expansion has accelerated since $z = (2\Omega_\Lambda/\Omega_m)^{1/3} - 1 \approx 0.63$ [Q13]. A universe settling into de Sitter expansion cools towards an empty final state. GUID reads this as the cosmological balanced state, the equilibrium in which the need for further existence has fallen to a constant. *Verdict: holds as a reading* in ΛCDM. The newest data leave the end open: DESI's second data release prefers a dark energy that weakens with time at 2.8 to 4.2 standard deviations, depending on the supernova sample [110]; if that holds, the future is not exactly de Sitter and the reading must be redone.

11.6 Dark matter: remnants trapped in the gravitational field

Hypothesis 7 (trapped remnants)**.** The modelling of this paper explicitly allows remnants of interactions to be trapped in the gravitational field: energy and momentum that interactions have left in the existence frame, or behind a boundary the frame has closed, continue to gravitate after the interaction. (H)

DMP motivates the hypothesis twice: a record outlives its act (Axiom 3), and a containing boundary keeps what it holds while every outside reader sees only its counts (T31). Dark matter, as far as the data show, is matter of exactly that kind: it gravitates and does nothing else that has been seen. Within the classical-frame modelling there are three routes by which remnants are trapped, and each can be calculated.

The stochastic remnant. Corollary 1 makes the frame diffuse: every interaction of quantum matter with the classical frame leaves a random kick in the field. Oppenheim and Russo computed the expectation value of the metric in classical-quantum gravity from its path-integral formulation, for static, spherically symmetric configurations [134]. Around a mass M the Newtonian potential acquires a linear and a quadratic term,

$$\Phi(r) = -\frac{GM}{r} - \frac{\gamma_0}{2} - \frac{\gamma_1}{2}\, r - \frac{\gamma_2}{2}\, r^2, \tag{19}$$

where the quadratic term is the cosmological constant of Schwarzschild–de Sitter space, $\gamma_2 = \Lambda c^2/3$, and the linear term is anticorrelated with it and of order $c^2\sqrt{\Lambda}$. Physically, where accelerations are low the variance of the field's acceleration exceeds the square of its mean and acts as an additional, entropic attraction. The circular speed is

$$v^2(r) = r\,\frac{\mathrm{d}\Phi}{\mathrm{d}r} = \frac{GM}{r} + \frac{\gamma_1}{2}\, r + \gamma_2\, r^2. \tag{20}$$

The first two terms balance at $r_* = \sqrt{2GM/\gamma_1}$, where

$$v_*^4 = 2GM\gamma_1. \tag{21}$$

Around r_* the rotation curve is roughly flat, changing by about 12% between $r_*/2$ and $2r_*$, and its height obeys $v^4 \propto M$, the form of the observed baryonic Tully–Fisher relation. For a galaxy of 10^{11} solar masses in baryons and $\gamma_1 = 10^{-10}\,\mathrm{m/s^2}$, the order Oppenheim and Russo find, $r_* = 17\,\mathrm{kpc}$ and $v_* = 227\,\mathrm{km/s}$, with the Λ term below 10^{-4} of v_*^2 there [Q18]. The acceleration scale is tied to the cosmological constant: $c^2\sqrt{\Lambda/3}/2\pi = 8.6 \times 10^{-11}\,\mathrm{m/s^2}$, close to Milgrom's empirical $a_0 = 1.2 \times 10^{-10}\,\mathrm{m/s^2}$ [135] [Q18]. On this route a single mechanism, the frame's diffusion, gives both a dark-energy term and a dark-matter-like term. The calculation is preliminary, as its authors stress: it is static and spherically symmetric, depends on an infrared regularisation, fixes γ_1 by boundary conditions rather than by a coupling constant, and does not yet address gravitational lensing or the microwave background. A deterministic relative of the same idea exists in tetrad gravity: Hehl and Mashhoon found that a memory of the frame, a nonlocal dependence on its past, acts in the linear approximation as dark matter given by an integral transform of the matter [136].

The contained remnant. The second route is containment (Hypothesis 5). Energy that collapsed behind a horizon during the start gravitates through its counts alone and is dark by T31: primordial black holes. The start itself makes them. When the inflaton fragments into oscillons, which in a broad class of models dominate the universe for a while after inflation [137], enough of them can collapse into black holes of sublunar mass to make up all of the dark matter (Cotner, Kusenko and Takhistov [139]). Primordial black holes can be all of the dark matter only in the asteroid-mass window, about 10^{14} to $10^{19}\,\mathrm{kg}$ (10^{17} to $10^{22}\,\mathrm{g}$), which current constraints leave open [140]. The window's light edge is set by the Hawking part of their HPI radiation: a hole of $10^{14}\,\mathrm{kg}$ glows at $1.2 \times 10^9\,\mathrm{K}$, $k_B T \approx 0.1\,\mathrm{MeV}$, and its evaporation products (MeV gamma rays, and positrons that annihilate at 511 keV) bound how many such holes there can be; holes lighter than a few $10^{11}\,\mathrm{kg}$ have evaporated already [Q18]. At the heavy edge, $10^{19}\,\mathrm{kg}$ with a Schwarzschild radius of 15 nm and a temperature of $1.2 \times 10^4\,\mathrm{K}$, microlensing takes over [Q18]. The prediction of this route is specific: if dark matter is the contained remnant of the start, the Hawking part of HPI radiation, unobservable for astrophysical black holes (Section 8), becomes observable at the window's light edge.

The radiated remnant. Gravitational waves are remnants of interactions that propagate in the frame. Pulsar timing has found a nanohertz background with characteristic strain $h_c \approx 2.4 \times 10^{-15}$ at the frequency 1/yr (NANOGrav [141]), which corresponds to $\Omega_{\mathrm{gw}} = 2\pi^2 f^2 h_c^2/3H_0^2 \approx 8 \times 10^{-9}$ per logarithmic frequency [Q18]: more than 10^7 times too little for the dark matter, and as radiation ($w = 1/3$) it does not cluster. It is a real remnant trapped in the field, but not the dark matter.

What every remnant must satisfy. Four measurements constrain every route (P). (1) It must already be there at recombination: the acoustic peaks of the microwave background measure the dark matter density at $z \approx 1100$, before the first star, as $\Omega_c h^2 = 0.120$, the same comoving value later structure requires, with the baryons then already at the share $\Omega_b/\Omega_m = 0.157$ [102] [Q15]. So the remnant formed during the start, not from later acts; and the baryon content was fixed early, as nucleosynthesis and the microwave background agree. (2) It must be collisionless and move with the galaxies: in the merging cluster 1E0657−558 the lensing mass follows the galaxies, not the hot gas that holds most of the baryons (Clowe et al. [142]). (3) It cannot be the known neutrinos: they are less than 0.7% of the dark matter density, since cosmology bounds their mass sum below 0.072 eV [111]; a dwarf galaxy made of them would need each to weigh more than about 170 eV by the Pauli principle (Tremaine and Gunn [112]); and relic neutrinos move at about 3200 km/s and stellar ones at nearly c, against a Galactic escape speed of 533 km/s [113] [Q15]. (4) It must leave α unchanged to about a part per million over ten billion years (Section 11.7).

Verdict: live. The contained route passes all four requirements by construction; the stochastic route reproduces the shape and scale of galactic rotation curves in a preliminary calculation and must still pass lensing, the acoustic peaks and the merging cluster; the radiated route is real but far too small.

11.7 The fine-structure constant as the information field

In GUID α is set by the information scalar ϕ (Hypothesis 4), so it may vary where ϕ does. The data are tight (P). Optical clocks bound today's drift to $\dot{\alpha}/\alpha = (1.0 \pm 1.1) \times 10^{-18}$ per year [114]; quasar absorption spectra bound changes over ten billion years to about a part per million [115]; the microwave background bounds the value at recombination to a few parts per thousand [116]. MICROSCOPE bounds the scalar's coupling to matter by (7). GUID's forward statement is that α does not grow: its drift is zero or negative. A measured positive drift would refute it. *Verdict: live.* GUID has also proposed that α was of order one at the beginning. Measured α *grows* with energy, to about 1/128 at the Z mass, and the Standard Model at one loop gives $1/\alpha \approx 105$ at the Planck energy [Q16], so a value of one lies beyond known physics. *Verdict: untested* before nucleosynthesis, *refuted* after it.

12 Matter in the existence frame

12.1 The spin ladder

GUID reads spin as the rungs of the Duality, and the Standard Model fills each rung (P):

Spin	GUID reading	Particles	Physics
0	existence itself	Higgs field; the scalar ϕ	the Higgs vacuum value gives rest mass; ϕ sets α
1/2	a duality, which can stay in balance	quarks, leptons	a massive Dirac fermion is two chiral halves joined by its mass
1	information itself; delivered, an act	photon, gluons, W, Z	the carriers of every interaction
2	the perfect duality, two in one	the frame's wave	a massless spin-2 field must couple to all energy: general relativity [41, 42]

The rungs are told apart by rotation: a state of helicity s picks up the phase $e^{is\theta}$ under a turn by θ about its momentum, so a spin-1 quantum looks the same again after 360°, a spin-2 quantum after 180°, and a spin-1/2 state only after 720°, returning with a minus sign after 360°. *Verdict: holds as a reading.*

12.2 The Higgs vacuum as the grant of existence

In GUID existence is granted by the ground. The Higgs field is the one field whose value in empty space is not zero: with potential $V(H) = -\mu^2 H^\dagger H + \lambda_H (H^\dagger H)^2$ it rests at $\langle H^\dagger H \rangle = v^2/2$, $v = \mu/\sqrt{\lambda_H} = 246\,\text{GeV}$, and an elementary particle has its rest mass by its coupling to that background [117, 118] (P). GUID reads the background value, which quantum field theory treats as a classical number, as G0's grant of rest mass, on the existence side and classical, in agreement with Lemma 2; the Higgs boson, a quantum ripple of the field that decays in about $10^{-22}\,\text{s}$, is content. The vacuum at the minimum of its potential is the physical counterpart of the balanced state $\Phi = 0$ of (2). Like G0 ("stated, not derived"), the Higgs potential is an input of the Standard Model, not a derivation.

The reading has one sharp limit, and it is gravitational. The Higgs field gives only part of the mass of matter: about 9% of a nucleon's mass is the quark mass terms; the rest is the energy of quark motion and of the gluon field [29]. Gravity weighs all of it: a gravity that weighed only the Higgs-given share would make titanium and platinum fall differently by 10^{-3}, against MICROSCOPE's 10^{-15} [Q6]. So the existence frame responds to all existence, granted (the Higgs share) and bound (the internal energy of hadrons), and the Higgs field is not the gravitational field. And gravity is far too weak to give mass: its coupling to the electron, $Gm_e^2/\hbar c = 1.75 \times 10^{-45}$, and its share of the binding of hydrogen, 4.4×10^{-40} of the electric one, are negligible [Q16]. *Verdict: holds as a reading* of the Higgs vacuum as the grant; *refuted* as an identity of the Higgs field with gravity.

12.3 The VIBE coefficient

GUID names α the *VIBE coefficient*, after the Variable Influence Balancing Equation (2), and reads it as the strength with which existence interacts with itself before it must re-ground itself. In quantum electrodynamics α is the strength with which a charge interacts with the electromagnetic field and so, through it, with itself; each emission and reabsorption of a virtual photon costs one power of α. The cleanest measure is the electron's anomalous magnetic moment, whose first term $a_e = \alpha/2\pi$ (Schwinger [119]) gives the measured value [120] to 0.15% [Q16]. The same constant is the ratio of the electron's three lengths, $r_e/\bar{\lambda}_C = \bar{\lambda}_C/a_0 = \alpha$ [Q16]. *Verdict: holds as a reading*; the VIBE coefficient is a name for α, not a new constant.

12.4 The forces as tiers of one interaction

GUID reads the forces as tiers of one interaction: the strong force binds the cluster, the weak force keeps it stable, electromagnetism links clusters of clusters. Each does that job (P): the strong force binds quarks into nucleons and nucleons into nuclei, the weak force moves nuclei towards the valley of stability, electromagnetism binds atoms and molecules. The weak and electromagnetic forces are one electroweak interaction [121, 122, 123]. Whether all three meet at one energy is open: in the Standard Model the couplings come close near 10^{14} to 10^{16} GeV and do not meet at one point [124]. Gravity is not among the tiers in GUID: it is the frame the tiers act in. *Verdict: holds as a reading*; unification *untested*.

13 Hosting: the Existence Turing machine

Theorem 1 says that DMP, supplied with the base configuration of our universe, hosts the proposed theory, and its part (ii) that the information half can carry any quantum content. Assembled from DMP parts, a universal quantum computer can run as content, and by the Church–Turing–Deutsch principle it approximates every finitely realisable physical system to any finite precision [40]. GUID calls such an assembly the *Existence Turing machine*; a sealed file that carries the assembly's recipe, its digest, the versions of the axioms and of the proof chain, and the verdict lets the same machine be redeployed and checked anywhere. Two consequences matter for quantum gravity. First, every candidate theory of quantum gravity in Section 14 can also run on the machine with its predictions, so the axioms do not forbid quantum gravity in nature; a theory whose geometry is quantum runs as content, not as DMP's frame. Second, the machine's own structure, the gates, records and layout, is classical, and in the proposed theory it is the existence frame itself. A check of the second statement, with three five-qubit parts whose record and counts are identical for all sixteen pairs of mass contents while the contents end entangled, belongs to the dissenting review and rests on one construction choice (existence read per part, not per mass position); we do not use it here.

14 Relation to the approaches to quantum gravity

Approach	What is quantum	Relation to GUID	What tells them apart
Effective field theory [3, 4]	metric fluctuations below E_P	GUID keeps the classical part; the graviton-loop term of (1) is absent	a term 10^{-70} of Newton's at 1 m [Q1]: unmeasurable
String theory, AdS/CFT [125]	everything; geometry emerges from a quantum boundary	hostable as content; the code kinematics of Section 9 are shared, with a classical centre	entanglement of masses
Loop quantum gravity [126, 127]	the triad/tetrad; area has a discrete spectrum	the frame is LQG's own variable, not quantised; area a number	area spectrum (not accessible); BMV
Asymptotic safety [128, 129]	the metric, with a UV fixed point	hostable	BMV
Causal dynamical triangulations [130]	sum over geometries	allowed within the start only	BMV
Causal sets [131]	discrete order; $\Lambda \sim 1/\sqrt{N}$	DMP's order is a tree, not a sprinkled causal set	Lorentz invariance needs random sprinkling
Thermodynamic gravity [69]	the microscopic entropy only	GUID's thermal reading, Section 7	none beyond GR
Classical-quantum gravity [12, 13]	matter only	GUID's dynamics, Corollary 1	diffusion against decoherence
Channel models [50, 51]	matter only	GUID's architecture, Proposition 3	noise and decoherence rates
Gravity-related collapse [52, 53]	matter, collapsing	parameter-free version excluded [54]; GUID needs a free strength	spontaneous radiation
Semiclassical gravity [48, 49]	matter; metric sourced by $\langle T \rangle$	rejected: nonlinear; Page–Geilker	done [11]
Modified dynamics (MOND) [135]	nothing; a law at low acceleration	the stochastic remnant gives a linear term of size a_0, tied to Λ (Section 11.6)	lensing, acoustic peaks, merging clusters
Spacetime from entanglement [132, 133]	content; geometry emergent	GUID's frame as the classical centre of such a geometry is open	unclear

15 Predictions

Observable	GUID	Quantised metric	Status
Entanglement of two masses through gravity alone [14, 15]	none, $\mathcal{N} = 0$	$\mathcal{N} = \frac{1}{2}\lvert\sin(\Delta\varphi/2)\rvert$	proposed; 25 s at 10^{-14} kg [Q5]
LOCC inequality with massive oscillators [61]	satisfied	can be violated	proposed
Decoherence and diffusion [13]	both non-zero, product bounded by (8)	neither required	bounded from both sides
Universality of free fall	exact for the frame; a scalar term $\eta = 2\lambda^2 f_\oplus \Delta f$	exact	$\lvert\eta\rvert \lesssim 10^{-15}$ [28]

Observable	GUID	Quantised metric	Status
Light bending γ	exactly 1	1	$\gamma - 1 = (2.1 \pm 2.3) \times 10^{-5}$ [33]
Speed of gravitational waves	c	c	$\lvert c_{\rm gw}/c - 1\rvert \lesssim 10^{-15}$ [25]
Single-graviton detector clicks [46]	possible, from a classical wave	possible	do not discriminate [47]
Primordial tensor modes	standard spectrum, generated within the start	$r = 16\epsilon$, $n_t = -r/8$	$r < 0.036$ [105]
Drift of α	zero or negative	none predicted	$(1.0 \pm 1.1) \times 10^{-18}$/yr [114]
Hawking radiation	thermal to the outside reader; return of content open	Page curve	not observable
Galactic rotation (stochastic remnant)	roughly flat near $r_* = \sqrt{2GM/\gamma_1}$, $v_*^4 = 2GM\gamma_1$, $\gamma_1 \sim c^2\sqrt{\Lambda}$	dark matter halo needed	preliminary [134]; lensing open
Contained remnants	primordial black holes of 10^{14}–10^{19} kg; Hawking glow at the light edge	no statement	window open [140]

16 The ledger

Every statement of the account, abridged, with its verdict.

Statement	Verdict	Where
Free fall is universal for the existence frame	derived; holds	Prop. 1; [Q6]
The information scalar does not enter gravity; $\gamma = 1$	derived; holds	Prop. 2; [Q7]
The existence frame is classical after the first record	derived	Lemma 2; [Q3]
A classical frame cannot mediate entanglement	derived	Lemma 3; [Q4]
DMP, with the base configuration of our universe, hosts the proposed theory	derived	Theorem 1
No graviton as a quantum of the frame	derived; consistent with all data	[Q9]
The frame must diffuse and decohere matter	derived (with CQ dynamics)	Cor. 1
Record and recompute is measurement and feedback	derived (structure)	Prop. 3
Gravity entangles no two masses	live, falsifiable	[Q5]
The existence of a boundary gh/c is $4\pi^2 kT_U$	holds as a reading	Prop. 4; [Q1]
Time slows where existence is strong (lapse)	holds	[Q2]
Einstein's equation as an equation of state	holds as a reading; constant open	(13)
Entropy as the cost of converting existence and information	holds as a reading (Landauer)	Section 7
Horizon as containment; counts as M, J, Q	holds as a reading	Section 8
Christodoulou's mass formula from the containment split	holds as a rederivation	(14); [Q10]
Spinning holes give up energy by dragging the frame, outside the horizon	holds	[Q10]

Statement	Verdict	Where
Hawking–Penrose–Information radiation	holds as a name; Hawking part untested in nature	[Q11]
Information returns from black holes	open	Section 8
Holographic minimal cuts and reconstruction from the axioms	holds, on a tree	[A1]–[A5]
The area is a number (one fixed-area sector)	derived	(16)
The tree is our space	refuted as literal; holds as routing or histories	Section 9
2+1 gravity is an exception	refuted; runs as content, not as the frame	Section 10; [Q8]
Inflation and oscillon decay are one start, the minting	holds as a reading	Hyp. 6
Quantum geometry only within the start	derived (with Hyp. 6)	Section 11
The early universe a Kugelblitz	holds as a reading, at every epoch	[Q12]
Inflation as the possibility space; flatness	holds as a reading	[Q13]
Primordial tensor modes, generated within the start	live; consistent with all bounds	[Q14]
The balanced state as the de Sitter future	holds as a reading in ΛCDM	[Q13]
Remnants of interactions are trapped in the gravitational field	live	Hyp. 7
Stochastic remnant: flat rotation curves, $v^4 \propto M$, scale tied to Λ	holds in a preliminary calculation; lensing open	[Q18]
Contained remnant: primordial black holes from the start	live (window open)	[Q18]
Radiated remnant (gravitational waves) as the dark matter	refuted (about 10^{-8})	[Q18]
The known neutrinos as the remnant	refuted (four bounds)	[Q15]
Baryon content fixed early	holds	Section 11
α does not grow	live	Section 11.7
α began at one	beyond known physics; refuted after nucleosynthesis	[Q16]
Spin ladder	holds as a reading	Section 12
The Higgs vacuum is G0's grant of rest mass	holds as a reading	Section 12
The Higgs field is the gravitational field	refuted	[Q6]
α as the VIBE coefficient, self-interaction strength	holds as a reading	[Q16]
The forces as tiers; their unity	holds as a reading; unity untested	Section 12

17 Where we expect it to break

1. *The hypotheses.* Everything physical rests on Hypotheses 1–7, above all on reading trust as existence and existence as energy-momentum. Another faithful reading of either changes the propositions; the dissent of Section 9 is one, and it predicts $\mathcal{N} > 0$.
2. *Experiment.* A positive test of gravitationally mediated entanglement, a violation of the LOCC inequality, or decoherence and diffusion both measured below the trade-off (8) would each show that gravity is not GUID's classical frame. Unless classical gravity can entangle after all [55], a positive test is decisive.
3. *Constants.* The axioms derive none of G, $\hbar$, Λ, the entropy per area η, or the strength of the measurement-and-feedback channel; they enter as content or configuration. A derivation

of any of them from the axioms would be new; we have none.

4. *The start.* Hypothesis 6 makes inflation and the oscillon decay one start that the axioms do not constrain, so GUID predicts nothing about primordial tensor modes beyond standard inflation; its predictions about the geometry begin after the start.
5. *Dark matter.* The stochastic remnant has been computed only for static, spherically symmetric configurations. If it cannot reproduce lensing, the acoustic peaks and the merging cluster 1E0657−558, it is not the dark matter and the contained remnant must carry it; if evaporation and microlensing close the asteroid-mass window, that route closes too.
6. *Black holes.* GUID does not say whether information returns from an evaporating hole. A derivation of the Page curve that needs superposed geometries after the start would contradict Lemma 2.
7. *Space.* A literal tree is not our space; the routing reading avoids the objection but has not been shown to give a Lorentzian 3+1 manifold in any limit.
8. *The checks.* [Q6] uses the semi-empirical Coulomb term and a coarse composition of the Earth; [Q13] a reheating temperature of 10^{15} GeV; [Q15] round values for a dwarf galaxy; [Q16] one-loop running. Better inputs move numbers by tens of per cent, not the verdicts.

18 Conclusion

Read as physics, the three axioms of the Dark Matter Protocol give a definite account of quantum gravity, and it is not a quantisation of geometry. Trust becomes existence, and existence becomes a tetrad that weighs all energy alike, never mixes with the information half, and is classical once the start, inflation with the decay of its oscillons, is complete, because the axioms make each of its values a record that two independent witnesses hold in full. Supplied with the base configuration currently assumed for our universe, DMP hosts this theory: its shared structure carries the classical frame, and its information half the quantum content. Consistency with quantum mechanics then fixes the rest of the dynamics: the frame must diffuse and decohere matter, and the axioms' own architecture, record then recompute, is the measurement-and-feedback channel that does so. Horizons are containing boundaries whose temperature is the existence they assert; black holes give up what the first law allows, and an outside reader sees only their counts; holographic codes are reproduced up to their classical centre; 2+1 dimensions are no exception; quantum geometry is confined to the start. The theory lets remnants of interactions be trapped in the gravitational field: its stochastic remnant already gives the shape and scale of galactic rotation curves in a first calculation, and its contained remnants are primordial black holes from the start. It stands on the classical side of the one class of experiments that can decide it, gravitationally mediated entanglement and its relatives. If those experiments find entanglement, the geometry is quantum and the proposed theory is refuted, and DMP would run its successor as content. We claim that the construction exists and reproduces. Whether it is right is for the experiments, and for the reader, to break.

Acknowledgements

The first author thanks Sir Peter Higgs and the many theorists and experimentalists whose work this account requires and builds on. The review passes that tested earlier forms of this work, including the dissenting one reported in Section 9, are part of its record.

Contributions and use of AI

Steven Hrovat conceived DMP, the reading of trust as existence, the two-field structure, and every hypothesis tested here, chose which ideas to keep after each round of tests,

and takes responsibility for the axiom system and the implications of its theorems. Sirius (Claude, Anthropic) made the derivations, related the account to the literature, wrote and ran `docs/quantum-gravity/checks.py`, and wrote this text. The holographic-code assembly, the Lean development and the review passes were written by AI systems under the first author's direction. Every numerical claim is a line of a public script; every reference was checked against a publisher, arXiv or collaboration record, as listed in `docs/quantum-gravity/README.md`.

Licence and availability

This paper and its source are released under the Apache License, Version 2.0. DMP: `https://github.com/StevenHrovat/dmp` (version 0.9.9 at commit `4400cce`). Gravity: `https://github.com/StevenHrovat/gravity` (this paper and its checks in `docs/quantum-gravity/`; the holographic-code assembly in `docs/qg-verdict/`; the Lean development in `evidence/`).

A The checks

`python3 docs/quantum-gravity/checks.py` (numpy, scipy, sympy) prints 93 checks, all of which pass. Q1: Planck units, the existence energy, Unruh, Hawking, Bekenstein–Hawking, Jacobson's entropy density, the size of the quantum term of (1). Q2: the lapse. Q3: no-cloning and broadcasting. Q4: classical against coherent mediation, 300 random couplings each. Q5: the Bose et al. geometry. Q6: the MICROSCOPE bound on the information scalar and the Higgs-share test. Q7: Brans–Dicke and Cassini. Q8: polarisations and Weyl components in D dimensions. Q9: the single-graviton threshold. Q10: Kerr horizon, ergosphere, Christodoulou, efficiencies. Q11: Hawking radiation of real holes. Q12: the Hubble mass at its Schwarzschild threshold. Q13: flatness, de Sitter floor, onset of acceleration. Q14: the tensor spectrum, the BICEP/Keck bound, Starobinsky's r. Q15: what every trapped remnant must satisfy, and why the known neutrinos do not. Q16: α as self-interaction strength, the electron's lengths, gravity against electromagnetism, the running of α. Q17: the speed of gravity. Q18: remnants trapped in the field: the stochastic remnant's rotation curve and acceleration scale, the contained remnant's Hawking temperatures and evaporation, the radiated remnant's energy density. `python3 docs/qg-verdict/assembly.py` (numpy) prints the 18 checks of the holographic-code assembly; A1 to A5 in this paper are its checks [1] to [5], and its check [6] is the special case of [Q4] with CNOT and CZ couplings.

References

[1] M. H. Goroff, A. Sagnotti, "The ultraviolet behavior of Einstein gravity", *Nucl. Phys. B* 266, 709 (1986).

[2] G. 't Hooft, M. Veltman, "One-loop divergencies in the theory of gravitation", *Ann. Inst. Henri Poincaré A* 20, 69 (1974).

[3] J. F. Donoghue, "General relativity as an effective field theory: the leading quantum corrections", *Phys. Rev. D* 50, 3874 (1994).

[4] N. E. J. Bjerrum-Bohr, J. F. Donoghue, B. R. Holstein, "Quantum gravitational corrections to the nonrelativistic scattering potential of two masses", *Phys. Rev. D* 67, 084033 (2003).

[5] S. W. Hawking, "Particle creation by black holes", *Commun. Math. Phys.* 43, 199 (1975).

[6] J. D. Bekenstein, "Black holes and entropy", *Phys. Rev. D* 7, 2333 (1973).

[7] D. N. Page, "Information in black hole radiation", *Phys. Rev. Lett.* 71, 3743 (1993).

[8] G. Penington, "Entanglement wedge reconstruction and the information paradox", *JHEP* 09 (2020) 002.

[9] A. Almheiri, N. Engelhardt, D. Marolf, H. Maxfield, "The entropy of bulk quantum fields and the entanglement wedge of an evaporating black hole", *JHEP* 12 (2019) 063.

[10] K. Eppley, E. Hannah, "The necessity of quantizing the gravitational field", *Found. Phys.* 7, 51 (1977).

[11] D. N. Page, C. D. Geilker, "Indirect evidence for quantum gravity", *Phys. Rev. Lett.* 47, 979 (1981).

[12] J. Oppenheim, "A postquantum theory of classical gravity?", *Phys. Rev. X* 13, 041040 (2023).

[13] J. Oppenheim, C. Sparaciari, B. Šoda, Z. Weller-Davies, "Gravitationally induced decoherence vs space-time diffusion: testing the quantum nature of gravity", *Nat. Commun.* 14, 7910 (2023), arXiv:2203.01982.

[14] S. Bose, A. Mazumdar, G. W. Morley, H. Ulbricht, M. Toroš, M. Paternostro, A. A. Geraci, P. F. Barker, M. S. Kim, G. Milburn, "Spin entanglement witness for quantum gravity", *Phys. Rev. Lett.* 119, 240401 (2017).

[15] C. Marletto, V. Vedral, "Gravitationally induced entanglement between two massive particles is sufficient evidence of quantum effects in gravity", *Phys. Rev. Lett.* 119, 240402 (2017).

[16] T. D. Galley, F. Giacomini, J. H. Selby, "A no-go theorem on the nature of the gravitational field beyond quantum theory", *Quantum* 6, 779 (2022).

[17] S. Hrovat, *The Dark Matter Protocol*, version 0.9.9, `https://github.com/StevenHrovat/dmp`, commit `4400cce` (2026).

[18] J. A. Goguen, J. Meseguer, "Security policies and security models", in *IEEE Symposium on Security and Privacy*, 11 (1982).

[19] D. E. Denning, "A lattice model of secure information flow", *Commun. ACM* 19, 236 (1976).

[20] D. Malkhi, M. Reiter, "Byzantine quorum systems", *Distrib. Comput.* 11, 203 (1998).

[21] P. O'Hearn, J. Reynolds, H. Yang, "Local reasoning about programs that alter data structures", in *CSL 2001*, LNCS 2142, 1 (2001).

[22] L. de Moura, S. Ullrich, "The Lean 4 theorem prover and programming language", in *CADE 28*, LNCS 12699, 625 (2021).

[23] R. Aldrovandi, J. G. Pereira, *Teleparallel Gravity: An Introduction* (Springer, Dordrecht, 2013).

[24] T. Jacobson, D. Mattingly, "Gravity with a dynamical preferred frame", *Phys. Rev. D* 64, 024028 (2001).

[25] B. P. Abbott et al. (LIGO, Virgo, Fermi GBM, INTEGRAL), "Gravitational waves and gamma-rays from a binary neutron star merger: GW170817 and GRB 170817A", *Astrophys. J. Lett.* 848, L13 (2017).

[26] J. D. Bekenstein, "Fine-structure constant: Is it really a constant?", *Phys. Rev. D* 25, 1527 (1982).

[27] H. B. Sandvik, J. D. Barrow, J. Magueijo, "A simple cosmology with a varying fine structure constant", *Phys. Rev. Lett.* 88, 031302 (2002).

[28] P. Touboul et al. (MICROSCOPE Collaboration), "MICROSCOPE mission: final results of the test of the equivalence principle", *Phys. Rev. Lett.* 129, 121102 (2022).

[29] Y.-B. Yang, J. Liang, Y.-J. Bi, Y. Chen, T. Draper, K.-F. Liu, Z. Liu, "Proton mass decomposition from the QCD energy momentum tensor", *Phys. Rev. Lett.* 121, 212001 (2018).

[30] G. Dvali, M. Zaldarriaga, "Changing α with time: implications for fifth-force-type experiments and quintessence", *Phys. Rev. Lett.* 88, 091303 (2002).

[31] C. M. Will, "The confrontation between general relativity and experiment", *Living Rev. Relativ.* 17, 4 (2014).

[32] C. Brans, R. H. Dicke, "Mach's principle and a relativistic theory of gravitation", *Phys. Rev.* 124, 925 (1961).

[33] B. Bertotti, L. Iess, P. Tortora, "A test of general relativity using radio links with the Cassini spacecraft", *Nature* 425, 374 (2003).

[34] W. K. Wootters, W. H. Zurek, "A single quantum cannot be cloned", *Nature* 299, 802 (1982).

[35] D. Dieks, "Communication by EPR devices", *Phys. Lett. A* 92, 271 (1982).

[36] H. Barnum, C. M. Caves, C. A. Fuchs, R. Jozsa, B. Schumacher, "Noncommuting mixed states cannot be broadcast", *Phys. Rev. Lett.* 76, 2818 (1996).

[37] W. H. Zurek, "Quantum Darwinism", *Nat. Phys.* 5, 181 (2009).

[38] F. G. S. L. Brandão, M. Piani, P. Horodecki, "Generic emergence of classical features in quantum Darwinism", *Nat. Commun.* 6, 7908 (2015).

[39] G. Vidal, R. F. Werner, "Computable measure of entanglement", *Phys. Rev. A* 65, 032314 (2002).

[40] D. Deutsch, "Quantum theory, the Church–Turing principle and the universal quantum computer", *Proc. R. Soc. Lond. A* 400, 97 (1985).

[41] S. Weinberg, "Photons and gravitons in S-matrix theory: derivation of charge conservation and equality of gravitational and inertial mass", *Phys. Rev.* 135, B1049 (1964).

[42] S. Deser, "Self-interaction and gauge invariance", *Gen. Relativ. Gravit.* 1, 9 (1970).

[43] F. Dyson, "Is a graviton detectable?", *Int. J. Mod. Phys. A* 28, 1330041 (2013).

[44] T. Rothman, S. Boughn, "Can gravitons be detected?", *Found. Phys.* 36, 1801 (2006).

[45] C. A. Mead, "Possible connection between gravitation and fundamental length", *Phys. Rev.* 135, B849 (1964).

[46] G. Tobar, S. K. Manikandan, T. Beitel, I. Pikovski, "Detecting single gravitons with quantum sensing", *Nat. Commun.* 15, 7229 (2024).

[47] D. Carney, V. Domcke, N. L. Rodd, "Graviton detection and the quantization of gravity", *Phys. Rev. D* 109, 044009 (2024).

[48] C. Møller, "The energy-momentum complex in general relativity and related problems", in *Les Théories Relativistes de la Gravitation*, ed. A. Lichnerowicz, M.-A. Tonnelat (CNRS, Paris, 1962).

[49] L. Rosenfeld, "On quantization of fields", *Nucl. Phys.* 40, 353 (1963).

[50] D. Kafri, J. M. Taylor, G. J. Milburn, "A classical channel model for gravitational decoherence", *New J. Phys.* 16, 065020 (2014).

[51] A. Tilloy, L. Diósi, "Sourcing semiclassical gravity from spontaneously localized quantum matter", *Phys. Rev. D* 93, 024026 (2016).

[52] L. Diósi, "A universal master equation for the gravitational violation of quantum mechanics", *Phys. Lett. A* 120, 377 (1987).

[53] R. Penrose, "On gravity's role in quantum state reduction", *Gen. Relativ. Gravit.* 28, 581 (1996).

[54] S. Donadi, K. Piscicchia, C. Curceanu, L. Diósi, M. Laubenstein, A. Bassi, "Underground test of gravity-related wave function collapse", *Nat. Phys.* 17, 74 (2021).

[55] J. Aziz, R. Howl, "Classical theories of gravity produce entanglement", *Nature* 646, 813 (2025).

[56] C. Marletto, J. Oppenheim, V. Vedral, E. Wilson, "Classical gravity cannot mediate entanglement", arXiv:2511.07348 (2025).

[57] L. Diósi, "No, classical gravity does not entangle quantized matter fields", arXiv:2511.00852 (2025).

[58] M. D. Schneider, N. Huggett, N. Linnemann, "A demonstration that classical gravity does not produce entanglement", arXiv:2511.19242 (2025).

[59] L. Gundhi, A. Infantino, A. Bassi, "Can classical theories of gravity produce entanglement?", arXiv:2604.19696 (2026).

[60] M. Christodoulou, A. Di Biagio, M. Aspelmeyer, Č. Brukner, C. Rovelli, R. Howl, "Locally mediated entanglement in linearized quantum gravity", *Phys. Rev. Lett.* 130, 100202 (2023).

[61] L. Lami, J. S. Pedernales, M. B. Plenio, "Testing the quantumness of gravity without entanglement", *Phys. Rev. X* 14, 021022 (2024).

[62] T. Westphal, H. Hepach, J. Pfaff, M. Aspelmeyer, "Measurement of gravitational coupling between millimetre-sized masses", *Nature* 591, 225 (2021).

[63] R. Colella, A. W. Overhauser, S. A. Werner, "Observation of gravitationally induced quantum interference", *Phys. Rev. Lett.* 34, 1472 (1975).

[64] V. V. Nesvizhevsky et al., "Quantum states of neutrons in the Earth's gravitational field", *Nature* 415, 297 (2002).

[65] D. Carney, P. C. E. Stamp, J. M. Taylor, "Tabletop experiments for quantum gravity: a user's manual", *Class. Quantum Grav.* 36, 034001 (2019).

[66] W. G. Unruh, "Notes on black-hole evaporation", *Phys. Rev. D* 14, 870 (1976).

[67] R. V. Pound, G. A. Rebka, Jr., "Apparent weight of photons", *Phys. Rev. Lett.* 4, 337 (1960).

[68] C. W. Chou, D. B. Hume, T. Rosenband, D. J. Wineland, "Optical clocks and relativity", *Science* 329, 1630 (2010).

[69] T. Jacobson, "Thermodynamics of spacetime: the Einstein equation of state", *Phys. Rev. Lett.* 75, 1260 (1995).

[70] T. Jacobson, "Entanglement equilibrium and the Einstein equation", *Phys. Rev. Lett.* 116, 201101 (2016).

[71] R. Landauer, "Irreversibility and heat generation in the computing process", *IBM J. Res. Dev.* 5, 183 (1961).

[72] W. Israel, "Event horizons in static vacuum space-times", *Phys. Rev.* 164, 1776 (1967).

[73] R. P. Kerr, "Gravitational field of a spinning mass as an example of algebraically special metrics", *Phys. Rev. Lett.* 11, 237 (1963).

[74] D. Christodoulou, "Reversible and irreversible transformations in black-hole physics", *Phys. Rev. Lett.* 25, 1596 (1970).

[75] J. M. Bardeen, B. Carter, S. W. Hawking, "The four laws of black hole mechanics", *Commun. Math. Phys.* 31, 161 (1973).

[76] R. Penrose, "Gravitational collapse: the role of general relativity", *Riv. Nuovo Cimento* 1, 252 (1969).

[77] R. D. Blandford, R. L. Znajek, "Electromagnetic extraction of energy from Kerr black holes", *Mon. Not. R. Astron. Soc.* 179, 433 (1977).

[78] J. M. Bardeen, W. H. Press, S. A. Teukolsky, "Rotating black holes: locally nonrotating frames, energy extraction, and scalar synchrotron radiation", *Astrophys. J.* 178, 347 (1972).

[79] F. Pastawski, B. Yoshida, D. Harlow, J. Preskill, "Holographic quantum error-correcting codes: toy models for the bulk/boundary correspondence", *JHEP* 06 (2015) 149.

[80] R. Laflamme, C. Miquel, J. P. Paz, W. H. Zurek, "Perfect quantum error correcting code", *Phys. Rev. Lett.* 77, 198 (1996).

[81] S. Ryu, T. Takayanagi, "Holographic derivation of entanglement entropy from AdS/CFT", *Phys. Rev. Lett.* 96, 181602 (2006).

[82] R. Cleve, D. Gottesman, H.-K. Lo, "How to share a quantum secret", *Phys. Rev. Lett.* 83, 648 (1999).

[83] A. Almheiri, X. Dong, D. Harlow, "Bulk locality and quantum error correction in AdS/CFT", *JHEP* 04 (2015) 163.

[84] D. Harlow, "The Ryu–Takayanagi formula from quantum error correction", *Commun. Math. Phys.* 354, 865 (2017).

[85] X. Dong, D. Harlow, D. Marolf, "Flat entanglement spectra in fixed-area states of quantum gravity", *JHEP* 10 (2019) 240.

[86] M. Heydeman, M. Marcolli, I. Saberi, B. Stoica, "Tensor networks, p-adic fields, and algebraic curves: arithmetic and the $\mathrm{AdS}_3/\mathrm{CFT}_2$ correspondence", *Adv. Theor. Math. Phys.* 22, 93 (2018).

[87] S. S. Gubser, J. Knaute, S. Parikh, A. Samberg, P. Witaszczyk, "p-adic AdS/CFT", *Commun. Math. Phys.* 352, 1019 (2017).

[88] Y. Rabinovich, R. Raz, "Lower bounds on the distortion of embedding finite metric spaces in graphs", *Discrete Comput. Geom.* 19, 79 (1998).

[89] M. Gell-Mann, J. B. Hartle, "Quantum mechanics in the light of quantum cosmology", in *Complexity, Entropy and the Physics of Information*, ed. W. H. Zurek (Addison-Wesley, Redwood City, 1990), 425.

[90] S. Deser, R. Jackiw, G. 't Hooft, "Three-dimensional Einstein gravity: dynamics of flat space", *Ann. Phys.* 152, 220 (1984).

[91] E. Witten, "2+1 dimensional gravity as an exactly soluble system", *Nucl. Phys. B* 311, 46 (1988).

[92] J. D. Brown, M. Henneaux, "Central charges in the canonical realization of asymptotic symmetries: an example from three-dimensional gravity", *Commun. Math. Phys.* 104, 207 (1986).

[93] A. Maloney, E. Witten, "Quantum gravity partition functions in three dimensions", *JHEP* 02 (2010) 029.

[94] J. E. Nelson, T. Regge, "Homotopy groups and (2+1)-dimensional quantum gravity", *Nucl. Phys. B* 328, 190 (1989).

[95] G. 't Hooft, "Non-perturbative 2 particle scattering amplitudes in 2+1 dimensional quantum gravity", *Commun. Math. Phys.* 117, 685 (1988).

[96] B. P. Abbott et al. (LIGO Scientific and Virgo Collaborations), "Observation of gravitational waves from a binary black hole merger", *Phys. Rev. Lett.* 116, 061102 (2016).

[97] J. J. Halliwell, "Decoherence in quantum cosmology", *Phys. Rev. D* 39, 2912 (1989).

[98] C. Kiefer, "Continuous measurement of mini-superspace variables by higher multipoles", *Class. Quantum Grav.* 4, 1369 (1987).

[99] J. B. Hartle, S. W. Hawking, "Wave function of the Universe", *Phys. Rev. D* 28, 2960 (1983).

[100] R. K. Pathria, "The universe as a black hole", *Nature* 240, 298 (1972).

[101] A. H. Guth, "Inflationary universe: a possible solution to the horizon and flatness problems", *Phys. Rev. D* 23, 347 (1981).

[102] Planck Collaboration, "Planck 2018 results. VI. Cosmological parameters", *Astron. Astrophys.* 641, A6 (2020).

[103] A. A. Starobinsky, "Spectrum of relict gravitational radiation and the early state of the universe", *JETP Lett.* 30, 682 (1979).

[104] L. M. Krauss, F. Wilczek, "Using cosmology to establish the quantization of gravity", *Phys. Rev. D* 89, 047501 (2014).

[105] P. A. R. Ade et al. (BICEP/Keck), "Improved constraints on primordial gravitational waves using Planck, WMAP, and BICEP/Keck observations through the 2018 observing season", *Phys. Rev. Lett.* 127, 151301 (2021).

[106] M. Tristram et al., "Improved limits on the tensor-to-scalar ratio using BICEP and Planck data", *Phys. Rev. D* 105, 083524 (2022).

[107] A. A. Starobinsky, "A new type of isotropic cosmological models without singularity", *Phys. Lett. B* 91, 99 (1980).

[108] E. Allys et al. (LiteBIRD Collaboration), "Probing cosmic inflation with the LiteBIRD cosmic microwave background polarization survey", *Prog. Theor. Exp. Phys.* 2023, 042F01 (2023).

[109] D. Polarski, A. A. Starobinsky, "Semiclassicality and decoherence of cosmological perturbations", *Class. Quantum Grav.* 13, 377 (1996).

[110] DESI Collaboration, "DESI DR2 results II: measurements of baryon acoustic oscillations and cosmological constraints", *Phys. Rev. D* 112, 083515 (2025).

[111] DESI Collaboration, "DESI 2024 VI: cosmological constraints from the measurements of baryon acoustic oscillations", *JCAP* 02 (2025) 021.

[112] S. Tremaine, J. E. Gunn, "Dynamical role of light neutral leptons in cosmology", *Phys. Rev. Lett.* 42, 407 (1979).

[113] T. Piffl et al., "The RAVE survey: the Galactic escape speed and the mass of the Milky Way", *Astron. Astrophys.* 562, A91 (2014).

[114] R. Lange et al., "Improved limits for violations of local position invariance from atomic clock comparisons", *Phys. Rev. Lett.* 126, 011102 (2021).

[115] M. T. Murphy et al., "Fundamental physics with ESPRESSO: precise limit on variations in the fine-structure constant towards the bright quasar HE 0515−4414", *Astron. Astrophys.* 658, A123 (2022).

[116] Planck Collaboration, "Planck intermediate results. XXIV. Constraints on variations in fundamental constants", *Astron. Astrophys.* 580, A22 (2015).

[117] P. W. Higgs, "Broken symmetries and the masses of gauge bosons", *Phys. Rev. Lett.* 13, 508 (1964).

[118] F. Englert, R. Brout, "Broken symmetry and the mass of gauge vector mesons", *Phys. Rev. Lett.* 13, 321 (1964).

[119] J. Schwinger, "On quantum-electrodynamics and the magnetic moment of the electron", *Phys. Rev.* 73, 416 (1948).

[120] X. Fan, T. G. Myers, B. A. D. Sukra, G. Gabrielse, "Measurement of the electron magnetic moment", *Phys. Rev. Lett.* 130, 071801 (2023).

[121] S. L. Glashow, "Partial-symmetries of weak interactions", *Nucl. Phys.* 22, 579 (1961).

[122] S. Weinberg, "A model of leptons", *Phys. Rev. Lett.* 19, 1264 (1967).

[123] A. Salam, "Weak and electromagnetic interactions", in *Elementary Particle Theory*, ed. N. Svartholm (Almqvist and Wiksell, Stockholm, 1968), 367.

[124] U. Amaldi, W. de Boer, H. Fürstenau, "Comparison of grand unified theories with electroweak and strong coupling constants measured at LEP", *Phys. Lett. B* 260, 447 (1991).

[125] J. M. Maldacena, "The large N limit of superconformal field theories and supergravity", *Adv. Theor. Math. Phys.* 2, 231 (1998).

[126] A. Ashtekar, "New variables for classical and quantum gravity", *Phys. Rev. Lett.* 57, 2244 (1986).

[127] C. Rovelli, L. Smolin, "Discreteness of area and volume in quantum gravity", *Nucl. Phys. B* 442, 593 (1995).

[128] S. Weinberg, "Ultraviolet divergences in quantum theories of gravitation", in *General Relativity: An Einstein Centenary Survey*, ed. S. W. Hawking, W. Israel (Cambridge University Press, 1979), 790.

[129] M. Reuter, "Nonperturbative evolution equation for quantum gravity", *Phys. Rev. D* 57, 971 (1998).

[130] J. Ambjørn, A. Görlich, J. Jurkiewicz, R. Loll, "Planckian birth of the quantum de Sitter universe", arXiv:0712.2485 (2007).

[131] L. Bombelli, J. Lee, D. Meyer, R. D. Sorkin, "Space-time as a causal set", *Phys. Rev. Lett.* 59, 521 (1987).

[132] M. Van Raamsdonk, "Building up spacetime with quantum entanglement", *Gen. Relativ. Gravit.* 42, 2323 (2010).

[133] J. Maldacena, L. Susskind, "Cool horizons for entangled black holes", *Fortschr. Phys.* 61, 781 (2013).

[134] J. Oppenheim, A. Russo, "Anomalous contribution to galactic rotation curves due to stochastic spacetime", arXiv:2402.19459 (2024).

[135] M. Milgrom, "A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis", *Astrophys. J.* 270, 365 (1983).

[136] F. W. Hehl, B. Mashhoon, "Nonlocal gravity simulates dark matter", arXiv:0812.1059 (2008).

[137] M. A. Amin, R. Easther, H. Finkel, R. Flauger, M. P. Hertzberg, "Oscillons after inflation", *Phys. Rev. Lett.* 108, 241302 (2012).

[138] L. Kofman, A. Linde, A. A. Starobinsky, "Reheating after inflation", *Phys. Rev. Lett.* 73, 3195 (1994).

[139] E. Cotner, A. Kusenko, V. Takhistov, "Primordial black holes from inflaton fragmentation into oscillons", *Phys. Rev. D* 98, 083513 (2018).

[140] M. Gorton, A. M. Green, "How open is the asteroid-mass primordial black hole window?", *SciPost Phys.* 17, 032 (2024).

[141] G. Agazie et al. (NANOGrav Collaboration), "The NANOGrav 15 yr data set: evidence for a gravitational-wave background", *Astrophys. J. Lett.* 951, L8 (2023).

[142] D. Clowe, M. Bradač, A. H. Gonzalez, M. Markevitch, S. W. Randall, C. Jones, D. Zaritsky, "A direct empirical proof of the existence of dark matter", *Astrophys. J. Lett.* 648, L109 (2006).