

# Data Science for Biological, Medical and Health Research: Notes for 432

Thomas E. Love, Ph.D.

Built 2021-01-28 12:54:59



# Contents

|  |           |
|--|-----------|
| <b>Introduction</b>  | <b>5</b>  |
| <b>R Packages used in these notes</b>                                    | <b>7</b>  |
| General Theme for <code>ggplot</code> work . . . . .                     | 8         |
| Data used in these notes . . . . .                                       | 8         |
| <b>1 Building Table 1</b>  | <b>9</b>  |
| 1.1 Data load . . . . .  | 9         |
| 1.2 Two examples from the <i>New England Journal of Medicine</i> . . . . | 10        |
| 1.3 The MR CLEAN trial . . . . .   | 12        |
| 1.4 Simulated <code>fakestroke</code> data . . . . .                     | 14        |
| 1.5 Building Table 1 for <code>fakestroke</code> : Attempt 1 . . . . .   | 15        |
| 1.6 <code>fakestroke</code> Table 1: Attempt 2 . . . . .                 | 18        |
| 1.7 Obtaining a more detailed Summary . . . . .                          | 21        |
| 1.8 Exporting the Completed Table 1 from R to Excel or Word . . . .      | 24        |
| 1.9 A Controlled Biological Experiment - The Blood-Brain Barrier . .     | 26        |
| 1.10 The <code>bloodbrain.csv</code> file . . . . .                      | 26        |
| 1.11 A Table 1 for <code>bloodbrain</code> . . . . .                     | 27        |
| <b>2 BRFSS SMART Data</b>  | <b>35</b> |
| 2.1 Key resources . . . . .  | 35        |
| 2.2 Ingesting the Raw Data . . . . .                                     | 36        |
| 2.3 Ingesting from our CSV file . . . . .                                | 37        |
| 2.4 What does the raw data look like? . . . . .                          | 37        |
| 2.5 Cleaning the BRFSS Data . . . . .                                    | 38        |
| 2.6 Imputing Age and Income as Quantitative from Thin Air . . . .        | 101       |
| 2.7 Clean Data in the State of Ohio . . . . .                            | 105       |
| 2.8 Clean Cleveland-Elyria Data . . . . .                                | 106       |



# Introduction

These Notes provide a series of examples using R to work through issues that are likely to come up in PQHS/CRSP/MPHP 432.

While these Notes share some of the features of a textbook, they are neither comprehensive nor completely original. The main purpose is to give students in 432 a set of common materials on which to draw during the course. In class, we will sometimes:

- reiterate points made in this document,
- amplify what is here,
- simplify the presentation of things done here,
- use new examples to show some of the same techniques,
- refer to issues not mentioned in this document,

but what we don't (always) do is follow these notes very precisely. We assume instead that you will read the materials and try to learn from them, just as you will attend classes and try to learn from them. We welcome feedback of all kinds on this document or anything else via Piazza.

What you will mostly find are brief explanations of a key idea or summary, accompanied (most of the time) by R code and a demonstration of the results of applying that code.

Everything you see here is available to you as HTML or PDF. You will also have access to the R Markdown files, which contain the code which generates everything in the document, including all of the R results. We will demonstrate the use of R Markdown (this document is generated with the additional help of an R package called bookdown) and R Studio (the “program” which we use to interface with the R language) in class.

To download the data and R code related to these notes, visit the appropriate link at the 432 course website.



# R Packages used in these notes

Here, we'll load in some packages used in these notes. The list of R Packages we will use in 432 is more extensive, and is available on our course website.

```
library(here)
library(janitor)
library(magrittr)

library(tableone)

library(broom)
library(haven)
library(janitor)
library(patchwork)
library(Hmisc)
library(rms)

library(visdat)
library(naniar)
library(caret)
library(simputation)
library(car)
library(mice)
library(leaps)
library(lars)
library(Epi)
library(pROC)
library(ROCR)
library(VGAM)
library(ggribes)
library(pander)
library(arm)
```

```
library(survival)
library(survminer)
library(kableExtra)

## and of course, we conclude with...

library(tidymodels)
library(tidyverse)

specify_decimal <- function(x, k) format(round(x, k), nsmall=k)
```

## General Theme for ggplot work

```
theme_set(theme_bw())
```

## Data used in these notes

All data sets used in these notes are available on our Data and Code website.

Dr. Love is in the process of moving all of the data loads below to their individual chapters.

```
prost <- read_csv("data/prost.csv")
pollution <- read_csv("data/pollution.csv")

bonding <- read_csv("data/bonding.csv")
cortisol <- read_csv("data/cortisol.csv")
emphysema <- read_csv("data/emphysema.csv")
resect <- read_csv("data/resect.csv")
colscr <- read_csv("data/screening.csv")
colscr2 <- read_csv("data/screening2.csv")
authorship <- read_csv("data/authorship.csv")
hem <- read_csv("data/hem.csv")
leukem <- read_csv("data/leukem.csv")
```



# Chapter 1

## Building Table 1

Many scientific articles involve direct comparison of results from various exposures, perhaps treatments. In 431, we studied numerous methods, including various sorts of hypothesis tests, confidence intervals, and descriptive summaries, which can help us to understand and compare outcomes in such a setting. One common approach is to present what's often called Table 1. Table 1 provides a summary of the characteristics of a sample, or of groups of samples, which is most commonly used to help understand the nature of the data being compared.

### 1.1 Data load

Let's load two data sets for this Chapter. All data sets used in these notes are available on our Data and Code website.

```
fakestroke <- read_csv("data/fakestroke.csv")
```

```
-- Column specification -----  
cols(  
  studyid = col_character(),  
  trt = col_character(),  
  age = col_double(),  
  sex = col_character(),  
  nihss = col_double(),  
  location = col_character(),  
  hx.isch = col_character(),  
  afib = col_double(),  
  dm = col_double(),  
  mrankin = col_character(),  
  sbp = col_double(),
```

```

    iv.altep = col_character(),
    time.iv = col_double(),
    aspects = col_double(),
    ia.occlus = col_character(),
    extra.ica = col_double(),
    time.rand = col_double(),
    time.punc = col_double()
  )
bloodbrain <- read_csv("data/bloodbrain.csv")

```

```

-- Column specification -----
cols(
  case = col_double(),
  brain = col_double(),
  liver = col_double(),
  tlratio = col_double(),
  solution = col_character(),
  sactime = col_double(),
  postin = col_double(),
  sex = col_character(),
  wt.init = col_double(),
  wt.loss = col_double(),
  wt.tumor = col_double()
)

```

## 1.2 Two examples from the *New England Journal of Medicine*

### 1.2.1 A simple Table 1

Table 1 is especially common in the context of clinical research. Consider the excerpt below, from a January 2015 article in the *New England Journal of Medicine* (Tolaney et al. 2015).

## 1.2. TWO EXAMPLES FROM THE NEW ENGLAND JOURNAL OF MEDICINE<sup>11</sup>

**Table 1. Baseline Characteristics of the Patients.\***

| Characteristic | Patients (N = 406) |
|----------------|--------------------|
|                | no. (%)            |
| Age group      |                    |
| <50 yr         | 132 (32.5)         |
| 50–59 yr       | 137 (33.7)         |
| 60–69 yr       | 96 (23.6)          |
| ≥70 yr         | 41 (10.1)          |
| Sex            |                    |
| Female         | 405 (99.8)         |
| Male           | 1 (0.2)            |
| Race†          |                    |
| White          | 351 (86.5)         |
| Black          | 28 (6.9)           |
| Asian          | 11 (2.7)           |
| Other          | 16 (3.9)           |

This (partial) table reports baseline characteristics on age group, sex and race, describing 406 patients with HER2-positive<sup>1</sup> invasive breast cancer that began the protocol therapy. Age, sex and race (along with severity of illness) are the most commonly identified characteristics in a Table 1.

In addition to the measures shown in this excerpt, the full Table also includes detailed information on the primary tumor for each patient, including its size, nodal status and histologic grade. Footnotes tell us that the percentages shown are subject to rounding, and may not total 100, and that the race information was self-reported.

### 1.2.2 A group comparison

A more typical Table 1 involves a group comparison, for example in this excerpt from Roy et al. (2008). This Table 1 describes a multi-center randomized clinical trial comparing two different approaches to caring for patients with heart failure and atrial fibrillation<sup>2</sup>.

<sup>1</sup>HER2 = human epidermal growth factor receptor type 2. Over-expression of this occurs in 15–20% of invasive breast cancers, and has been associated with poor outcomes.

<sup>2</sup>The complete Table 1 appears on pages 2668–2669 of Roy et al. (2008), but I have only reproduced the first page and the footnote in this excerpt.

| Table 1. Baseline Characteristics of the Patients.* |                                 |                               |
|---|---------------------------------|-------------------------------|
| Variable  | Rhythm-Control Group<br>(N=682) | Rate-Control Group<br>(N=694) |
| Male sex (%)  | 78                              | 85                            |
| Age (yr)  | 66±11                           | 67±11                         |
| Body-mass index†                                    | 27.8±5.4                        | 28.0±5.1                      |
| Nonwhite race (%)‡                                  | 16                              | 13                            |
| NYHA class III or IV (%)                            |                                 |                               |
| At baseline   | 32                              | 31                            |
| During previous 6 mo                                | 76                              | 76                            |
| Predominant cardiac diagnosis (%)§                  |                                 |                               |
| Coronary artery disease                             | 48                              | 48                            |
| Valvular heart disease                              | 5                               | 5                             |
| Nonischemic cardiomyopathy                          | 36                              | 39                            |
| Congenital heart disease                            | 1                               | 1                             |
| Hypertensive heart disease                          | 10                              | 7                             |

The article provides percentages, means and standard deviations across groups, but note that it does not provide p values for the comparison of baseline characteristics. This is a common feature of NEJM reports on randomized clinical trials, where we anticipate that the two groups will be well matched at baseline. Note that the patients in this study were *randomly* assigned to either the rhythm-control group or to the rate-control group, using blocked randomization stratified by study center.

### 1.3 The MR CLEAN trial

Berkhemer et al. (2015) reported on the MR CLEAN trial, involving 500 patients with acute ischemic stroke caused by a proximal intracranial arterial occlusion. The trial was conducted at 16 medical centers in the Netherlands, where 233 were randomly assigned to the intervention (intraarterial treatment plus usual care) and 267 to control (usual care alone.) The primary outcome was the modified Rankin scale score at 90 days; this categorical scale measures functional outcome, with scores ranging from 0 (no symptoms) to 6 (death). The fundamental conclusion of Berkhemer et al. (2015) was that in patients with acute ischemic stroke caused by a proximal intracranial occlusion of the anterior circulation, intraarterial treatment administered within 6 hours after stroke onset was effective and safe.

Here's the Table 1 from Berkhemer et al. (2015).

**Table 1. Baseline Characteristics of the 500 Patients.\***

| Characteristic  | Intervention<br>(N = 233) | Control<br>(N = 267) |
|---|---------------------------|----------------------|
| Age — yr  |                           |                      |
| Median  | 65.8                      | 65.7                 |
| Interquartile range   | 54.5–76.0                 | 55.5–76.4            |
| Male sex — no. (%)  | 135 (57.9)                | 157 (58.8)           |
| NIHSS score†  |                           |                      |
| Median (interquartile range)                                  | 17 (14–21)                | 18 (14–22)           |
| Range   | 3–30                      | 4–38                 |
| Location of stroke in left hemisphere — no. (%)               | 116 (49.8)                | 153 (57.3)           |
| History of ischemic stroke — no. (%)                          | 29 (12.4)                 | 25 (9.4)             |
| Atrial fibrillation — no. (%)                                 | 66 (28.3)                 | 69 (25.8)            |
| Diabetes mellitus — no. (%)                                   | 34 (14.6)                 | 34 (12.7)            |
| Prestroke modified Rankin scale score — no. (%)‡              |                           |                      |
| 0   | 190 (81.5)                | 214 (80.1)           |
| 1   | 21 (9.0)                  | 29 (10.9)            |
| 2   | 12 (5.2)                  | 13 (4.9)             |
| >2  | 10 (4.3)                  | 11 (4.1)             |
| Systolic blood pressure — mm Hg§                              | 146±26.0                  | 145±24.4             |
| Treatment with IV alteplase — no. (%)                         | 203 (87.1)                | 242 (90.6)           |
| Time from stroke onset to start of IV alteplase — min         |                           |                      |
| Median  | 85                        | 87                   |
| Interquartile range   | 67–110                    | 65–116               |
| ASPECTS — median (interquartile range)¶                       | 9 (7–10)                  | 9 (8–10)             |
| Intracranial arterial occlusion — no./total no. (%)           |                           |                      |
| Intracranial ICA  | 1/233 (0.4)               | 3/266 (1.1)          |
| ICA with involvement of the M1 middle cerebral artery segment | 59/233 (25.3)             | 75/266 (28.2)        |
| M1 middle cerebral artery segment                             | 154/233 (66.1)            | 165/266 (62.0)       |
| M2 middle cerebral artery segment                             | 18/233 (7.7)              | 21/266 (7.9)         |
| A1 or A2 anterior cerebral artery segment                     | 1/233 (0.4)               | 2/266 (0.8)          |
| Extracranial ICA occlusion — no./total no. (%)  **            | 75/233 (32.2)             | 70/266 (26.3)        |
| Time from stroke onset to randomization — min††               |                           |                      |
| Median  | 204                       | 196                  |
| Interquartile range   | 152–251                   | 149–266              |
| Time from stroke onset to groin puncture — min                |                           |                      |
| Median  | 260                       | NA                   |
| Interquartile range   | 210–313                   |                      |

The Table was accompanied by the following notes.

\* The intervention group was assigned to intraarterial treatment plus usual care, and the control group was assigned to usual care alone. Plus-minus values are means ±SD. ICA denotes internal carotid artery, IV intravenous, and NA not applicable.

† Scores on the National Institutes of Health Stroke Scale (NIHSS) range from 0 to 42, with higher scores indicating more severe neurologic deficits. The NIHSS is a 15-item scale, and values for 30 of the 7500 items were missing (0.4%). The highest number of missing items for a single patient was 6.

‡ Scores on the modified Rankin scale of functional disability range from 0 (no symptoms) to 6 (death). A score of 2 or less indicates functional independence.

§ Data on systolic blood pressure at baseline were missing for one patient assigned to the control group.

¶ The Alberta Stroke Program Early Computed Tomography Score (ASPECTS) is a measure of the extent of stroke. Scores range from 0 to 10, with higher scores indicating fewer early ischemic changes. Scores were not available for four patients assigned to the control group: noncontrast computed tomography was not performed in one patient, and three patients had strokes in the territory of the anterior cerebral artery.

|| Vessel imaging was not performed in one patient in the control group, so the level of occlusion was not known.

\*\* Extracranial ICA occlusions were reported by local investigators.

†† Data were missing for two patients in the intervention group.

## 1.4 Simulated `fakestroke` data

Consider the simulated data, available on our Data and Code website in the `fakestroke.csv` file, which I built to let us mirror the Table 1 for MR CLEAN (Berkhemer et al. 2015). The `fakestroke.csv` file contains the following 18 variables for 500 patients.

| Variable               | Description  |
|------------------------|--|
| <code>studyid</code>   | Study ID # (z001 through z500)   |
| <code>trt</code>       | Treatment group (Intervention or Control)  |
| <code>age</code>       | Age in years   |
| <code>sex</code>       | Male or Female   |
| <code>nihss</code>     | NIH Stroke Scale Score (can range from 0-42; higher scores indicate more severe neurological deficits)   |
| <code>location</code>  | Stroke Location - Left or Right Hemisphere   |
| <code>hx.isch</code>   | History of Ischemic Stroke (Yes/No)  |
| <code>afib</code>      | Atrial Fibrillation (1 = Yes, 0 = No)  |
| <code>dm</code>        | Diabetes Mellitus (1 = Yes, 0 = No)  |
| <code>mraink</code>    | Pre-stroke modified Rankin scale score (0, 1, 2 or > 2) indicating functional disability - complete range is 0 (no symptoms) to 6 (death)                |
| <code>sbp</code>       | Systolic blood pressure, in mm Hg  |
| <code>iv.altep</code>  | Treatment with IV alteplase (Yes/No)   |
| <code>time.iv</code>   | Time from stroke onset to start of IV alteplase (minutes) if <code>iv.altep=Yes</code>   |
| <code>aspects</code>   | Alberta Stroke Program Early Computed Tomography score, which measures extent of stroke from 0 - 10; higher scores indicate fewer early ischemic changes |
| <code>ia.occlus</code> | Intracranial arterial occlusion, based on vessel imaging - five categories <sup>3</sup>  |
| <code>extra.ica</code> | Extracranial ICA occlusion (1 = Yes, 0 = No)   |
| <code>time.rand</code> | Time from stroke onset to study randomization, in minutes  |
| <code>time.punc</code> | Time from stroke onset to groin puncture, in minutes (only if Intervention)  |

Here's a quick look at the simulated data in `fakestroke`.

```
fakestroke
```

```
# A tibble: 500 x 18
```

```
  studyid trt   age sex  nihss location hx.isch afib   dm mraink  sbp
  <chr>   <chr> <dbl> <chr> <dbl> <chr>   <chr> <dbl> <dbl> <chr> <dbl>
```

<sup>3</sup>The five categories are Intracranial ICA, ICA with involvement of the M1 middle cerebral artery segment, M1 middle cerebral artery segment, M2 middle cerebral artery segment, A1 or A2 anterior cerebral artery segment

```

1 z001    Cont~    53 Male    21 Right    No          0      0 2      127
2 z002    Inte~    51 Male    23 Left     No          1      0 0      137
3 z003    Cont~    68 Fema~   11 Right    No          0      0 0      138
4 z004    Cont~    28 Male    22 Left     No          0      0 0      122
5 z005    Cont~    91 Male    24 Right    No          0      0 0      162
6 z006    Cont~    34 Fema~   18 Left     No          0      0 2      166
7 z007    Inte~    75 Male    25 Right    No          0      0 0      140
8 z008    Cont~    89 Fema~   18 Right    No          0      0 0      157
9 z009    Cont~    75 Male    25 Left     No          1      0 2      129
10 z010   Inte~    26 Fema~   27 Right    No          0      0 0      143
# ... with 490 more rows, and 7 more variables: iv.altep <chr>, time.iv <dbl>,
#   aspects <dbl>, ia.occlus <chr>, extra.ica <dbl>, time.rand <dbl>,
#   time.punc <dbl>

```

## 1.5 Building Table 1 for fakestroke: Attempt 1

Our goal, then, is to take the data in `fakestroke.csv` and use it to generate a Table 1 for the study that compares the 233 patients in the Intervention group to the 267 patients in the Control group, on all of the other variables (except study ID #) available. I'll use the `tableone` package of functions available in R to help me complete this task. We'll make a first attempt, using the `CreateTableOne` function in the `tableone` package. To use the function, we'll need to specify:

- the `vars` or variables we want to place in the rows of our Table 1 (which will include just about everything in the `fakestroke` data except the `studyid` code and the `trt` variable for which we have other plans, and the `time.punc` which applies only to subjects in the Intervention group.)
  - A useful trick here is to use the `dput` function, specifically something like `dput(names(fakestroke))` can be used to generate a list of all of the variables included in the `fakestroke` tibble, and then this can be copied and pasted into the `vars` specification, saving some typing.
- the `strata` which indicates the levels want to use in the columns of our Table 1 (for us, that's `trt`)

```

fs.vars <- c("age", "sex", "nihss", "location",
             "hx.isch", "afib", "dm", "mrankin", "sbp",
             "iv.altep", "time.iv", "aspects",
             "ia.occlus", "extra.ica", "time.rand")

fs.trt <- c("trt")

att1 <- CreateTableOne(data = fakestroke,
                      vars = fs.vars,
                      strata = fs.trt)

print(att1)

```

|                       | Stratified by trt |                |       |      |
|-----------------------|-------------------|----------------|-------|------|
|                       | Control           | Intervention   | p     | test |
| n                     | 267               | 233            |       |      |
| age (mean (SD))       | 65.38 (16.10)     | 63.93 (18.09)  | 0.343 |      |
| sex = Male (%)        | 157 (58.8)        | 135 (57.9)     | 0.917 |      |
| nihss (mean (SD))     | 18.08 (4.32)      | 17.97 (5.04)   | 0.787 |      |
| location = Right (%)  | 114 (42.7)        | 117 (50.2)     | 0.111 |      |
| hx.isch = Yes (%)     | 25 ( 9.4)         | 29 (12.4)      | 0.335 |      |
| afib (mean (SD))      | 0.26 (0.44)       | 0.28 (0.45)    | 0.534 |      |
| dm (mean (SD))        | 0.13 (0.33)       | 0.12 (0.33)    | 0.923 |      |
| mrankin (%)           |                   |                | 0.922 |      |
| > 2                   | 11 ( 4.1)         | 10 ( 4.3)      |       |      |
| 0                     | 214 (80.1)        | 190 (81.5)     |       |      |
| 1                     | 29 (10.9)         | 21 ( 9.0)      |       |      |
| 2                     | 13 ( 4.9)         | 12 ( 5.2)      |       |      |
| sbp (mean (SD))       | 145.00 (24.40)    | 146.03 (26.00) | 0.647 |      |
| iv.altep = Yes (%)    | 242 (90.6)        | 203 (87.1)     | 0.267 |      |
| time.iv (mean (SD))   | 87.96 (26.01)     | 98.22 (45.48)  | 0.003 |      |
| aspects (mean (SD))   | 8.65 (1.47)       | 8.35 (1.64)    | 0.033 |      |
| ia.occlus (%)         |                   |                | 0.795 |      |
| A1 or A2              | 2 ( 0.8)          | 1 ( 0.4)       |       |      |
| ICA with M1           | 75 (28.2)         | 59 (25.3)      |       |      |
| Intracranial ICA      | 3 ( 1.1)          | 1 ( 0.4)       |       |      |
| M1                    | 165 (62.0)        | 154 (66.1)     |       |      |
| M2                    | 21 ( 7.9)         | 18 ( 7.7)      |       |      |
| extra.ica (mean (SD)) | 0.26 (0.44)       | 0.32 (0.47)    | 0.150 |      |
| time.rand (mean (SD)) | 213.88 (70.29)    | 202.51 (57.33) | 0.051 |      |

### 1.5.1 Some of this is very useful, and other parts need to be fixed.

1. The 1/0 variables (`afib`, `dm`, `extra.ica`) might be better if they were treated as the factors they are, and reported as the Yes/No variables are reported, with counts and percentages rather than with means and standard deviations.
2. In some cases, we may prefer to re-order the levels of the categorical (factor) variables, particularly the `mrankin` variable, but also the `ia.occlus` variable. It would also be more typical to put the Intervention group to the left and the Control group to the right, so we may need to adjust our `trt` variable's levels accordingly.
3. For each of the quantitative variables (`age`, `nihss`, `sbp`, `time.iv`, `aspects`, `extra.ica`, `time.rand` and `time.punc`) we should make a decision whether a summary with mean and standard deviation is appropriate, or whether we should instead summarize with, say, the median and quartiles. A mean and standard deviation really only yields an appropriate summary when



the data are least approximately Normally distributed. This will make the  $p$  values a bit more reasonable, too. The `test` column in the first attempt will soon have something useful to tell us.

4. If we'd left in the `time.punc` variable, we'd get some warnings, having to do with the fact that `time.punc` is only relevant to patients in the Intervention group.

### 1.5.2 fakestroke Cleaning Up Categorical Variables

Let's specify each of the categorical variables as categorical explicitly. This helps the `CreateTableOne` function treat them appropriately, and display them with counts and percentages. This includes all of the 1/0, Yes/No and multi-categorical variables.

```
fs.factorvars <- c("sex", "location", "hx.isch", "afib", "dm",
                  "mrankin", "iv.altep", "ia.occlus", "extra.ica")
```

Then we simply add a `factorVars = fs.factorvars` call to the `CreateTableOne` function.

We also want to re-order some of those categorical variables, so that the levels are more useful to us. Specifically, we want to:

- place Intervention before Control in the `trt` variable,
- reorder the `mrankin` scale as 0, 1, 2, > 2, and
- rearrange the `ia.occlus` variable to the order<sup>4</sup> presented in Berkhemer et al. (2015).

To accomplish this, we'll use the `fct_relevel` function from the `forcats` package (loaded with the rest of the core `tidyverse` packages) to reorder our levels manually.

```
fakestroke <- fakestroke %>%
  mutate(trt = fct_relevel(trt, "Intervention", "Control"),
         mrankin = fct_relevel(mrankin, "0", "1", "2", "> 2"),
         ia.occlus = fct_relevel(ia.occlus, "Intracranial ICA",
                                "ICA with M1", "M1", "M2",
                                "A1 or A2")
  )
```

---

<sup>4</sup>We might also have considered reordering the `ia.occlus` factor by its frequency, using the `fct_infreq` function

## 1.6 fakestroke Table 1: Attempt 2

```
att2 <- CreateTableOne(data = fakestroke,
                        vars = fs.vars,
                        factorVars = fs.factorvars,
                        strata = fs.trt)
print(att2)
```

|                       | Stratified by trt |                |       |      |
|-----------------------|-------------------|----------------|-------|------|
|                       | Intervention      | Control        | p     | test |
| n                     | 233               | 267            |       |      |
| age (mean (SD))       | 63.93 (18.09)     | 65.38 (16.10)  | 0.343 |      |
| sex = Male (%)        | 135 (57.9)        | 157 (58.8)     | 0.917 |      |
| nihss (mean (SD))     | 17.97 (5.04)      | 18.08 (4.32)   | 0.787 |      |
| location = Right (%)  | 117 (50.2)        | 114 (42.7)     | 0.111 |      |
| hx.isch = Yes (%)     | 29 (12.4)         | 25 ( 9.4)      | 0.335 |      |
| afib = 1 (%)          | 66 (28.3)         | 69 (25.8)      | 0.601 |      |
| dm = 1 (%)            | 29 (12.4)         | 34 (12.7)      | 1.000 |      |
| mrankin (%)           |                   |                | 0.922 |      |
| 0                     | 190 (81.5)        | 214 (80.1)     |       |      |
| 1                     | 21 ( 9.0)         | 29 (10.9)      |       |      |
| 2                     | 12 ( 5.2)         | 13 ( 4.9)      |       |      |
| > 2                   | 10 ( 4.3)         | 11 ( 4.1)      |       |      |
| sbp (mean (SD))       | 146.03 (26.00)    | 145.00 (24.40) | 0.647 |      |
| iv.altep = Yes (%)    | 203 (87.1)        | 242 (90.6)     | 0.267 |      |
| time.iv (mean (SD))   | 98.22 (45.48)     | 87.96 (26.01)  | 0.003 |      |
| aspects (mean (SD))   | 8.35 (1.64)       | 8.65 (1.47)    | 0.033 |      |
| ia.occlus (%)         |                   |                | 0.795 |      |
| Intracranial ICA      | 1 ( 0.4)          | 3 ( 1.1)       |       |      |
| ICA with M1           | 59 (25.3)         | 75 (28.2)      |       |      |
| M1                    | 154 (66.1)        | 165 (62.0)     |       |      |
| M2                    | 18 ( 7.7)         | 21 ( 7.9)      |       |      |
| A1 or A2              | 1 ( 0.4)          | 2 ( 0.8)       |       |      |
| extra.ica = 1 (%)     | 75 (32.2)         | 70 (26.3)      | 0.179 |      |
| time.rand (mean (SD)) | 202.51 (57.33)    | 213.88 (70.29) | 0.051 |      |

The categorical data presentation looks much improved.

### 1.6.1 What summaries should we show?

Now, we'll move on to the issue of making a decision about what type of summary to show for the quantitative variables. Since the **fakestroke** data are just simulated and only match the summary statistics of the original results, not the details, we'll adopt the decisions made by Berkhemer et al. (2015), which

were to use medians and interquartile ranges to summarize the distributions of all of the continuous variables **except** systolic blood pressure.

- Specifying certain quantitative variables as *non-normal* causes R to show them with medians and the 25th and 75th percentiles, rather than means and standard deviations, and also causes those variables to be tested using non-parametric tests, like the Wilcoxon signed rank test, rather than the t test. The **test** column indicates this with the word **nonnorm**.
  - In real data situations, what should we do? The answer is to look at the data. I would not make the decision as to which approach to take without first plotting (perhaps in a histogram or a Normal Q-Q plot) the observed distributions in each of the two samples, so that I could make a sound decision about whether Normality was a reasonable assumption. If the means and medians are meaningfully different from each other, this is especially important.
  - To be honest, though, if the variable in question is a relatively unimportant covariate and the *p* values for the two approaches are nearly the same, I'd say that further investigation is rarely important.
- Specifying *exact* tests for certain categorical variables (we'll try this for the **location** and **mrarkin** variables) can be done, and these changes will be noted in the **test** column, as well.
  - In real data situations, I would rarely be concerned about this issue, and often choose Pearson (approximate) options across the board. This is reasonable so long as the number of subjects falling in each category is reasonably large, say above 10. If not, then an exact test may be a tiny improvement.
  - Paraphrasing Rosenbaum (2017), having an exact rather than an approximate test result is about as valuable as having a nice crease in your trousers.

To finish our Table 1, then, we need to specify which variables should be treated as non-Normal in the **print** statement - notice that we don't need to redo the **CreateTableOne** for this change.

```
print(att2,
      nonnormal = c("age", "nihss", "time.iv", "aspects", "time.rand"),
      exact = c("location", "mrarkin"))
```

|                      | Stratified by trt    |                      |
|----------------------|----------------------|----------------------|
|                      | Intervention         | Control              |
| n                    | 233                  | 267                  |
| age (median [IQR])   | 65.80 [54.50, 76.00] | 65.70 [55.75, 76.20] |
| sex = Male (%)       | 135 (57.9)           | 157 (58.8)           |
| nihss (median [IQR]) | 17.00 [14.00, 21.00] | 18.00 [14.00, 22.00] |
| location = Right (%) | 117 (50.2)           | 114 (42.7)           |
| hx.isch = Yes (%)    | 29 (12.4)            | 25 (9.4)             |
| afib = 1 (%)         | 66 (28.3)            | 69 (25.8)            |

|                          |                         |                         |
|--------------------------|-------------------------|-------------------------|
| dm = 1 (%)               | 29 (12.4)               | 34 (12.7)               |
| mrankin (%)              |                         |                         |
| 0                        | 190 (81.5)              | 214 (80.1)              |
| 1                        | 21 ( 9.0)               | 29 (10.9)               |
| 2                        | 12 ( 5.2)               | 13 ( 4.9)               |
| > 2                      | 10 ( 4.3)               | 11 ( 4.1)               |
| sbp (mean (SD))          | 146.03 (26.00)          | 145.00 (24.40)          |
| iv.altep = Yes (%)       | 203 (87.1)              | 242 (90.6)              |
| time.iv (median [IQR])   | 85.00 [67.00, 110.00]   | 87.00 [65.00, 116.00]   |
| aspects (median [IQR])   | 9.00 [7.00, 10.00]      | 9.00 [8.00, 10.00]      |
| ia.occlus (%)            |                         |                         |
| Intracranial ICA         | 1 ( 0.4)                | 3 ( 1.1)                |
| ICA with M1              | 59 (25.3)               | 75 (28.2)               |
| M1                       | 154 (66.1)              | 165 (62.0)              |
| M2                       | 18 ( 7.7)               | 21 ( 7.9)               |
| A1 or A2                 | 1 ( 0.4)                | 2 ( 0.8)                |
| extra.ica = 1 (%)        | 75 (32.2)               | 70 (26.3)               |
| time.rand (median [IQR]) | 204.00 [152.00, 249.50] | 196.00 [149.00, 266.00] |
| Stratified by trt        |                         |                         |
|                          | p                       | test                    |
| n                        |                         |                         |
| age (median [IQR])       | 0.579                   | nonnorm                 |
| sex = Male (%)           | 0.917                   |                         |
| nihss (median [IQR])     | 0.453                   | nonnorm                 |
| location = Right (%)     | 0.106                   | exact                   |
| hx.isch = Yes (%)        | 0.335                   |                         |
| afib = 1 (%)             | 0.601                   |                         |
| dm = 1 (%)               | 1.000                   |                         |
| mrankin (%)              | 0.917                   | exact                   |
| 0                        |                         |                         |
| 1                        |                         |                         |
| 2                        |                         |                         |
| > 2                      |                         |                         |
| sbp (mean (SD))          | 0.647                   |                         |
| iv.altep = Yes (%)       | 0.267                   |                         |
| time.iv (median [IQR])   | 0.596                   | nonnorm                 |
| aspects (median [IQR])   | 0.075                   | nonnorm                 |
| ia.occlus (%)            | 0.795                   |                         |
| Intracranial ICA         |                         |                         |
| ICA with M1              |                         |                         |
| M1                       |                         |                         |
| M2                       |                         |                         |
| A1 or A2                 |                         |                         |
| extra.ica = 1 (%)        | 0.179                   |                         |
| time.rand (median [IQR]) | 0.251                   | nonnorm                 |

## 1.7 Obtaining a more detailed Summary

If this was a real data set, we'd want to get a more detailed description of the data to make decisions about things like potentially collapsing categories of a variable, or whether or not a normal distribution was useful for a particular continuous variable, etc. You can do this with the `summary` command applied to a created Table 1, which shows, among other things, the effect of changing from normal to non-normal  $p$  values for continuous variables, and from approximate to "exact"  $p$  values for categorical factors.

Again, as noted above, in a real data situation, we'd want to plot the quantitative variables (within each group) to make a smart decision about whether a  $t$  test or Wilcoxon approach is more appropriate.

Note in the summary below that we have some missing values here. Often, we'll present this information within the Table 1, as well.

```
summary(att2)
```

```
### Summary of continuous variables ###
```

```
trt: Intervention
```

|           | n   | miss | p.miss | mean | sd | median | p25 | p75 | min | max | skew  | kurt  |
|-----------|-----|------|--------|------|----|--------|-----|-----|-----|-----|-------|-------|
| age       | 233 | 0    | 0.0    | 64   | 18 | 66     | 54  | 76  | 23  | 96  | -0.34 | -0.52 |
| nihss     | 233 | 0    | 0.0    | 18   | 5  | 17     | 14  | 21  | 10  | 28  | 0.48  | -0.74 |
| sbp       | 233 | 0    | 0.0    | 146  | 26 | 146    | 129 | 164 | 78  | 214 | -0.07 | -0.22 |
| time.iv   | 233 | 30   | 12.9   | 98   | 45 | 85     | 67  | 110 | 42  | 218 | 1.03  | 0.08  |
| aspects   | 233 | 0    | 0.0    | 8    | 2  | 9      | 7   | 10  | 5   | 10  | -0.56 | -0.98 |
| time.rand | 233 | 2    | 0.9    | 203  | 57 | 204    | 152 | 250 | 100 | 300 | 0.01  | -1.16 |

```
-----
```

```
trt: Control
```

|           | n   | miss | p.miss | mean | sd | median | p25 | p75 | min | max | skew   | kurt  |
|-----------|-----|------|--------|------|----|--------|-----|-----|-----|-----|--------|-------|
| age       | 267 | 0    | 0.0    | 65   | 16 | 66     | 56  | 76  | 24  | 94  | -0.296 | -0.28 |
| nihss     | 267 | 0    | 0.0    | 18   | 4  | 18     | 14  | 22  | 11  | 25  | 0.017  | -1.24 |
| sbp       | 267 | 1    | 0.4    | 145  | 24 | 145    | 128 | 161 | 82  | 231 | 0.156  | 0.08  |
| time.iv   | 267 | 25   | 9.4    | 88   | 26 | 87     | 65  | 116 | 44  | 130 | 0.001  | -1.32 |
| aspects   | 267 | 4    | 1.5    | 9    | 1  | 9      | 8   | 10  | 5   | 10  | -1.071 | 0.36  |
| time.rand | 267 | 0    | 0.0    | 214  | 70 | 196    | 149 | 266 | 120 | 360 | 0.508  | -0.93 |

```
p-values
```

|           | pNormal     | pNonNormal |
|-----------|-------------|------------|
| age       | 0.342813660 | 0.57856976 |
| nihss     | 0.787487252 | 0.45311695 |
| sbp       | 0.647157646 | 0.51346132 |
| time.iv   | 0.003073372 | 0.59641104 |
| aspects   | 0.032662901 | 0.07464683 |
| time.rand | 0.050803672 | 0.25134327 |

Standardize mean differences

1 vs 2  
age 0.08478764  
nihss 0.02405390  
sbp 0.04100833  
time.iv 0.27691223  
aspects 0.19210662  
time.rand 0.17720957

=====  
### Summary of categorical variables ###

trt: Intervention

| var       | n   | miss | p.miss | level            | freq | percent | cum.percent |
|-----------|-----|------|--------|------------------|------|---------|-------------|
| sex       | 233 | 0    | 0.0    | Female           | 98   | 42.1    | 42.1        |
|           |     |      |        | Male             | 135  | 57.9    | 100.0       |
| location  | 233 | 0    | 0.0    | Left             | 116  | 49.8    | 49.8        |
|           |     |      |        | Right            | 117  | 50.2    | 100.0       |
| hx.isch   | 233 | 0    | 0.0    | No               | 204  | 87.6    | 87.6        |
|           |     |      |        | Yes              | 29   | 12.4    | 100.0       |
| afib      | 233 | 0    | 0.0    | 0                | 167  | 71.7    | 71.7        |
|           |     |      |        | 1                | 66   | 28.3    | 100.0       |
| dm        | 233 | 0    | 0.0    | 0                | 204  | 87.6    | 87.6        |
|           |     |      |        | 1                | 29   | 12.4    | 100.0       |
| mrankin   | 233 | 0    | 0.0    | 0                | 190  | 81.5    | 81.5        |
|           |     |      |        | 1                | 21   | 9.0     | 90.6        |
|           |     |      |        | 2                | 12   | 5.2     | 95.7        |
|           |     |      |        | > 2              | 10   | 4.3     | 100.0       |
| iv.altep  | 233 | 0    | 0.0    | No               | 30   | 12.9    | 12.9        |
|           |     |      |        | Yes              | 203  | 87.1    | 100.0       |
| ia.occlus | 233 | 0    | 0.0    | Intracranial ICA | 1    | 0.4     | 0.4         |
|           |     |      |        | ICA with M1      | 59   | 25.3    | 25.8        |
|           |     |      |        | M1               | 154  | 66.1    | 91.8        |
|           |     |      |        | M2               | 18   | 7.7     | 99.6        |
|           |     |      |        | A1 or A2         | 1    | 0.4     | 100.0       |
| extra.ica | 233 | 0    | 0.0    | 0                | 158  | 67.8    | 67.8        |

|   |    |      |       |
|---|----|------|-------|
| 1 | 75 | 32.2 | 100.0 |
|---|----|------|-------|

-----  
trt: Control

| var       | n   | miss | p.miss | level            | freq | percent | cum.percent |
|-----------|-----|------|--------|------------------|------|---------|-------------|
| sex       | 267 | 0    | 0.0    | Female           | 110  | 41.2    | 41.2        |
|           |     |      |        | Male             | 157  | 58.8    | 100.0       |
| location  | 267 | 0    | 0.0    | Left             | 153  | 57.3    | 57.3        |
|           |     |      |        | Right            | 114  | 42.7    | 100.0       |
| hx.isch   | 267 | 0    | 0.0    | No               | 242  | 90.6    | 90.6        |
|           |     |      |        | Yes              | 25   | 9.4     | 100.0       |
| afib      | 267 | 0    | 0.0    | 0                | 198  | 74.2    | 74.2        |
|           |     |      |        | 1                | 69   | 25.8    | 100.0       |
| dm        | 267 | 0    | 0.0    | 0                | 233  | 87.3    | 87.3        |
|           |     |      |        | 1                | 34   | 12.7    | 100.0       |
| mrankin   | 267 | 0    | 0.0    | 0                | 214  | 80.1    | 80.1        |
|           |     |      |        | 1                | 29   | 10.9    | 91.0        |
|           |     |      |        | 2                | 13   | 4.9     | 95.9        |
|           |     |      |        | > 2              | 11   | 4.1     | 100.0       |
| iv.altep  | 267 | 0    | 0.0    | No               | 25   | 9.4     | 9.4         |
|           |     |      |        | Yes              | 242  | 90.6    | 100.0       |
| ia.occlus | 267 | 1    | 0.4    | Intracranial ICA | 3    | 1.1     | 1.1         |
|           |     |      |        | ICA with M1      | 75   | 28.2    | 29.3        |
|           |     |      |        | M1               | 165  | 62.0    | 91.4        |
|           |     |      |        | M2               | 21   | 7.9     | 99.2        |
|           |     |      |        | A1 or A2         | 2    | 0.8     | 100.0       |
| extra.ica | 267 | 1    | 0.4    | 0                | 196  | 73.7    | 73.7        |
|           |     |      |        | 1                | 70   | 26.3    | 100.0       |

p-values

|          | pApprox   | pExact    |
|----------|-----------|-----------|
| sex      | 0.9171387 | 0.8561188 |
| location | 0.1113553 | 0.1056020 |
| hx.isch  | 0.3352617 | 0.3124683 |
| afib     | 0.6009691 | 0.5460206 |
| dm       | 1.0000000 | 1.0000000 |
| mrankin  | 0.9224798 | 0.9173657 |

```
iv.altep 0.2674968 0.2518374
ia.occlus 0.7945580 0.8189090
extra.ica 0.1793385 0.1667574
```

```
Standardize mean differences
      1 vs 2
sex      0.017479025
location 0.151168444
hx.isch  0.099032275
afib     0.055906317
dm       0.008673478
mrankin  0.062543164
iv.altep 0.111897009
ia.occlus 0.117394890
extra.ica 0.129370206
```

In this case, I have simulated the data to mirror the results in the published Table 1 for this study. In no way have I captured the full range of the real data, or any of the relationships in that data, so it's more important here to see what's available in the analysis, rather than to interpret it closely in the clinical context.

## 1.8 Exporting the Completed Table 1 from R to Excel or Word

Once you've built the table and are generally satisfied with it, you'll probably want to be able to drop it into Excel or Word for final cleanup.

### 1.8.1 Approach A: Save and open in Excel

One option is to **save the Table 1** to a `.csv` file within our `data` subfolder (note that the `data` folder must already exist), which you can then open directly in Excel. This is the approach I generally use. Note the addition of some `quote`, `noSpaces` and `printToggle` selections here.

```
fs.table1save <- print(att2,
  nonnormal = c("age", "nihss", "time.iv", "aspects", "time.rand"),
  exact = c("location", "mrankin"),
  quote = FALSE, noSpaces = TRUE, printToggle = FALSE)

write.csv(fs.table1save, file = "data/fs-table1.csv")
```

When I then open the `fs-table1.csv` file in Excel, it looks like this:



## 1.8. EXPORTING THE COMPLETED TABLE 1 FROM R TO EXCEL OR WORD<sup>25</sup>

|    | A                        | B                       | C                       | D     | E       |
|----|--------------------------|-------------------------|-------------------------|-------|---------|
| 1  |                          | Intervention            | Control                 | p     | test    |
| 2  | n                        | 233                     | 267                     |       |         |
| 3  | age (median [IQR])       | 65.80 [54.50, 76.00]    | 65.70 [55.75, 76.20]    | 0.579 | nonnorm |
| 4  | sex = Male (%)           | 135 (57.9)              | 157 (58.8)              | 0.917 |         |
| 5  | nihss (median [IQR])     | 17.00 [14.00, 21.00]    | 18.00 [14.00, 22.00]    | 0.453 | nonnorm |
| 6  | location = Right (%)     | 117 (50.2)              | 114 (42.7)              | 0.111 |         |
| 7  | hx.isch = Yes (%)        | 29 (12.4)               | 25 (9.4)                | 0.335 |         |
| 8  | afib = 1 (%)             | 66 (28.3)               | 69 (25.8)               | 0.601 |         |
| 9  | dm = 1 (%)               | 29 (12.4)               | 34 (12.7)               | 1     |         |
| 10 | mrarkin (%)              |                         |                         | 0.922 |         |
| 11 |                          | 0 190 (81.5)            | 214 (80.1)              |       |         |
| 12 |                          | 1 21 (9.0)              | 29 (10.9)               |       |         |
| 13 |                          | 2 12 (5.2)              | 13 (4.9)                |       |         |
| 14 | > 2                      | 10 (4.3)                | 11 (4.1)                |       |         |
| 15 | sbp (mean (sd))          | 146.03 (26.00)          | 145.00 (24.40)          | 0.647 |         |
| 16 | iv.altep = Yes (%)       | 203 (87.1)              | 242 (90.6)              | 0.267 |         |
| 17 | time.iv (median [IQR])   | 85.00 [67.00, 110.00]   | 87.00 [65.00, 116.00]   | 0.596 | nonnorm |
| 18 | aspects (median [IQR])   | 9.00 [7.00, 10.00]      | 9.00 [8.00, 10.00]      | 0.075 | nonnorm |
| 19 | ia.occlus (%)            |                         |                         | 0.795 |         |
| 20 | Intracranial ICA         | 1 (0.4)                 | 3 (1.1)                 |       |         |
| 21 | ICA with M1              | 59 (25.3)               | 75 (28.2)               |       |         |
| 22 | M1                       | 154 (66.1)              | 165 (62.0)              |       |         |
| 23 | M2                       | 18 (7.7)                | 21 (7.9)                |       |         |
| 24 | A1 or A2                 | 1 (0.4)                 | 2 (0.8)                 |       |         |
| 25 | extra.ica = 1 (%)        | 75 (32.2)               | 70 (26.3)               | 0.179 |         |
| 26 | time.rand (median [IQR]) | 204.00 [152.00, 249.50] | 196.00 [149.00, 266.00] | 0.251 | nonnorm |
| 27 | time.punc (median [IQR]) | 260.00 [212.00, 313.00] | NA [NA, NA]             | NA    | nonnorm |

And from here, I can either drop it directly into Word, or present it as is, or start tweaking it to meet formatting needs.

### 1.8.2 Approach B: Produce the Table so you can cut and paste it

```
print(att2,
      nonnormal = c("age", "nihss", "time.iv", "aspects", "time.rand"),
      exact = c("location", "mrarkin"),
      quote = TRUE, noSpaces = TRUE)
```

This will look like a mess by itself, but if you:

1. copy and paste that mess into Excel
2. select Text to Columns from the Data menu
3. select Delimited, then Space and select Treat consecutive delimiters as one

you should get something usable again.

Or, in Word,

1. insert the text
2. select the text with your mouse
3. select Insert ... Table ... Convert Text to Table
4. place a quotation mark in the “Other” area under Separate text at ...

After dropping blank columns, the result looks pretty good.

## 1.9 A Controlled Biological Experiment - The Blood-Brain Barrier

My source for the data and the following explanatory paragraph is page 307 from Ramsey and Schafer (2002). The original data come from Barnett et al. (1995).

The human brain (and that of rats, coincidentally) is protected from the bacteria and toxins that course through the bloodstream by something called the blood-brain barrier. After a method of disrupting the barrier was developed, researchers tested this new mechanism, as follows. A series of 34 rats were inoculated with human lung cancer cells to induce brain tumors. After 9-11 days they were infused with either the barrier disruption (BD) solution or, as a control, a normal saline (NS) solution. Fifteen minutes later, the rats received a standard dose of a particular therapeutic antibody (L6-F(ab')<sub>2</sub>). The key measure of the effectiveness of transmission across the brain-blood barrier is the ratio of the antibody concentration in the brain tumor to the antibody concentration in normal tissue outside the brain. The rats were then sacrificed, and the amounts of antibody in the brain tumor and in normal tissue from the liver were measured. The study's primary objective is to determine whether the antibody concentration in the tumor increased when the blood-barrier disruption infusion was given, and if so, by how much?

### 1.10 The `bloodbrain.csv` file

Consider the data, available on our Data and Code website in the `bloodbrain.csv` file, which includes the following variables:

| Variable             | Description                                       |
|----------------------|---|
| <code>case</code>    | identification number for the rat (1 - 34)        |
| <code>brain</code>   | an outcome: Brain tumor antibody count (per gram) |
| <code>liver</code>   | an outcome: Liver antibody count (per gram)       |
| <code>tlratio</code> | an outcome: tumor / liver concentration ratio     |

| Variable        | Description   |
|-----------------|---|
| <b>solution</b> | the treatment: BD (barrier disruption) or NS (normal saline)        |
| <b>sactime</b>  | a design variable: Sacrifice time (hours; either 0.5, 3, 24 or 72)  |
| <b>postin</b>   | covariate: Days post-inoculation of lung cancer cells (9, 10 or 11) |
| <b>sex</b>      | covariate: M or F   |
| <b>wt.init</b>  | covariate: Initial weight (grams)                                   |
| <b>wt.loss</b>  | covariate: Weight loss (grams)                                      |
| <b>wt.tumor</b> | covariate: Tumor weight ( $10^{-4}$ grams)                          |

And here's what the data look like in R.

```
bloodbrain

# A tibble: 34 x 11
  case brain liver tlratio solution sactime postin sex wt.init wt.loss
<dbl> <dbl> <dbl> <dbl> <chr>      <dbl> <dbl> <chr> <dbl> <dbl>
1     1  41081 1.46e6 0.0282 BD         0.5     10 F     239    5.9
2     2  44286 1.60e6 0.0276 BD         0.5     10 F     225     4
3     3 102926 1.60e6 0.0642 BD         0.5     10 F     224   -4.9
4     4  25927 1.78e6 0.0146 BD         0.5     10 F     184    9.8
5     5  42643 1.35e6 0.0316 BD         0.5     10 F     250     6
6     6  31342 1.79e6 0.0175 NS         0.5     10 F     196    7.7
7     7  22815 1.63e6 0.0140 NS         0.5     10 F     200    0.5
8     8  16629 1.62e6 0.0103 NS         0.5     10 F     273     4
9     9  22315 1.57e6 0.0142 NS         0.5     10 F     216    2.8
10    10  77961 1.06e6 0.0735 BD         3       10 F     267    2.6
# ... with 24 more rows, and 1 more variable: wt.tumor <dbl>
```

## 1.11 A Table 1 for bloodbrain

Barnett et al. (1995) did not provide a Table 1 for these data, so let's build one to compare the two **solutions** (BD vs. NS) on the covariates and outcomes, plus the natural logarithm of the tumor/liver concentration ratio (**tlratio**). We'll opt to treat the sacrifice time (**sactime**) and the days post-inoculation of lung cancer cells (**postin**) as categorical rather than quantitative variables.

```
bloodbrain <- bloodbrain %>%
  mutate(logTL = log(tlratio))

dput(names(bloodbrain))

c("case", "brain", "liver", "tlratio", "solution", "sactime",
  "postin", "sex", "wt.init", "wt.loss", "wt.tumor", "logTL")
```

OK - there's the list of variables we'll need. I'll put the outcomes at the bottom of the table.

```
bb.vars <- c("sactime", "postin", "sex", "wt.init", "wt.loss",
            "wt.tumor", "brain", "liver", "tlratio", "logTL")

bb.factors <- c("sactime", "sex", "postin")

bb.att1 <- CreateTableOne(data = bloodbrain,
                        vars = bb.vars,
                        factorVars = bb.factors,
                        strata = c("solution"))

summary(bb.att1)
```

### Summary of continuous variables ###

solution: BD

|          | n  | miss | p.miss | mean   | sd    | median | p25    | p75   | min    | max   | skew  |
|----------|----|------|--------|--------|-------|--------|--------|-------|--------|-------|-------|
| wt.init  | 17 | 0    | 0      | 243    | 3e+01 | 2e+02  | 2e+02  | 3e+02 | 2e+02  | 3e+02 | -0.39 |
| wt.loss  | 17 | 0    | 0      | 3      | 5e+00 | 4e+00  | 1e+00  | 6e+00 | -5e+00 | 1e+01 | -0.10 |
| wt.tumor | 17 | 0    | 0      | 157    | 8e+01 | 2e+02  | 1e+02  | 2e+02 | 2e+01  | 4e+02 | 0.53  |
| brain    | 17 | 0    | 0      | 56043  | 3e+04 | 5e+04  | 4e+04  | 8e+04 | 6e+03  | 1e+05 | 0.29  |
| liver    | 17 | 0    | 0      | 672577 | 7e+05 | 6e+05  | 2e+04  | 1e+06 | 2e+03  | 2e+06 | 0.35  |
| tlratio  | 17 | 0    | 0      | 2      | 3e+00 | 1e-01  | 6e-02  | 3e+00 | 1e-02  | 9e+00 | 1.58  |
| logTL    | 17 | 0    | 0      | -1     | 2e+00 | -2e+00 | -3e+00 | 1e+00 | -4e+00 | 2e+00 | 0.08  |

kurt

|          |      |
|----------|------|
| wt.init  | 0.7  |
| wt.loss  | 0.2  |
| wt.tumor | 1.0  |
| brain    | -0.6 |
| liver    | -1.7 |
| tlratio  | 1.7  |
| logTL    | -1.7 |

-----

solution: NS

|          | n  | miss | p.miss | mean   | sd    | median | p25    | p75    | min    | max   | skew  |
|----------|----|------|--------|--------|-------|--------|--------|--------|--------|-------|-------|
| wt.init  | 17 | 0    | 0      | 240    | 3e+01 | 2e+02  | 2e+02  | 3e+02  | 2e+02  | 3e+02 | 0.33  |
| wt.loss  | 17 | 0    | 0      | 4      | 4e+00 | 3e+00  | 2e+00  | 7e+00  | -4e+00 | 1e+01 | -0.09 |
| wt.tumor | 17 | 0    | 0      | 209    | 1e+02 | 2e+02  | 2e+02  | 3e+02  | 3e+01  | 5e+02 | 0.63  |
| brain    | 17 | 0    | 0      | 23887  | 1e+04 | 2e+04  | 1e+04  | 3e+04  | 1e+03  | 5e+04 | 0.30  |
| liver    | 17 | 0    | 0      | 664975 | 7e+05 | 7e+05  | 2e+04  | 1e+06  | 9e+02  | 2e+06 | 0.40  |
| tlratio  | 17 | 0    | 0      | 1      | 2e+00 | 5e-02  | 3e-02  | 9e-01  | 1e-02  | 7e+00 | 2.27  |
| logTL    | 17 | 0    | 0      | -2     | 2e+00 | -3e+00 | -3e+00 | -7e-02 | -5e+00 | 2e+00 | 0.27  |

kurt

|         |       |
|---------|-------|
| wt.init | -0.48 |
| wt.loss | 0.08  |

```
wt.tumor  0.77
brain     -0.35
liver     -1.56
tlratio   4.84
logTL     -1.61
```

## p-values

|          | pNormal     | pNonNormal  |
|----------|-------------|-------------|
| wt.init  | 0.807308940 | 0.641940278 |
| wt.loss  | 0.683756156 | 0.876749808 |
| wt.tumor | 0.151510151 | 0.190482094 |
| brain    | 0.001027678 | 0.002579901 |
| liver    | 0.974853609 | 0.904045603 |
| tlratio  | 0.320501715 | 0.221425879 |
| logTL    | 0.351633525 | 0.221425879 |

## Standardize mean differences

|          | 1 vs 2     |
|----------|------------|
| wt.init  | 0.08435244 |
| wt.loss  | 0.14099823 |
| wt.tumor | 0.50397184 |
| brain    | 1.23884159 |
| liver    | 0.01089667 |
| tlratio  | 0.34611465 |
| logTL    | 0.32420504 |

```
#####
### Summary of categorical variables ###
```

## solution: BD

| var     | n  | miss | p.miss | level | freq | percent | cum.percent |
|---------|----|------|--------|-------|------|---------|-------------|
| sactime | 17 | 0    | 0.0    | 0.5   | 5    | 29.4    | 29.4        |
|         |    |      |        | 3     | 4    | 23.5    | 52.9        |
|         |    |      |        | 24    | 4    | 23.5    | 76.5        |
|         |    |      |        | 72    | 4    | 23.5    | 100.0       |
| postin  | 17 | 0    | 0.0    | 9     | 1    | 5.9     | 5.9         |
|         |    |      |        | 10    | 14   | 82.4    | 88.2        |
|         |    |      |        | 11    | 2    | 11.8    | 100.0       |
| sex     | 17 | 0    | 0.0    | F     | 13   | 76.5    | 76.5        |
|         |    |      |        | M     | 4    | 23.5    | 100.0       |

```
-----
solution: NS
```

| var     | n  | miss | p.miss | level | freq | percent | cum.percent |
|---------|----|------|--------|-------|------|---------|-------------|
| sactime | 17 | 0    | 0.0    | 0.5   | 4    | 23.5    | 23.5        |
|         |    |      |        | 3     | 5    | 29.4    | 52.9        |
|         |    |      |        | 24    | 4    | 23.5    | 76.5        |
|         |    |      |        | 72    | 4    | 23.5    | 100.0       |
| postin  | 17 | 0    | 0.0    | 9     | 2    | 11.8    | 11.8        |
|         |    |      |        | 10    | 13   | 76.5    | 88.2        |
|         |    |      |        | 11    | 2    | 11.8    | 100.0       |
| sex     | 17 | 0    | 0.0    | F     | 13   | 76.5    | 76.5        |
|         |    |      |        | M     | 4    | 23.5    | 100.0       |

p-values

|         | pApprox   | pExact |
|---------|-----------|--------|
| sactime | 0.9739246 | 1      |
| postin  | 0.8309504 | 1      |
| sex     | 1.0000000 | 1      |

Standardize mean differences

|         | 1 vs 2    |
|---------|-----------|
| sactime | 0.1622214 |
| postin  | 0.2098877 |
| sex     | 0.0000000 |

Note that, in this particular case, the decisions we make about normality vs. non-normality (for quantitative variables) and the decisions we make about approximate vs. exact testing (for categorical variables) won't actually change the implications of the  $p$  values. Each approach gives similar results for each variable. Of course, that's not always true.

### 1.11.1 Generate final Table 1 for bloodbrain

I'll choose to treat `tlratio` and its logarithm as non-Normal, but otherwise, use `t` tests, but admittedly, that's an arbitrary decision, really.

```
print(bb.att1, nonnormal = c("tlratio", "logTL"))
```

| n           | Stratified by solution |          |
|-------------|------------------------|----------|
|             | BD                     | NS       |
| sactime (%) | 17                     | 17       |
| 0.5         | 5 (29.4)               | 4 (23.5) |
| 3           | 4 (23.5)               | 5 (29.4) |
| 24          | 4 (23.5)               | 4 (23.5) |

|                        |                       |                       |
|------------------------|-----------------------|-----------------------|
| 72                     | 4 (23.5)              | 4 (23.5)              |
| postin (%)             |                       |                       |
| 9                      | 1 ( 5.9)              | 2 (11.8)              |
| 10                     | 14 (82.4)             | 13 (76.5)             |
| 11                     | 2 (11.8)              | 2 (11.8)              |
| sex = M (%)            | 4 (23.5)              | 4 (23.5)              |
| wt.init (mean (SD))    | 242.82 (27.23)        | 240.47 (28.54)        |
| wt.loss (mean (SD))    | 3.34 (4.68)           | 3.94 (3.88)           |
| wt.tumor (mean (SD))   | 157.29 (84.00)        | 208.53 (116.68)       |
| brain (mean (SD))      | 56043.41 (33675.40)   | 23887.18 (14610.53)   |
| liver (mean (SD))      | 672577.35 (694479.58) | 664975.47 (700773.13) |
| tlratio (median [IQR]) | 0.12 [0.06, 2.84]     | 0.05 [0.03, 0.94]     |
| logTL (median [IQR])   | -2.10 [-2.74, 1.04]   | -2.95 [-3.41, -0.07]  |
| Stratified by solution |                       |                       |
|                        | p                     | test                  |
| n                      |                       |                       |
| sactime (%)            | 0.974                 |                       |
| 0.5                    |                       |                       |
| 3                      |                       |                       |
| 24                     |                       |                       |
| 72                     |                       |                       |
| postin (%)             | 0.831                 |                       |
| 9                      |                       |                       |
| 10                     |                       |                       |
| 11                     |                       |                       |
| sex = M (%)            | 1.000                 |                       |
| wt.init (mean (SD))    | 0.807                 |                       |
| wt.loss (mean (SD))    | 0.684                 |                       |
| wt.tumor (mean (SD))   | 0.152                 |                       |
| brain (mean (SD))      | 0.001                 |                       |
| liver (mean (SD))      | 0.975                 |                       |
| tlratio (median [IQR]) | 0.221 nonnorm         |                       |
| logTL (median [IQR])   | 0.221 nonnorm         |                       |

Or, we can get an Excel-readable version placed in a `data` subfolder, using

```
bb.t1 <- print(bb.att1, nonnormal = c("tlratio", "logTL"), quote = FALSE,
               noSpaces = TRUE, printToggle = FALSE)
write.csv(bb.t1, file = "data/bb-table1.csv")
```

which, when dropped into Excel, will look like this:

|    | A                      | B                     | C                     | D             | E    |
|----|------------------------|-----------------------|-----------------------|---------------|------|
| 1  |                        | BD                    | NS                    | p             | test |
| 2  | n                      | 17                    | 17                    |               |      |
| 3  | sex = M (%)            | 4 (23.5)              | 4 (23.5)              | 1             |      |
| 4  | sactime (%)            |                       |                       | 0.974         |      |
| 5  | 0.5 5 (29.4)           |                       | 4 (23.5)              |               |      |
| 6  | 3 4 (23.5)             |                       | 5 (29.4)              |               |      |
| 7  | 24 4 (23.5)            |                       | 4 (23.5)              |               |      |
| 8  | 72 4 (23.5)            |                       | 4 (23.5)              |               |      |
| 9  | postin (%)             |                       |                       | 0.831         |      |
| 10 | 9 1 (5.9)              |                       | 2 (11.8)              |               |      |
| 11 | 10 14 (82.4)           |                       | 13 (76.5)             |               |      |
| 12 | 11 2 (11.8)            |                       | 2 (11.8)              |               |      |
| 13 | wt.init (mean (sd))    | 242.82 (27.23)        | 240.47 (28.54)        | 0.807         |      |
| 14 | wt.loss (mean (sd))    | 3.34 (4.68)           | 3.94 (3.88)           | 0.684         |      |
| 15 | wt.tumor (mean (sd))   | 157.29 (84.00)        | 208.53 (116.68)       | 0.152         |      |
| 16 | brain (mean (sd))      | 56043.41 (33675.40)   | 23887.18 (14610.53)   | 0.001         |      |
| 17 | liver (mean (sd))      | 672577.35 (694479.58) | 664975.47 (700773.13) | 0.975         |      |
| 18 | tlratio (median [IQR]) | 0.12 [0.06, 2.84]     | 0.05 [0.03, 0.94]     | 0.221 nonnorm |      |
| 19 | logTL (median [IQR])   | -2.10 [-2.74, 1.04]   | -2.95 [-3.41, -0.07]  | 0.221 nonnorm |      |
| 20 |                        |                       |                       |               |      |

One thing I would definitely clean up here, in practice, is to change the presentation of the  $p$  value for **sex** from 1 to  $> 0.99$ , or just omit it altogether. I'd also drop the **computer-ese** where possible, add units for the measures, round **a lot**, identify the outcomes carefully, and use notes to indicate deviations from the main approach.



## 1.11.2 A More Finished Version (after Cleanup in Word)

**Table 1. Comparing Rats Receiving BD to those Receiving NS on Available Covariates and Design Variables, and Key Outcomes**

|   | Barrier Disruption<br>(BD: treatment) | Normal Saline<br>(NS: control) | p     |
|---|---------------------------------------|--------------------------------|-------|
| # of Rats   | 17                                    | 17                             |       |
| Sex = Male  | 4 (23.5)                              | 4 (23.5)                       | -     |
| Sacrifice Time (hours)                                  |                                       |                                | 0.97  |
| 0.5   | 5 (29.4)                              | 4 (23.5)                       |       |
| 3   | 4 (23.5)                              | 5 (29.4)                       |       |
| 24  | 4 (23.5)                              | 4 (23.5)                       |       |
| 72  | 4 (23.5)                              | 4 (23.5)                       |       |
| Days post-inoculation of<br>lung cancer cells           |                                       |                                | 0.83  |
| 9   | 1 (5.9)                               | 2 (11.8)                       |       |
| 10  | 14 (82.4)                             | 13 (76.5)                      |       |
| 11  | 2 (11.8)                              | 2 (11.8)                       |       |
| Initial Weight (g)                                      | 243 (27)                              | 240 (29)                       | 0.81  |
| Weight Loss (g)   | 3.3 (4.7)                             | 3.9 (3.9)                      | 0.68  |
| Tumor Weight (10 <sup>-4</sup> g)                       | 157.3 (84.0)                          | 208.5 (116.7)                  | 0.15  |
| Key Outcomes: mean (sd) unless otherwise indicated      |                                       |                                |       |
| Brain Tumor Antibody Count (per g)                      | 56,043 (33,675)                       | 23,887 (14,611)                | 0.001 |
| Liver Antibody Count (per g)                            | 672,577 (694,480)                     | 664,975 (700,773)              | 0.98  |
| Tumor/Liver Ratio<br>(median [Q25, Q75])                | 0.12<br>[0.06, 2.84]                  | 0.05<br>[0.03, 0.94]           | 0.22  |
| Natural Log of Tumor/Liver Ratio<br>(median [Q25, Q75]) | -2.10<br>[-2.74, 1.04]                | -2.95<br>[-3.41, -0.07]        | 0.22  |

Table 1 Notes:

- Categorical variables are summarized with counts, percentages and p values based on approximate chi-square tests.
- Continuous variables, unless otherwise indicated, are summarized with means, standard deviations and p values based on t tests.
- The Tumor / Liver ratio and its natural logarithm are summarized with the median and quartiles and a p value from a non-parametric (Wilcoxon signed rank) test.



## Chapter 2

# BRFSS SMART Data

The Centers for Disease Control analyzes Behavioral Risk Factor Surveillance System (BRFSS) survey data for specific metropolitan and micropolitan statistical areas (MMSAs) in a program called the Selected Metropolitan/Micropolitan Area Risk Trends of BRFSS (SMART BRFSS.)

In this work, we will focus on data from the 2017 SMART, and in particular on data from the state of Ohio, and from the Cleveland-Elyria, OH, Metropolitan Statistical Area. The purpose of this survey is to provide localized health information that can help public health practitioners identify local emerging health problems, plan and evaluate local responses, and efficiently allocate resources to specific needs.

In this chapter, I describe some cleaning of the BRFSS SMART data, and break it out into national, statewide, and local samples.

The data files produced by this chapter include:

- `smart_ohio.Rds` which includes data on approximately 100 variables for over 7000 subjects in six MMSAs that are at least partially located in the state of Ohio.
- `smart_cle.Rds` which includes data on those same variables for a little over 1000 subjects in the Cleveland-Elyria-Lorain OH MMSA.

### 2.1 Key resources

- the “raw” data, in the form of the 2017 SMART BRFSS MMSA Data, found in a zipped SAS Transport Format file. The data were released in October 2018.
- the MMSA Variable Layout which simply lists the variables included in the data file

- the Calculated Variables PDF which describes the risk factors by data variable names - there is also an online summary matrix of these calculated variables.
- the lengthy 2017 Survey Questions PDF which lists all questions asked as part of the BRFSS in 2017
- the enormous Codebook for the 2017 BRFSS Survey PDF which identifies the variables by name for us.

Also, for each subject, we are also provided with a sampling weight, in `_MMSAWT`, which will help us incorporate the sampling design later. These weights are at the MMSA level, and are used for generating MMSA-level estimates for variables in the data set. Details on the weighting methodology are available at this PDF.

## 2.2 Ingesting the Raw Data

To create the data files we'll use, I used the `read_xpt` function from the `haven` package to bring in the SAS XPT data file that is provided by CDC. The codes I used (but won't use in these Notes) were:

```
smart_raw <- read_xpt("MMSA2017/MMSA2017.xpt")
```

This gives the nationwide data, which has 230,875 rows and 177 columns.

But for the purposes of putting these Notes online, I needed to crank down the sample size enormously. To that end, I created a new data file, which I developed by

- importing the MMSA2017.xpt file as above
- filtering away all observations except those from MMSAs which include Ohio in their name, and
- saving the result, which now has 7,412 rows and 177 columns.

The code (again, not run here) that I used to filter to the OH-based MMSAs was:

```
smart_ohio_raw <- smart_raw %>%  
  filter(str_detect(MMSANAME, "OH"))  
  
write_csv(smart_ohio_raw, "data/smart_ohio_raw.csv")
```

So, for purposes of these notes, our complete data set is actually coming from `smart_ohio_raw.csv` and consists only of the 7,412 observations associated with the six MMSAs that include Ohio in their names.

## 2.3 Ingesting from our CSV file

Note that the `smart_ohio_raw.csv` and other data files we're developing in this Chapter are available on our Data and Code website

```
smart_ohio_raw <- read_csv("data/smart_ohio_raw.csv")

dim(smart_ohio_raw)
```

```
[1] 7412 177
```

## 2.4 What does the raw data look like?

Here is a list of all variable names included in this file. We're not going to use all of those variables, but this will give you a sense of what is available.

```
names(smart_ohio_raw)
```

```
[1] "DISPCODE" "STATERE1" "SAFETIME" "HHADULT" "GENHLTH" "PHYSHLTH"
[7] "MENTHLTH" "POORHLTH" "HLTHPLN1" "PERSDOC2" "MEDCOST" "CHECKUP1"
[13] "BPHIGH4" "BPMEDS" "CHOLCHK1" "TOLDHI2" "CHOLMED1" "CVDINFR4"
[19] "CVDCRHD4" "CVDSTRK3" "ASTHMA3" "ASTHNOW" "CHCSCNCR" "CHCOCNCR"
[25] "CHCCOPD1" "HAVARTH3" "ADDEPEV2" "CHCKIDNY" "DIABETE3" "DIABAGE2"
[31] "LMTJOIN3" "ARTHDIS2" "ARTHSOCL" "JOINPAI1" "SEX" "MARITAL"
[37] "EDUCA" "RENTHOM1" "NUMHHOL2" "NUMPHON2" "CPDEMO1A" "VETERAN3"
[43] "EMPLOY1" "CHILDREN" "INCOME2" "INTERNET" "WEIGHT2" "HEIGHT3"
[49] "PREGNANT" "DEAF" "BLIND" "DECIDE" "DIFFWALK" "DIFFDRES"
[55] "DIFFALON" "SMOKE100" "SMOKDAY2" "STOPSMK2" "LASTSMK2" "USENOW3"
[61] "ECIGARET" "ECIGNOW" "ALCDAY5" "AVEDRNK2" "DRNK3GE5" "MAXDRNKS"
[67] "FRUIT2" "FRUITJU2" "FVGREEN1" "FRENCHF1" "POTATOE1" "VEGETAB2"
[73] "EXERANY2" "EXTRACT11" "EXEROFT1" "EXERHMM1" "EXTRACT21" "EXEROFT2"
[79] "EXERHMM2" "STRENGTH" "SEATBELT" "FLUSHOT6" "FLSHTMY2" "PNEUVAC3"
[85] "SHINGLE2" "HIVTST6" "HIVTSTD3" "HIVRISK5" "CASTHDX2" "CASTHNO2"
[91] "CALLBCKZ" "WDUSENOW" "WDINFTRK" "WDHOWOFT" "WDSHARE" "NAMTRIBE"
[97] "NAMOTHR" "_URBNRRL" "_STSTR" "_IMPSEX" "_RFHLTH" "_PHYS14D"
[103] "_MENT14D" "_HCVU651" "_RFHYPE5" "_CHOLCH1" "_RFCHOL1" "_MICHD"
[109] "_LTASTH1" "_CASTHM1" "_ASTHMS1" "_DRDXAR1" "_LMTACT1" "_LMTWRK1"
[115] "_LMTSCL1" "_PRACE1" "_MRACE1" "_HISPANC" "_RACE" "_RACEG21"
[121] "_RACEGR3" "_AGEG5YR" "_AGE65YR" "_AGE80" "_AGE_G" "WTKG3"
[127] "_BMI5" "_BMI5CAT" "_RFBMI5" "_EDUCAG" "_INCOMG" "_SMOKER3"
[133] "_RFSMOK3" "_ECIGSTS" "_CURECIG" "DRNKANY5" "_RFBING5" "_DRNKWEK"
[139] "_RFDRHV5" "FTJUDA2_" "FRUTDA2_" "GREND1_" "FRNCHDA_" "POTADA1_"
[145] "VEGEDA2_" "_MISFRT1" "_MISVEG1" "_FRTRES1" "_VEGRES1" "_FRUTSU1"
[151] "_VEGESU1" "_FRTL1A" "_VEGLT1A" "_FRT16A" "_VEG23A" "_FRUITE1"
[157] "_VEGETE1" "_TOTINDA" "_MINAC11" "_MINAC21" "_PACAT1" "_PAINDX1"
```

```
[163] "_PA150R2" "_PA300R2" "_PA30021" "_PASTRNG" "_PAREC1" "_PASTAE1"
[169] "_RFSEAT2" "_RFSEAT3" "_FLSHOT6" "_PNEUMO2" "_AIDTST3" "_MMSA"
[175] "_MMSAWT" "SEQNO" "MMSANAME"
```

## 2.5 Cleaning the BRFSS Data

### 2.5.1 Identifying Information

The identifying variables for each subject are gathered in `SEQNO`, which I'll leave alone.

- Each statistical (geographic) area is identified by a `_MMSA` variable, which I'll rename `mmsa_code`, and by an `MMSANAME` which I'll rename as `mmsa_name`
- For each subject, we are also provided with a sampling weight, in `_MMSAWT`, which will help us incorporate the sampling design later in the semester. We'll rename this as `mmsa_wt`. Details on the weighting methodology are available at [https://www.cdc.gov/brfss/annual\\_data/2017/pdf/2017\\_SMART\\_BRFSS\\_MMSA\\_Methodology-508.pdf](https://www.cdc.gov/brfss/annual_data/2017/pdf/2017_SMART_BRFSS_MMSA_Methodology-508.pdf)

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(mmsa_code = `_MMSA`,
         mmsa_name = `MMSANAME`,
         mmsa_wt = `_MMSAWT`)

smart_ohio_raw %>% count(mmsa_code, mmsa_name)

# A tibble: 6 x 3
  mmsa_code mmsa_name                                     n
  <dbl> <chr>                                     <int>
1 17140 Cincinnati, OH-KY-IN, Metropolitan Statistical Area 1737
2 17460 Cleveland-Elyria, OH, Metropolitan Statistical Area 1133
3 18140 Columbus, OH, Metropolitan Statistical Area 2033
4 19380 Dayton, OH, Metropolitan Statistical Area 587
5 26580 Huntington-Ashland, WV-KY-OH, Metropolitan Statistical Area 1156
6 45780 Toledo, OH, Metropolitan Statistical Area 766
```

Those names are very long. I'll build some shorter ones, by dropping everything after the comma.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(mmsa = str_replace_all(string = mmsa_name, pattern="\\,.*$", replacement=" "))


smart_ohio_raw %>% count(mmsa, mmsa_name)

# A tibble: 6 x 3
  mmsa mmsa_name n
```

|   | <chr>               | <chr>  | <int> |
|---|---------------------|--|-------|
| 1 | "Cincinnati "       | Cincinnati, OH-KY-IN, Metropolitan Statistical Area  | 1737  |
| 2 | "Cleveland-Elyria " | Cleveland-Elyria, OH, Metropolitan Statistical Area  | 1133  |
| 3 | "Columbus "         | Columbus, OH, Metropolitan Statistical Area          | 2033  |
| 4 | "Dayton "           | Dayton, OH, Metropolitan Statistical Area            | 587   |
| 5 | "Huntington-Ashlan~ | Huntington-Ashland, WV-KY-OH, Metropolitan Statisti~ | 1156  |
| 6 | "Toledo "           | Toledo, OH, Metropolitan Statistical Area            | 766   |

And here are the sampling weights for the subjects in the Cleveland-Elyria MSA.

```
smart_ohio_raw %>%
  filter(mmsa_code == 17460) %>%
  ggplot(., aes(x = mmsa_wt)) +
  geom_histogram(bins = 30, fill = "blue", col = "white")
```



## 2.5.2 Survey Method

### 2.5.2.1 DISPCODE and its cleanup to completed

DISPCODE which is 1100 if the subject completed the interview, and 1200 if they partially completed the interview. We'll create a variable called **completed** that indicates (1 = complete, 0 = not) whether the subject completed the interview.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(completed = 12 - (DISPCODE/100))
```

```
smart_ohio_raw %>% count(DISPCODE, completed)
```

```
# A tibble: 2 x 3
  DISPCODE completed     n
  <dbl>      <dbl> <int>
1    1100          1  6277
2    1200          0  1135
```

### 2.5.2.2 STATERE1 and SAFETIME and their reduction to landline

BRFSSS is conducted by telephone. The next two variables help us understand whether the subject was contacted via land line or via cellular phone.

- **STATERE1** is 1 if the subject is a resident of the state (only asked of people in the land line version of the survey).
- **SAFETIME** is 1 if this is a safe time to talk (only asked of people in the cell phone version of the survey).
- We'll use **STATERE1** and **SAFETIME** to create an indicator variable **landline** that specifies how the respondent was surveyed (1 = land line, 0 = cell phone), as follows...

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(landline = replace_na(STATERE1, 0))

smart_ohio_raw %>% count(STATERE1, SAFETIME, landline)
```

```
# A tibble: 2 x 4
  STATERE1 SAFETIME landline     n
  <dbl>      <dbl>   <dbl> <int>
1         1        NA         1  3649
2        NA         1         0  3763
```

### 2.5.2.3 HHADULT and its cleanup to hhadults

- **HHADULT** is the response to “How many members of your household, including yourself, are 18 years of age or older?”
  - The permitted responses range from 1-76, with special values 77 for Don't Know/Not Sure and 99 for refused, with BLANK for missing or not asked.
  - So we should change all numerical values above 76 to NA for our analyses (the blanks are already regarded as NAs by R in the ingestion process.)

```
smart_ohio_raw %>% tabyl(HHADULT)
```



| HHADULT | n    | percent      | valid_percent |
|---------|------|--------------|---------------|
| 1       | 274  | 0.0369670804 | 0.236206897   |
| 2       | 603  | 0.0813545602 | 0.519827586   |
| 3       | 170  | 0.0229357798 | 0.146551724   |
| 4       | 73   | 0.0098488937 | 0.062931034   |
| 5       | 28   | 0.0037776579 | 0.024137931   |
| 6       | 4    | 0.0005396654 | 0.003448276   |
| 7       | 3    | 0.0004047491 | 0.002586207   |
| 8       | 1    | 0.0001349164 | 0.000862069   |
| 10      | 1    | 0.0001349164 | 0.000862069   |
| 11      | 1    | 0.0001349164 | 0.000862069   |
| 99      | 2    | 0.0002698327 | 0.001724138   |
| NA      | 6252 | 0.8434970318 | NA            |

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(hhadults = HHADULT,
         hhadults = replace(hhadults, hhadults > 76, NA))

smart_ohio_raw %>% count(HHADULT, hhadults) %>% tail()
```

```
# A tibble: 6 x 3
  HHADULT hhadults      n
  <dbl>    <dbl> <int>
1      7        7      3
2      8        8      1
3     10       10      1
4     11       11      1
5     99       NA      2
6     NA       NA    6252
```

### 2.5.3 Health Status (1 item)

The next variable describes relate to the subject's health status.

#### 2.5.3.1 GENHLTH and its cleanup to genhealth

- GENHLTH, the General Health variable, which is the response to “Would you say that in general your health is ...”
  - 1 = Excellent
  - 2 = Very good
  - 3 = Good
  - 4 = Fair
  - 5 = Poor
  - 7 = Don't know/Not sure
  - 9 = Refused

– BLANK = Not asked or missing

To clean up the GENHLTH data into a new variable called `genhealth` we'll need to - convince R that the 7 and 9 values are in fact best interpreted as NA, - and perhaps change the variable to a factor and incorporate the names into the levels.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(genhealth = fct_recode(factor(GENHLTH),
                                "1_Excellent" = "1",
                                "2_VeryGood" = "2",
                                "3_Good" = "3",
                                "4_Fair" = "4",
                                "5_Poor" = "5",
                                NULL = "7",
                                NULL = "9"))

smart_ohio_raw %>% count(GENHLTH, genhealth)
```

```
# A tibble: 7 x 3
  GENHLTH genhealth      n
  <dbl> <fct>      <int>
1      1 1_Excellent 1057
2      2 2_VeryGood 2406
3      3 3_Good     2367
4      4 4_Fair     1139
5      5 5_Poor      428
6      7 <NA>        10
7      9 <NA>         5
```

### 2.5.4 Healthy Days - Health-Related Quality of Life (3 items)

The next three variables describe the subject's health-related quality of life.

#### 2.5.4.1 PHYSHLTH and its cleanup to physhealth

PHYSHLTH, the Number of Days Physical Health Not Good variable, which is the response to “Now thinking about your physical health, which includes physical illness and injury, for how many days during the past 30 days was your physical health not good?”

- Values of 1-30 are numeric and reasonable.
- A value of 88 indicates “none” and should be recoded to 0.
- 77 is the code for Don't know/Not sure
- 99 is the code for Refused

- BLANK indicates Not asked or missing, and R recognizes this as NA properly.

To clean up PHYSHLTH to a new variable called `physhealth`, we'll need: - to convince R that the 77 and 99 values are in fact best interpreted as NA, and - to convince R that the 88 should be interpreted as 0.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(physhealth = PHYSHLTH,
         physhealth = replace(physhealth, physhealth %in% c(77, 99), NA),
         physhealth = replace(physhealth, physhealth == 88, 0))

smart_ohio_raw %>% count(PHYSHLTH, physhealth) %>% tail()
```

```
# A tibble: 6 x 3
  PHYSHLTH physhealth     n
    <dbl>      <dbl> <int>
1      28          28     12
2      29          29     14
3      30          30    677
4      77          NA    123
5      88           0   4380
6      99          NA     15
```

Note that we present the `tail` of the counts in this case so we can see what happens to the key values (77, 88, 99) of our original variable PHYSHLTH.

#### 2.5.4.2 MENTHLTH and its cleanup to menthealth

MENTHLTH, the Number of Days Mental Health Not Good variable, which is the response to “Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?”

- This is coded just like the PHYSHLTH variable, so we need to do the same cleaning we did there.

To clean up MENTHLTH to a new variable called `menthealth`, we'll need: - to convince R that the 77 and 99 values are in fact best interpreted as NA, and - to convince R that the 88 should be interpreted as 0.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(menthealth = MENTHLTH,
         menthealth = replace(menthealth, menthealth %in% c(77, 99), NA),
         menthealth = replace(menthealth, menthealth == 88, 0))

smart_ohio_raw %>% count(MENTHLTH, menthealth) %>% tail()
```

```
# A tibble: 6 x 3
  MENTHLTH menthealth      n
    <dbl>      <dbl> <int>
1      28         28     7
2      29         29    10
3      30         30   475
4      77         NA    86
5      88          0  4823
6      99         NA    28
```

### 2.5.4.3 POORHLTH and its cleanup to poorhealth

POORHLTH, the Poor Physical or Mental Health variable, which is the response to “During the past 30 days, for about how many days did poor physical or mental health keep you from doing your usual activities, such as self-care, work, or recreation?”

- Again, we recode just like the PHYSHLTH variable.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(poorhealth = POORHLTH,
         poorhealth = replace(poorhealth, poorhealth %in% c(77, 99), NA),
         poorhealth = replace(poorhealth, poorhealth == 88, 0))

smart_ohio_raw %>% count(POORHLTH, poorhealth) %>% tail()
```

```
# A tibble: 6 x 3
  POORHLTH poorhealth      n
    <dbl>      <dbl> <int>
1      29         29     4
2      30         30   382
3      77         NA    64
4      88          0  2194
5      99         NA    11
6      NA         NA  3337
```

There’s a lot more missingness in the `poorhealth` counts than in the other health-related quality of life measures. There’s also a strong mode at 0, and a smaller mode at 30 in each variable.

```
p1 <- ggplot(smart_ohio_raw, aes(x = physhealth)) +
  geom_histogram(binwidth = 1, fill = "orange") +
  labs(title = paste0("Bad Physical Health Days (",
                     sum(is.na(smart_ohio_raw$physhealth)),
                     " NA)"))

p2 <- ggplot(smart_ohio_raw, aes(x = menthealth)) +
```

```


geom_histogram(binwidth = 1, fill = "blue") +
labs(title = paste0("Bad Mental Health Days (",
                     sum(is.na(smart_ohio_raw$menthealth)),
                     " NA)"))

p3 <- ggplot(smart_ohio_raw, aes(x = poorhealth)) +
  geom_histogram(binwidth = 1, fill = "red") +
  labs(title = paste0("Unable to Do Usual Activities Days (",
                     sum(is.na(smart_ohio_raw$poorhealth)),
                     " NA)"))

(p1 + p2) / p3 +
  plot_annotation(title = "Health Related Quality of Life Measures in BRFSS/SMART (Ohio MMSAs)"

```

Health Related Quality of Life Measures in BRFSS/SMART (Ohio MMSAs)



### 2.5.5 Health Care Access (4 items)

The next four variables relate to the subject's health care access.

#### 2.5.5.1 HLTHPLN1 and its cleanup to healthplan

HLTHPLN1, the Have any health care coverage variable, is the response to "Do you have any kind of health care coverage, including health insurance, prepaid

plans such as HMOs, or government plans such as Medicare, or Indian Health Service?”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused

To clean up the `HLTHPLN1` data into a new variable called `healthplan` we’ll - convince R that the 7 and 9 values are in fact best interpreted as `NA`, - and turn it into an indicator variable, e.g., we will leave the variable as numeric, but change the values to 1 = Yes and 0 = No.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(healthplan = HLTHPLN1,
         healthplan = replace(healthplan, healthplan %in% c(7, 9), NA),
         healthplan = replace(healthplan, healthplan == 2, 0))

smart_ohio_raw %>% count(HLTHPLN1, healthplan)
```

```
# A tibble: 4 x 3
  HLTHPLN1 healthplan     n
  <dbl>      <dbl> <int>
1       1         1  6994
2       2         0   398
3       7        NA    10
4       9        NA    10
```

### 2.5.5.2 PERSDOC2 and its cleanup to `hasdoc` and to `numdocs2`

`PERSDOC2`, the Multiple Health Care Professionals variable, is the response to “Do you have one person you think of as your personal doctor or health care provider?” where if the response is “No,” the survey then asks “Is there more than one or is there no person who you think of as your personal doctor or health care provider?”

- 1 = Yes, only one
- 2 = More than one
- 3 = No
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

To clean up the `PERSDOC2` data into a new variable called `hasdoc` we’ll - convince R that the 7 and 9 values are in fact best interpreted as `NA`, - and turn it into an indicator variable, e.g., we will leave the variable as numeric, but change the values to 1 = Yes and 0 = No, so that the original 1 and 2 become 1, and the original 3 becomes 0.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(hasdoc = PERSDOC2,
         hasdoc = replace(hasdoc, hasdoc %in% c(7, 9), NA),
         hasdoc = replace(hasdoc, hasdoc %in% c(1, 2), 1),
         hasdoc = replace(hasdoc, hasdoc == 3, 0))

smart_ohio_raw %>% count(PERSDOC2, hasdoc)
```

```
# A tibble: 5 x 3
  PERSDOC2 hasdoc      n
    <dbl>   <dbl> <int>
1         1     1  5784
2         2     1   623
3         3     0   990
4         7    NA    14
5         9    NA     1
```

### 2.5.5.3 MEDCOST and its cleanup to costprob

MEDCOST, the Could Not See Doctor Because of Cost variable, is the response to “Was there a time in the past 12 months when you needed to see a doctor but could not because of cost?”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

This is just like HLTHPLAN.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(costprob = MEDCOST,
         costprob = replace(costprob, costprob %in% c(7, 9), NA),
         costprob = replace(costprob, costprob == 2, 0))

smart_ohio_raw %>% count(MEDCOST, costprob)
```

```
# A tibble: 4 x 3
  MEDCOST costprob      n
    <dbl>   <dbl> <int>
1         1     1   714
2         2     0  6680
3         7    NA    14
4         9    NA     4
```

### 2.5.5.4 CHECKUP1 and its cleanup to t\_checkup

CHECKUP1, the Length of time since last routine checkup variable, is the response to “About how long has it been since you last visited a doctor for a routine checkup? [A routine checkup is a general physical exam, not an exam for a specific injury, illness, or condition.]”

- 1 = Within past year (anytime less than 12 months ago)
- 2 = Within past 2 years (1 year but less than 2 years ago)
- 3 = Within past 5 years (2 years but less than 5 years ago)
- 4 = 5 or more years ago
- 7 = Don’t know/Not sure
- 8 = Never
- 9 = Refused
- BLANK = Not asked or missing

To clean up the CHECKUP1 data into a new variable called `t_checkup` we’ll - convince R that the 7 and 9 values are in fact best interpreted as NA, - relabel options 1, 2, 3, 4 and 8 while turning the variable into a factor.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(t_checkup = fct_recode(factor(CHECKUP1),
                                     "1_In-past-year" = "1",
                                     "2_1-to-2-years" = "2",
                                     "3_2-to-5-years" = "3",
                                     "4_5_plus_years" = "4",
                                     "8_Never" = "8",
                                     NULL = "7",
                                     NULL = "9"))

smart_ohio_raw %>% count(CHECKUP1, t_checkup)
```

```
# A tibble: 7 x 3
  CHECKUP1 t_checkup      n
  <dbl> <fct>      <int>
1      1 1_In-past-year 5803
2      2 2_1-to-2-years  714
3      3 3_2-to-5-years  413
4      4 4_5_plus_years  376
5      7 <NA>          68
6      8 8_Never        32
7      9 <NA>          6
```



## 2.5.6 Blood Pressure (2 measures)

### 2.5.6.1 BPHIGH4 and its cleanup to bp\_high

BPHIGH4 is asking about awareness of a hypertension diagnosis. It's the response to the question: "Have you EVER been told by a doctor, nurse or other health professional that you have high blood pressure?" In addition, if the answer was "Yes" and the respondent is female, they were then asked "Was this only when you were pregnant?"

The available codes are:

- 1 = Yes
- 2 = Yes, but female told only during pregnancy
- 3 = No
- 4 = Told borderline high or pre-hypertensive
- 7 = Don't know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

To clean up the BPHIGH4 data into a new variable called `bp_high` we'll - convince R that the 7 and 9 values are in fact best interpreted as NA, - relabel (and re-order) options 1, 2, 3, 4 while turning the variable into a factor.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(bp_high = fct_recode(factor(BPHIGH4),
                                "0_No" = "3",
                                "1_Yes" = "1",
                                "2_Only_while_pregnant" = "2",
                                "4_Borderline" = "4",
                                NULL = "7",
                                NULL = "9"),
         bp_high = fct_relevel(bp_high,
                                "0_No", "1_Yes",
                                "2_Only_while_pregnant",
                                "4_Borderline"))
```

```
smart_ohio_raw %>% count(BPHIGH4, bp_high)
```

```
# A tibble: 6 x 3
```

|   | BPHIGH4 | bp_high               | n     |
|---|---------|-----------------------|-------|
|   | <dbl>   | <fct>                 | <int> |
| 1 | 1       | 1_Yes                 | 3161  |
| 2 | 2       | 2_Only_while_pregnant | 67    |
| 3 | 3       | 0_No                  | 4114  |
| 4 | 4       | 4_Borderline          | 49    |
| 5 | 7       | <NA>                  | 19    |
| 6 | 9       | <NA>                  | 2     |

### 2.5.6.2 BPMEDS and its cleanup to bp\_meds

BPMEDS is the response to the question “Are you currently taking medicine for your high blood pressure?”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

To clean up the BPMEDS data into a new variable called `bp_meds` we’ll treat it just as we did with `HLTHPLN1` and - convince R that the 7 and 9 values are in fact best interpreted as NA, - and turn it into an indicator variable, e.g., we will leave the variable as numeric, but change the values to 1 = Yes and 0 = No.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(bp_meds = BPMEDS,
         bp_meds = replace(bp_meds, bp_meds %in% c(7, 9), NA),
         bp_meds = replace(bp_meds, bp_meds == 2, 0))

smart_ohio_raw %>% count(BPMEDS, bp_meds)
```

```
# A tibble: 5 x 3
  BPMEDS bp_meds     n
  <dbl>   <dbl> <int>
1     1     1 2675
2     2     0  481
3     7    NA    4
4     9    NA    1
5    NA    NA 4251
```

What is the relationship between our two blood pressure variables? Only the people with `bp_meds = “1_Yes”` were asked the `bp_meds` question.

```
smart_ohio_raw %>% tabyl(bp_high, bp_meds)
```

```
      bp_high  0  1  NA_
0_No        0  0 4114
1_Yes      481 2675    5
2_Only_while_pregnant 0  0  67
4_Borderline  0  0  49
<NA>        0  0  21
```

## 2.5.7 Cholesterol (3 items)

### 2.5.7.1 CHOLCHK1 and its cleanup to t\_chol

CHOLCHK1, the Length of time since cholesterol was checked, is the response to “Blood cholesterol is a fatty substance found in the blood. About how long has it been since you last had your blood cholesterol checked?”

- 1 = Never
- 2 = Within past year (anytime less than 12 months ago)
- 3 = Within past 2 years (1 year but less than 2 years ago)
- 4 = Within past 5 years (2 years but less than 5 years ago)
- 5 = 5 or more years ago
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

To clean up the CHOLCHK1 data into a new variable called `t_chol` we’ll - convince R that the 7 and 9 values are in fact best interpreted as NA, - relabel options 1, 2, 3, 4 and 8 while turning the variable into a factor.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(t_chol = fct_recode(factor(CHOLCHK1),
                                "1_Never" = "1",
                                "2_In-past-year" = "2",
                                "3_1-to-2-years" = "3",
                                "4_2-to-5-years" = "4",
                                "5_5-plus-years" = "5",
                                NULL = "7",
                                NULL = "9"))

smart_ohio_raw %>% count(CHOLCHK1, t_chol)
```

```
# A tibble: 8 x 3
  CHOLCHK1 t_chol      n
  <dbl> <fct>    <int>
1       1 1_Never    424
2       2 2_In-past-year 5483
3       3 3_1-to-2-years  559
4       4 4_2-to-5-years  289
5       5 5_5-plus-years  272
6       7 <NA>      376
7       9 <NA>       8
8      NA <NA>       1
```

The next two measures are not gathered from the people who answered “Never” to this question.

### 2.5.7.2 TOLDHI2 and its cleanup to chol\_high

TOLDHI2 is asking about awareness of a diagnosis of high cholesterol. It's the response to the question: "Have you EVER been told by a doctor, nurse or other health professional that your blood cholesterol is high?"

The available codes are:

- 1 = Yes
- 2 = No
- 7 = Don't know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

To clean up the TOLDHI2 data into a new variable called `chol_high` we'll treat it like `BPMEDS` and `HLTHPLN1` - convince R that the 7 and 9 values are in fact best interpreted as NA, - and turn it into an indicator variable, e.g., we will leave the variable as numeric, but change the values to 1 = Yes and 0 = No.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(chol_high = TOLDHI2,
         chol_high = replace(chol_high, chol_high %in% c(7, 9), NA),
         chol_high = replace(chol_high, chol_high == 2, 0))

smart_ohio_raw %>% count(TOLDHI2, chol_high)
```

```
# A tibble: 5 x 3
  TOLDHI2 chol_high      n
  <dbl>    <dbl> <int>
1      1         1  2612
2      2         0  4286
3      7        NA    70
4      9        NA     4
5     NA        NA   440
```

### 2.5.7.3 CHOLMED1 and its cleanup to chol\_meds

CHOLMED1 is the response to the question "Are you currently taking medicine prescribed by a doctor or other health professional for your blood cholesterol?"

- 1 = Yes
- 2 = No
- 7 = Don't know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

To clean up the CHOLMED1 data into a new variable called `chol_meds` we'll treat it just as we did with `HLTHPLN1` and - convince R that the 7 and 9 values are in

fact best interpreted as NA, - and turn it into an indicator variable, e.g., we will leave the variable as numeric, but change the values to 1 = Yes and 0 = No.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(chol_meds = CHOLMED1,
         chol_meds = replace(chol_meds, chol_meds %in% c(7, 9), NA),
         chol_meds = replace(chol_meds, chol_meds == 2, 0))

smart_ohio_raw %>% count(CHOLMED1, chol_meds)
```

```
# A tibble: 4 x 3
  CHOLMED1 chol_meds      n
  <dbl>      <dbl> <int>
1         1         1  1781
2         2         0   826
3         7        NA     5
4        NA        NA  4800
```

## 2.5.8 Chronic Health Conditions (14 items)

### 2.5.8.1 Self-reported diagnosis history (11 items)

The next few variables describe whether or not the subject meets a particular standard, and are all coded in the raw data the same way:

- 1 = Yes
- 2 = No
- 7 = Don't know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

and we'll recode them all to 1 = Yes, 0 = No, otherwise NA, as we've done previously.

The questions are all started with "Has a doctor, nurse, or other health professional ever told you that you had any of the following? For each, tell me Yes, No, or you're Not sure."

| Original | Revised    | Details  |
|----------|------------|--|
| CVDINFR4 | hx_mi      | (Ever told) you had a heart attack, also called a myocardial infarction? |
| CVDCRHD4 | hx_chd     | (Ever told) you had angina or coronary heart disease?                    |
| CVDSTRK3 | hx_stroke  | (Ever told) you had a stroke?  |
| ASTHMA3  | hx_asthma  | (Ever told) you had asthma?  |
| ASTHNOW  | now_asthma | Do you still have asthma? (only asked of those with Yes in ASTHMA3)      |

| Original | Revised    | Details  |
|----------|------------|--|
| CHCSCNCR | hx_skinc   | (Ever told) you had skin cancer?   |
| CHCOCNCR | hx_otherc  | (Ever told) you had any other types of cancer?   |
| CHCCOPD1 | hx_copd    | (Ever told) you have Chronic Obstructive Pulmonary Disease or COPD, emphysema or chronic bronchitis?   |
| HAVARTH3 | hx_arthr   | (Ever told) you have some form of arthritis, rheumatoid arthritis, gout, lupus, or fibromyalgia? (Arthritis diagnoses include: rheumatism, polymyalgia rheumatica; osteoarthritis (not osteoporosis); tendonitis, bursitis, bunion, tennis elbow; carpal tunnel syndrome, tarsal tunnel syndrome; joint infection, etc.) |
| ADDEPEV2 | hx_depress | (Ever told) you that you have a depressive disorder, including depression, major depression, dysthymia, or minor depression?   |
| CHCKIDNY | hx_kidney  | (Ever told) you have kidney disease? Do NOT include kidney stones, bladder infection or incontinence.  |

```

smart_ohio_raw <- smart_ohio_raw %>%
  mutate(hx_mi = CVDINFR4,
    hx_mi = replace(hx_mi, hx_mi %in% c(7, 9), NA),
    hx_mi = replace(hx_mi, hx_mi == 2, 0),
    hx_chd = CVDCRHD4,
    hx_chd = replace(hx_chd, hx_chd %in% c(7, 9), NA),
    hx_chd = replace(hx_chd, hx_chd == 2, 0),
    hx_stroke = CVDSTRK3,
    hx_stroke = replace(hx_stroke, hx_stroke %in% c(7, 9), NA),
    hx_stroke = replace(hx_stroke, hx_stroke == 2, 0),
    hx_asthma = ASTHMA3,
    hx_asthma = replace(hx_asthma, hx_asthma %in% c(7, 9), NA),
    hx_asthma = replace(hx_asthma, hx_asthma == 2, 0),
    now_asthma = ASTHNOW,
    now_asthma = replace(now_asthma, now_asthma %in% c(7, 9), NA),
    now_asthma = replace(now_asthma, now_asthma == 2, 0),
    hx_skinc = CHCSCNCR,
    hx_skinc = replace(hx_skinc, hx_skinc %in% c(7, 9), NA),
    hx_skinc = replace(hx_skinc, hx_skinc == 2, 0),
    hx_otherc = CHCOCNCR,
    hx_otherc = replace(hx_otherc, hx_otherc %in% c(7, 9), NA),
    hx_otherc = replace(hx_otherc, hx_otherc == 2, 0),
    hx_copd = CHCCOPD1,
    hx_copd = replace(hx_copd, hx_copd %in% c(7, 9), NA),
    hx_copd = replace(hx_copd, hx_copd == 2, 0),
    hx_arthr = HAVARTH3,
    hx_arthr = replace(hx_arthr, hx_arthr %in% c(7, 9), NA),

```

```

    hx_arthr = replace(hx_arthr, hx_arthr == 2, 0),
    hx_depress = ADDEPEV2,
    hx_depress = replace(hx_depress, hx_depress %in% c(7, 9), NA),
    hx_depress = replace(hx_depress, hx_depress == 2, 0),
    hx_kidney = CHCKIDNY,
    hx_kidney = replace(hx_kidney, hx_kidney %in% c(7, 9), NA),
    hx_kidney = replace(hx_kidney, hx_kidney == 2, 0))

```

We definitely should have written a function to do that, of course.

### 2.5.8.2 `_ASTHMS1` and its cleanup to `asthma`

`_ASTHMS1` categorizes subjects by asthma status as:

- 1 = Current
- 2 = Former
- 3 = Never
- 9 = Don't Know / Not Sure / Refused / Missing

We'll turn this into a factor with appropriate levels and NA information.

```

smart_ohio_raw <- smart_ohio_raw %>%
  mutate(asthma = fct_recode(
    factor(`_ASTHMS1`),
    "Current" = "1",
    "Former" = "2",
    "Never" = "3",
    NULL = "9"))

smart_ohio_raw %>% count(`_ASTHMS1`, asthma)

```

```

# A tibble: 4 x 3
  `_ASTHMS1` asthma      n
    <dbl> <fct>    <int>
1         1 Current    734
2         2 Former    248
3         3 Never    6376
4         9 <NA>      54

```

### 2.5.8.3 `DIABETE3` and its cleanup to `hx_diabetes` and `dm_status`

`DIABETE3`, the (Ever told) you have diabetes variable, is the response to “(Ever told) you have diabetes (If Yes and respondent is female, ask Was this only when you were pregnant?. If Respondent says pre-diabetes or borderline diabetes, use response code 4.)”

- 1 = Yes
- 2 = Yes, but female told only during pregnancy
- 3 = No
- 4 = No, pre-diabetes or borderline diabetes
- 7 = Don't know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

I'll create one variable called `hx_diabetes` which is 1 if `DIABETE3` = 1, and 0 otherwise, with appropriate NAs, like our other variables. Then I'll create `dm_status` to include all of this information in a factor, but again recode the missing values properly.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(hx_diabetes = DIABETE3,
         hx_diabetes = replace(hx_diabetes, hx_diabetes %in% c(7, 9), NA),
         hx_diabetes = replace(hx_diabetes, hx_diabetes %in% 2:4, 0),
         dm_status = fct_recode(factor(DIABETE3),
                                "Diabetes" = "1",
                                "Pregnancy-Induced" = "2",
                                "No-Diabetes" = "3",
                                "Pre-Diabetes" = "4",
                                NULL = "7",
                                NULL = "9"),
         dm_status = fct_relevel(dm_status,
                                "No-Diabetes",
                                "Pre-Diabetes",
                                "Pregnancy-Induced",
                                "Diabetes"))

smart_ohio_raw %>% count(DIABETE3, hx_diabetes, dm_status)
```

```
# A tibble: 6 x 4
  DIABETE3 hx_diabetes dm_status      n
    <dbl>    <dbl> <dbl> <fct>    <int>
1       1         1         1 Diabetes    1098
2       2         0         0 Pregnancy-Induced    67
3       3         0         0 No-Diabetes    6100
4       4         0         0 Pre-Diabetes    133
5       7         NA      NA <NA>         12
6       9         NA      NA <NA>          2
```

#### 2.5.8.4 DIABAGE2 and its cleanup to `dm_age`

DIABAGE2, the Age When Told Diabetic variable, is the response to “How old were you when you were told you have diabetes?” It is asked only of people with



DIABETE3 = 1 (Yes).

- The response is 1-97, with special values 98 for Don't Know/Not Sure and 99 for refused, with BLANK for missing or not asked. People 97 years of age and above were listed as 97.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(dm_age = DIABAGE2,
         dm_age = replace(dm_age, dm_age > 97, NA))

smart_ohio_raw %>% count(DIABAGE2, dm_age) %>% tail()
```

```
# A tibble: 6 x 3
  DIABAGE2 dm_age      n
  <dbl>   <dbl> <int>
1      84     84      1
2      85     85      2
3      90     90      1
4      98    NA     61
5      99    NA      4
6     NA     NA    6314
```

### 2.5.9 Arthritis Burden (4 items)

The first two measures are only asked of people with `hx_arthr = 1`, and are coded as:

- 1 = Yes
- 2 = No
- 7 = Don't know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

and we'll recode them to 1 = Yes, 0 = No, otherwise NA, as we've done previously.

#### 2.5.9.1 LMTJOIN3 (Limited because of joint symptoms), and its cleanup to `arth_lims`

This is the response to “Are you now limited in any way in any of your usual activities because of arthritis or joint symptoms?”

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(arth_lims = LMTJOIN3,
         arth_lims = replace(arth_lims, arth_lims %in% c(7, 9), NA),
         arth_lims = replace(arth_lims, arth_lims == 2, 0))

smart_ohio_raw %>% count(hx_arthr, LMTJOIN3, arth_lims)
```

```
# A tibble: 6 x 4
  hx_arthr LMTJOIN3 arth_lims      n
  <dbl>    <dbl>    <dbl> <int>
1      0      NA      NA  4587
2      1      1      1  1378
3      1      2      0  1388
4      1      7      NA   17
5      1      9      NA    2
6     NA     NA     NA   40
```

### 2.5.9.2 ARTHDIS2 (Does Arthritis Affect Whether You Work), and its cleanup to arth\_work

This is the response to “Do arthritis or joint symptoms now affect whether you work, the type of work you do or the amount of work you do?”

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(arth_work = ARTHDIS2,
         arth_work = replace(arth_work, arth_work %in% c(7, 9), NA),
         arth_work = replace(arth_work, arth_work == 2, 0))

smart_ohio_raw %>% count(ARTHDIS2, arth_work)
```

```
# A tibble: 5 x 3
  ARTHDIS2 arth_work      n
  <dbl>    <dbl> <int>
1      1      1    925
2      2      0  1808
3      7     NA    42
4      9     NA    10
5     NA     NA  4627
```

### 2.5.9.3 ARTHSOCL (Social Activities Limited Because of Joint Symptoms) and its cleanup to arth\_soc

This is the response to “During the past 30 days, to what extent has your arthritis or joint symptoms interfered with your normal social activities, such as going shopping, to the movies, or to religious or social gatherings?”

The responses are:

- 1 = A lot
- 2 = A little
- 3 = Not at all
- 7 = Don’t know/Not sure
- 9 = Refused

- BLANK = Not asked or missing

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(arth_soc = fct_recode(factor(ARTHSOCL),
                                "A lot" = "1",
                                "A little" = "2",
                                "Not at all" = "3",
                                NULL = "7",
                                NULL = "9"))

smart_ohio_raw %>% count(ARTHSOCL, arth_soc)
```

```
# A tibble: 6 x 3
  ARTHSOCL arth_soc      n
    <dbl> <fct>    <int>
1       1 A lot      606
2       2 A little    734
3       3 Not at all 1427
4       7 <NA>        15
5       9 <NA>         3
6      NA <NA>     4627
```

#### 2.5.9.4 JOINPAI1 (How Bad Was Joint Pain - scale of 0-10) and its cleanup to joint\_pain

This is the response to the following question: “Please think about the past 30 days, keeping in mind all of your joint pain or aching and whether or not you have taken medication. On a scale of 0 to 10 where 0 is no pain or aching and 10 is pain or aching as bad as it can be, DURING THE PAST 30 DAYS, how bad was your joint pain ON AVERAGE?”

The available values are 0-10, plus codes 77 (Don’t Know / Not Sure), 99 (Refused) and BLANK.

To clean up JOINPAI1 to a new variable called `joint_pain`, we’ll need to convince R that the 77 and 99 values are, like BLANK, in fact best interpreted as NA.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(joint_pain = JOINPAI1,
         joint_pain = replace(joint_pain, joint_pain %in% c(77, 99), NA))

smart_ohio_raw %>% count(JOINPAI1, joint_pain) %>% tail()
```

```
# A tibble: 6 x 3
  JOINPAI1 joint_pain      n
    <dbl>    <dbl> <int>
1       8         8    277
```

|   |    |    |      |
|---|----|----|------|
| 2 | 9  | 9  | 72   |
| 3 | 10 | 10 | 158  |
| 4 | 77 | NA | 28   |
| 5 | 99 | NA | 5    |
| 6 | NA | NA | 4627 |

## 2.5.10 Demographics (25 items)

### 2.5.10.1 \_AGEG5YR, which we'll edit into agegroup

The `_AGEG5YR` variable is a calculated variable (by CDC) obtained from the subject's age. Since the `age` data are not available, we instead get these groupings, which we'll rearrange into the `agegroup` factor.

| <code>_AGEG5YR</code> | Age range                      | <code>agegroup</code> |
|-----------------------|--------------------------------|-----------------------|
| 1                     | 18 <= AGE <= 24                | 18-24                 |
| 2                     | 25 <= AGE <= 29                | 25-29                 |
| 3                     | 30 <= AGE <= 34                | 30-34                 |
| 4                     | 35 <= AGE <= 39                | 35-39                 |
| 5                     | 40 <= AGE <= 44                | 40-44                 |
| 6                     | 45 <= AGE <= 49                | 45-49                 |
| 7                     | 50 <= AGE <= 54                | 50-54                 |
| 8                     | 55 <= AGE <= 59                | 55-59                 |
| 9                     | 60 <= AGE <= 64                | 60-64                 |
| 10                    | 65 <= AGE <= 69                | 65-69                 |
| 11                    | 70 <= AGE <= 74                | 70-74                 |
| 12                    | 75 <= AGE <= 79                | 75-79                 |
| 13                    | AGE >= 80                      | 80plus                |
| 14                    | Don't Know, Refused or Missing | NA                    |

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(agegroup = fct_recode(factor(`_AGEG5YR`),
    "18-24" = "1",
    "25-29" = "2",
    "30-34" = "3",
    "35-39" = "4",
    "40-44" = "5",
    "45-49" = "6",
    "50-54" = "7",
    "55-59" = "8",
    "60-64" = "9",
    "65-69" = "10",
    "70-74" = "11",
```



```

"Amer Indian or Alaskan" = "3",
"Asian" = "4",
"Hawaiian or Pac Island" = "5",
"Other Race" = "6",
"Multiracial" = "7",
NULL = "77",
NULL = "99"))

```

```
smart_ohio_raw %>% count(`_MRACE1`, race)
```

```

# A tibble: 9 x 3
  `_MRACE1` race          n
    <dbl> <fct>         <int>
1         1 White        6177
2         2 Black or African A 739
3         3 Amer Indian or Alaskan 66
4         4 Asian        115
5         5 Hawaitian or Pac Island 5
6         6 Other Race      43
7         7 Multiracial    153
8        77 <NA>          14
9        99 <NA>         100

```

### 2.5.10.3 \_HISPANC recoded to hispanic

The `_HISPANC` variable specifies whether or not the respondent is of Hispanic or Latinx origin. The available responses are:

- 1 = Hispanic, Latinx or Spanish origin
- 2 = Not of Hispanic, Latinx or Spanish origin
- 9 = Don't Know, Refused, or Missing

We'll turn the 9s into NA, and create an indicator variable (1 = Hispanic or Latinx, 0 = not)

```

smart_ohio_raw <- smart_ohio_raw %>%
  mutate(hispanic = 2 - `_HISPANC`,
         hispanic = replace(hispanic, hispanic < 0, NA))

smart_ohio_raw %>% count(`_HISPANC`, hispanic)

```

```

# A tibble: 3 x 3
  `_HISPANC` hispanic    n
    <dbl>    <dbl> <int>
1         1         1  146
2         2         0 7217

```

3            9            NA        49

#### 2.5.10.4 `_RACEGR3` recoded to `race_eth`

The `_RACEGR3` variable is a five-level combination of race and ethnicity. The responses are:

- 1 = White non-Hispanic
- 2 = Black non-Hispanic
- 3 = Other race non-Hispanic
- 4 = Multiracial non-Hispanic
- 5 = Hispanic
- 9 = Don't Know / Not Sure / Refused

We'll create a factor out of this information, with appropriate level names.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(race_eth = fct_recode(
    factor(`_RACEGR3`),
    "White non-Hispanic" = "1",
    "Black non-Hispanic" = "2",
    "Other race non-Hispanic" = "3",
    "Multiracial non-Hispanic" = "4",
    "Hispanic" = "5",
    NULL = "9"))

smart_ohio_raw %>% count(`_RACEGR3`, race_eth)
```

```
# A tibble: 6 x 3
  ` _RACEGR3` race_eth      n
    <dbl> <fct>          <int>
1         1 1 White non-Hispanic 6086
2         2 2 Black non-Hispanic  725
3         3 3 Other race non-Hispanic 193
4         4 4 Multiracial non-Hispanic 143
5         5 5 Hispanic        146
6         9 9 <NA>            119
```

#### 2.5.10.5 `SEX` recoded to `female`

The available levels of `SEX` are:

- 1 = Male
- 2 = Female
- 9 = Refused

We'll recode that to `female = 1` for Female, 0 Male, otherwise NA. Note the trick here is to subtract one from the coded `SEX` to get the desired `female`, but this requires that we move 9 to NA, rather than 9.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(female = SEX - 1,
         female = replace(female, female == 8, NA))

smart_ohio_raw %>% count(SEX, female)
```

```
# A tibble: 2 x 3
  SEX female     n
  <dbl> <dbl> <int>
1     1     0 3136
2     2     1 4276
```

#### 2.5.10.6 MARITAL status, revised to marital

The available levels of `MARITAL` are:

- 1 = Married
- 2 = Divorced
- 3 = Widowed
- 4 = Separated
- 5 = Never married
- 6 = A member of an unmarried couple
- 9 = Refused
- BLANK = Not asked or missing

We'll just turn this into a factor, and move 9 to NA.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(marital = fct_recode(factor(MARITAL),
                                "Married" = "1",
                                "Divorced" = "2",
                                "Widowed" = "3",
                                "Separated" = "4",
                                "Never_Married" = "5",
                                "Unmarried_Couple" = "6",
                                NULL = "9"))

smart_ohio_raw %>% count(MARITAL, marital)
```

```
# A tibble: 7 x 3
  MARITAL marital     n
  <dbl> <fct>         <int>
1     1 1 Married     3668
```



|   |                    |      |
|---|--------------------|------|
| 2 | 2 Divorced         | 1110 |
| 3 | 3 Widowed          | 978  |
| 4 | 4 Separated        | 142  |
| 5 | 5 Never_Married    | 1248 |
| 6 | 6 Unmarried_Couple | 208  |
| 7 | 9 <NA>             | 58   |

### 2.5.10.7 EDUCA recoded to educgroup

The available levels of EDUCA (Education Level) are responses to: “What is the highest grade or year of school you completed?”

- 1 = Never attended school or only kindergarten
- 2 = Grades 1 through 8 (Elementary)
- 3 = Grades 9 through 11 (Some high school)
- 4 = Grade 12 or GED (High school graduate)
- 5 = College 1 year to 3 years (Some college or technical school)
- 6 = College 4 years or more (College graduate)
- 9 = Refused
- BLANK = Not asked or missing

We'll just turn this into a factor, and move 9 to NA.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(educgroup = fct_recode(factor(EDUCA),
                                "Kindergarten" = "1",
                                "Elementary" = "2",
                                "Some_HS" = "3",
                                "HS_Grad" = "4",
                                "Some_College" = "5",
                                "College_Grad" = "6",
                                NULL = "9"))

smart_ohio_raw %>% count(EDUCA, educgroup)
```

```
# A tibble: 7 x 3
  EDUCA educgroup      n
  <dbl> <fct>      <int>
1     1 Kindergarten     3
2     2 Elementary    117
3     3 Some_HS       332
4     4 HS_Grad      2209
5     5 Some_College  2079
6     6 College_Grad  2646
7     9 <NA>          26
```

### 2.5.10.8 RENTHOM1 recoded to home\_own

The available levels of RENTHOM1 (Own or Rent Home) are responses to: “Do you own or rent your home? (Home is defined as the place where you live most of the time/the majority of the year.)”

- 1 = Own
- 2 = Rent
- 3 = Other Arrangement
- 7 = Don’t know/Not Sure
- 9 = Refused
- BLANK = Not asked or missing

We’ll recode as `home_own` = 1 if they own their home, and 0 otherwise, and dealing with missingness properly.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(home_own = RENTHOM1,
         home_own = replace(home_own, home_own %in% c(7,9), NA),
         home_own = replace(home_own, home_own %in% c(2,3), 0))

smart_ohio_raw %>% count(RENTHOM1, home_own)
```

```
# A tibble: 5 x 3
  RENTHOM1 home_own      n
    <dbl>     <dbl> <int>
1         1         1  5216
2         2         0  1793
3         3         0   348
4         7        NA    28
5         9        NA    27
```

### 2.5.10.9 CPDEM01A and its cleanup to cell\_own

CPDEM01A is the response to “Including phones for business and personal use, do you have a cell phone for personal use?”

Available responses are:

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

and we’ll recode them to 1 = Yes, 0 = No, otherwise NA, as we’ve done previously.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(cell_own = 2 - CPDEMO1A,
         cell_own = replace(cell_own, cell_own < 0, NA))

smart_ohio_raw %>% count(CPDEMO1A, cell_own)
```

```
# A tibble: 5 x 3
  CPDEMO1A cell_own      n
    <dbl>    <dbl> <int>
1         1         1  2930
2         2         0   698
3         7        NA     2
4         9        NA    19
5        NA        NA  3763
```

#### 2.5.10.10 VETERAN3 and its cleanup to veteran

VETERAN3, the Are You A Veteran variable, is the response to “Have you ever served on active duty in the United States Armed Forces, either in the regular military or in a National Guard or military reserve unit? (Active duty does not include training for the Reserves or National Guard, but DOES include activation, for example, for the Persian Gulf War.)”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(veteran = VETERAN3,
         veteran = replace(veteran, veteran %in% c(7, 9), NA),
         veteran = replace(veteran, veteran == 2, 0))

smart_ohio_raw %>% count(VETERAN3, veteran)
```

```
# A tibble: 3 x 3
  VETERAN3 veteran      n
    <dbl>    <dbl> <int>
1         1         1   927
2         2         0  6479
3         9        NA     6
```

### 2.5.10.11 EMPLOY1 and its cleanup to employment

EMPLOY1, the Employment Status variable, is the response to “Are you currently ... ?”

- 1 = Employed for wages
- 2 = Self-employed
- 3 = Out of work for 1 year or more
- 4 = Out of work for less than 1 year
- 5 = A homemaker
- 6 = A student
- 7 = Retired
- 8 = Unable to work
- 9 = Refused
- BLANK = Not asked or missing

We’ll just turn this into a factor, and move 9 to NA.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(employment = fct_recode(factor(EMPLOY1),
    "Employed_for_wages" = "1",
    "Self-employed" = "2",
    "Outofwork_1yearormore" = "3",
    "Outofwork_lt1year" = "4",
    "Homemaker" = "5",
    "Student" = "6",
    "Retired" = "7",
    "Unable_to_work" = "8",
    NULL = "9"))

smart_ohio_raw %>% count(EMPLOY1, employment)
```

```
# A tibble: 9 x 3
  EMPLOY1 employment      n
  <dbl> <fct>         <int>
1      1 Employed_for_wages 3119
2      2 Self-employed    466
3      3 Outofwork_1yearormore 254
4      4 Outofwork_lt1year  134
5      5 Homemaker        411
6      6 Student          190
7      7 Retired         2202
8      8 Unable_to_work    603
9      9 <NA>             33
```

**2.5.10.12 CHILDREN and its cleanup to kids**

CHILDREN, the Number of Children in Household variable, is the response to “How many children less than 18 years of age live in your household?”

- 1-87 = legitimate responses
- 88 = None
- 99 = Refused
- BLANK = Not asked or missing

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(kids = CHILDREN,
         kids = replace(kids, kids == 99, NA),
         kids = replace(kids, kids == 88, 0))

smart_ohio_raw %>% count(CHILDREN, kids) %>% tail()
```

```
# A tibble: 6 x 3
  CHILDREN kids      n
    <dbl> <dbl> <int>
1         6     6     7
2         7     7     5
3         8     8     2
4        12    12     1
5        88     0  5449
6        99    NA     43
```

**2.5.10.13 INCOME2 to incomegroup**

The available levels of INCOME2 (Income Level) are responses to: “Is your annual household income from all sources ...”

- 1 = Less than \$10,000
- 2 = \$10,000 to less than \$15,000
- 3 = \$15,000 to less than \$20,000
- 4 = \$20,000 to less than \$25,000
- 5 = \$25,000 to less than \$35,000
- 6 = \$35,000 to less than \$50,000
- 7 = \$50,000 to less than \$75,000
- 8 = \$75,000 or more
- 77 = Don’t know/Not sure
- 99 = Refused
- BLANK = Not asked or missing

We’ll just turn this into a factor, and move 77 and 99 to NA.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(incomegroup = fct_recode(factor(`INCOME2`),
    "0-9K" = "1",
    "10-14K" = "2",
    "15-19K" = "3",
    "20-24K" = "4",
    "25-34K" = "5",
    "35-49K" = "6",
    "50-74K" = "7",
    "75K+" = "8",
    NULL = "77",
    NULL = "99"))

smart_ohio_raw %>% count(`INCOME2`, incomegroup)
```

```
# A tibble: 11 x 3
  INCOME2 incomegroup    n
  <dbl> <fct>         <int>
1     1 0-9K           285
2     2 10-14K          306
3     3 15-19K          477
4     4 20-24K          589
5     5 25-34K          685
6     6 35-49K          922
7     7 50-74K          928
8     8 75K+         1910
9    77 <NA>           610
10    99 <NA>           678
11    NA <NA>            22
```

#### 2.5.10.14 INTERNET and its cleanup to internet30

INTERNET, the Internet use in the past 30 days variable, is the response to “Have you used the internet in the past 30 days?”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(internet30 = INTERNET,
    internet30 = replace(internet30, internet30 %in% c(7, 9), NA),
    internet30 = replace(internet30, internet30 == 2, 0))
```

```
smart_ohio_raw %>% count(INTERNET, internet30)
```

```
# A tibble: 5 x 3
  INTERNET internet30     n
  <dbl>      <dbl> <int>
1       1         1  6020
2       2         0  1335
3       7        NA    10
4       9        NA    10
5      NA        NA    37
```

#### 2.5.10.15 WTKG3 is weight\_kg

WTKG3 is computed by CDC, as the respondent's weight in kilograms with two implied decimal places. We calculate the actual weight in kg, with the following:

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(weight_kg = WTKG3/100)

smart_ohio_raw %>% count(WTKG3, weight_kg) %>% tail()
```

```
# A tibble: 6 x 3
  WTKG3 weight_kg     n
  <dbl>      <dbl> <int>
1 19051      191.     1
2 19278      193.     1
3 19504      195.     1
4 20412      204.     2
5 20865      209.     1
6    NA        NA    462
```

#### 2.5.10.16 HEIGHT3 is replaced with height\_m

HEIGHT3 is strangely gathered to allow people to specify their height in either feet and inches or in meters and centimeters.

- 200-711 indicates height in feet (first digit) and inches (second two digits)
- 9000 - 9998 indicates height in meters (second digit) and centimeters (last two digits)
- 7777 = Don't know/Not sure
- 9999 = Refused

Note that there is one impossible value of 575 in the data set. We'll make that an NA, and we'll also make NA any heights below 3 feet, or above 2.24 meters. Specifically, we calculate the actual height in meters, with the following:

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(height_m = case_when(
    HEIGHT3 >= 300 & HEIGHT3 <= 511 ~ round((12*floor(HEIGHT3/100) + (HEIGHT3 - 100)/12), 2),
    HEIGHT3 >= 600 & HEIGHT3 <= 711 ~ round((12*floor(HEIGHT3/100) + (HEIGHT3 - 100)/12), 2),
    HEIGHT3 >= 9000 & HEIGHT3 <= 9224 ~ ((HEIGHT3 - 9000)/100)))

smart_ohio_raw %>% count(HEIGHT3, height_m) %>% tail()
```

```
# A tibble: 6 x 3
  HEIGHT3 height_m     n
  <dbl>    <dbl> <int>
1     607      2.01     2
2     608      2.03     6
3     609      2.06     1
4    7777      NA     27
5    9999      NA     86
6      NA      NA     67
```

#### 2.5.10.17 bmi is calculated from height\_m and weight\_kg

We'll calculate body-mass index from height and weight.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(bmi = round(weight_kg/(height_m^2), 2))

smart_ohio_raw %>% count(height_m, weight_kg, bmi) # %>% tail()
```

```
# A tibble: 1,806 x 4
  height_m weight_kg  bmi     n
  <dbl>    <dbl> <dbl> <int>
1     1.35     39.0  21.4     1
2     1.35     52.2  28.6     1
3     1.4     89.8  45.8     1
4     1.42     31.8  15.8     1
5     1.42     45.4  22.5     1
6     1.42     55.8  27.7     1
7     1.42     58.5  29.0     1
8     1.42     59.9  29.7     1
9     1.42     60.8  30.1     1
10    1.42     71.2  35.3     1
# ... with 1,796 more rows
```

#### 2.5.10.18 bmgrouper is calculated from bmi

We'll then divide the respondents into adult BMI categories, in the usual way.



- BMI < 18.5 indicates underweight
- BMI from 18.5 up to 25 indicates normal weight
- BMI from 25 up to 30 indicates overweight
- BMI of 30 and higher indicates obesity

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(bmigroup = factor(cut2(as.numeric(bmi),
                                cuts = c(18.5, 25.0, 30.0))))
```

```
smart_ohio_raw %>% count(bmigroup)
```

```
# A tibble: 5 x 2
  bmigroup      n
* <fct>      <int>
1 [13.3,18.5)  119
2 [18.5,25.0) 2017
3 [25.0,30.0) 2445
4 [30.0,75.5] 2338
5 <NA>         493
```

#### 2.5.10.19 PREGNANT and its cleanup to pregnant

PREGNANT, the Pregnancy Status variable, is the response to “To your knowledge, are you now pregnant?”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing (includes SEX = male)

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(pregnant = PREGNANT,
         pregnant = replace(pregnant, pregnant %in% c(7, 9), NA),
         pregnant = replace(pregnant, pregnant == 2, 0))
```

```
smart_ohio_raw %>% count(PREGNANT, pregnant)
```

```
# A tibble: 5 x 3
  PREGNANT pregnant      n
  <dbl>      <dbl> <int>
1      1          1    41
2      2          0  1329
3      7         NA     3
4      9         NA     3
5     NA         NA  6036
```

### 2.5.10.20 DEAF and its cleanup to deaf

DEAF, the Are you deaf or do you have serious difficulty hearing variable, is the response to “Are you deaf or do you have serious difficulty hearing?”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(deaf = DEAF,
         deaf = replace(deaf, deaf %in% c(7, 9), NA),
         deaf = replace(deaf, deaf == 2, 0))

smart_ohio_raw %>% count(DEAF, deaf)
```

```
# A tibble: 5 x 3
  DEAF deaf    n
  <dbl> <dbl> <int>
1     1     1  708
2     2     0 6551
3     7    NA   15
4     9    NA    4
5    NA    NA  134
```

### 2.5.10.21 BLIND and its cleanup to blind

BLIND, the Blind or Difficulty seeing variable, is the response to “Are you blind or do you have serious difficulty seeing, even when wearing glasses?”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(blind = BLIND,
         blind = replace(blind, blind %in% c(7, 9), NA),
         blind = replace(blind, blind == 2, 0))

smart_ohio_raw %>% count(BLIND, blind)
```

```
# A tibble: 5 x 3
  BLIND blind    n
```

|   | <dbl> | <dbl> | <int> |
|---|-------|-------|-------|
| 1 | 1     | 1     | 415   |
| 2 | 2     | 0     | 6834  |
| 3 | 7     | NA    | 14    |
| 4 | 9     | NA    | 1     |
| 5 | NA    | NA    | 148   |

#### 2.5.10.22 DECIDE and its cleanup to decide

DECIDE, the Difficulty Concentrating or Remembering variable, is the response to “Because of a physical, mental, or emotional condition, do you have serious difficulty concentrating, remembering, or making decisions?”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(decide = DECIDE,
         decide = replace(decide, decide %in% c(7, 9), NA),
         decide = replace(decide, decide == 2, 0))

smart_ohio_raw %>% count(DECIDE, decide)
```

```
# A tibble: 5 x 3
  DECIDE decide     n
  <dbl>   <dbl> <int>
1     1     1     870
2     2     0    6348
3     7    NA     30
4     9    NA      2
5    NA    NA    162
```

#### 2.5.10.23 DIFFWALK and its cleanup to diffwalk

DIFFWALK, the Difficulty Walking or Climbing Stairs variable, is the response to “Do you have serious difficulty walking or climbing stairs?”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(diffwalk = DIFFWALK,
         diffwalk = replace(diffwalk, diffwalk %in% c(7, 9), NA),
         diffwalk = replace(diffwalk, diffwalk == 2, 0))

smart_ohio_raw %>% count(DIFFWALK, diffwalk)
```

```
# A tibble: 5 x 3
  DIFFWALK diffwalk     n
    <dbl>     <dbl> <int>
1       1         1  1482
2       2         0  5738
3       7        NA    19
4       9        NA     2
5      NA        NA   171
```

#### 2.5.10.24 DIFFDRES and its cleanup to diffdress

DIFFDRES, the Difficulty Dressing or Bathing variable, is the response to “Do you have difficulty dressing or bathing?”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(diffdress = DIFFDRES,
         diffdress = replace(diffdress, diffdress %in% c(7, 9), NA),
         diffdress = replace(diffdress, diffdress == 2, 0))

smart_ohio_raw %>% count(DIFFDRES, diffdress)
```

```
# A tibble: 5 x 3
  DIFFDRES diffdress     n
    <dbl>     <dbl> <int>
1       1         1   352
2       2         0 6868
3       7        NA    12
4       9        NA     1
5      NA        NA   179
```

**2.5.10.25 DIFFALON and its cleanup to diffalone**

DIFFALON, the Difficulty Doing Errands Alone variable, is the response to “Because of a physical, mental, or emotional condition, do you have difficulty doing errands alone such as visiting a doctor’s office or shopping?”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(diffalone = DIFFALON,
         diffalone = replace(diffalone, diffalone %in% c(7, 9), NA),
         diffalone = replace(diffalone, diffalone == 2, 0))

smart_ohio_raw %>% count(DIFFALON, diffalone)
```

```
# A tibble: 5 x 3
  DIFFALON diffalone      n
    <dbl>      <dbl> <int>
1       1          1   636
2       2          0  6560
3       7         NA    15
4       9         NA     4
5      NA         NA   197
```

**2.5.11 Tobacco Use (2 items)****2.5.11.1 SMOKE100 and its cleanup to smoke100**

SMOKE100, the Smoked at Least 100 Cigarettes variable, is the response to “Have you smoked at least 100 cigarettes in your entire life? [Note: 5 packs = 100 cigarettes]”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(smoke100 = SMOKE100,
         smoke100 = replace(smoke100, smoke100 %in% c(7, 9), NA),
         smoke100 = replace(smoke100, smoke100 == 2, 0))

smart_ohio_raw %>% count(SMOKE100, smoke100)
```

```
# A tibble: 5 x 3
  SMOKE100 smoke100     n
    <dbl>    <dbl> <int>
1         1         1 3294
2         2         0 3881
3         7        NA   31
4         9        NA    4
5        NA        NA  202
```

#### 2.5.11.2 `_SMOKER3` and its cleanup to `smoker`

`_SMOKER3`, is a calculated variable which categorizes subjects by their smoking status:

- 1 = Current smoker who smokes daily
- 2 = Current smoker but not every day
- 3 = Former smoker
- 4 = Never smoked
- 9 = Don't Know / Refused / Missing

We'll reclassify this as a factor with appropriate labels and NAs.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(smoker = fct_recode(factor(`_SMOKER3`),
    "Current_daily" = "1",
    "Current_not_daily" = "2",
    "Former" = "3",
    "Never" = "4",
    NULL = "9"))

smart_ohio_raw %>% count(`_SMOKER3`, smoker)
```

```
# A tibble: 5 x 3
  `_SMOKER3` smoker           n
    <dbl> <fct>           <int>
1         1 Current_daily     990
2         2 Current_not_daily  300
3         3 Former          1999
4         4 Never           3881
5         9 <NA>             242
```

## 2.5.12 E-Cigarettes (2 items)

### 2.5.12.1 ECIGARET and its cleanup to ecig\_ever

ECIGARET, the Ever used an e-cigarette variable, is the response to “Have you ever used an e-cigarette or other electronic vaping product, even just one time, in your entire life?”

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(ecig_ever = ECIGARET,
         ecig_ever = replace(ecig_ever, ecig_ever %in% c(7, 9), NA),
         ecig_ever = replace(ecig_ever, ecig_ever == 2, 0))

smart_ohio_raw %>% count(ECIGARET, ecig_ever)
```

```
# A tibble: 5 x 3
  ECIGARET ecig_ever     n
  <dbl>     <dbl> <int>
1       1         1  1354
2       2         0  5799
3       7        NA     9
4       9        NA     3
5      NA        NA    247
```

### 2.5.12.2 \_ECIGSTS and its cleanup to ecigs

\_ECIGSTS, is a calculated variable which categorizes subjects by their smoking status:

- 1 = Current and uses daily
- 2 = Current user but not every day
- 3 = Former user
- 4 = Never used e-cigarettes
- 9 = Don’t Know / Refused / Missing

We’ll reclassify this as a factor with appropriate labels and NAs.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(ecigs = fct_recode(factor(`_ECIGSTS`),
                                "Current_daily" = "1",
                                "Current_not_daily" = "2",
                                "Former" = "3",
```

```

"Never" = "4",
NULL = "9"))

smart_ohio_raw %>% count(`_ECIGSTS`, ecigs)

```

```

# A tibble: 5 x 3
  `_ECIGSTS` ecigs      n
    <dbl> <fct>    <int>
1         1 Current_daily    102
2         2 Current_not_daily 165
3         3 Former      1085
4         4 Never       5799
5         9 <NA>         261

```

### 2.5.13 Alcohol Consumption (6 items)

#### 2.5.13.1 ALCDAY5 and its cleanup to alcdays

ALCDAY5, the Days in past 30 had alcoholic beverage variable, is the response to “During the past 30 days, how many days per week or per month did you have at least one drink of any alcoholic beverage such as beer, wine, a malt beverage or liquor?”

- 101-107 = # of days per week (101 = 1 day per week, 107 = 7 days per week)
- 201-230 = # of days in past 30 days (201 = 1 day in last 30, 230 = 30 days in last 30)
- 777 = Don’t know/Not sure
- 888 = No drinks in past 30 days
- 999 = Refused
- BLANK = Not asked or Missing

We’re going to convert this to a single numeric value. Answers in days per week (in the past 7 days) will be converted (after rounding) to days in the past 30. This is a little bit of a mess, really, but we can do it.

```

smart_ohio_raw <- smart_ohio_raw %>%
  mutate(alcdays = as.numeric(ALCDAY5)) %>%
  mutate(alcdays = replace(alcdays, alcdays == 888, 0),
         alcdays = replace(alcdays, alcdays %in% c(777, 999), NA)) %>%
  mutate(alcdays = case_when(ALCDAY5 > 199 & ALCDAY5 < 231 ~ ALCDAY5 - 200,
                             ALCDAY5 > 100 & ALCDAY5 < 108 ~ round((ALCDAY5 - 100)*30),
                             TRUE ~ alcdays))

smart_ohio_raw %>% count(ALCDAY5, alcdays)

```



```
# A tibble: 39 x 3
  ALCDAY5 alcdays      n
  <dbl>    <dbl> <int>
1     101         4   263
2     102         9   197
3     103        13   142
4     104        17    76
5     105        21    53
6     106        26    18
7     107        30   114
8     201         1   621
9     202         2   448
10    203         3   233
# ... with 29 more rows
```

### 2.5.13.2 AVEDRNK2 and its cleanup to avgdrinks

AVEDRNK2, the Avg alcoholic drinks per day in past 30 variable, is the response to “One drink is equivalent to a 12-ounce beer, a 5-ounce glass of wine, or a drink with one shot of liquor. During the past 30 days, on the days when you drank, about how many drinks did you drink on the average? (A 40 ounce beer would count as 3 drinks, or a cocktail drink with 2 shots would count as 2 drinks.)”

- 1-76 = # of drinks per day
- 77 = Don’t know/Not sure
- 99 = Refused
- BLANK = Not asked or Missing (always happens when ALCDAY5 = 777, 888 or 999)

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(avgdrinks = AVEDRNK2,
         avgdrinks = replace(avgdrinks, avgdrinks > 76, NA))

smart_ohio_raw %>% count(AVEDRNK2, avgdrinks) %>% tail()
```

```
# A tibble: 6 x 3
  AVEDRNK2 avgdrinks      n
  <dbl>    <dbl> <int>
1      42         42     1
2      60         60     2
3      76         76     1
4      77         NA    46
5      99         NA     5
6      NA         NA  3876
```

### 2.5.13.3 MAXDRNKS and its cleanup to maxdrinks

MAXDRNKS, the most drinks on a single occasion in the past 30 days variable, is the response to “During the past 30 days, what is the largest number of drinks you had on any occasion?”

- 1-76 = # of drinks
- 77 = Don’t know/Not sure
- 99 = Refused
- BLANK = Not asked or Missing (always happens when ALCDAY5 = 777, 888 or 999)

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(maxdrinks = MAXDRNKS,
         maxdrinks = replace(maxdrinks, maxdrinks > 76, NA))

smart_ohio_raw %>% count(MAXDRNKS, maxdrinks) %>% tail()
```

```
# A tibble: 6 x 3
  MAXDRNKS maxdrinks      n
    <dbl>      <dbl> <int>
1      42         42      1
2      48         48      1
3      76         76      2
4      77         NA     94
5      99         NA     11
6      NA         NA    3899
```

### 2.5.13.4 \_RFBING5 and its cleanup to binge

\_RFBING5 identifies binge drinkers (males having five or more drinks on one occasion, females having four or more drinks on one occasion in the past 30 days)

The values are

- 1 = No
- 2 = Yes
- 9 = Don’t Know / Refused / Missing

People who reported no alcdays are reported here as “No,” so we’ll adjust this into an indicator variable, and create the necessary NAs.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(binge = `_RFBING5` - 1,
         binge = replace(binge, binge > 1, NA))

smart_ohio_raw %>% count(`_RFBING5`, binge)
```

```
# A tibble: 3 x 3
  ` _RFBING5` binge      n
      <dbl> <dbl> <int>
1         1     0  6035
2         2     1  1000
3         9    NA   377
```

#### 2.5.13.5 `_DRNKWEK` and its cleanup to `drinks_wk`

`_DRNKWEK` provides the computed number of alcoholic drinks per week, with two implied decimal places. The code 99900 is used for “Don’t know / Not sure / Refused / Missing” so we’ll fix that, and also divide by 100 to get an average with a decimal point.

**Note:** We’re also going to treat all results of 100 or more drinks per week as incorrect, and thus indicate them as missing data here.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(drinks_wk = `_DRNKWEK` / 100,
         drinks_wk = replace(drinks_wk, drinks_wk > 99, NA))

smart_ohio_raw %>% count(`_DRNKWEK`, drinks_wk) %>% tail(12)
```

```
# A tibble: 12 x 3
  `_DRNKWEK` drinks_wk      n
      <dbl>      <dbl> <int>
1      9333      93.3     2
2     10000      NA      1
3     10500      NA      2
4     11667      NA      1
5     14000      NA      2
6     16800      NA      2
7     17500      NA      1
8     18200      NA      1
9     28000      NA      1
10    29400      NA      1
11    53200      NA      1
12    99900      NA     379
```

#### 2.5.13.6 `_RFDRHV5` and its cleanup to `drink_heavy`

`_RFDRHV5` identifies heavy drinkers (males having 14 or more drinks per week, females having 7 or more drinks per week)

The values are

- 1 = No
- 2 = Yes
- 9 = Don't Know / Refused / Missing

People who reported no `alcdays` are reported here as “No,” so we’ll adjust this into an indicator variable, and create the necessary NAs.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(drink_heavy = `_RFDRHV5` - 1,
         drink_heavy = replace(drink_heavy, drink_heavy > 1, NA))

smart_ohio_raw %>% count(`_RFDRHV5`, drink_heavy)
```

```
# A tibble: 3 x 3
  `_RFDRHV5` drink_heavy    n
    <dbl>         <dbl> <int>
1         1             0  6607
2         2             1   426
3         9            NA   379
```

## 2.5.14 Fruits and Vegetables (8 items)

### 2.5.14.1 `\_FRUTSU1` and its cleanup to `fruit_day`

`_FRUTSU1` provides the computed number of fruit servings consumed per day, with two implied decimal places. We’ll divide by 100 to insert the decimal point.

**Note:** We’re also going to treat all results exceeding 16 servings per day as implausible, and thus indicate them as missing data here, following some CDC procedures.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(fruit_day = `_FRUTSU1` / 100,
         fruit_day = replace(fruit_day, fruit_day > 16, NA))

smart_ohio_raw %>% count(`_FRUTSU1`, fruit_day) %>% tail()
```

```
# A tibble: 6 x 3
  `_FRUTSU1` fruit_day    n
    <dbl>         <dbl> <int>
1      913         9.13     1
2     1000         10      4
3     1400         14      1
4     3000         NA       1
5     7600         NA       1
6        NA         NA     555
```

#### 2.5.14.2 `_VEGESU1` and its cleanup to `veg_day`

`_VEGESU1` provides the computed number of vegetable servings consumed per day, with two implied decimal places. We'll divide by 100 to insert the decimal point.

**Note:** We're also going to treat all results exceeding 23 servings per day as implausible, and thus indicate them as missing data here, following some CDC procedures.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(veg_day = `_VEGESU1` / 100,
         veg_day = replace(veg_day, veg_day > 23, NA))

smart_ohio_raw %>% count(`_VEGESU1`, veg_day) %>% tail()
```

```
# A tibble: 6 x 3
  `_VEGESU1` veg_day      n
    <dbl>    <dbl> <int>
1      1414      14.1     1
2      1603      16.0     1
3      1891      18.9     1
4      2167      21.7     1
5      3150      NA      1
6         NA      NA    666
```

#### 2.5.14.3 `FTJUDA2_` and its cleanup to `eat_juice`

`FTJUDA2_` provides the servings of fruit juice consumed per day, with two implied decimal places. We'll divide by 100 to insert the decimal point.

**Note:** We're also going to treat all results exceeding 16 servings per day as implausible, and thus indicate them as missing data here.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(eat_juice = `FTJUDA2_` / 100,
         eat_juice = replace(eat_juice, eat_juice > 16, NA))

smart_ohio_raw %>% count(`FTJUDA2_`, eat_juice) %>% tail()
```

```
# A tibble: 6 x 3
  FTJUDA2_ eat_juice      n
    <dbl>    <dbl> <int>
1      500         5     6
2      600         6     1
3      700         7     1
4     1200        12     1
```

|   |      |    |     |
|---|------|----|-----|
| 5 | 7500 | NA | 1   |
| 6 | NA   | NA | 469 |

#### 2.5.14.4 FRUTDA2\_ and its cleanup to eat\_fruit

FRUTDA2\_ provides the servings of fruit consumed per day, with two implied decimal places. We'll divide by 100 to insert the decimal point.

**Note:** We're also going to treat all results exceeding 16 servings per day as implausible, and thus indicate them as missing data here.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(eat_fruit = `FRUTDA2_` / 100,
         eat_fruit = replace(eat_fruit, eat_fruit > 16, NA))

smart_ohio_raw %>% count(`FRUTDA2_`, eat_fruit) %>% tail()
```

```
# A tibble: 6 x 3
  FRUTDA2_ eat_fruit      n
    <dbl>     <dbl> <int>
1     700         7     5
2     800         8     3
3     900         9     1
4    1000        10     1
5    3000        NA     1
6      NA        NA    456
```

#### 2.5.14.5 GREENDA1\_ and its cleanup to eat\_greenveg

GREENDA1\_ provides the servings of dark green vegetables consumed per day, with two implied decimal places. We'll divide by 100 to insert the decimal point.

**Note:** We're also going to treat all results exceeding 16 servings per day as implausible, and thus indicate them as missing data here.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(eat_greenveg = `GREENDA1_` / 100,
         eat_greenveg = replace(eat_greenveg, eat_greenveg > 16, NA))

smart_ohio_raw %>% count(`GREENDA1_`, eat_greenveg) %>% tail()
```

```
# A tibble: 6 x 3
  GREENDA1_ eat_greenveg      n
    <dbl>     <dbl> <int>
1     700         7     4
2     786        7.86     1
3     800         8     2
```

|   |      |    |     |
|---|------|----|-----|
| 4 | 2000 | NA | 1   |
| 5 | 3000 | NA | 1   |
| 6 | NA   | NA | 447 |

#### 2.5.14.6 FRNCHDA\_ and its cleanup to eat\_fries

FRNCHDA\_ provides the servings of french fries consumed per day, with two implied decimal places. We'll divide by 100 to insert the decimal point.

**Note:** We're also going to treat all results exceeding 16 servings per day as implausible, and thus indicate them as missing data here.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(eat_fries = `FRNCHDA_` / 100,
         eat_fries = replace(eat_fries, eat_fries > 16, NA))

smart_ohio_raw %>% count(`FRNCHDA_`, eat_fries) %>% tail()
```

```
# A tibble: 6 x 3
  FRNCHDA_ eat_fries      n
  <dbl>     <dbl> <int>
1     300         3       9
2     314        3.14       1
3     400         4       3
4     500         5       1
5     700         7       1
6      NA        NA     453
```

#### 2.5.14.7 POTADA1\_ and its cleanup to eat\_potato

POTADA1\_ provides the servings of potatoes consumed per day, with two implied decimal places. We'll divide by 100 to insert the decimal point.

**Note:** We're also going to treat all results exceeding 16 servings per day as implausible, and thus indicate them as missing data here.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(eat_potato = `POTADA1_` / 100,
         eat_potato = replace(eat_potato, eat_potato > 16, NA))

smart_ohio_raw %>% count(`POTADA1_`, eat_potato) %>% tail()
```

```
# A tibble: 6 x 3
  POTADA1_ eat_potato      n
  <dbl>     <dbl> <int>
1     314        3.14       1
2     329        3.29       1
```

|   |     |      |     |
|---|-----|------|-----|
| 3 | 400 | 4    | 3   |
| 4 | 471 | 4.71 | 1   |
| 5 | 700 | 7    | 1   |
| 6 | NA  | NA   | 501 |

#### 2.5.14.8 VEGEDA2\_ and its cleanup to eat\_otherveg

VEGEDA2\_ provides the servings of other vegetables consumed per day, with two implied decimal places. We'll divide by 100 to insert the decimal point.

**Note:** We're also going to treat all results exceeding 16 servings per day as implausible, and thus indicate them as missing data here.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(eat_otherveg = `VEGEDA2_` / 100,
         eat_otherveg = replace(eat_otherveg, eat_otherveg > 16, NA))

smart_ohio_raw %>% count(`VEGEDA2_`, eat_otherveg) %>% tail()
```

```
# A tibble: 6 x 3
  VEGEDA2_ eat_otherveg     n
  <dbl>      <dbl> <int>
1     600          6      3
2     700          7     11
3     800          8      1
4    1000         10      2
5    1100         11      1
6      NA         NA     509
```

### 2.5.15 Exercise and Physical Activity (8 items)

#### 2.5.15.1 \_TOTINDA and its cleanup to exerany

\_TOTINDA, the Exercise in Past 30 Days variable, is the response to “During the past month, other than your regular job, did you participate in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise?”

- 1 = Yes
- 2 = No
- 7 = Don't know/Not sure
- 9 = Refused
- BLANK = Not asked or missing

This is just like HLTHPLAN.



```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(exerany = `_TOTINDA`,
         exerany = replace(exerany, exerany %in% c(7, 9), NA),
         exerany = replace(exerany, exerany == 2, 0))

smart_ohio_raw %>% count(`_TOTINDA`, exerany)

# A tibble: 3 x 3
#   `_TOTINDA` exerany      n
#   <dbl>    <dbl> <int>
1         1        1  4828
2         2        0  2137
3         9       NA   447
```

### 2.5.15.2 `\_PACAT1` and its cleanup to activity

`\_PACAT1` contains physical activity categories, estimated from responses to the BRFSS. The categories are:

- 1 = Highly Active
- 2 = Active
- 3 = Insufficiently Active
- 4 = Inactive
- 9 = Don't Know / Not Sure / Refused / Missing

So we'll create a factor.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(activity = factor(`_PACAT1`,
                          activity = fct_recode(activity,
                                                  "Highly_Active" = "1",
                                                  "Active" = "2",
                                                  "Insufficiently_Active" = "3",
                                                  "Inactive" = "4",
                                                  NULL = "9"))

smart_ohio_raw %>% count(`_PACAT1`, activity)

# A tibble: 5 x 3
#   `_PACAT1` activity      n
#   <dbl>    <fct>    <int>
1         1 1 Highly_Active  2053
2         2 2 Active       1132
3         3 3 Insufficiently_Active 1293
4         4 4 Inactive     2211
5         9 9 <NA>         723
```

### 2.5.15.3 `_PAINDX1` and its cleanup to `rec_aerobic`

`_PAINDX1` indicates whether the respondent's stated levels of physical activity meet recommendations for aerobic activity. The responses are:

- 1 = Yes
- 2 = No
- 9 = Don't know/Not sure/Refused/Missing

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(rec_aerobic = 2 - `_PAINDX1`,
         rec_aerobic = replace(rec_aerobic, rec_aerobic < 0, NA))

smart_ohio_raw %>% count(`_PAINDX1`, rec_aerobic)
```

```
# A tibble: 3 x 3
  `_PAINDX1` rec_aerobic    n
    <dbl>         <dbl> <int>
1         1             1 3228
2         2             0 3504
3         9            NA  680
```

### 2.5.15.4 `_PASTRNG` and its cleanup to `rec_strength`

`_PASTRNG` indicates whether the respondent's stated levels of physical activity meet recommendations for strength-building activity. The responses are:

- 1 = Yes
- 2 = No
- 9 = Don't know/Not sure/Refused/Missing

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(rec_strength = 2 - `_PASTRNG`,
         rec_strength = replace(rec_strength, rec_strength < 0, NA))

smart_ohio_raw %>% count(`_PASTRNG`, rec_strength)
```

```
# A tibble: 3 x 3
  `_PASTRNG` rec_strength    n
    <dbl>         <dbl> <int>
1         1             1 1852
2         2             0 5004
3         9            NA  556
```

**2.5.15.5 EXTRACT11 and its cleanup to exer1\_type**

Respondents are asked “What type of physical activity or exercise did you spend the most time doing during the past month?” and these responses are gathered into a set of 76 named categories, including an “other” category. Codes 77 (Don’t Know / Not Sure) and 99 (Refused) are dropped into NA in my code below, and Code 98 (“Other type of activity”) remains. Then I went through the tedious work of converting the factor levels from numbers to names, following the value labels provided by BRFSS.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(exer1_type = factor(EXTRACT11),
         exer1_type = fct_recode(
           exer1_type,
           "Active Gaming Devices" = "1",
           "Aerobics video or class" = "2",
           "Backpacking" = "3",
           "Badminton" = "4",
           "Basketball" = "5",
           "Bicycling machine" = "6",
           "Bicycling" = "7",
           "Boating" = "8",
           "Bowling" = "9",
           "Boxing" = "10",
           "Calisthenics" = "11",
           "Canoeing" = "12",
           "Carpentry" = "13",
           "Dancing" = "14",
           "Elliptical machine" = "15",
           "Fishing" = "16",
           "Frisbee" = "17",
           "Gardening" = "18",
           "Golf with cart" = "19",
           "Golf without cart" = "20",
           "Handball" = "21",
           "Hiking" = "22",
           "Hockey" = "23",
           "Horseback riding" = "24",
           "Hunting large game" = "25",
           "Hunting small game" = "26",
           "Inline skating" = "27",
           "Jogging" = "28",
           "Lacrosse" = "29",
           "Mountain climbing" = "30",
           "Mowing lawn" = "31",
           "Paddleball" = "32",
```

```
"Painting house" = "33",  
"Pilates" = "34",  
"Racquetball" = "35",  
"Raking lawn" = "36",  
"Running" = "37",  
"Rock climbing" = "38",  
"Rope skipping" = "39",  
"Rowing machine" = "40",  
"Rugby" = "41",  
"Scuba diving" = "42",  
"Skateboarding" = "43",  
"Skating" = "44",  
"Sledding" = "45",  
"Snorkeling" = "46",  
"Snow blowing" = "47",  
"Snow shoveling" = "48",  
"Snow skiing" = "49",  
"Snowshoeing" = "50",  
"Soccer" = "51",  
"Softball/Baseball" = "52",  
"Squash" = "53",  
"Stair Climbing" = "54",  
"Stream fishing" = "55",  
"Surfing" = "56",  
"Swimming" = "57",  
"Swimming in laps" = "58",  
"Table tennis" = "59",  
"Tai Chi" = "60",  
"Tennis" = "61",  
"Touch football" = "62",  
"Volleyball" = "63",  
"Walking" = "64",  
"Waterskiing" = "66",  
"Weight lifting" = "67",  
"Wrestling" = "68",  
"Yoga" = "69",  
"Child Care" = "71",  
"Farm Work" = "72",  
"Household Activities" = "73",  
"Martial Arts" = "74",  
"Upper Body Cycle" = "75",  
"Yard Work" = "76",  
"Other Activities" = "98",  
NULL = "77",  
NULL = "99")
```

```
)
```

```
Warning: Problem with `mutate()` input `exer1_type`.
i Unknown levels in `f`: 3, 17, 21, 32, 36, 41, 42, 45, 47, 53, 55, 56, 59
i Input `exer1_type` is `fct_recode(...)`.
```

The warning generated here is caused by the fact that some of the available types of exercise were not mentioned by people in our sample. Looking at the last few results, we can see how many people fell into several categories.

```
smart_ohio_raw %>% count(EXTRACT11, exer1_type) %>% tail()
```

```
# A tibble: 6 x 3
  EXTRACT11 exer1_type      n
  <dbl> <fct>      <int>
1      75 Upper Body Cycle    6
2      76 Yard Work          78
3      77 <NA>              10
4      98 Other Activities   276
5      99 <NA>               4
6      NA <NA>            2588
```

The most common activities are:

```
smart_ohio_raw %>% count(exer1_type, sort = TRUE) %>% head(10)
```

```
# A tibble: 10 x 2
  exer1_type      n
  <fct>      <int>
1 Walking    2605
2 <NA>       2602
3 Running    324
4 Other Activities 276
5 Gardening  242
6 Weight lifting 189
7 Aerobics video or class 103
8 Bicycling machine 103
9 Bicycling    96
10 Golf with cart 90
```

#### 2.5.15.6 EXTRACT21 and its cleanup to exer2\_type

As a follow-up, respondents are asked “What other type of physical activity gave you the next most exercise during the past month?” and these responses are also gathered into the same set of 76 named categories, including an “other” category, but now also adding a “No Other Activity” category (code 88). Codes 77 (Don’t Know / Not Sure) and 99 (Refused) are dropped into NA in my code below, and

Code 98 (“Other type of activity”) remains. Then I went through the tedious work of converting the factor levels from numbers to names, following the value labels provided by BRFSS. I’m sure there’s a better way to do this.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(exer2_type = factor(EXTRACT21),
         exer2_type = fct_recode(
           exer2_type,
           "Active Gaming Devices" = "1",
           "Aerobics video or class" = "2",
           "Backpacking" = "3",
           "Badminton" = "4",
           "Basketball" = "5",
           "Bicycling machine" = "6",
           "Bicycling" = "7",
           "Boating" = "8",
           "Bowling" = "9",
           "Boxing" = "10",
           "Calisthenics" = "11",
           "Canoeing" = "12",
           "Carpentry" = "13",
           "Dancing" = "14",
           "Elliptical machine" = "15",
           "Fishing" = "16",
           "Frisbee" = "17",
           "Gardening" = "18",
           "Golf with cart" = "19",
           "Golf without cart" = "20",
           "Handball" = "21",
           "Hiking" = "22",
           "Hockey" = "23",
           "Horseback riding" = "24",
           "Hunting large game" = "25",
           "Hunting small game" = "26",
           "Inline skating" = "27",
           "Jogging" = "28",
           "Lacrosse" = "29",
           "Mountain climbing" = "30",
           "Mowing lawn" = "31",
           "Paddleball" = "32",
           "Painting house" = "33",
           "Pilates" = "34",
           "Racquetball" = "35",
           "Raking lawn" = "36",
           "Running" = "37",
           "Rock climbing" = "38",
```

```

    "Rope skipping" = "39",
    "Rowing machine" = "40",
    "Rugby" = "41",
    "Scuba diving" = "42",
    "Skateboarding" = "43",
    "Skating" = "44",
    "Sledding" = "45",
    "Snorkeling" = "46",
    "Snow blowing" = "47",
    "Snow shoveling" = "48",
    "Snow skiing" = "49",
    "Snowshoeing" = "50",
    "Soccer" = "51",
    "Softball/Baseball" = "52",
    "Squash" = "53",
    "Stair Climbing" = "54",
    "Stream fishing" = "55",
    "Surfing" = "56",
    "Swimming" = "57",
    "Swimming in laps" = "58",
    "Table tennis" = "59",
    "Tai Chi" = "60",
    "Tennis" = "61",
    "Touch football" = "62",
    "Volleyball" = "63",
    "Walking" = "64",
    "Waterskiing" = "66",
    "Weight lifting" = "67",
    "Wrestling" = "68",
    "Yoga" = "69",
    "Child Care" = "71",
    "Farm Work" = "72",
    "Household Activities" = "73",
    "Martial Arts" = "74",
    "Upper Body Cycle" = "75",
    "Yard Work" = "76",
    "No Other Activity" = "88",
    "Other Activities" = "98",
    NULL = "77",
    NULL = "99")
)

```

```

Warning: Problem with `mutate()` input `exer2_type`.
i Unknown levels in `f`: 3, 21, 30, 39, 41, 46, 50, 62
i Input `exer2_type` is `fct_recode(...)`

```

```
smart_ohio_raw %>% count(EXTRACT21, exer2_type) %>% tail()
```

```
# A tibble: 6 x 3
  EXTRACT21 exer2_type      n
    <dbl> <fct>      <int>
1         76 Yard Work      153
2         77 <NA>           26
3         88 No Other Activity 1854
4         98 Other Activities   246
5         99 <NA>           19
6         NA <NA>          2627
```

The most common activity types in this group are:

```
smart_ohio_raw %>% count(exer2_type, sort = TRUE) %>% head(10)
```

```
# A tibble: 10 x 2
  exer2_type      n
    <fct>      <int>
1 <NA>          2672
2 No Other Activity 1854
3 Walking         629
4 Weight lifting   272
5 Other Activities  246
6 Gardening        202
7 Household Activities 169
8 Yard Work        153
9 Running          148
10 Bicycling        118
```

#### 2.5.15.7 `_MINAC11` and its cleanup to `exer1_min`

`_MINAC11` is minutes of physical activity per week for the first activity (listed as `exer1_type` above.) Since there are only about 10,080 minutes in a typical week, we'll treat as implausible any values larger than 4200 minutes (which would indicate 70 hours per week.)

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(exer1_min = `_MINAC11`,
         exer1_min = replace(exer1_min, exer1_min > 4200, NA))

smart_ohio_raw %>% count(`_MINAC11`, exer1_min) %>% tail()
```

```
# A tibble: 6 x 3
  `_MINAC11` exer1_min      n
    <dbl>      <dbl> <int>
1      3780      3780      8
```



|   |       |      |      |
|---|-------|------|------|
| 2 | 3959  | 3959 | 1    |
| 3 | 3960  | 3960 | 1    |
| 4 | 4193  | 4193 | 6    |
| 5 | 27000 | NA   | 1    |
| 6 | NA    | NA   | 2760 |

#### 2.5.15.8 `_MINAC21` and its cleanup to `exer2_min`

`_MINAC21` is minutes of physical activity per week for the second activity (listed as `exer2_type` above.) Again, we'll treat as implausible any values larger than 4200 minutes (which would indicate 70 hours per week.)

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(exer2_min = `_MINAC21`,
         exer2_min = replace(exer2_min, exer2_min > 4200, NA))

smart_ohio_raw %>% count(`_MINAC21`, exer2_min) %>% tail()
```

```
# A tibble: 6 x 3
  `_MINAC21` exer2_min      n
    <dbl>      <dbl> <int>
1     3360      3360      3
2     3780      3780      7
3     4193      4193      3
4     6120         NA      1
5     8400         NA      1
6         NA         NA    2770
```

### 2.5.16 Seatbelt Use (1 item)

#### 2.5.16.1 SEATBELT and its cleanup to `seatbelt`

This question asks “How often do you use seat belts when you drive or ride in a car?” Possible responses are:

- 1 = Always
- 2 = Nearly always
- 3 = Sometimes
- 4 = Seldom
- 5 = Never
- 7 = Don't know / Not sure
- 8 = Never drive or ride in a car
- 9 = Refused

We'll treat codes 7, 8 and 9 as NA, and turn this into a factor.

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(seatbelt = fct_recode(factor(SEATBELT),
                                "Always" = "1",
                                "Nearly_always" = "2",
                                "Sometimes" = "3",
                                "Seldom" = "4",
                                "Never" = "5",
                                NULL = "7",
                                NULL = "8",
                                NULL = "9"))

smart_ohio_raw %>% count(SEATBELT, seatbelt)

# A tibble: 9 x 3
  SEATBELT seatbelt      n
  <dbl> <fct>      <int>
1      1 Always      6047
2      2 Nearly_always 409
3      3 Sometimes     191
4      4 Seldom        81
5      5 Never        148
6      7 <NA>          7
7      8 <NA>         21
8      9 <NA>          2
9     NA <NA>        506
```

## 2.5.17 Immunization (3 items)

### 2.5.17.1 FLUSHOT6 and its cleanup to vax\_flu

FLUSHOT6 gives the response to “During the past 12 months, have you had either a flu shot or a flu vaccine that was sprayed in your nose?” The responses are:

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(vax_flu = 2 - FLUSHOT6,
         vax_flu = replace(vax_flu, vax_flu < 0, NA))

smart_ohio_raw %>% count(FLUSHOT6, vax_flu)

# A tibble: 5 x 3
  FLUSHOT6 vax_flu      n
```

|   | <dbl> | <dbl> | <int> |
|---|-------|-------|-------|
| 1 | 1     | 1     | 3453  |
| 2 | 2     | 0     | 3410  |
| 3 | 7     | NA    | 26    |
| 4 | 9     | NA    | 3     |
| 5 | NA    | NA    | 520   |

### 2.5.17.2 PNEUVAC3 and its cleanup to vax\_pneumo

PNEUVAC3 gives the response to “A pneumonia shot or pneumococcal vaccine is usually given only once or twice in a person’s lifetime and is different from the flu shot. Have you ever had a pneumonia shot?” The responses are:

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(vax_pneumo = 2 - PNEUVAC3,
         vax_pneumo = replace(vax_pneumo, vax_pneumo < 0, NA))

smart_ohio_raw %>% count(PNEUVAC3, vax_pneumo)
```

```
# A tibble: 5 x 3
  PNEUVAC3 vax_pneumo     n
  <dbl>      <dbl> <int>
1       1         1  3112
2       2         0  3262
3       7        NA   509
4       9        NA     3
5      NA        NA   526
```

### 2.5.17.3 SHINGLE2 and its cleanup to vax\_shingles

SHINGLE2 gives the response to “Have you ever had the shingles or zoster vaccine?” The responses are:

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(vax_shingles = 2 - SHINGLE2,
         vax_shingles = replace(vax_shingles, vax_shingles < 0, NA))
```

```
smart_ohio_raw %>% count(SHINGLE2, vax_shingles)
```

```
# A tibble: 4 x 3
  SHINGLE2 vax_shingles     n
    <dbl>      <dbl> <int>
1       1          1  1503
2       2          0  2979
3       7         NA    78
4      NA         NA  2852
```

## 2.5.18 HIV/AIDS (2 items)

### 2.5.18.1 HIVTST6 and its cleanup to hiv\_test

HIVTST6 gives the response to “Have you ever been tested for HIV? Do not count tests you may have had as part of a blood donation. Include testing fluid from your mouth.” The responses are:

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(hiv_test = 2 - HIVTST6,
         hiv_test = replace(hiv_test, hiv_test < 0, NA))

smart_ohio_raw %>% count(HIVTST6, hiv_test)
```

```
# A tibble: 5 x 3
  HIVTST6 hiv_test     n
    <dbl>      <dbl> <int>
1       1          1  2017
2       2          0  4565
3       7         NA   260
4       9         NA   14
5      NA         NA  556
```

### 2.5.18.2 HIVRISK5 and its cleanup to hiv\_risk

HIVRISK5 gives the response to “I am going to read you a list. When I am done, please tell me if any of the situations apply to you. You do not need to tell me which one. You have injected any drug other than those prescribed for you in the past year. You have been treated for a sexually transmitted disease or STD

## 2.6. IMPUTING AGE AND INCOME AS QUANTITATIVE FROM THIN AIR101

in the past year. You have given or received money or drugs in exchange for sex in the past year.” The responses are:

- 1 = Yes
- 2 = No
- 7 = Don’t know/Not sure
- 9 = Refused

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(hiv_risk = 2 - HIVRISK5,
         hiv_risk = replace(hiv_risk, hiv_risk < 0, NA))

smart_ohio_raw %>% count(HIVRISK5, hiv_risk)
```

```
# A tibble: 5 x 3
  HIVRISK5 hiv_risk      n
  <dbl>     <dbl> <int>
1       1         1   277
2       2         0  6537
3       7        NA     2
4       9        NA    17
5      NA        NA   579
```

## 2.6 Imputing Age and Income as Quantitative from Thin Air

This section is purely for teaching purposes. I would never use the variables created in this section for research work.

### 2.6.1 age\_imp: Imputing Age Data

I want a quantitative age variable, so I’m going to create an imputed `age_imp` value for each subject based on their `agegroup`. For each age group, I will assume that each of the ages represented by a value in that age group will be equally likely, and will draw from the relevant uniform distribution to impute age.

```
set.seed(2020432002)

smart_ohio_raw <- smart_ohio_raw %>%
  mutate(age_low = as.numeric(str_sub(as.character(agegroup), 1, 2))) %>%
  mutate(age_high = as.numeric(str_sub(as.character(agegroup), 4, 5))) %>%
  rowwise() %>%
  mutate(age_imp = ifelse(!is.na(agegroup),
                          round(runif(1, min = age_low, max = age_high), 0),
```

```

NA))

smart_ohio_raw %>% count(agegroup, age_imp) %>% tail()

```

```

# A tibble: 80 x 3
# Rowwise:
  agegroup age_imp     n
  <fct>      <dbl> <int>
1 18-24      18     46
2 18-24      19     75
3 18-24      20     76
4 18-24      21     82
5 18-24      22     80
6 18-24      23     54
7 18-24      24     35
8 25-29      25     42
9 25-29      26     93
10 25-29      27     77
# ... with 70 more rows

```


Here is a histogram of the `age_imp` variable.

```

ggplot(smart_ohio_raw, aes(x = age_imp)) +
  geom_histogram(fill = "navy", col = "white",
                 binwidth = 1) +
  scale_x_continuous(breaks = c(18, 25, 35, 45, 55, 65, 75, 85, 96)) +
  labs(x = "Imputed Age in Years",
       title = paste0("Imputed Income: ",
                      sum(is.na(smart_ohio_raw$age_imp)),
                      " respondents have missing age group"))

```

## 2.6. IMPUTING AGE AND INCOME AS QUANTITATIVE FROM THIN AIR103



### 2.6.2 inc\_imp: Imputing Income Data

I want a quantitative income variable, so I'm going to create an imputed `inc_imp` value for each subject based on their `incomegroup`. For most income groups, I will assume that each of the incomes represented by a value in that income group will be equally likely, and will draw from the relevant uniform distribution to impute income. The exception is the highest income group, where I will impute a value drawn from a distribution that places all values at \$75,000 or more, but has a substantial right skew and long tail.

```
set.seed(2020432001)
```

```
smart_ohio_raw <- smart_ohio_raw %>%
  mutate(inc_imp = case_when(
    incomegroup == "0-9K" ~ round(runif(1, min = 100, max = 9999)),
    incomegroup == "10-14K" ~ round(runif(1, min = 10000, max = 14999)),
    incomegroup == "15-19K" ~ round(runif(1, min = 15000, max = 19999)),
    incomegroup == "20-24K" ~ round(runif(1, min = 20000, max = 24999)),
    incomegroup == "25-34K" ~ round(runif(1, min = 25000, max = 34999)),
    incomegroup == "35-49K" ~ round(runif(1, min = 35000, max = 49999)),
    incomegroup == "50-74K" ~ round(runif(1, min = 50000, max = 74999)),
    incomegroup == "75K+" ~ round((rnorm(n = 1, mean = 0, sd = 300)^2) + 74999)))

smart_ohio_raw %>% count(incomegroup, inc_imp) %>% tail()
```

```
# A tibble: 6 x 3
# Rowwise:
  incomegroup inc_imp      n
  <fct>        <dbl> <int>
1 75K+         774009      1
2 75K+         798174      1
3 75K+         806161      1
4 75K+         847758      1
5 75K+        1085111      1
6 <NA>          NA     1310
```

Here are density plots of the `inc_imp` variable. The top picture shows the results on a linear scale, and the bottom shows them on a log (base 10) scale.


```
p1 <- ggplot(smart_ohio_raw, aes(x = inc_imp/1000)) +
  geom_density(fill = "darkgreen", col = "white") +
  labs(x = "Imputed Income in Thousands of Dollars",
       title = "Imputed Income on the Linear scale") +
  scale_x_continuous(breaks = c(25, 75, 250, 1000))

p2 <- ggplot(smart_ohio_raw, aes(x = inc_imp/1000)) +
  geom_density(fill = "darkgreen", col = "white") +
  labs(x = "Imputed Income in Thousands of Dollars",
       title = "Imputed Income on the Log (base 10) scale") +
  scale_x_log10(breaks = c(0.1, 1, 5, 25, 75, 250, 1000))

p1 / p2 +
  plot_annotation(title =
    paste0("Imputed Income: ", sum(is.na(smart_ohio_raw$inc_imp))),
```



Imputed Income: 1310 respondents have missing income group



## 2.7 Clean Data in the State of Ohio

There are six MMSAs associated with the state of Ohio. We're going to create a `smart_ohio` that includes each of them. First, I'll ungroup the data that I created earlier, so I get a clean tibble.

```
smart_ohio_raw <- smart_ohio_raw %>% ungroup()
```

Next, I'll select the variables I want to retain (they are the ones I created, plus `SEQNO`.)

```
smart_ohio <- smart_ohio_raw %>%
  select(SEQNO, mmsa, mmsa_code, mmsa_name, mmsa_wt, completed,
         landline, hhadults,
         genhealth, physhealth, menthealth, poorhealth,
         agegroup, age_imp, race, hispanic, race_eth,
         female, marital, kids, educgroup, home_own,
         veteran, employment, incomegroup, inc_imp,
         cell_own, internet30,
         weight_kg, height_m, bmi, bmigroup,
         pregnant, deaf, blind, decide,
         diffwalk, diffdress, diffalone,
         smoke100, smoker, ecig_ever, ecigs,
         healthplan, hasdoc, costprob, t_checkup,
```

```

    bp_high, bp_meds,
    t_chol, chol_high, chol_meds,
    asthma, hx_asthma, now_asthma,
    hx_mi, hx_chd, hx_stroke, hx_skinc, hx_otherc,
    hx_copd, hx_depress, hx_kidney,
    hx_diabetes, dm_status, dm_age,
    hx_arthr, arth_lims, arth_work, arth_soc,
    joint_pain, alcdays, avgdrinks, maxdrinks,
    binge, drinks_wk, drink_heavy,
    fruit_day, veg_day, eat_juice, eat_fruit,
    eat_greenveg, eat_fries, eat_potato,
    eat_otherveg, exerany, activity, rec_aerobic,
    rec_strength, exer1_type, exer2_type,
    exer1_min, exer2_min, seatbelt,
    vax_flu, vax_pneumo, vax_shingles,
    hiv_test, hiv_risk)

saveRDS(smart_ohio, "data/smart_ohio.Rds")

write_csv(smart_ohio, "data/smart_ohio.csv")

```

The `smart_ohio` file should contain 99 variables, describing 7412 respondents.

## 2.8 Clean Cleveland-Elyria Data

### 2.8.1 Cleveland - Elyria Data

The `mmsa_name` variable is probably the simplest way for us to filter our data down to the MMSA we are interested in. Here, I'm using the `str_detect` function to identify the values of `mmsa_name` that contain the text "Cleveland."

```

smart_cle <- smart_ohio %>%
  filter(str_detect(mmsa_name, 'Cleveland'))

saveRDS(smart_cle, "data/smart_cle.Rds")

```

In the Cleveland-Elyria MSA, we have 1133 observations on the same 99 variables.

We'll build a variety of smaller subsets from these data, eventually.

Barnett, Peggy A., Simon Roman-Golstein, Fred Ramsey, and others. 1995. "Differential Permeability and Quantitative MR Imaging of a Human Lung Carcinoma Brain Xenograft in the Nude Rat." *American Journal of Pathology* 146(2): 436–49. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1869863/>.

- Berkhemer, Olvert A., Puck S. S. Fransen, Debbie Buemer, and others. 2015. "A Randomized Trial of Intraarterial Treatment for Acute Ischemic Stroke." *New England Journal of Medicine* 372: 11–20. <http://www.nejm.org/doi/full/10.1056/NEJMoa1411587>.
- Ramsey, Fred L., and Daniel W. Schafer. 2002. *The Statistical Sleuth: A Course in Methods of Data Analysis*. Second Edition. Pacific Grove, CA: Duxbury.
- Rosenbaum, Paul R. 2017. *Observation and Experiment: An Introduction to Causal Inference*. Cambridge, MA: Harvard University Press.
- Roy, Denis, Mario Talajic, Stanley Nattel, and others. 2008. "Rhythm Control Versus Rate Control for Atrial Fibrillation and Heart Failure." *New England Journal of Medicine* 358: 2667–77. <http://www.nejm.org/doi/full/10.1056/NEJMoa0708789>.
- Tolaney, Sara M, William T. Barry, T. Dang Chau, and others. 2015. "Adjuvant Paclitaxel and Trastuzumab for Node-Negative, Her2-Positive Breast Cancer." *New England Journal of Medicine* 372: 134–41. <http://www.nejm.org/doi/full/10.1056/NEJMoa1406281>.