

# Maximum time-dependent space-charge limited diode currents

Cite as: Phys. Plasmas **23**, 014502 (2016); <https://doi.org/10.1063/1.4939607>

Submitted: 06 October 2015 • Accepted: 23 December 2015 • Published Online: 08 January 2016

M. E. Griswold and  N. J. Fisch



View Online



Export Citation



CrossMark

## ARTICLES YOU MAY BE INTERESTED IN

[100 years of the physics of diodes](#)

Applied Physics Reviews **4**, 011304 (2017); <https://doi.org/10.1063/1.4978231>

[An upper bound to time-averaged space-charge limited diode currents](#)

Physics of Plasmas **17**, 114503 (2010); <https://doi.org/10.1063/1.3503661>

[A coordinate system invariant formulation for space-charge limited current in vacuum](#)

Applied Physics Letters **115**, 054101 (2019); <https://doi.org/10.1063/1.5115261>

Physics of Plasmas

Papers from 62nd Annual Meeting of the  
APS Division of Plasma Physics

Read now!

# Maximum time-dependent space-charge limited diode currents

M. E. Griswold<sup>1</sup> and N. J. Fisch<sup>2</sup>

<sup>1</sup>Tri Alpha Energy, Inc., Rancho Santa Margarita, California 92688, USA

<sup>2</sup>Princeton Plasma Physics Laboratory, Princeton University, Princeton, New Jersey 08543, USA

(Received 6 October 2015; accepted 23 December 2015; published online 8 January 2016)

Recent papers claim that a one dimensional (1D) diode with a time-varying voltage drop can transmit current densities that exceed the Child-Langmuir (CL) limit on average, apparently contradicting a previous conjecture that there is a hard limit on the average current density across any 1D diode, as  $t \rightarrow \infty$ , that is equal to the CL limit. However, these claims rest on a different definition of the CL limit, namely, a comparison between the time-averaged diode current and the adiabatic average of the expression for the stationary CL limit. If the current were considered as a function of the maximum applied voltage, rather than the average applied voltage, then the original conjecture would not have been refuted. © 2016 AIP Publishing LLC.

[<http://dx.doi.org/10.1063/1.4939607>]

This short research note points out that a conjecture regarding the Child-Langmuir (CL) limiting current density, while yet to be demonstrated, has not been refuted. The Child-Langmuir limit [Eq. (1)] gives the maximum current density of charged particles that can be forced across an un-neutralized gap by a certain voltage. It applies to a wide range of devices and natural phenomena such as ion beam sources, vacuum diodes, and plasma sheaths. There has been recent interest in generalizing this limit to situations where the current emission is a function of time. For example, the photoemission electron microscopy requires a series of electron pulses whose time interval can be distorted if the parameters violate a certain space charge limit.<sup>1</sup> In other cases, time-dependent current emission can arise spontaneously in low-voltage diodes even though the voltage drop is constant, with the cathode emitting intermittent bursts of single electrons<sup>2,3</sup> or sheets of electrons.<sup>4</sup>

In a previous paper, we investigated whether a current source with time-dependent emissivity could drive more current across a fixed-voltage gap, on average, than steady-state emission would allow. We were not able to beat this limit through empirical investigations, leading us to conjecture that there is a hard limit on the average current density across any one dimensional (1D) planar diode (as  $t \rightarrow \infty$ ) that is equal to the CL limit<sup>5</sup>

$$J_{CL} = \frac{4}{9} \epsilon_0 \sqrt{\frac{2q}{m}} \frac{V^{3/2}}{d^2}, \quad (1)$$

where  $q$  and  $m$  are the charge and mass of the current-carrying particles,  $V$  is the voltage across the diode,  $d$  is the width of the gap between electrodes, and  $\epsilon_0$  is the free space permittivity.

Since then, there have been several attempts to refute the conjecture, leading to refinements of the original conjecture. For example, in the regime where space charge effects limit the current across the diode to only a few electrons at a time, the average of the spontaneously time-varying current across a constant-voltage diode was shown to exceed  $J_{CL}$  by

13%.<sup>2</sup> Here, the increase in average current was not caused by the time-dependence of the current, but rather by special boundary conditions that pertain when the space charge electric field from a single electron is not negligible compared to the vacuum electric field in the diode.<sup>3</sup> The refined conjecture then ruled out discreteness effects, by excluding the case when the electric field at the cathode turns negative.

More recently, it has been claimed that even current in a traditional macroscopic diode could exceed  $J_{CL}$  if the voltage drop across the diode varied in time.<sup>6,7</sup> The point hinges on the question of how to define “exceeding the CL limit” for a diode whose voltage varies in time, because  $J_{CL}$  is only defined for diodes with constant voltage. In References 6 and 7, a new figure of merit,  $J_1$ , is defined, which is equal to the adiabatic average of the expression for  $J_{CL}$  [Eq. (2)]. A time-varying diode is then considered to exceed the CL limit when its average current exceeds  $J_1$ , namely,

$$J_1 = \frac{4}{9} \epsilon_0 \sqrt{\frac{2q}{m}} \frac{1}{d^2} \cdot \frac{1}{T} \int (V_T(t))^{\frac{3}{2}} dt, \quad (2)$$

where  $V_T$  is the time-varying voltage across the diode, and the adiabatic average is taken over the time interval  $T$ . It is then shown<sup>6</sup> that current in time-varying diodes can exceed  $J_1$  by as much as 50%. Note that exceeding this current does not refute the original conjecture, since what is called the CL limit is redefined.

However, the new definition represented by Eq. (2) raises the question of whether this redefined current limit might represent a meaningful current to exceed. First, note that the re-definition does not really pertain to the CL limit because the CL limit is not defined as a function of time. Second, note that a more physically meaningful measure, and the intent of our original conjecture, would be to compare the performance of a time-varying diode, with voltage  $V_T$ , to some stationary diode that has a constant voltage  $V_S$ . But what is the proper choice for  $V_S$ ? In our previous paper,<sup>5</sup> only the current emission varied in time and  $V_T$  was constant, so it was straightforward to set  $V_S = V_T$ . The choice is less clear when  $V_T$  is a function of time.

The definition of  $J_1$  is equivalent to comparing the average time-varying current to the CL limit of a stationary diode with constant voltage  $V_{S,1}$  defined as follows:

$$V_{S,1} = \left[ \frac{1}{T} \int (V_T(t))^{\frac{3}{2}} dt \right]^{2/3}. \quad (3)$$

The physical significance of this comparison is not clear. Practical considerations generally constrain the maximum voltage that drives charged particle current, not the time-average of  $V^{3/2}$ . Accordingly, the test for exceeding the Child-Langmuir law should be a comparison to a stationary diode with voltage  $V_{S,2}$  equal to the maximum voltage reached by the time-varying diode

$$V_{S,2} = \text{Maximum}[V_T(t)], \quad (4)$$

$$J_2 = \frac{4}{9} \epsilon_0 \sqrt{\frac{2q}{m}} \frac{1}{d^2} \{ \text{Maximum}[V_T(t)] \}^{3/2}.$$

Stated another way, the time-varying diode “exceeds the CL limit” when its average current is larger than  $J_2$ . Note that  $J_2$  is always larger than  $J_1$ , so it is harder to exceed the CL limit in this formulation and neither Reference 6 or 7 exceeded the CL limit using this stricter criteria. Thus, the original conjecture, with the more standard definition of the CL limiting current, captures succinctly the physical case and remains to be either refuted or proved.

<sup>1</sup>Y. L. Liu, P. Zhang, S. H. Chen, and L. K. Ang, *Phys. Plasmas* **22**, 084504 (2015).

<sup>2</sup>Y. Zhu and L. K. Ang, *Appl. Phys. Lett.* **98**, 051502 (2011).

<sup>3</sup>M. E. Griswold, N. J. Fisch, and J. S. Wurtele, *Phys. Plasmas* **19**, 024502 (2012).

<sup>4</sup>A. Pedersen, A. Manolescu, and A. Valfells, *Phys. Rev. Lett.* **104**, 175002 (2010).

<sup>5</sup>M. E. Griswold, N. J. Fisch, and J. S. Wurtele, *Phys. Plasmas* **17**, 114503 (2010).

<sup>6</sup>A. Rokhlenko, *Phys. Plasmas* **22**, 022126 (2015).

<sup>7</sup>R. E. Caflisch and M. S. Rosin, *Phys. Rev. E* **85**, 056408 (2012).