

Partial Differential Equations

Edited by Stewart Nash

2026

Preface

This material is taken from “Schaum’s Outline of Theory and Problems of Partial Differential Equations” by Paul DuChateau and David W. Zachmann. The footnotes are taken from content gathered from Wikipedia.

Contents

1	Difference Methods for Parabolic Equations	1
1.1	Difference Equations	1
2	Difference Methods for Hyperbolic Equations	3
2.1	One-Dimensional Wave Equation	3
3	Variational Formulation of Boundary Value Problems	5
3.1	Introduction	5
4	Variational Approximation Methods	7
4.1	The Rayleigh-Ritz Procedure	7
4.2	The Galerkin Procedure	8
5	The Finite Element Method: An Introduction	9

Chapter 1

Difference Methods for Parabolic Equations

1.1 Difference Equations

The various partial derivatives of a function $u(x, t)$ can be expressed as a *difference quotient* plus a truncation error (T.E.).

$$u_t(x, t) = \frac{u(x, t+k) - u(x, t)}{k} + \text{T.E.} \quad (1.1)$$

$$\text{T.E.} = -\frac{k}{2}u_{tt}(x, \bar{t}) \quad (t < \bar{t} < t+k) \quad (\text{Forward Difference for } u_t)$$

Chapter 2

Difference Methods for Hyperbolic Equations

2.1 One-Dimensional Wave Equation

Methods similar to those given in Section 9.4 may be used to approximate smooth solutions to

$$u_{tt} = c^2 u_{xx} \tag{2.1}$$

Let $(x_n, t_j) = (nh, jk)$ ($n, j = 0, 1, 2, \dots$) and write $s \equiv k/h$; we have as representatives of the two sorts of methods:

Chapter 3

Variational Formulation of Boundary Value Problems

3.1 Introduction

In certain cases, the solution of a boundary value problem for a PDE is also a solution of an associated calculus of variations problem.

Chapter 4

Variational Approximation Methods

This chapter presents some techniques for constructing approximate solutions to boundary value problems. These techniques are based on the ideas of Chapter 12 and differ markedly from the finite-difference methods of Chapters 9, 10, and 11.

4.1 The Rayleigh-Ritz Procedure

This approximation procedure is limited to boundary value problems admitting the variational formulation “Find u^* in \mathcal{A} such that functional $J[u]$ is minimized over \mathcal{A} by u^* .” It was seen in Chapter 12 that such boundary value problems arise in connection with self-adjoint elliptic PDEs.

We suppose \mathcal{A} to be some subset of $L^2(\Omega)$, where Ω denotes a bounded set in \mathbb{R}^n with smooth boundary S consisting of complementary portion S_1 and S_2 . Specifically, we take $\mathcal{A} = \{u \text{ in } H^1(\Omega) : u = g \text{ on } S_1\}$ for a given g in $C(\bar{\Omega})$; the associated subspace of comparison functions is taken as $\mathcal{M} = \{v \text{ in } H^1(\Omega) : v = 0 \text{ on } S_1\}$.

Let ϕ_0 denote an arbitrary function from \mathcal{A} (e.g., $\phi_0 = g$) and let ϕ_1, \dots, ϕ_N denote N linearly independent functions \mathcal{M} . Then,

$$u_N(\mathbf{x}) \equiv \phi_0(\mathbf{x}) + \sum_{j=1}^N c_j \phi_j(\mathbf{x}) \quad (4.1)$$

belongs to \mathcal{A} for *all* choices of the constants c_1, \dots, c_N ; we denote by \mathcal{A}_N the subset of \mathcal{A} consisting of all such functions u_N . Let u_N^* denote the function

in \mathcal{A}_N that minimizes J over \mathcal{A}_N . It can be shown that u_N^* represents the best approximation, in the least-squares sense, from \mathcal{A}_N to the exact solution u^* . This function u_N^* is called the *Rayleigh-Ritz approximation* to the solution of the boundary value problem.

The minimization of J over \mathcal{A}_N is tantamount to the minimization over all \mathbf{c} in \mathbb{R}^N of the ordinary function

$$H(c_1, \dots, c_N) \equiv J \left[\phi_0 + \sum_{j=1}^N c_j \phi_j \right] \quad (4.2)$$

The minimizing constants $c_1^*, c_2^*, \dots, c_N^*$ must satisfy

$$\frac{\partial H}{\partial c_m}(c_1, \dots, c_N) = 0 \quad (m = 1, \dots, N) \quad (4.3)$$

which is a system of N equations in N unknowns.

The Rayleigh-Ritz procedure may also be applied to eigenvalue problems of the sort treated in Section 12.4. In any eigenvalue problem the boundary conditions are all homogeneous. Hence, we take $\phi_0 \equiv 0$ in (13.1) and minimize the Rayleigh quotient J over the subspace $\mathcal{A}_N = \mathcal{M}_N$.

4.2 The Galerkin Procedure

This approximation method is employed when the boundary value problem admits only a weak formulation; e.g., in the case of a linear elliptic PDE containing odd-order derivatives. In the event that conditions (12.24) hold in the weak formulation, so that a variational formulation also exists, it can be shown (see Problem 13.4(b)) that the Galerkin and Rayleigh-Ritz procedure coincide.

Consider, then, a boundary value problem with the weak formulation “Find u^* in \mathcal{A} such that $K[u^*, v] = F[v]$ for all v in \mathcal{M} .” Here we suppose that Ω , S , \mathcal{A} , and \mathcal{M} are as described in Section 13.1. Let ϕ_1, \dots, ϕ_N denote N linearly independent *trial functions* in \mathcal{M} and let ϕ_0 denote an arbitrary function in \mathcal{A} . As in the Rayleigh-Ritz procedure, we seek an approximation u_N^* to the weak solution u^* of the form (13.1). In addition, let ψ_1, \dots, ψ_N denote N linearly independent *weight functions* in \mathcal{M} ; these may or may not be the same as the trial functions. The Galerkin approximate solution is required to satisfy

$$K[u_N^*, \psi_j] = F[\psi_j] \quad (j = 1, \dots, N) \quad (4.4)$$

This is a set of N equations in the N unknowns c_1^*, \dots, c_N^* .

Chapter 5

The Finite Element Method: An Introduction

The success of the approximation methods presented in Chapter 13 is largely dependent on the selection of an effective collection of trial functions ϕ_j and/or weight functions ψ_j . If these functions are chosen from certain families of piecewise polynomials, called *finite element spaces*, the following advantages are realized:

- (i) It is possible to deal in a systematic fashion with regions Ω having curved boundaries of rather arbitrary shape.
- (ii) One can systematically estimate the accuracy of the approximate solution in terms of the adjustable parameters associated with the finite element family.
- (iii) The coefficient matrix and data vector for the system of algebraic equations defining the approximate solution can be efficiently generated by computer.