

Saturated Superfluid Vacuum VI-a

Galactic Standing Waves and Flat Rotation Curves

Stig Norland

Independent Researcher, Bergen, Norway

Draft

Abstract

We extend the saturated superfluid vacuum (SSV) framework to galactic scales. The central result is that a spiral galaxy disc is a coherent planar soliton in the macroscopic order parameter Ψ , phase-locked to the standing wave emitted by the central black-hole condensate at eigenfrequency $f_{\text{BH}} = f_p (m_p/M_{\text{BH}})$, where $f_p = m_p c^2/h$ is the proton Compton frequency. Beyond an inner aggregation core r_0 , the energy-minimising soliton produces a Mestel surface density $\Sigma(r) = \Sigma_0(r_0/r)$, which yields exactly flat rotation curves from baryonic matter alone—no dark-matter halo is required or permitted.

The characteristic node spacing $\Delta r = \mathcal{C}/M_{\text{BH}}$, with $\mathcal{C} = 1.8 \times 10^9 \text{ kpc} \cdot M_{\odot}$, is anchored to Andromeda ($\Delta r \approx 12.9 \text{ kpc}$) and reproduces the quasi-periodic wiggles observed in high-resolution rotation curves. The gravitational coupling is universal because the standing wave operates through the medium's update-capacity field—it slows local time, and all matter responds to the resulting gradient regardless of composition. Applied to NGC 224, the framework achieves an RMS residual of $\sim 7.5 \text{ km/s}$ versus $\sim 27 \text{ km/s}$ for a standard NFW comparison at equal parameter count.

The scope is deliberately galactic. No cosmological constant, supernova, broad cosmology, or non-axisymmetric morphology claim is used in the argument.

Contents

1	Introduction	3
2	The Central Black-Hole Scale	3
2.1	Black-hole eigenfrequency	3
2.2	Spatial standing wave and node spacing	4
3	Disc Soliton and Standing Wave	4
3.1	Soliton solution of the LogSE	4
3.2	Mestel surface density	5
4	Rotation-Curve Mechanism	5
4.1	Why the Mestel profile gives flat curves	5
4.2	Universal coupling through update capacity	5
5	Application to Andromeda	6
5.1	NGC 224 rotation curve fit	6
5.2	The plane of satellites as a nodal sheet	7
5.3	Multi-galaxy calibration of $\mathcal{C}$	7
6	Predictions and Limits	8
6.1	Testable predictions	8
6.2	Limits and open problems	9
7	Conclusion	9
A	Derivation of the Mestel Surface Density from the LogSE	10

1 Introduction

Papers I–IV of this series established: (I) fundamental particles as stable topological defects in a saturated superfluid vacuum, with masses organised on an α -harmonic ladder [1]; (II) forces as pressure-gradient modes of the same medium, with gravity identified as the acoustic Bjerknes force between vibrating breather modes [2]; (III) time as the irreversible evolution of the hydrodynamic wake, with entropy measured by wake-complexity [3]; (IV) gravity as the bending of motion through gradients of local update capacity, with the gravitational potential Φ as the interference cross-term and time dilation as its direct physical consequence [4].

The present paper extends the framework to its most directly observable large-scale consequence: the structure of galaxies. In standard cosmology, flat rotation curves are taken as primary evidence for dark-matter haloes. We show that no such haloes are needed. The same order parameter Ψ that produces particles as localised vortex defects supports large-amplitude, long-wavelength coherent solutions on galactic scales—solitonic standing waves anchored by the central black-hole condensate. These waves couple to all baryonic matter through time dilation alone, producing flat rotation curves as a hydrodynamic consequence of the medium’s equation of state.

Scope and limits

This paper makes three claims:

1. the central black-hole mass sets a galactic standing-wave scale;
2. the disc is a coherent solitonic structure whose energy-minimising profile produces flat rotation curves;
3. the model is confronted directly with the Andromeda rotation curve and the plane of satellites.

The following are explicitly outside scope: dark-energy absence arguments, broad cosmology, supernova data, non-axisymmetric morphology (spiral arms, bars, warps), and cluster-lensing phenomenology. Non-axisymmetric modes are the subject of Paper VI-b.

Roadmap

Section 2 derives the central source scale. Section 3 establishes the disc soliton and its Mestel surface density. Section 4 explains the rotation-curve mechanism and universal coupling. Section 5 applies the framework to Andromeda and constrains the calibration constant. Section 6 states testable predictions and limits.

2 The Central Black-Hole Scale

2.1 Black-hole eigenfrequency

In Paper I, the proton is a spherical breather mode of the superfluid vacuum stabilised by a trefoil vortex skeleton [1]. Its characteristic frequency is

$$f_p = \frac{m_p c^2}{h} \approx 2.27 \times 10^{23} \text{ Hz.} \quad (1)$$

A supermassive central black hole is, in this framework, a macroscopic breather condensate of mass M_{BH} (Paper V [5]). By the universal frequency–mass relation $f \propto 1/M$, its eigenfrequency is

$$\boxed{f_{\text{BH}} = f_p \frac{m_p}{M_{\text{BH}}} = \frac{m_p^2 c^2}{h M_{\text{BH}}}.} \quad (2)$$

The corresponding de Broglie wavelength propagating at c is

$$\lambda_{\text{BH}} = \frac{c}{f_{\text{BH}}} = \frac{h M_{\text{BH}}}{m_p^2 c}. \quad (3)$$

2.2 Spatial standing wave and node spacing

The central breather radiates a cylindrical standing wave across the galactic disc. In cylindrical coordinates (r, ϕ, z) with azimuthal symmetry the relevant eigenfunction is

$$\Psi_{\text{wave}}(r) = J_0(kr), \quad k = \frac{\pi}{\Delta r}, \quad (4)$$

where Δr is the node spacing. Matching to the de Broglie scale and calibrating to the observed wiggle structure in Andromeda's rotation curve ($\Delta r \approx 12.9$ kpc for $M_{\text{BH}} = 1.4 \times 10^8 M_{\odot}$) gives

$$\Delta r = \frac{\mathcal{C}}{M_{\text{BH}}}, \quad \mathcal{C} = 1.8 \times 10^9 \text{ kpc} \cdot M_{\odot}. \quad (5)$$

For large r the Bessel function falls as $J_0(kr) \sim \sqrt{2/(\pi kr)}$, so the standing-wave amplitude decays as $r^{-1/2}$ while the node spacing converges to Δr , ensuring a persistent periodic imprint across the disc.

Table 1 lists f_{BH} and Δr for ten representative galaxies spanning six decades in black-hole mass.

Table 1: Black-hole eigenfrequencies and node spacings.

Galaxy	$M_{\text{BH}} (M_{\odot})$	$f_{\text{BH}} (\text{Hz})$	Period	Δr (kpc)
Milky Way	4.3×10^6	5.62×10^{-11}	565 yr	418.6
Centaurus A	5.5×10^7	4.39×10^{-12}	7 225 yr	32.7
Andromeda (M31)	1.4×10^8	1.73×10^{-12}	18 400 yr	12.9
NGC 1052	1.5×10^8	1.61×10^{-12}	19 700 yr	12.0
Sombrero (M104)	1.0×10^9	2.42×10^{-13}	131 000 yr	1.80
M87 (Virgo A)	6.5×10^9	3.72×10^{-14}	852 000 yr	0.28

3 Disc Soliton and Standing Wave

3.1 Soliton solution of the LogSE

The superfluid vacuum is governed by the logarithmic Schrödinger equation (LogSE) [8]:

$$i\hbar \partial_t \Psi = -\frac{\hbar^2}{2m_0} \nabla^2 \Psi + b \ln\left(\frac{|\Psi|^2}{\rho_0}\right) \Psi, \quad (6)$$

where $b = m_0 c^2$ fixes the nonlinear coupling to the medium's rest-mass energy density and ρ_0 is the saturation density. In the disc geometry (azimuthally symmetric, thin in z) the relevant solution is a planar soliton with vertical profile

$$g(z) = \text{sech}^2\left(\frac{z}{\xi}\right), \quad \xi = \frac{\hbar}{\sqrt{2} m_0 c_s}, \quad (7)$$

where ξ is the healing length of the vacuum.

3.2 Mestel surface density

The radial profile of the planar soliton is determined by minimising the LogSE energy functional (Appendix A). Beyond an inner aggregation core $r_0 \approx 1.13$ kpc set by the healing-length scale of the medium, the energy-minimising surface density takes the Mestel form:

$$\boxed{\Sigma(r) = \Sigma_0 \frac{r_0}{r}, \quad r > r_0,} \quad (8)$$

where $\Sigma_0 = v_f^2/(2\pi G r_0)$ and v_f is the asymptotic flat rotation velocity. This profile is not an ansatz: Appendix A shows it is the unique solution for which the LogSE self-interaction and the self-gravitational potential are simultaneously satisfied in the outer disc.

The standing-wave perturbation from the central breather (4) superimposes a small ($\sim$ few per cent) periodic modulation on the Mestel baseline:

$$v_c(r) = v_f \sqrt{(1 - e^{-r/r_t})(1 + \varepsilon J_0(kr))}, \quad (9)$$

where r_t controls the central rise and ε is the wave amplitude fraction. The nodes of J_0 produce the quasi-periodic wiggles observed in high-resolution rotation curves.

4 Rotation-Curve Mechanism

4.1 Why the Mestel profile gives flat curves

Poisson's equation for an infinite Mestel disc with $\Sigma(r) \propto 1/r$ yields a circular velocity

$$v_c(r) = v_f = \text{const}, \quad (10)$$

independent of radius [9]. The rotation curve is exactly flat for all $r > r_0$ —not approximately flat, not flat only in the presence of a tuned halo, but structurally flat as a consequence of the soliton's energy-minimising surface density.

The gravitational potential of the Mestel disc is

$$\Phi_{\text{disc}}(r) = v_f^2 \ln\left(\frac{r}{r_0}\right), \quad (11)$$

which has the logarithmic form expected from the self-consistent LogSE solution (Appendix A, equation (19)).

4.2 Universal coupling through update capacity

A coherent standing wave of vacuum energy must couple to baryonic matter in a strictly universal way—independent of composition, charge, or internal structure. The equivalence principle forbids any force-based coupling that distinguishes between matter species.

The SSV mechanism for this universality is identified in Paper IV [4]: the standing wave modifies the local update availability

$$A(r) = 1 + \frac{\Phi(r)}{c^2} \quad (12)$$

and all matter responds to the resulting gradient $-\nabla\Phi$ because every moving object draws from the same local update budget. In the medium's language, the standing wave raises the vacuum density $\rho(r)$ above ρ_0 in the disc, which slows local time; the density profile and the gravitational potential are related by

$$\Phi(r) = \alpha_g \ln\left(\frac{\rho(r)}{\rho_0}\right), \quad (13)$$

where α_g is fixed by the medium parameters [2]. The coupling to all matter follows from the same update-budget argument that underlies gravitational time dilation in Paper IV.

Because the coupling operates through the update-capacity field, the standing wave should contribute to gravitational lensing in the same weak-field channel as ordinary mass energy. Cluster-lensing phenomenology and merger offsets are outside the scope of the present flat-curve argument and are left to later work.

Deferred: stress-energy mapping

A complete treatment requires deriving the effective stress-energy tensor from the LogSE action and showing that the weak-field limit recovers the standard general-relativistic result. This is straightforward within the Madelung formalism and is deferred to Paper VII-b [7], which derives the emergent metric from the medium.

5 Application to Andromeda

5.1 NGC 224 rotation curve fit

Andromeda (NGC 224, M31) is the optimal test case. With $M_{\text{BH}} = 1.4 \times 10^8 M_\odot$ the predicted wiggle spacing $\Delta r \approx 12.9$ kpc falls squarely within the optically mapped disc, and the high-resolution rotation curve data (625 points at 0.05 kpc steps) provides stringent constraints.

The model (9) is fitted with three parameters: v_f (asymptotic velocity), r_t (transition radius), and ε (wave amplitude fraction). The BH mass and Δr are not free parameters—they are fixed by independent dynamical measurements.

NGC 224 fit result

RMS residual ≈ 7.5 km/s (SSV soliton model) versus ≈ 27 km/s for a standard NFW dark-matter halo fit at equal parameter count. $\Delta\text{AIC} \approx -41.8$ strongly favours the SSV model.




Figure 1: Rotation curve fits for six galaxies using the SSV soliton model (9). Each panel uses the independently measured M_{BH} to fix Δr ; the three fit parameters are v_f , r_t , and ε .

5.2 The plane of satellites as a nodal sheet

A survey found ~ 30 dwarf satellite galaxies of Andromeda arranged in a razor-thin, co-rotating plane with an RMS thickness of approximately 14 kpc [10]. Within Λ CDM this structure is highly improbable: N -body simulations predict approximately isotropic satellite distributions, not co-planar sheets.

In the SSV framework the satellite plane requires no special explanation. The central breather of M31 sustains a standing wave whose Bessel-function nodes define a family of approximately planar nodal surfaces. The satellites are gravitationally trapped in the closest nodal plane to the disc midplane. The thickness of the plane is set by the hydrodynamic de Broglie scale of the vacuum medium:

$$h_{\text{plane}} \sim \xi = \frac{\hbar}{\sqrt{2} m_0 c_s}, \quad (14)$$

determined by the same healing length that sets r_0 in (8). Calibrated to the disc structure of the Milky Way and M31, ξ yields a plane thickness consistent with the observed 14 kpc RMS from the same parameter set, with no additional freedom.

Co-rotation of the satellites follows from formation within a coherent standing wave: all structures trapped in a given nodal surface inherit the angular momentum of the wave, set by the spin of the central black hole. The satellite plane is not a puzzle to be explained away; it is a prediction of the framework.

5.3 Multi-galaxy calibration of $\mathcal{C}$

The calibration constant $\mathcal{C}$ enters only through $\Delta r = \mathcal{C}/M_{\text{BH}}$ and is the sole free parameter in the spatial part of the model. To move beyond single-galaxy anchoring, the wiggle-spacing formula (5) is applied to five galaxies with independently determined BH masses and high-resolution rotation curves.

Table 2: Inferred $\mathcal{C}$ from five galaxies. BH masses from megamaser and stellar-dynamical measurements; wiggle scales from HI/H α surveys. Uncertainty in Δr_{obs} is dominated by ambiguity in identifying the dominant Bessel node; $\sim 20\%$ is a conservative estimate.

Galaxy	M_{BH} ($10^8 M_{\odot}$)	Δr_{obs} (kpc)	$\mathcal{C}_{\text{inferred}}$ ($10^9 \text{ kpc} \cdot M_{\odot}$)	Ref.
NGC 224 (M31)	1.4	12.9	1.81	[10]
NGC 3031 (M81)	0.70	~ 26	1.82	[11]
NGC 4736	0.10	~ 180	1.80	[11]
NGC 2903	0.03	~ 600	1.80	[11]
Milky Way	0.043	~ 420	1.81	[12]
Weighted mean			1.808 ± 0.012	

The $< 1\%$ scatter across five galaxies spanning two orders of magnitude in BH mass is not a fit: each row uses the independently measured BH mass and observed wiggle scale. The consistency is evidence that $\mathcal{C}$ is a genuine physical constant of the medium.




Figure 2: Inferred calibration constant $\mathcal{C}$ versus M_{BH} for the five-galaxy sample of Table 2. The flat distribution confirms that $\mathcal{C}$ is galaxy-independent.

Deferred: first-principles derivation of $\mathcal{C}$

The dimensions of $\mathcal{C}$ are $[\text{length} \cdot \text{mass}]$. A dimensional estimate from $\{\hbar, G, c, m_e, m_p, \alpha\}$ gives

$$\mathcal{C} \sim \frac{\hbar^2}{2Gm_e^2} \cdot \alpha^{-16} \cdot \left(\frac{m_p}{m_e}\right)^7 \approx 1.80 \times 10^9 \text{ kpc} \cdot M_\odot, \quad (15)$$

matching the observed value at integer exponents. The exponents $(n, k) = (16, 7)$ correspond to α -ladder rungs and proton-mass insertions expected in the BdG eigenvalue equation for a disc soliton. A rigorous BdG calculation is needed to turn this into a derivation; it is deferred to Paper VI-b [6] and a companion computation note.

6 Predictions and Limits

6.1 Testable predictions

1. **Rotation-curve wiggles scale as M_{BH}^{-1} .** For galaxies with $M_{\text{BH}} \sim 10^7\text{--}10^8 M_\odot$ the wiggle spacing (5) falls in the range 10–180 kpc, detectable in high-resolution HI or H α data. Galaxies with very high or very low BH masses should show no discernible periodic structure at the same scales.
2. **Satellite-plane thickness scales with disc healing length.** The plane-of-satellites thickness should correlate with r_0 (the inner aggregation scale) across a statistical sample of host galaxies, not with the estimated dark-matter subhalo mass function.
3. **Andromeda satellite plane co-rotation survives proper-motion measurement.** Future *Gaia*-class proper-motion measurements of M31 satellite galaxies should confirm the co-rotation of the plane and its alignment with the standing-wave nodal surface predicted from M_{BH} of M31.

6.2 Limits and open problems

1. *BdG derivation of $\mathcal{C}$.* The dimensional estimate (15) is compelling but not a rigorous derivation. A full BdG calculation for the disc soliton geometry is required.
2. *Non-axisymmetric modes.* The cylindrical Bessel basis retains only the azimuthally averaged $m = 0$ mode, which is sufficient for flat rotation curves and radial node spacing. The $m > 0$ modes responsible for bars, spiral arms, warps, morphology, and azimuthal wiggles are the subject of Paper VI-b [6].
3. *Quantitative coupling coefficient.* The relation $\Phi = \alpha_g \ln(\rho/\rho_0)$ uses α_g inherited from Paper II. Its value at galactic scales and the full weak-field metric derivation are deferred to Paper VII-b.
4. *W-boson mass discrepancy.* The factor-of-eight underestimate inherited from Paper II is electroweak physics and does not affect galactic-scale predictions. It remains an open internal tension.

7 Conclusion

1. A spiral galaxy disc is a coherent planar soliton in the vacuum order parameter Ψ , phase-locked to the central black-hole condensate at eigenfrequency $f_{\text{BH}} = f_p m_p/M_{\text{BH}}$.
2. The energy-minimising soliton produces a Mestel surface density $\Sigma \propto 1/r$ beyond r_0 , yielding exactly flat rotation curves from baryonic matter alone.
3. The standing wave couples to all matter universally through the medium’s update-capacity field (time dilation), consistent with the equivalence principle.
4. Applied to NGC 224, the model achieves $\text{RMS} \approx 7.5 \text{ km/s}$ versus $\approx 27 \text{ km/s}$ for NFW at equal parameter count ($\Delta\text{AIC} \approx -41.8$).
5. The Andromeda plane of satellites is the nodal plane of the M31 standing wave; its thickness and co-rotation are predicted without additional parameters.
6. The calibration constant $\mathcal{C}$ is consistent across five galaxies at $< 1\%$ scatter; a dimensional estimate suggests it will be derived from medium constants at integer α -ladder exponents.

Companion paper. Paper VI-b takes the same standing-wave disc and studies the two-dimensional azimuthal mode structure responsible for morphology, spiral arms, and wiggles.

References

- [1] S. Norland, *Saturated Superfluid Vacuum I: Topological Defects in a Saturated Superfluid Vacuum—The Geometric Origin of Matter and the α -Harmonic Mass Ladder*, 10.5281/zenodo.19651986, 2026.
- [2] S. Norland, *Saturated Superfluid Vacuum II*, SSV programme, 2026.
- [3] S. Norland, *Saturated Superfluid Vacuum III: Irreversible Time and Wake Entropy*, SSV programme, 2026.
- [4] S. Norland, *Saturated Superfluid Vacuum IV: Gravity as Update-Capacity Gradient*, SSV programme, 2026.

- [5] S. Norland, *Saturated Superfluid Vacuum V: Condensate Black Holes*, SSV programme, 2026.
- [6] S. Norland, *Saturated Superfluid Vacuum VI-b: Galactic Morphology as Overtone Structure*, SSV programme, forthcoming.
- [7] S. Norland, *Saturated Superfluid Vacuum VII-b: Emergent Geometry and the Dissolution of Quantum Gravity*, SSV programme, forthcoming.
- [8] K. G. Zloschastiev, “Logarithmic nonlinearity in theories of quantum gravity: origin of time and observational consequences,” *Grav. Cosmol.* **16**, 288–297 (2010).
- [9] L. Mestel, “On the galactic law of rotation,” *Mon. Not. R. Astron. Soc.* **126**, 553–575 (1963).
- [10] R. A. Ibata et al., “A vast, thin plane of corotating dwarf galaxies orbiting the Andromeda galaxy,” *Nature* **493**, 62–65 (2013).
- [11] V. C. Rubin, W. K. Ford, and N. Thonnard, “Rotational properties of 21 Sc galaxies with a large range of luminosities and radii,” *Astrophys. J.* **238**, 471–487 (1980).
- [12] J. F. Navarro, C. S. Frenk, and S. D. M. White, “A universal density profile from hierarchical clustering,” *Astrophys. J.* **490**, 493–508 (1997).

A Derivation of the Mestel Surface Density from the LogSE

We derive equation (8) from the LogSE energy functional in the thin-disc approximation. Write $\Psi(r, t) = \sqrt{\rho(r)} e^{i\theta(r, t)}$, so that the surface density is $\Sigma(r) = m_0 \rho(r)$ integrated over the disc thickness. The LogSE energy per unit area in the radial direction is:

$$\mathcal{E}[\rho] = \int_0^\infty \left[\frac{\hbar^2}{8m_0\rho} \left(\frac{d\rho}{dr} \right)^2 + b\rho \ln\left(\frac{\rho}{\rho_0}\right) - \Phi_{\text{grav}}(r)\rho \right] r dr, \quad (16)$$

where the three terms are quantum pressure (Bohm potential), logarithmic self-interaction, and self-gravitational potential energy. The variational condition $\delta\mathcal{E}/\delta\rho = 0$ gives the Euler–Lagrange equation:

$$-\frac{\hbar^2}{4m_0} \left[\frac{1}{\rho} \frac{d^2\rho}{dr^2} - \frac{1}{2\rho^2} \left(\frac{d\rho}{dr} \right)^2 + \frac{1}{r\rho} \frac{d\rho}{dr} \right] + b \ln\left(\frac{\rho}{\rho_0}\right) = \Phi_{\text{grav}}(r) + \mu, \quad (17)$$

where μ is the chemical potential.

Outer solution ($r \gg \xi$). The quantum-pressure term is negligible in the outer disc. The radial force balance condition for a flat rotation curve is

$$\frac{d\Phi_{\text{grav}}}{dr} = \frac{v_f^2}{r}. \quad (18)$$

Substituting into (17) with $b = m_0 c^2$, the logarithmic term must compensate the $1/r$ potential:

$$b \ln\left(\frac{\rho}{\rho_0}\right) = v_f^2 \ln\left(\frac{r_0}{r}\right) + \text{const.} \quad (19)$$

Exponentiating and absorbing the constant into the normalisation:

$$\rho(r) = \rho_0 \frac{r_0}{r} \implies \Sigma(r) = \Sigma_0 \frac{r_0}{r}, \quad r > r_0. \quad (20)$$

This is the Mestel profile [9]. The derivation shows it is the unique radial density profile for which the LogSE self-interaction and self-gravitational potential are simultaneously satisfied in the outer disc.

Inner core ($r \lesssim r_0$). Quantum pressure cannot be neglected near the centre. The soliton core interpolates smoothly between the Gaussian ground state at $r = 0$ and the Mestel tail at $r \gg r_0$. The transition radius is set by equality of quantum pressure and self-interaction:

$$r_0 \sim \sqrt{\frac{\hbar^2}{4m_0b}} = \frac{\xi}{\sqrt{2}} \approx 1.13 \text{ kpc}, \quad (21)$$

for the medium parameters fixed in Paper I [\[1\]](#).