

Minimal Models in Mixed Characteristic

Liam Stigant

Imperial College London

Abstract

Recent work of Bhatt et al.[1] has established the bulk of the Minimal Model Program for klt threefolds over suitable mixed characteristic bases. This poster focuses on subsequent work to extend and apply these results. In particular on the Abundance and Finiteness of Minimal Models results and their applications.

Aims of the Minimal Model Program (MMP):

1. To show there is a distinguished class of representatives for a given birational class of varieties
2. To show these representatives have 'nice' geometry

In dimension 2 these representatives have some clear minimality properties, hence the name. In the modern formulation if we start with a variety X then we seek a birational model X' with 'good' singularities such that either

1. X' is a **Minimal Model** - it has nef canonical divisor $K_{X'}$
2. X' is a **Mori Fibre Space** - it admits a fibration $X' \rightarrow Z$ such that $-K_{X'}$ is ample over Z

The output is **not unique** but its nature is determined by the curvature of X . Either every possible output is a Mori Fibre Space or they are all Minimal Models.

In practice we work with **pairs** (X, B) consisting of a variety, X and a divisor (called a **boundary**) B . We then use $K_X + B$ instead of just K_X .

Here 'good' singularities means **klt** and \mathbb{Q} -factorial. A rather backwards definition is that (X, B) is a klt pair if it comes from an MMP for some log smooth pair (X', B') where B' is a divisor with coefficients less than 1.

Bibliography

- [1] Bhargav Bhatt et al. "Globally+-regular varieties and the minimal model program for threefolds in mixed characteristic". In: (2020).
- [2] Christopher D Hacon and James McKernan. "The Sarkisov program". In: (2009).

Abundance Conjecture

When (X, B) is a minimal model we expect it to have some quite specific geometry. Probably the most important conjecture here is the **Abundance Conjecture**. When $B = 0$ it says, roughly speaking, that X looks like a fibration of varieties with flat curvature over a lower dimensional one. E.g. An elliptic fibration.

More precisely, it says that if $K_X + B$ is nef, then it should be semiample.

If $\pi : (X, B) \dashrightarrow (X, B')$ is a minimal model and $\phi : X' \rightarrow Z$ is induced by abundance then $\phi \circ \pi$ is the (unique) **ample model** of (X, B) .


Theorem: Abundance in Mixed Characteristic

Let R be an excellent, Noetherian ring admitting a dualising complex with residue fields of characteristic $p > 5$. Let (X, B) be a klt threefold R -pair with \mathbb{R} -boundary. If $K_X + B$ is nef, then it is semiample.

Geography of Ample Models - Example


Let S be the blowup of \mathbb{P}_κ^2 at 2 points $p_1 \neq p_2$. Let E_1, E_2 be the exceptional curves of $S \rightarrow \mathbb{P}_\kappa^2$ and L the strict transform of a line through p_1, p_2 , so L, E_1, E_2 span $\text{Pic}(S)$. Choose $A \sim -K_S$ with $(S, A + E_1 + E_2 + L)$ log smooth.

Let T be the triangle formed by L, E_1, E_2 . Then for $B \in T$ the **ample model** of $K_S + A + B$ corresponds to polygon in a decomposition of T as below.



The decomposition of T describes the geometry of the Mori fibre spaces.

1. Triangles inside T with a side along the boundary correspond to Mori fibre spaces
2. Shared sides of triangles correspond to blowups
3. All the morphisms (blowups and Mori fibrations) are induced by abundance for pairs on the corresponding polygon



Geography of Ample Models - Definitions

Fix a \mathbb{Q} -divisor $A \geq 0$. Let V be a finite dimensional, rational affine subspace of $WDiv_{\mathbb{R}}(X)$ containing no components of A .

We have the following.

$$V_A = \{A + B : B \in V\}$$

$$\mathcal{L}_A(V) = \{\Delta = A + B \in V_A : (X, \Delta) \text{ is klt}\}$$

If $C \subseteq \mathcal{L}_A(V)$ is a **rational polytope** then we have

$$\mathcal{E}(C) = \{\Delta \in C : K_X + \Delta \text{ is pseudoeffective}\}$$

Given a birational contraction $\phi : X \dashrightarrow Y$ we also define

$$\mathcal{W}_\phi(C) = \{\Delta \in \mathcal{E}(C) : \phi \text{ is a weak log canonical (wlc) model of } (X, \Delta)\}$$

and given a rational map $\psi : X \dashrightarrow Z$ we have its ample class

$$\mathcal{A}_\phi(C) = \{\Delta \in \mathcal{E}(C) : \phi \text{ is the ample model of } (X, \Delta)\}$$

Roughly speaking a **wlc model** is an output of the MMP and an **ample model** is the map induced by abundance from a wlc model. On the diagram to the left the wlc models correspond to the interior triangles and the ample models to unions of their faces. The theorem below essentially says that these decompositions always exist.

These definitions ignore some of the mixed characteristic specific difficulties. Rather than working with klt pairs, we have to work with **rlt pairs**. Essentially these are pairs which can be replaced by a klt pair locally over a base. This is due to the lack of global Bertini type theorems. For example if A is ample on X , (X, A) need not be klt but it is always rlt.

Theorem: Finiteness of Minimal Models

Let R be an excellent ring admitting a dualising complex with residue fields of characteristic $p > 5$. Let X be a threefold over R . Let C be a polytope inside $\mathcal{L}_A(V)$. Suppose there is a boundary $A + B \in \mathcal{L}_A(V)$ such that $(X, A + B)$ is a klt pair. Then the following hold:

1. There are finitely many birational contractions $\phi_i : X \dashrightarrow Y_i$ such that

$$\mathcal{E}(C) = \bigcup \mathcal{W}_i = \mathcal{W}_{\phi_i}(C)$$

where each \mathcal{W}_i is a rational polytope. Moreover if $\phi : X \rightarrow Y$ is a wlc model for any choice of $\Delta \in \mathcal{E}(C)$ then $\phi = \phi_i$ for some i , up to composition with an isomorphism.


2. There are finitely many rational maps $\psi_j : X \dashrightarrow Z_j$ which partition $\mathcal{E}(C)$ into subsets $\mathcal{A}_{\psi_j}(C) = \mathcal{A}_i$.

3. For each W_i there is a j such that we can find a morphism $f_{i,j} : Y_i \rightarrow Z_j$ and $W_i \subseteq \overline{A_j}$.

4. $\mathcal{E}(C)$ is a rational polytope and A_j is a union of the interiors of finitely many rational polytopes.

Application: Sarkisov Program

If $f : X \rightarrow Z, g : Y \rightarrow W$ are two Mori Fibre Spaces, a **Sarkisov link** $s : X \dashrightarrow Y$ is one of the following.



where

- The horizontal map is a sequence of flops (codimension 2 birational transformations)
- Every vertical morphism is a contraction
- If the target of a vertical morphism is X or Y then it is an extremal divisorial contraction
- Either p, q are both Mori Fibre Spaces (this is type IV_m) or they are both small contractions (type IV_s)

The idea here is exactly the same as the example. We find a roof T dominating X, Y and a polygon $C \subseteq \mathcal{L}_A(V)$ for suitable A, V on T such that X, Y correspond to ample classes its decomposition (called a **geography**). The Sarkisov links then correspond to the boundaries between ample classes. On the left, π_1 is type III and $\pi_2 \circ \pi_3^{-1}$ is type II .

The **Sarkisov program** says that every Mori fibre space outputs of an MMP are related by Sarkisov links. That Finiteness of Minimal Models implies the Sarkisov Program is essentially due to [2], though a few modifications are needed in mixed characteristic.