

Patient ID: PA02	Patient Name: Anonymous
Age: 27 Years	Sex: F
Accession Number:	Modality: MR
Referring Physician:	Study: MRI LUMBAR SPINE +WSS
Study Date:	

MRI LUMBAR SPINE + WSS

PROTOCOL:-

Axial T1 Spin Echo and T2 Fast Spin Echo.

Sagittal T1 Spin Echo and T2 Fast Spin Echo.

Coronal STIR.

OBSERVATIONS:-

Lumbar lordosis straightened

Marginal osteophytes are seen at multiple levels

Mild Diffuse disc bulge with posterocentral disc is seen at the L5-S1 level indenting the thecal sac with mild narrowing of bilateral lateral recess and neural foramina

No subluxation / dislocation / spondylolisthesis is seen.

Vertebral body, pedicle, transverse process, spinous process, lamina and articular surfaces are normal. Pars interarticularis appear normal.

Facetal joints appear normal.

No altered signal intensity noted in the lumbar vertebral body.

The intervertebral disc seen at lumbar vertebra shows normal signal intensity.

There is no altered signal intensity noted in the visualised part of spinal cord.

Cord ends at L1 vertebral body and show normal signal intensity.

Cauda equina & filum terminale appear normal.

The iliopsoas and perivertebral soft tissues are unremarkable. Sacroiliac joint appear normal.

The sagittal diameters of the lumbar spinal canal are as follows (in mm):

Level :	L1-2	L2-3	L3-4	L4-5	L5-S1
(mm):	15.4	14.5	13.5	12.8	13.2

Patient ID: PA02	Patient Name:
Age: 27 Years	Sex: F
Accession Number:	Modality: MR
Referring Physician:	Study: MRI LUMBAR SPINE +WSS
Study Date:	

CERVICAL SPINE:

There is reversal of cervical curvature with mild diffuse disc bulges at C3-C4 to C6 C7 levels

The cervical vertebrae reveal normal signal intensity and alignment.

The intervertebral discs are of normal signal intensity on T2 weighted images.

No disc bulge is noted. The facet joints are unremarkable.

The cord is of normal signal intensity.

No abnormal pre or para vertebral soft tissue is seen.

DORSAL SPINE:

The dorsal vertebrae reveal normal signal intensity and alignment.

The intervertebral discs are of normal signal intensity on T2 weighted images.

No disc bulge is noted. The facet joints are unremarkable.

The cord is of normal signal intensity. No abnormality is detected in the conus medullaris.

No abnormal pre or para vertebral soft tissue is seen.

IMPRESSION: -

- **Mild Diffuse disc bulge with posterocentral disc is seen at the L5-S1 level indenting the thecal sac with mild narrowing of bilateral lateral recess and neural foramina**