

NAME : PA05	DATE : 14.10.2024
AGE/ SEX : 20years/Female	REF. BY:

Obstetric Fetal well Being Colour Doppler Study and BPP

LMP: 03.03.2024

EDD BY LMP: 08.12.2024

Gestational Age: 32wks 1day

EDD BY USG: 11.12.2024

FETAL. NO.	Single
PRESENTATION	Cephalic
LIQUOR	is adequate (AFI~10)
CARDIAC ACTIVITY	Normal - 136bpm/min
PLACENTA	is anterior, well above internal os, Grade- I, not low lying.

BPD	8.06cm	32weeks 3days
HC	28.57cm	31weeks 3days
FL	5.94cm	31weeks 0days
AC	27.90cm	32weeks 0days

EFW- 1805gms +/- 264gms

DOPPLER

	PSV	EDV	PI	RI	S/D
Umbilical artery	38.1	12.4	1.29	0.67	3.1
Fetal MCA	45.2	7.09	2.40	0.84	6.4
Right Uterine artery	72.2	37.6	0.66	0.48	1.9
left uterine artery	129	62.6	0.75	0.52	2.1

Fetal MCA and umbilical artery are normal, show colour flow and flow pattern.
The SD Ratio of umbilical artery is (Which is normal).

B/L Uterine arteries show normal colour flow & spectral trace.

Fetal thoracic aorta show normal colour flow and flow pattern.

BIOPHYSICAL PROFILE SCORING

Biophysical variable	Normal	Score
Fetal posture and tone (FT)	2 episodes of active extension with return to flexion of fetal limb(s) or trunk; opening and closing of hand considered normal tone.	2
Fetal movement (FM)	3 discrete body/limb movements in 30 min or less (episodes of active continuous movement considered as single movement).	2
Fetal breathing movement (FBM)	3-4 episodes of >30 s of FBM in 30min or less.	2
Amniotic fluid volume (AFV)	Adequate (AFI ~10).	2

Total Score = 8/8

No loop of cord seen around fetal neck in present scan.

IMPRESSION

- Single live intrauterine fetus of approx GA 31week's 5days +/- 2weeks in cephalic presentation.
- Normal colour Doppler study.
- BPP – 8/8

Suggested: Interval growth scans.

I, [Doctor Name] declare that while conducting Ultrasonography/image scanning on **[Patient Name]** I have neither detected nor disclosed the sex of her fetus to anybody in any manner.

(Not all the anomalies can be diagnosed on USG. Difficulty in assessment increases with increasing gestation age. Some anomalies may be missed due to the position of the fetus at the time of examination, not allowing proper visualizations. Subtle defects (cleft lip, anomalies of hand and feet, ear etc.) may not be detected in a 2D scan.)

Consultant MD Radiologist