

Lab 3: Exploring Discrete Probability Distributions

75694189 16257626 13341225 49442423

Chen Wang, Junke Wang, Zhuozhi Xiong, Titas Dutta

due: Oct 16, 2019 5:00 pm

Bernoulli and Binomial Distributions

Functions `dbinom`, `pbinom`, `qbinom`, `rbino`

Binomial

Taken from Open Intro Exercise 4.17:

Data collected by the Substance Abuse and Mental Health Services Administration (SAMSHA) suggests that 69.7% of 18-20 year olds consumed alcoholic beverages in any given year. A random sample of 10 people age 18-20 was taken.

Write text below:

1. What distribution does the number of 18-20 that consumed alcohol follow?

Binomial distribution.

2. What is the expected value of this distribution?

```
n1 <- 10
p1 <- 0.697
```

```
n1*p1
```

```
## [1] 6.97
```

6.97.

3. What is the variance of this distribution?

```
n1*p1*(1-p1)
```

```
## [1] 2.11191
```

The variance is 2.11.

Write code below:

4. What is the probability that exactly 5 out of 10 18-20 year olds have consumed an alcoholic beverage?

```
dbinom(x=5,size=10,prob=0.697)
```

```
## [1] 0.1058711
```

The probability is 0.11.

5. What is the probability that at most 3 out of 10 randomly sampled 18-20 year olds have consumed alcoholic beverages?

```
pbinom(q=3,size=10,prob = 0.697)
```

```
## [1] 0.01123721
```

The probability is 0.01.

6. What is the probability that at least 6 out of 10 randomly sampled 18-20 year olds have consumed alcoholic beverages?

```
1-pbinom(q=5, size=10, prob=0.697)
```

```
## [1] 0.844538
```

The probability is 0.84

7. What is the median of this distribution?

```
qbinom(p=0.5, size=10, prob=0.697)
```

```
## [1] 7
```

The median is 7.

You can also randomly generate samples using a binomial distribution.

```
rbinom(n=100,size = 20,prob = 0.5)
```

```
## [1] 12 12 11 13 13 5 8 6 10 10 9 10 10 10 11 9 8 13 12 11 8 11 10
## [24] 12 6 10 10 5 12 13 9 13 13 9 8 9 10 10 8 12 7 8 9 12 12 7
## [47] 12 10 13 10 9 9 9 9 10 6 12 9 9 11 11 11 10 8 11 8 6 6 9
## [70] 12 10 9 12 11 10 10 9 11 15 12 9 5 13 7 9 8 4 13 12 14 12 9
## [93] 13 8 12 12 12 7 7 8
```

Geometric Distribution

Functions `dgeom()`, `pgeom()`, `qgeom()`, `rgeom()`

In R, these functions model differently than the book. This looks at the number of failures until the first success.

The probability of a defective lightbulb at a certain factor is 0.30. Write text here:

8. What is distribution would the number of bulbs until the first defective is found follow?

Geometric

9. What is the expected number of lightbulbs that are checked before finding a defective bulb?

```
1/0.3
```

```
## [1] 3.333333
```

The expected number is 3.33.

10. What is the probability that the tenth lightbulb is the first defective bulb?

```
dgeom(x=9, prob = 0.3)
```

```
## [1] 0.01210608
```

11. What is the probability that the first defective bulb is found after the first 3 bulbs are checked?

```
1-pgeom(q=3,prob=0.3)
```

```
## [1] 0.2401
```

12. Again we can draw a random sample:

```
dataset <- rgeom(100,0.3)
dataset
```

```
## [1] 0 9 2 1 0 1 1 0 6 0 9 2 1 1 2 2 4 2 1 2 1 2 4
## [24] 1 1 2 0 2 1 4 5 2 3 1 13 3 0 1 3 3 0 1 1 7 6 1
## [47] 2 6 1 1 1 6 0 2 1 0 1 2 3 4 0 2 4 0 0 1 1 0 0
## [70] 0 3 1 0 8 0 1 3 0 0 0 0 4 3 3 5 1 0 0 5 7 0 4
## [93] 0 1 0 0 5 0 4 0
```

```
mean(dataset)
```

```
## [1] 2.1
```

```
hist(dataset)
```

Histogram of dataset

