EWRC for DY processes: dedicated generators

 $pp \to W^\pm \to l\nu$

 $-l_+l \leftarrow Z_* \downarrow \leftarrow dd$

➤ Pole approximation
 Wackeroth, Hollik, Phys. Rev. D55 (1997) 6788;

Baur, Keller, Wackeroth, PRD59 (1999) 013002;

 \star $\mathcal{O}(\alpha)$ photonic corrections Baur, Keller, Sakumoto, PRD57 (1998) 199; WGRAD

ZGRAD

 \star Complete $\mathcal{O}(\alpha)$ corrections U. Baur et al., Phys. Rev. **b65** (2002) 033007; V.A. Zykunov, Eur. P. J. C3 (2001) 9, Phys. Atom. Nucl. 69 (2006) 1522; Dittmaier, Krämer, Phys. Rev. D65 (2002) 073007; Brensing et al., Phys. Rev. D77 (2008) 073006; Baur, Wackeroth, Phys. Rev. D70 (2004) 073015; \star Complete $\mathcal{O}(\alpha)$ corrections

ZGRAD2

HORACE

C.M. Carloni Calame et al., JHEP 10 (2007) 190;

WGRAD2 A. Arbuzov et al., Eur. Phys. J. C46 (2006) 407; D. Bardin et al., arXiv:1207.4400 SANC C.M. Carloni Calame et al., JHEP 12 (2006) 016;

SANC

S. Dittmaier and M. Huber, JHEP 01 (2010) 060.

V.A. Zykunov, Phys. Rev. **D75** (2007) 073019; A. Arbuzov *et al.*, Eur. Phys. J **C54**:451-460, 2008; D. Bardin et al., arXiv:1207.4400

HORACE

F. Piccinini (INFN)