

# Sphere Packing in Lean

Maryna Viazovska, Sidharth Hariharan

August 16, 2024

## Abstract

This blueprint consists of an adaptation of Maryna Viazovska's Fields Medal-winning paper proving that no packing of unit balls in Euclidean space  $\mathbb{R}^8$  has density greater than that of the  $E_8$ -lattice packing. This blueprint is a work in progress, and will be frequently updated and restructured as the formalisation effort progresses. We recommend that you look at [this webpage](#) for the latest version.

## Contents

<b>1</b>	<b>Sphere packings</b>	<b>2</b>
1.1	The Setup	2
1.2	Scaling Sphere Packings	3
1.3	Lattices and Periodic packings	5
1.4	Density of periodic packings	6
1.5	Main Result	6
<b>2</b>	<b>The <math>E_8</math> lattice</b>	<b>6</b>
2.1	Definitions of $E_8$ lattice	6
2.2	Basic Properties of $E_8$ lattice	7
2.3	The $E_8$ sphere packing	8
<b>3</b>	<b>Facts from Fourier analysis</b>	<b>8</b>
<b>4</b>	<b>Cohn-Elkies linear programming bounds</b>	<b>9</b>
<b>5</b>	<b>Modular forms</b>	<b>11</b>
5.1	Modular forms and examples	11
5.2	Quasimodular forms and derivatives	17
<b>6</b>	<b>Fourier eigenfunctions with double zeroes at lattice points</b>	<b>23</b>

# 1 Sphere packings

The Sphere Packing problem is a classic optimisation problem with widespread applications that go well beyond mathematics. The task is to determine the “densest” possible arrangement of spheres in a given space. It remains unsolved in all but finitely many dimensions.

It was famously determined, in [12], that the optimal arrangement in  $\mathbb{R}^8$  is given by the  $E_8$  lattice. The result is strongly dependent on the Cohn-Elkies linear programming bound (Theorem 3.1 in [3]), which, if a  $\mathbb{R}^d \rightarrow \mathbb{R}$  function satisfying certain conditions exists, bounds the optimal density of sphere packings in  $\mathbb{R}^d$  in terms of it. The proof in [12] uses the theory of modular forms to construct a function that can be used to bound the density of all sphere packings in  $\mathbb{R}^8$  above by the density of the  $E_8$  lattice packing. This then allows us to conclude that no packing in  $\mathbb{R}^8$  can be denser than the  $E_8$  lattice packing.

## 1.1 The Setup

This subsection gives an overview for the setup of the problem, both informally and in Lean. Throughout this blueprint,  $\mathbb{R}^d$  will denote the Euclidean vector space equipped with distance  $\|\cdot\|$  and Lebesgue measure  $\text{Vol}(\cdot)$ . For any  $x \in \mathbb{R}^d$  and  $r \in \mathbb{R}_{>0}$ , we denote by  $B_d(x, r)$  the open ball in  $\mathbb{R}^d$  with center  $x$  and radius  $r$ . While we will give a more formal definition of a sphere packing below (and in Lean), the underlying idea is that it is a union of balls of equal radius centred at points that are far enough from each other that the balls do not overlap.

Arguably the most important definition in this subsection is that of *packing density*, which measures which portion of  $d$ -dimensional Euclidean space is covered by a given sphere packing. Taking the supremum over all packings gives what we refer to as the *sphere packing constant*, which is the quantity we are interested in optimising.

**Definition 1.1.** *Given a set  $X \subset \mathbb{R}^d$  and a real number  $r > 0$  (known as the separation radius) such that  $\|x - y\| \geq r$  for all distinct  $x, y \in X$ , we define the sphere packing  $\mathcal{P}(X)$  with centres at  $X$  to be the union of all open balls of radius  $r$  centred at points in  $X$ :*

$$\mathcal{P}(X) := \bigcup_{x \in X} B_d(x, r)$$

**Remark 1.2.** *Note that a sphere packing is uniquely defined from a given set of centres (which, in order to be a valid set of centres, must admit a corresponding separation radius). Therefore, as a conscious choice during the formalisation process, we will define everything that depends on sphere packings in terms of `SpherePacking`, a `structure` that bundles all the identifying information of*

a packing, but not the actual balls themselves. For the purposes of this blueprint, however, we will refrain from making this distinction.

We now define a notion of density for bounded regions of space by considering the density inside balls of finite radius.

**Definition 1.3.** *The finite density of a packing  $\mathcal{P}$  is defined as*

$$\Delta_{\mathcal{P}}(R) := \frac{\text{Vol}(\mathcal{P} \cap B_d(0, R))}{\text{Vol}(B_d(0, R))}, \quad R > 0.$$

As intuitive as it seems to take the density of a packing to be the limit of the finite densities as the radius of the ball goes to infinity, it is not immediately clear that this limit exists. Therefore, we define the density of a sphere packing as a limit superior instead.

**Definition 1.4.** *We define the density of a packing  $\mathcal{P}$  as the limit superior*

$$\Delta_{\mathcal{P}} := \limsup_{R \rightarrow \infty} \Delta_{\mathcal{P}}(R).$$

We may now define the sphere packing constant, the quantity that the sphere packing problem requires us to compute.

**Definition 1.5.** *The sphere packing constant is defined as supremum of packing densities over all possible packings:*

$$\Delta_d := \sup_{\substack{\mathcal{P} \subset \mathbb{R}^d \\ \text{sphere packing}}} \Delta_{\mathcal{P}}.$$

## 1.2 Scaling Sphere Packings

Given that the problem involves the *arrangement* of balls in space, it is intuitive not to worry about the radius of the balls (so long as they are all equal to each other). However, Definition 1.1 involves a choice of separation radius. In principle, we would want two sphere packing configurations that differ only in separation radii to ‘encode the same information’. In this brief subsection, we will describe how to change the separation radius of a sphere packing by *scaling* the packing by a positive real number and prove that this does not affect its density. This will give us the freedom to choose any separation radius we like when attempting to define the optimal sphere packing in  $\mathbb{R}^d$ .

**Definition 1.6.** *Given a sphere packing  $\mathcal{P}(X)$  with separation radius  $r$ , we defined the scaled packing with respect to a real number  $c > 0$  to be the packing  $\mathcal{P}(cX)$ , where  $cX = \{cx \in V \mid x \in X\}$  has separation radius  $cr$ .*

**Lemma 1.7.** *Let  $\mathcal{P}(X)$  be a sphere packing and  $c$  a positive real number. Then, for all  $R > 0$ ,*

$$\Delta_{\mathcal{P}(cX)}(cR) = \Delta_{\mathcal{P}(X)}(R).$$

*Proof.* The proof follows by direct computation:

$$\Delta_{\mathcal{P}(cX)}(cR) = \frac{\text{Vol}(\mathcal{P}(cX) \cap B_d(0, cR))}{\text{Vol}(B_d(0, cR))} = \frac{c^d \cdot \text{Vol}(\mathcal{P}(X) \cap B_d(0, R))}{c^d \cdot \text{Vol}(B_d(0, R))} = \Delta_{\mathcal{P}(X)}(R)$$

where the second equality follows from applying the fact that scaling a (measurable) set by a factor of  $c$  scales its volume by a factor of  $c^d$  to the fact that  $\mathcal{P}(cX) \cap B_d(0, cR) = c \cdot (\mathcal{P}(X) \cap B_d(0, R))$ .  $\square$

**Lemma 1.8.** *Let  $\mathcal{P}(X)$  be a sphere packing and  $c$  a positive real number. Then, the density of the scaled packing  $\mathcal{P}(cX)$  is equal to the density of the original packing  $\mathcal{P}(X)$ .*

*Proof.* One can show, using relatively unsophisticated real analysis, that

$$\limsup_{R \rightarrow \infty} \Delta_{\mathcal{P}(cX)}(R) = \limsup_{cR \rightarrow \infty} \Delta_{\mathcal{P}(cX)}(cR)$$

Lemma 1.7 tells us that  $\Delta_{\mathcal{P}(cX)}(cR) = \Delta_{\mathcal{P}(X)}(R)$  for every  $R > 0$ . Therefore,

$$\limsup_{cR \rightarrow \infty} \Delta_{\mathcal{P}(cX)}(cR) = \limsup_{cR \rightarrow \infty} \Delta_{\mathcal{P}(X)}(R) = \limsup_{R \rightarrow \infty} \Delta_{\mathcal{P}(X)}(R)$$

where the second equality is the result of a similar change of variables to the one done above.  $\square$

Therefore, as expected, we do not need to worry about the separation radius when constructing sphere packings. This will be useful when we attempt to construct the optimal sphere packing in  $\mathbb{R}^8$ —and even more so when attempting to *formalise* this construction—because the underlying structure of the packing is given by a set known as the  $E_8$  lattice, which has separation radius  $\sqrt{2}$ .

We can also use Lemma 1.8 to simplify the computation of the sphere packing constant by taking the supremum not over *all* sphere packings but only over those with density 1.

**Lemma 1.9.**

$$\Delta_d = \sup_{\substack{\mathcal{P} \subset \mathbb{R}^d \\ \text{sphere packing} \\ \text{sep. rad.}=1}} \Delta_{\mathcal{P}}$$

*Proof.* That the supremum over packings of unit density is at most the sphere packing constant is obvious. For the reverse inequality, let  $\mathcal{P}(X)$  be any sphere packing with separation radius  $r$ . We know, from Lemma 1.8, that the density of  $\mathcal{P}(X)$  is equal to that of the scaled packing  $\mathcal{P}(\frac{X}{r})$ . Since the scaled packing has separation radius 1, its density is naturally at most the supremum over all packings of unit density, meaning that the same is true of  $\mathcal{P}(X)$ .  $\square$

### 1.3 Lattices and Periodic packings

**Definition 1.10.** We say that an additive subgroup  $\Lambda \leq \mathbb{R}^d$  is a lattice if it is discrete and its  $\mathbb{R}$ -span contains all the elements of  $\mathbb{R}^d$ .

**Definition 1.11.** We say that a sphere packing  $\mathcal{P}(X)$  is  $(\Lambda)$ -periodic if there exists a lattice  $\Lambda \subset \mathbb{R}^d$  such that for all  $x \in X$  and  $y \in \Lambda$ ,  $x + y \in X$  (ie,  $X$  is  $\Lambda$ -periodic).

**Lemma 1.12.** Every periodic sphere packing is a sphere packing.

*Proof.* Mathematically, this lemma is hardly worth mentioning. We only do so to underscore the automatically constructed forgetful map `PeriodicSpherePacking.toSpherePacking` in Lean.  $\square$

**Lemma 1.13.** If  $\mathcal{P}(X)$  is a  $\Lambda$ -periodic sphere packing, then  $\Lambda$  acts on  $X$  by translation.

*Proof.* This is immediate from the definition of a periodic sphere packing.  $\square$

**Definition 1.14.** If  $\Lambda$  is a lattice, we call the  $\Lambda$ -periodic packing  $\mathcal{P}(\Lambda)$  with centres at points in  $\Lambda$  a lattice packing.

**Lemma 1.15.** If  $X \subseteq \mathbb{R}^d$  is a set of sphere packing centres that is periodic with respect to some lattice  $\Lambda$ , then the density of the corresponding (periodic) sphere packing is given by

$$\frac{|X/\Lambda|}{\text{Vol}(\mathbb{R}^d/\Lambda)} \cdot \text{Vol}(B_d(0,1))$$

where the quotients in the numerator and denominator correspond to the orbits of the action by translation of  $\Lambda$  on  $X$  and  $\mathbb{R}^d$  respectively.

**Remark 1.16.** This can be thought of as the “volume of spheres per fundamental domain”: the number of spheres per fundamental domain is  $|X/\Lambda|$ , and the volume of each sphere is  $\text{Vol}(B_d(0,1))$ .

**Definition 1.17.** The periodic sphere packing constant is defined to be

$$\Delta_d^{\text{periodic}} := \sup_{\substack{P \subset \mathbb{R}^d \\ \text{periodic packing}}} \Delta_P$$

**Theorem 1.18.** For all  $d$ , the periodic sphere packing constant in  $\mathbb{R}^d$  is equal to the sphere packing constant in  $\mathbb{R}^d$ .

*Proof.* **State this in Lean (ready). Fill in proof here (see [3, Appendix A])**  $\square$

In other words, it suffices to compute and optimise the periodic sphere packing constant.

## 1.4 Density of periodic packings

In this subsection, we build up results about the density of periodic packings. In particular, the density of a periodic packing, defined as the limit of the periodic packing intersected with a growing ball centered at the origin, is equal to the density within any fundamental region of the period lattice.

## 1.5 Main Result

With the terminologies above, we can state the main theorem of this project.

**Theorem 1.19.** *All periodic packing  $\mathcal{P} \subseteq \mathbb{R}^8$  has density satisfying  $\Delta_{\mathcal{P}} \leq \Delta_{E_8}$ , the density of the  $E_8$  sphere packing (see Definition ??).*

*Proof.* We will prove this theorem over the course of this blueprint. □

**Corollary 1.20.** *All packing  $\mathcal{P} \subseteq \mathbb{R}^8$  has density satisfying  $\Delta_{\mathcal{P}} \leq \Delta_{E_8}$ .*

*Proof.* This is a direct consequence of Theorem 1.18 and Theorem ??. □

**Corollary 1.21.**  $\Delta_8 = \Delta_{E_8}$ .

*Proof.* By definition,  $\Delta_{E_8} \leq \Delta_8$ , while Corollary ?? shows  $\Delta_8 = \sup_{\text{packing } \mathcal{P}} \Delta_{\mathcal{P}} \leq \Delta_{E_8}$ , and the result follows. □

## 2 The $E_8$ lattice

### 2.1 Definitions of $E_8$ lattice

There are several equivalent definitions of the  $E_8$  lattice. Below, we formalise two of them, and prove they are equivalent.

**Definition 2.1.**  *$E_8$ -lattice, Definition 1* We define the  $E_8$ -lattice (as a subset of  $\mathbb{R}^8$ ) to be

$$\Lambda_8 = \{(x_i) \in \mathbb{Z}^8 \cup (\mathbb{Z} + \frac{1}{2})^8 \mid \sum_{i=1}^8 x_i \equiv 0 \pmod{2}\}.$$

**Definition 2.2.** *We define the scaled  $E_8$ -lattice (by a real number  $c$ ) as*

$$c\Lambda_8 = \{c \cdot \vec{v} : \vec{v} \in \Lambda_8\}$$

.

**Definition 2.3.**  *$E_8$ -lattice, Definition 2* We define the  $E_8$  basis vectors to be the set of vectors

$$\mathcal{B}_8 = \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ -1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \\ -1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ -1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1/2 \\ -1/2 \\ -1/2 \\ -1/2 \\ -1/2 \\ -1/2 \\ -1/2 \\ -1/2 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ -1 \end{bmatrix} \right\}$$

**Definition 2.4.** We define the scaled  $E_8$  basis vectors (by a real number  $c$ ) to be  $c\mathcal{B}_8 = \{c \cdot \vec{v} : \vec{v} \in \mathcal{B}_8\}$ .

**Theorem 2.5.** The two definitions above coincide, i.e.  $c\Lambda_8 = \text{span}_{\mathbb{Z}}(c\mathcal{B}_8)$ .

*Proof.* We prove each side contains the other side.

For a vector  $\vec{v} \in \Lambda_8 \subseteq \mathbb{R}^8$ , we have  $\sum_i \vec{v}_i \equiv 0 \pmod{2}$  and  $\vec{v}_i$  being either all integers or all half-integers. After some modulo arithmetic, it can be seen that  $\mathcal{B}_8^{-1}\vec{v}$  as integer coordinates (i.e. it is congruent to 0 modulo 1). Hence,  $\vec{v} \in \text{span}_{\mathbb{Z}}(\mathcal{B}_8)$ .

For the opposite direction, we write the vector as  $\vec{v} = \sum_i c_i \mathcal{B}_8^i \in \text{span}_{\mathbb{Z}}(\mathcal{B}_8)$  with  $c_i$  being integers and  $\mathcal{B}_8^i$  being the  $i$ -th basis vector. Expanding the definition then gives  $\vec{v} = (c_1 - \frac{1}{2}c_7, -c_1 + c_2 - \frac{1}{2}c_7, \dots, -\frac{1}{2}c_7)$ . Again, after some modulo arithmetic, it can be seen that  $\sum_i \vec{v}_i$  is indeed 0 modulo 2 and are all either integers and half-integers, concluding the proof.

(Note: this proof doesn't depend on that  $\mathcal{B}_8$  is linearly independent.) □

## 2.2 Basic Properties of $E_8$ lattice

In this section, we establish basic properties of the  $E_8$  lattice and the  $\mathcal{B}_8$  vectors.

**Lemma 2.6.** For nonzero real numbers  $c$ , the set  $c\mathcal{B}_8$  is a  $\mathbb{R}$ -basis of  $\mathbb{R}^8$ .

*Proof.* It suffices to prove that  $\mathcal{B}_8 \in \text{GL}_8(\mathbb{R})$ . We prove this by explicitly defining the inverse matrix  $\mathcal{B}_8^{-1}$  and proving  $\mathcal{B}_8\mathcal{B}_8^{-1} = I_8$ , which implies that  $\det(\mathcal{B}_8)$  is nonzero. □

**Lemma 2.7.** For real numbers  $c$ ,  $c\Lambda_8$  is an additive subgroup of  $\mathbb{R}^8$ .

*Proof.* Trivially follows from that  $\Lambda_8 \subseteq \mathbb{R}^8$  is the  $\mathbb{Z}$ -span of  $\mathcal{B}_8$  and hence an additive group. □

**Lemma 2.8.** All vectors in  $\Lambda_8$  have norm of the form  $\sqrt{2n}$ , where  $n$  is a nonnegative integer.

*Proof.* Writing  $\vec{v} = \sum_i c_i \mathcal{B}_8^i$ , we have  $\|\vec{v}\|^2 = \sum_i \sum_j c_i c_j (\mathcal{B}_8^i \cdot \mathcal{B}_8^j)$ . Computing all 64 pairs of dot products and simplifying, we get a massive term that is a quadratic form in  $c_i$  with even integer coefficients, concluding the proof.  $\square$

**Lemma 2.9.** *For nonzero real numbers  $c$ ,  $c\Lambda_8$  is discrete, i.e. that the subspace topology induced by its inclusion into  $\mathbb{R}^8$  is the discrete topology.*

*Proof.* We prove this for  $c = 1$ . Since  $\Lambda_8$  is a topological group and  $+$  is continuous, it suffices to prove that  $\{0\}$  is open in  $\Lambda_8$ . This follows from the fact that there is an open ball  $\mathcal{B}(\sqrt{2}) \subseteq \mathbb{R}^8$  around it containing no other lattice points, since the shortest nonzero vector has norm  $\sqrt{2}$ .  $\square$

**Lemma 2.10.** *For nonzero real numbers  $c$ ,  $c\Lambda_8$  is a lattice, i.e. it is discrete and spans  $\mathbb{R}^8$  over  $\mathbb{R}$ .*

*Proof.* The first part is the above lemma. The second part follows from that  $\mathcal{B}_8$  is a basis and hence linearly independent over  $\mathbb{R}$ .  $\square$

**Prove  $E_8$  is unimodular. Prove  $E_8$  is positive-definite.**

## 2.3 The $E_8$ sphere packing

**Definition 2.11.** *The  $E_8$  sphere packing is the sphere packing with separation 1, whose set of centres is  $\frac{1}{\sqrt{2}}\Lambda_8$ .*

**Theorem 2.12.** *We have  $\Delta_{E_8} = \frac{\pi^4}{384}$ .*

*Proof.* **Finish proof. Preferably we want APIs about fundamental region of lattice, and use that to reduce this theorem to computation inside the fundamental region, and use formula for volume of ball.**  $\square$

## 3 Facts from Fourier analysis

In this section, we recall a few definitions from Fourier analysis.

**Definition 3.1.** *The Fourier transform of an  $L^1$ -function  $f : \mathbb{R}^d \rightarrow \mathbb{C}$  is defined as*

$$\mathcal{F}(f)(y) = \hat{f}(y) := \int_{\mathbb{R}^d} f(x) e^{-2\pi i \langle x, y \rangle} dx, \quad y \in \mathbb{R}^d$$

where  $\langle x, y \rangle = \frac{1}{2}\|x\|^2 + \frac{1}{2}\|y\|^2 - \frac{1}{2}\|x - y\|^2$  is the standard scalar product in  $\mathbb{R}^d$ .

**Definition 3.2.** *A  $C^\infty$  function  $f : \mathbb{R}^d \rightarrow \mathbb{C}$  is called a Schwartz function if it goes to zero as  $\|x\| \rightarrow \infty$  faster than any inverse power of  $\|x\|$ , and the same holds for all partial derivatives of  $f$ .*



**Definition 3.3.** *The set of all Schwartz functions is called a Schwartz space.*

**Lemma 3.4.** *The Fourier transform is an automorphism of the space of Schwartz functions.*

*Proof.* **Fill in proof.** □

**Lemma 3.5.**

$$\mathcal{F}(e^{\pi i \|x\|^2 z})(y) = z^{-4} e^{\pi i \|y\|^2 (\frac{-1}{z})}.$$

*Proof.* **Fill in proof.** □

**Theorem 3.6** ((Poisson summation formula)).

$$\sum_{\ell \in \Lambda} f(\ell) = \frac{1}{\text{Vol}(\mathbb{R}^d/\Lambda)} \sum_{m \in \Lambda^*} \widehat{f}(m).$$

*Proof.* **Fill in proof.** □

## 4 Cohn-Elkies linear programming bounds

In 2003 Cohn and Elkies [3] developed linear programming bounds that apply directly to sphere packings. The goal of this section is to formalize the Cohn–Elkies linear programming bound.

The following theorem is the key result of [3]. (The original theorem is stated for a class of functions more general than Schwartz functions)

**Theorem 4.1.** *(Cohn, Elkies [3]) Suppose that  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is a Schwartz function that is not identically zero and satisfies the following conditions:*

$$f(x) \leq 0 \text{ for } \|x\| \geq 1 \tag{1}$$

and

$$\widehat{f}(x) \geq 0 \text{ for all } x \in \mathbb{R}^d. \tag{2}$$

*Then the density of  $d$ -dimensional sphere packings is bounded above by*

$$\frac{f(0)}{\widehat{f}(0)} \cdot \text{vol}(B_d(0, 1/2)).$$

*Proof.* Here we reproduce the proof given in [3]. We will first prove the theorem for periodic packings.

Let  $X \subset \mathbb{R}^d$  be a discrete subset such that  $\|x - y\| \geq 1$  for any distinct  $x, y \in X$ . Suppose that  $X$  is  $\Lambda$ -periodic with respect to some lattice  $\Lambda \subset \mathbb{R}^d$ .

The inequality

$$\sharp(X/\Lambda) \cdot f(0) \geq \sum_{x \in X} \sum_{y \in X/\Lambda} f(x-y) = \sum_{x \in X/\Lambda} \sum_{y \in X/\Lambda} \sum_{\ell \in \Lambda} f(x-y+\ell) \quad (3)$$

follows from the condition (1) of the theorem and the assumption on the distances between points in  $X$ . The equality

$$\sum_{x \in X/\Lambda} \sum_{y \in X/\Lambda} \sum_{\ell \in \Lambda} f(x-y+\ell) = \sum_{x \in X/\Lambda} \sum_{y \in X/\Lambda} \frac{1}{\text{vol}(\mathbb{R}^d/\Lambda)} \sum_{m \in \Lambda^*} \widehat{f}(m) e^{2\pi i m(x-y)}.$$

follows from the Poisson summation formula. The right hand side of the above equation can be written as

$$\sum_{x \in X/\Lambda} \sum_{y \in X/\Lambda} \frac{1}{\text{vol}(\mathbb{R}^d/\Lambda)} \sum_{m \in \Lambda^*} \widehat{f}(m) e^{2\pi i m(x-y)} = \frac{1}{\text{vol}(\mathbb{R}^d/\Lambda)} \sum_{m \in \Lambda^*} \widehat{f}(m) \cdot \left| \sum_{x \in X/\Lambda} e^{2\pi i m x} \right|^2.$$

Note that  $\left| \sum_{x \in X/\Lambda} e^{2\pi i m x} \right|^2 \geq 0$  for all  $m \in \Lambda^*$ . Moreover, the term corresponding to  $m = 0$  satisfies  $\left| \sum_{x \in X/\Lambda} e^{2\pi i 0 x} \right|^2 = \sharp(X/\Lambda)^2$ . Now we use the condition (2) and estimate

$$\frac{1}{\text{vol}(\mathbb{R}^d/\Lambda)} \sum_{m \in \Lambda^*} \widehat{f}(m) \cdot \left| \sum_{x \in X/\Lambda} e^{2\pi i m(x-y)} \right|^2 \geq \frac{\sharp(X/\Lambda)^2}{\text{vol}(\mathbb{R}^d/\Lambda)} \cdot \widehat{f}(0). \quad (4)$$

Comparing inequalities (3) and (4) we arrive at

$$\frac{\sharp(X/\Lambda)}{\text{vol}(\mathbb{R}^d/\Lambda)} \leq \frac{f(0)}{\widehat{f}(0)}.$$

Now we see that the density of the periodic packing  $\mathcal{P}_X$  with balls of radius  $1/2$  is bounded by

$$\Delta(\mathcal{P}_X) = \frac{\sharp(X/\Lambda)}{\text{vol}(\mathbb{R}^d/\Lambda)} \cdot \text{vol}(B_d(0, 1/2)) \leq \frac{f(0)}{\widehat{f}(0)} \cdot \text{vol}(B_d(0, 1/2)).$$

This finishes the proof of the theorem for periodic packings. Theorem 1.18 implies the desired result for arbitrary packings.  $\square$

The main step in our proof of Theorem ?? is the explicit construction of an optimal function. It will be convenient for us to scale this function by  $\sqrt{2}$ .

**Theorem 4.2.** *There exists a radial Schwartz function  $g : \mathbb{R}^8 \rightarrow \mathbb{R}$  which satisfies:*

$$g(x) \leq 0 \text{ for } \|x\| \geq \sqrt{2} \quad (5)$$

$$\widehat{g}(x) \geq 0 \text{ for all } x \in \mathbb{R}^8 \quad (6)$$

$$g(0) = \widehat{g}(0) = 1. \quad (7)$$

Theorem 4.1 applied to the optimal function  $f(x) = g(x/\sqrt{2})$  immediately implies Theorem ??.

## 5 Modular forms

In this section, we recall and develop some theory of (quasi)modular forms.

### 5.1 Modular forms and examples

Let  $\mathfrak{H}$  be the upper half-plane  $\{z \in \mathbb{C} \mid \Im(z) > 0\}$ .

**Lemma 5.1.** *The modular group  $\Gamma_1 := \mathrm{SL}_2(\mathbb{Z})$  acts on  $\mathfrak{H}$  by linear fractional transformations*

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} z := \frac{az + b}{cz + d}.$$

Let  $N$  be a positive integer.

**Definition 5.2.** *The level  $N$  principal congruence subgroup of  $\Gamma_1$  is*

$$\Gamma(N) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_1 \mid \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \pmod{N} \right\}.$$

**Definition 5.3.** *A subgroup  $\Gamma \subset \Gamma_1$  is called a congruence subgroup if  $\Gamma(N) \subset \Gamma$  for some  $N \in \mathbb{N}$ .*

**Definition 5.4.** *Define the matrices*

$$S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \in \Gamma_1, T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \in \Gamma_1, \alpha = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} \in \Gamma_2 \subset \Gamma_1, \beta = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} \in \Gamma_2 \subset \Gamma_1.$$

*It is easily verifiable that  $\alpha = T^2$  and  $\beta = -S\alpha^{-1}S = -ST^{-2}S$ .*

The following two lemmas tell us the group structure of  $\Gamma(1) = \Gamma_1$  and  $\Gamma(2)$ , which we will use later on to define the theta forms.

**Lemma 5.5.** *We have  $\Gamma(1) = \langle S, T, -I \rangle$ .*

*Proof.* See [4, Exercise 1.1.1]. □

**Lemma 5.6.** *We have  $\Gamma(2) = \langle \alpha, \gamma, -I \rangle$ .*

*Proof.* See [4, Exercise 1.2.4]. □

Let  $z \in \mathfrak{H}$ ,  $k \in \mathbb{Z}$ , and  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ . We omit many of the proofs below when they exist in Mathlib already.

**Definition 5.7.** The automorphy factor of weight  $k$  is defined as

$$j_k(z, \begin{pmatrix} a & b \\ c & d \end{pmatrix}) := (cz + d)^{-k}.$$

**Lemma 5.8.** The automorphy factor satisfies the chain rule

$$j_k(z, \gamma_1 \gamma_2) = j_k(z, \gamma_1) j_k(\gamma_2 z, \gamma_1).$$

**Definition 5.9.** Let  $F$  be a function on  $\mathfrak{H}$  and  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ . Then the slash operator acts on  $F$  by

$$(F|_k \gamma)(z) := j_k(z, \gamma) F(\gamma z).$$

**Lemma 5.10.** The chain rule implies

$$F|_k \gamma_1 \gamma_2 = (F|_k \gamma_1)|_k \gamma_2.$$

In particular, this lemma implies that if  $\Gamma = \langle M_i \rangle_{i \in \mathcal{I}}$ , then the slash action  $F|_\gamma$  is uniquely determined by the action of generators, i.e.  $F|M_i$  and  $F|M_i^{-1}$ .

**Lemma 5.11.** For even  $k$ ,  $F|_k(-I) = F$ .

*Proof.* Follows from the definition of the slash operator:  $(F|_k(-I))(z) = (-1)^{-k} F((-I)z) = F(z)$ .  $\square$

**Definition 5.12.** A (holomorphic) modular form of integer weight  $k$  and congruence subgroup  $\Gamma$  is a holomorphic function  $f : \mathfrak{H} \rightarrow \mathbb{C}$  such that:

1. (Slash invariant)  $f|_k \gamma = f$  for all  $\gamma \in \Gamma$
2. (Holomorphic at  $i\infty$ ) for each  $\alpha \in \Gamma_1$   $f|_k \alpha$  has the Fourier expansion  $f|_k \alpha(z) = \sum_{n=0}^{\infty} c_f(\alpha, \frac{n}{n_\alpha}) e^{2\pi i \frac{n}{n_\alpha} z}$  for some  $n_\alpha \in \mathbb{N}$  and Fourier coefficients  $c_f(\alpha, m) \in \mathbb{C}$ .

**Definition 5.13.** Let  $M_k(\Gamma)$  be the space of modular forms of weight  $k$  and congruence subgroup  $\Gamma$ .

A key fact in the theory of modular forms is the following theorem:

**Theorem 5.14.** The spaces  $M_k(\Gamma)$  are finite dimensional.

*Proof.* In this project, we only require the theorem for  $\Gamma = \Gamma_1$  and  $\Gamma = \Gamma_2$ . For the proof, see [11].  $\square$

Let us consider several examples of modular forms.

**Definition 5.15.** For an even integer  $k \geq 4$  we define the weight  $k$  Eisenstein series as

$$E_k(z) := \frac{1}{2\zeta(k)} \sum_{(c,d) \in \mathbb{Z}^2 \setminus (0,0)} (cz + d)^{-k}. \quad (8)$$

**Lemma 5.16.** For all  $k$ ,  $E_k \in M_k(\Gamma_1)$ . Especially, we have

$$E_k\left(-\frac{1}{z}\right) = z^k E_k(z). \quad (9)$$

*Proof.* This follows from the fact that the sum converges absolutely. Now apply slash operator with  $\gamma = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$  gives (9).  $\square$

**Lemma 5.17.** The Eisenstein series possesses the Fourier expansion

$$E_k(z) = 1 + \frac{2}{\zeta(1-k)} \sum_{n=1}^{\infty} \sigma_{k-1}(n) e^{2\pi i n z}, \quad (10)$$

where  $\sigma_{k-1}(n) = \sum_{d|n} d^{k-1}$ . In particular, we have

$$\begin{aligned} E_4(z) &= 1 + 240 \sum_{n=1}^{\infty} \sigma_3(n) e^{2\pi i n z} \\ E_6(z) &= 1 - 504 \sum_{n=1}^{\infty} \sigma_5(n) e^{2\pi i n z}. \end{aligned}$$

The infinite sum (8) does not converge absolutely for  $k = 2$ . On the other hand, the expression (10) converges to a holomorphic function on the upper half-plane and we will take it as a definition of  $E_2$  (See Definition 5.34 below).

The discriminant form is a unique normalized cusp form of weight 12, which can be defined using  $E_4$  and  $E_6$ .

**Definition 5.18.** The discriminant form  $\Delta(z)$  is given by

$$\Delta(z) = \frac{E_4(z)^3 - E_6(z)^2}{1728} \quad (11)$$

**Lemma 5.19.**  $\Delta(z) \in M_{12}(\Gamma_1)$  and it vanishes at the unique cusp, i.e. it is a cusp form of level  $\Gamma_1$  and weight 12.

*Proof.* Being a modular form of desired weight and level directly follows from those of  $E_4$  and  $E_6$ . It is a cusp form since the constant terms of Fourier expansions of  $E_4$  and  $E_6$  are both 1.  $\square$

It also admits a product formula, which allow us to prove positivity of  $\Delta(it)$  for  $t > 0$  later.

**Lemma 5.20.** *We have*

$$\Delta(z) = e^{2\pi iz} \prod_{n \geq 1} (1 - e^{2\pi inz})^{24}. \quad (12)$$

*Proof.* There are several known proofs of this. One possible proof that we can formalize is from Kohnen [6], which prove

$$\frac{1}{2\pi iz} \frac{d}{dz} (\log \Delta) = 1 - 24 \sum_{n \geq 1} \frac{ne^{2\pi inz}}{1 - e^{2\pi inz}}. \quad (13)$$

by using a multiplicative analogue of the Hecke operator and the valence formula.  $\square$

Note that the RHS of (13) is equal to the  $E_2(z)$ . As a side note, we can also consider defining  $\Delta$  as (12), and prove that it coincides with (11). Such an argument can be found in [2, Section 2.4].

**Corollary 5.21.**  $\Delta(it) > 0$  for all  $t > 0$ .

*Proof.* By Lemma 5.20, we have

$$\Delta(it) = e^{-2\pi t} \prod_{n \geq 1} (1 - e^{-2\pi nt})^{24} > 0.$$

$\square$

Another examples of modular forms we would like to consider are *theta functions* [13, Section 3.1].

**Definition 5.22.** *We define three different theta functions (so called “Thetanullwerte”) as*

$$\begin{aligned} \Theta_2(z) &= \theta_{10}(z) = \sum_{n \in \mathbb{Z}} e^{\pi i(n + \frac{1}{2})^2 z}, \\ \Theta_3(z) &= \theta_{00}(z) = \sum_{n \in \mathbb{Z}} e^{\pi i n^2 z}, \\ \Theta_4(z) &= \theta_{01}(z) = \sum_{n \in \mathbb{Z}} (-1)^n e^{\pi i n^2 z} \end{aligned}$$

For convenience, we use the following notations for the fourth powers of the theta functions.

**Definition 5.23.** *Define*

$$H_2 = \Theta_2^4, \quad H_3 = \Theta_3^4, \quad H_4 = \Theta_4^4.$$

Note that we only need these fourth powers to define (6.13).

The group  $\Gamma_1$  is generated by the elements  $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ ,  $S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ , and  $-I = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$  (Lemma 5.5), and the transformation of functions under  $\Gamma(2)$  is determined by that under  $\Gamma_1$  (by Lemma 5.10). The

following lemma shows how the theta functions (and their powers) transform under the slash action of these matrices.

**Lemma 5.24.** *These elements act on the theta functions in the following way*

$$H_2|S = -H_4 \quad (14)$$

$$H_3|S = -H_3 \quad (15)$$

$$H_4|S = -H_2 \quad (16)$$

and

$$H_2|T = -H_2 \quad (17)$$

$$H_3|T = H_4$$

$$H_4|T = H_3$$

*Proof.* The last three identities easily follow from the definition. For example, (17) follows from

$$\begin{aligned} \Theta_2(z+1) &= \sum_{n \in \mathbb{Z}} e^{\pi i(n+\frac{1}{2})^2(z+1)} = \sum_{n \in \mathbb{Z}} e^{\pi i(n+\frac{1}{2})^2} e^{\pi i(n+\frac{1}{2})^2 z} \\ &= \sum_{n \in \mathbb{Z}} e^{\pi i(n^2+n+\frac{1}{4})} e^{\pi i(n+\frac{1}{2})^2 z} = \sum_{n \in \mathbb{Z}} (-1)^{n^2+n} e^{\pi i/4} e^{\pi i(n+\frac{1}{2})^2 z} \\ &= e^{\pi i/4} \Theta_2(z) \end{aligned}$$

and taking 4th power. (14) and (16) are equivalent under  $z \leftrightarrow -1/z$ , so it is enough to show (14) and (15). These identities follow from the identities of the *two-variable* Jacobi theta function, which is defined as (be careful for the variables, where we use  $\tau$  instead of  $z$ )

$$\theta(z, \tau) = \sum_{n \in \mathbb{Z}} e^{2\pi i n z + \pi i n^2 \tau}$$

and already formalized by David Loeffler. This function specialize to the theta functions as

$$\Theta_2(\tau) = e^{\pi i \tau/4} \theta(-\tau/2, \tau)$$

$$\Theta_3(\tau) = \theta(0, \tau)$$

$$\Theta_4(\tau) = \theta(1/2, \tau)$$

Possion summation formula gives

$$\theta(z, \tau) = \frac{1}{\sqrt{-i\tau}} e^{-\frac{\pi i z^2}{\tau}} \theta\left(\frac{z}{\tau}, -\frac{1}{\tau}\right)$$

and applying the specializations above yield the identities. For example, (16) follows from

$$\Theta_4(\tau) = \theta\left(\frac{1}{2}, \tau\right) = \frac{1}{\sqrt{-i\tau}} e^{-\frac{\pi i}{4\tau}} \theta\left(\frac{1}{2\tau}, -\frac{1}{\tau}\right) = \frac{1}{\sqrt{-i\tau}} \Theta_2\left(-\frac{1}{\tau}\right)$$

and taking 4th power. □

Using the above identities, we can prove that these are modular forms.

**Lemma 5.25.**  *$H_2$ ,  $H_3$ , and  $H_4$  are slash invariant under  $\Gamma(2)$ , i.e. for all  $\gamma \in \Gamma(2)$  and  $i \in \{2, 3, 4\}$ , we have  $H_i|\gamma = H_i|\gamma^{-1} = H_i$ .*

*Proof.* By Lemma 5.6 and Lemma 5.10, it suffices to show that the  $H_i$  are invariant under slash actions with respect to  $\alpha$ ,  $\beta$ , and  $-I$ . Invariance under  $-I$  follows from Lemma 5.11. The rest follows from Lemma 5.10, 5.24, and the matrix identities

$$\alpha = T^2, \quad \beta = -S\alpha^{-1}S = -ST^{-2}S.$$

For example, invariance for  $H_2$  can be proved by

$$\begin{aligned} H_2|\alpha &= H_2|T^2 = -H_2|T = H_2 \\ H_2|\beta &= H_2|(-S\alpha^{-1}S) = H_2|(S\alpha^{-1}S) = -H_4|(\alpha^{-1}S) = -H_4|S = H_2. \end{aligned}$$

□

**Lemma 5.26.**  *$H_2$ ,  $H_3$ , and  $H_4$  belong to  $M_2(\Gamma(2))$ .*

*Proof.* **Fill in this proof, in particular the analytic properties of  $H_i$ .**

We have to show that each  $H_i$  satisfies property (2) of Definition 5.12 □

We also have a nontrivial relation between these theta functions.

**Lemma 5.27.** *These three theta functions satisfy the Jacobi identity*

$$H_2 + H_4 = H_3 \Leftrightarrow \Theta_2^4 + \Theta_4^4 = \Theta_3^4. \tag{18}$$

*Proof.* Use the dimension formula of the space of modular forms of weight 2 and level  $\Gamma(2)$ . Especially, we have  $\dim M_2(\Gamma(2)) = 2$  with basis  $H_2$  and  $H_4$ . □

These are also related to  $E_4$ ,  $E_6$ , and  $\Delta$  as follows.



**Lemma 5.28.** *We have*

$$E_4 = \frac{1}{2}(H_2^2 + H_3^2 + H_4^2) = H_2^2 + H_2H_4 + H_4^2 \quad (19)$$

$$E_6 = \frac{1}{2}(H_2 + H_3)(H_3 + H_4)(H_4 - H_2) = \frac{1}{2}(H_2 + 2H_4)(2H_2 + H_4)(H_4 - H_2) \quad (20)$$

$$\Delta = \frac{1}{256}(H_2H_3H_4)^2. \quad (21)$$

*Proof.* We can prove these similarly as Lemma 5.27. Right hand sides of (19), (20), and (21) are all modular forms of level  $\Gamma_1$  and desired weights, where (21) is a cusp form since  $H_2$  is. Now the identities follow from the dimension calculations  $\dim M_4(\Gamma_1) = \dim M_6(\Gamma_1) = \dim S_{12}(\Gamma_1) = 1$  and comparing the first nonzero  $q$ -coefficients.  $\square$

The *strict* positivity of Jacobi theta functions might needed later.

**Lemma 5.29.** *All three functions  $t \mapsto H_2(it), H_3(it), H_4(it)$  are positive for  $t > 0$ .*

*Proof.* By Lemma 5.27 and the transformation law (14), it is enough to prove the positivity for  $\Theta_2(it)$ , which is clear from its definition:

$$\Theta_2(it) = \sum_{n \in \mathbb{Z}} e^{-\pi(n+\frac{1}{2})^2 t} > 0.$$

$\square$

## 5.2 Quasimodular forms and derivatives

Morally, quasimodular forms can be thought as *modular forms with differentiations*. It can be defined formally as follows.

**Definition 5.30** (Quasimodular form). *Let  $f : \mathfrak{H} \rightarrow \mathbb{C}$  be a holomorphic function, and let  $k$  and  $s \geq 0$  be integers. The function  $f$  is a quasimodular form of weight  $k$ , level  $\Gamma$ , and depth  $s$  if there exist holomorphic functions  $f_0, \dots, f_s : \mathfrak{H} \rightarrow \mathbb{C}$  such that*

$$(f|_k \gamma)(z) = (cz + d)^{-k} f\left(\frac{az + b}{cz + d}\right) = \sum_{j=0}^s f_j(z) \left(\frac{c}{cz + d}\right)^j \quad (22)$$

for all  $z \in \mathfrak{H}$  and  $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ .

By taking  $\gamma = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ , one can check that we should have  $f_0 = f$ . Thus, a quasimodular form of depth 0 is just a modular form of same weight and level.

**Definition 5.31.** *Let  $F$  be a quasimodular form. We define the (normalized) derivative of  $F$  as*

$$F' = DF := \frac{1}{2\pi i} \frac{d}{dz} F. \quad (23)$$

**Lemma 5.32.** *We have an equality of operators  $D = q \frac{d}{dq}$ . In particular, the  $q$ -series of the derivative of a quasimodular form  $F(z) = \sum_{n \geq n_0} a_n q^n$  is  $F'(z) = \sum_{n \geq n_0} n a_n q^n$ .*

*Proof.* Directly follows from the definition (5.31), where  $\frac{1}{2\pi i} \frac{d}{dz} e^{2\pi i n z} = n e^{2\pi i n z}$ .  $\square$

**Theorem 5.33.** *The space of quasimodular forms is closed under the derivative (23). If  $F$  is a quasimodular form of weight  $k$ , level  $\Gamma$ , and depth  $s$ , then  $F'$  is a quasimodular form of weight  $k + 2$ , level  $\Gamma$ , and depth  $s + 1$ .*

*Proof.* This follows from differentiating the definitional equation (22): we get

$$\begin{aligned} & -kc(cz+d)^{-k-1}F\left(\frac{az+b}{cz+d}\right) + (cz+d)^{-k-2}\frac{dF}{dz}\left(\frac{az+b}{cz+d}\right) \\ &= \sum_{j=0}^s \frac{dF_j}{dz}(z) \left(\frac{c}{cz+d}\right)^j + F_j(z) \cdot (-j) \left(\frac{c}{cz+d}\right)^{j+1} \\ &\Rightarrow (cz+d)^{-k-2}F'\left(\frac{az+b}{cz+d}\right) = \sum_{j=0}^{s+1} G_j(z) \left(\frac{c}{cz+d}\right)^j \end{aligned}$$

where

$$G_j(z) := F'_j(z) - \frac{ikc}{2\pi(cz+d)}F_j(z) + \frac{i(j-1)}{2\pi}F_{j-1}(z), \quad F_{-1} := 0$$

for  $0 \leq j \leq s+1$ , which are holomorphic.  $\square$

The most important quasimodular form is the weight 2 Eisenstein series  $E_2$ . We define it as a  $q$ -series, which gives a holomorphic function on  $\mathfrak{H}$ .

**Definition 5.34.** *We set*

$$E_2(z) := 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n) e^{2\pi i n z}.$$

**Lemma 5.35.** *This function is not modular, however it satisfies*

$$z^{-2} E_2\left(\frac{-1}{z}\right) = E_2(z) - \frac{6i}{\pi} \frac{1}{z}. \quad (24)$$

*More generally, we have*

$$(cz+d)^{-2} E_2\left(\frac{az+b}{cz+d}\right) = E_2(z) - \frac{6ic}{\pi(cz+d)}, \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}). \quad (25)$$

*Proof.* Use (13). Modularity of  $\Delta(z)$  gives  $(cz+d)^{-12} \Delta\left(\frac{az+b}{cz+d}\right) = \Delta(z)$  for  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_1$ , and by differentiating it we get

$$(cz+d)^{-14} \Delta'\left(\frac{az+b}{cz+d}\right) = \Delta'(z) - \frac{6ic}{\pi(cz+d)} \Delta(z).$$

Now, divide both sides with  $\Delta(z)$  proves (25).  $\square$

**Definition 5.36.** For  $k \in \mathbb{R}$ , define the weight  $k$  Serre derivative  $\partial_k$  of a modular form  $F$  as

$$\partial_k F := F' - \frac{k}{12} E_2 F.$$

**Theorem 5.37.** Let  $F$  be a modular form of weight  $k$  and level  $\Gamma$ . Then,  $\partial_k F$  is a modular form of weight  $k+2$  of the same level.

*Proof.* Let  $G = \partial_k F = F' - \frac{k}{12} E_2 F$ . It is enough to show that  $G$  is invariant under  $|_{k+2}\gamma$  for  $\gamma \in \Gamma$ . From  $F \in M_k(\Gamma)$ , we have

$$(F|_k \gamma)(z) := (cz+d)^{-k} F\left(\frac{az+b}{cz+d}\right) = F(z), \quad \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma.$$

By taking the derivative of the above equation, we get

$$\begin{aligned} & -kc(cz+d)^{-k-1} F\left(\frac{az+b}{cz+d}\right) + (cz+d)^{-k} (cz+d)^{-2} \frac{dF}{dz}\left(\frac{az+b}{cz+d}\right) = \frac{dF}{dz}(z) \\ \Leftrightarrow & (cz+d)^{-k-2} F'\left(\frac{az+b}{cz+d}\right) = F'(z) - \frac{ikc}{2\pi(cz+d)} F(z). \end{aligned}$$

Combined with (25), we get

$$\begin{aligned} ((\partial_k F)|_{k+2}\gamma)(z) &= (cz+d)^{-k-2} \left( F'\left(\frac{az+b}{cz+d}\right) - \frac{k}{12} E_2 \left(\frac{az+b}{cz+d}\right) F\left(\frac{az+b}{cz+d}\right) \right) \\ &= F'(z) - \frac{ikc}{2\pi(cz+d)} F(z) - \frac{k}{12} \left( E_2 - \frac{6c}{\pi(cz+d)} \right) F(z) \\ &= F'(z) - \frac{k}{12} E_2(z) F(z) = (\partial_k F)(z) \end{aligned}$$

so  $\partial_k F \in M_{k+2}(\Gamma)$ .  $\square$

**Remark 5.38.** More generally, the following theorem holds: if  $F$  is a quasimodular form of weight  $k$  and depth  $s$ , then  $\partial_{k-s} F$  is a quasimodular form of weight  $k+2$  and depth  $\leq s$  of the same level. We will not prove this here.

**Theorem 5.39.** We have

$$E'_2 = \frac{E_2^2 - E_4}{12} \tag{26}$$

$$E'_4 = \frac{E_2 E_4 - E_6}{3} \tag{27}$$

$$E'_6 = \frac{E_2 E_6 - E_4^2}{2} \tag{28}$$

*Proof.* In terms of Serre derivatives, these are equivalent to

$$\begin{aligned}\partial_1 E_2 &= -\frac{1}{12} E_4 \\ \partial_4 E_4 &= -\frac{1}{3} E_6 \\ \partial_6 E_6 &= -\frac{1}{2} E_4^2\end{aligned}$$

By Theorem 5.37, all the Serre derivatives are, in fact, modular. To be precise, the modularity of  $\partial_4 E_4$  and  $\partial_6 E_6$  directly follows from Theorem 5.37, and that of  $\partial_1 E_2$  follows from (25). Differentiating and squaring then gives us the following:

$$\begin{aligned}E_2'|_4\gamma &= E_2' - \frac{ic}{\pi(cz+d)} E_2 - \frac{3c^2}{\pi^2(cz+d)^2} \\ E_2^2|_4\gamma &= E_2^2 - \frac{12ic}{\pi(cz+d)} E_2 - \frac{36c^2}{\pi^2(cz+d)^2}\end{aligned}\tag{29}$$

Hence, (26)  $- \frac{1}{12}$  (29) is a modular form of weight 4. Since  $\dim M_k(\Gamma_1) = 1$  for  $k = 4, 6, 8$ , they should be multiples of  $E_4, E_6, E_8$ , and the proportionality constants can be determined by observing the constant terms of  $q$ -expansions.  $\square$

**Corollary 5.40.**

$$\Delta' = E_2 \Delta.\tag{30}$$

*Proof.* By Ramanujan's formula (27) and (28),

$$\Delta' = \frac{3E_4^2 E_4' - 2E_6 E_6'}{1728} = \frac{1}{1728} \left( 3E_4^2 \cdot \frac{E_2 E_4 - E_6}{3} - 2E_6 \cdot \frac{E_2 E_6 - E_4^2}{2} \right) = \frac{E_2(E_4^3 - E_6^2)}{1728} = E_2 \Delta.$$

$\square$

Similar argument allow us to compute (Serre) derivatives of  $H_2, H_3, H_4$ .

**Proposition 5.41.** *We have*

$$\begin{aligned}H_2' &= \frac{1}{6} (H_2^2 + 2H_2 H_4 + E_2 H_2) \\ H_3' &= \frac{1}{6} (H_2^2 - H_4^2 + E_2 H_3) \\ H_4' &= -\frac{1}{6} (2H_2 H_4 + H_4^2 - E_2 H_4)\end{aligned}$$

or equivalently,

$$\partial_2 H_2 = \frac{1}{6} (H_2^2 + 2H_2 H_4)\tag{31}$$

$$\partial_2 H_3 = \frac{1}{6} (H_2^2 - H_4^2)\tag{32}$$

$$\partial_2 H_4 = -\frac{1}{6}(2H_2 H_4 + H_4^2) \quad (33)$$

*Proof.* Equivalences are obvious from the definition of the Serre derivative. By Theorem 5.37, all the Serre derivatives are modular forms of weight 4 and level  $\Gamma(2)$ . We have  $\dim M_4(\Gamma(2)) = 3$  with basis  $H_2^2, H_2 H_4, H_4^2$ , and comparing the first three  $q$ -coefficients give (31), (32), and (33).  $\square$

**Theorem 5.42.** *The Serre derivative satisfies the following product rule: for any quasimodular forms  $F$  and  $G$ ,*

$$\partial_{w_1+w_2}(FG) = (\partial_{w_1}F)G + F(\partial_{w_2}G).$$

*Proof.* It follows from the definition:

$$\begin{aligned} \partial_{w_1+w_2}(FG) &= (FG)' - \frac{w_1+w_2}{12}E_2(FG) \\ &= F'G + FG' - \frac{w_1+w_2}{12}E_2(FG) \\ &= \left(F' - \frac{w_1}{12}E_2F\right)G + F\left(G' - \frac{w_2}{12}E_2G\right) \\ &= (\partial_{w_1}F)G + F(\partial_{w_2}G). \end{aligned}$$

$\square$

We also have the following useful theorem for proving positivity of quasimodular forms on the imaginary axis, which is [7, Proposition 3.5, Corollary 3.6].

**Theorem 5.43.** *Let  $F$  be a holomorphic quasimodular cusp form with real Fourier coefficients. Assume that there exists  $k$  such that  $(\partial_k F)(it) > 0$  for all  $t > 0$ . If the first Fourier coefficient of  $F$  is positive, then  $F(it) > 0$  for all  $t > 0$ .*

*Proof.* By (30), we have

$$\begin{aligned} \frac{d}{dt} \left( \frac{F(it)}{\Delta(it)^{\frac{k}{12}}} \right) &= (-2\pi) \frac{F'(it)\Delta(it)^{\frac{k}{12}} - F(it)\frac{k}{12}E_2(it)\Delta(it)^{\frac{k}{12}}}{\Delta(it)^{\frac{k}{6}}} \\ &= (-2\pi) \frac{(\partial_k F)(it)}{\Delta(it)^{\frac{k}{12}}} < 0, \end{aligned}$$

hence

$$t \mapsto \frac{F(it)}{\Delta(it)^{\frac{k}{12}}}$$

is monotone decreasing. Because of the assumption on the positivity of the first nonzero Fourier coefficient of  $F$ ,  $F(it) > 0$  for sufficiently large  $t$  since

$$F = \sum_{n \geq n_0} a_n q^n \Rightarrow e^{2\pi n_0 t} F(it) = a_{n_0} + e^{-2\pi t} \sum_{n \geq n_0+1} a_n e^{-2\pi(n-n_0-1)t}$$

and  $\lim_{t \rightarrow \infty} e^{2\pi n_0 t} F(it) = a_{n_0} > 0$ , hence the result follows.  $\square$

**Definition 5.44.** A weakly-holomorphic modular form of integer weight  $k$  and congruence subgroup  $\Gamma$  is a holomorphic function  $f : \mathfrak{H} \rightarrow \mathbb{C}$  such that:

1.  $f|_k \gamma = f$  for all  $\gamma \in \Gamma$
2. for each  $\alpha \in \Gamma_1$   $f|_k \alpha$  has Fourier expansion  $f|_k \alpha(z) = \sum_{n=n_0}^{\infty} c_f(\alpha, \frac{n}{n_\alpha}) e^{2\pi i \frac{n}{n_\alpha} z}$  for some  $n_0 \in \mathbb{Z}$  and  $n_\alpha \in \mathbb{N}$ .

For an  $m$ -periodic holomorphic function  $f$  and  $n \in \frac{1}{m}\mathbb{Z}$ , we will denote the  $n$ -th Fourier coefficient of  $f$  by  $c_f(n)$ , so that

$$f(z) = \sum_{n \in \frac{1}{m}\mathbb{Z}} c_f(n) e^{2\pi i n z}.$$

We denote the space of weakly-holomorphic modular forms of weight  $k$  and group  $\Gamma$  by  $M_k^!(\Gamma)$ . The spaces  $M_k^!(\Gamma)$  are infinite dimensional. Probably the most famous weakly-holomorphic modular form is the *elliptic  $j$ -invariant*

$$j := \frac{1728 E_4^3}{E_4^3 - E_6^2}.$$

This function belongs to  $M_0^!(\Gamma_1)$  and has the Fourier expansion

$$j(z) = q^{-1} + 744 + 196884q + 21493760q^2 + 864299970q^3 + 20245856256q^4 + O(q^5)$$

where  $q = e^{2\pi i z}$ . Using a simple computer algebra system, such as PARI GP or Mathematica, one can compute first hundred terms of this Fourier expansion in just a few seconds. An important question is to find an asymptotic formula for  $c_j(n)$ , the  $n$ -th Fourier coefficient of  $j$ . Using the Hardy-Ramanujan circle method [10] or the non-holomorphic Poincare series [9], one can show that

**Lemma 5.45.**

$$c_j(n) = \frac{2\pi}{n} \sum_{k=1}^{\infty} \frac{A_k(n)}{k} I_1\left(\frac{4\pi\sqrt{n}}{k}\right) \quad n \in \mathbb{Z}_{>0}$$

where

$$A_k(n) = \sum_{\substack{h \bmod k \\ (h,k)=1}} e^{\frac{-2\pi i}{k}(nh+h')}, \quad hh' \equiv -1 \pmod{k},$$

and  $I_\alpha(x)$  denotes the modified Bessel function of the first kind defined as in [1, Section 9.6].

A similar convergent asymptotic expansion holds for the Fourier coefficients of any weakly holomorphic modular form [5], [2, Propositions 1.10 and 1.12]. Such a convergent expansion implies effective estimates for the Fourier coefficients.

## 6 Fourier eigenfunctions with double zeroes at lattice points

In this section we construct two radial Schwartz functions  $a, b : \mathbb{R}^8 \rightarrow i\mathbb{R}$  such that

$$\mathcal{F}(a) = a \tag{34}$$

$$\mathcal{F}(b) = -b \tag{35}$$

which double zeroes at all  $\Lambda_8$ -vectors of length greater than  $\sqrt{2}$ . Recall that each vector of  $\Lambda_8$  has length  $\sqrt{2n}$  for some  $n \in \mathbb{N}_{\geq 0}$ . We define  $a$  and  $b$  so that their values are purely imaginary because this simplifies some of our computations. We will show in Section 7 that an appropriate linear combination of functions  $a$  and  $b$  satisfies conditions (5)–(7).

First, we will define function  $a$ . To this end we consider the following functions:

**Definition 6.1.**

$$\begin{aligned} \phi_{-4} &:= \frac{E_4^2}{\Delta} \\ \phi_{-2} &:= \frac{E_4(E_2E_4 - E_6)}{\Delta} \\ \phi_0 &:= \frac{(E_2E_4 - E_6)^2}{\Delta} \end{aligned}$$

**Lemma 6.2.** *These functions have the Fourier expansions*

$$\phi_{-4}(z) = q^{-1} + 504 + 73764q + 2695040q^2 + 54755730q^3 + O(q^4) \tag{36}$$

$$\phi_{-2}(z) = 720 + 203040q + 9417600q^2 + 223473600q^3 + 3566782080q^4 + O(q^5)$$

$$\phi_0(z) = 518400q + 31104000q^2 + 870912000q^3 + 15697152000q^4 + O(q^5) \tag{37}$$

where  $q = e^{2\pi iz}$ .

The function  $\phi_0(z)$  is not modular; however,

**Lemma 6.3.** *The identity 5.35 implies the following transformation rule:*

$$\phi_0\left(\frac{-1}{z}\right) = \phi_0(z) - \frac{12i}{\pi} \frac{1}{z} \phi_{-2}(z) - \frac{36}{\pi^2} \frac{1}{z^2} \phi_{-4}(z). \tag{38}$$

**Definition 6.4.** *For  $x \in \mathbb{R}^8$  we define*

$$\begin{aligned} a(x) &:= \int_{-1}^i \phi_0\left(\frac{-1}{z+1}\right) (z+1)^2 e^{\pi i \|x\|^2 z} dz + \int_1^i \phi_0\left(\frac{-1}{z-1}\right) (z-1)^2 e^{\pi i \|x\|^2 z} dz \\ &\quad - 2 \int_0^i \phi_0\left(\frac{-1}{z}\right) z^2 e^{\pi i \|x\|^2 z} dz + 2 \int_i^{i\infty} \phi_0(z) e^{\pi i \|x\|^2 z} dz. \end{aligned} \tag{39}$$

We observe that the contour integrals in (39) converge absolutely and uniformly for  $x \in \mathbb{R}^8$ . Indeed,  $\phi_0(z) = O(e^{-2\pi iz})$  as  $\Im(z) \rightarrow \infty$ . Therefore,  $a(x)$  is well defined. Now we prove that  $a$  satisfies condition (34).

**Proposition 6.5.**

*The function  $a$  defined by (39) belongs to the Schwartz space and satisfies*

$$\widehat{a}(x) = a(x).$$

*Proof.* First, we prove that  $a$  is a Schwartz function. From Lemma 5.17, Definition 5.34, and 5.45 we deduce that the Fourier coefficients of  $\phi_0$  satisfy

$$|c_{\phi_0}(n)| \leq 2 e^{4\pi\sqrt{n}} \quad n \in \mathbb{Z}_{>0}.$$

Thus, there exists a positive constant  $C$  such that

$$|\phi_0(z)| \leq C e^{-2\pi\Im z} \quad \text{for } \Im z > \frac{1}{2}.$$

We estimate the first summand in the right-hand side of (39). For  $r \in \mathbb{R}_{\geq 0}$  we have

$$\begin{aligned} \left| \int_{-1}^i \phi_0\left(\frac{-1}{z+1}\right) (z+1)^2 e^{\pi i r^2 z} dz \right| &= \left| \int_{i\infty}^{-1/(i+1)} \phi_0(z) z^{-4} e^{\pi i r^2 (-1/z-1)} dz \right| \leq \\ &C_1 \int_{1/2}^{\infty} e^{-2\pi t} e^{-\pi r^2/t} dt \leq C_1 \int_0^{\infty} e^{-2\pi t} e^{-\pi r^2/t} dt = C_2 r K_1(2\sqrt{2}\pi r) \end{aligned}$$

where  $C_1$  and  $C_2$  are some positive constants and  $K_\alpha(x)$  is the modified Bessel function of the second kind defined as in [1, Section 9.6]. This estimate also holds for the second and third summand in (39).

For the last summand we have

$$\left| \int_i^{i\infty} \phi_0(z) e^{\pi i r^2 z} dz \right| \leq C \int_1^{\infty} e^{-2\pi t} e^{-\pi r^2 t} dt = C_3 \frac{e^{\pi(r^2+2)}}{r^2+2}.$$

Therefore, we arrive at

$$|a(r)| \leq 4C_2 r K_1(2\sqrt{2}\pi r) + 2C_3 \frac{e^{-\pi(r^2+2)}}{r^2+2}.$$

It is easy to see that the left hand side of this inequality decays faster than any inverse power of  $r$ . Analogous estimates can be obtained for all derivatives  $\frac{d^k}{dr^k} a(r)$ .

Now we show that  $a$  is an eigenfunction of the Fourier transform. We recall that the Fourier transform



of a Gaussian function is

$$\mathcal{F}(e^{\pi i \|x\|^2 z})(y) = z^{-4} e^{\pi i \|y\|^2 (\frac{-1}{z})}. \quad (40)$$

Next, we exchange the contour integration with respect to  $z$  variable and Fourier transform with respect to  $x$  variable in (39). This can be done, since the corresponding double integral converges absolutely. In this way we obtain

$$\begin{aligned} \widehat{a}(y) &= \int_{-1}^i \phi_0\left(\frac{-1}{z+1}\right) (z+1)^2 z^{-4} e^{\pi i \|y\|^2 (\frac{-1}{z})} dz + \int_1^i \phi_0\left(\frac{-1}{z-1}\right) (z-1)^2 z^{-4} e^{\pi i \|y\|^2 (\frac{-1}{z})} dz \\ &\quad - 2 \int_0^i \phi_0\left(\frac{-1}{z}\right) z^2 z^{-4} e^{\pi i \|y\|^2 (\frac{-1}{z})} dz + 2 \int_i^{i\infty} \phi_0(z) z^{-4} e^{\pi i \|y\|^2 (\frac{-1}{z})} dz. \end{aligned}$$

Now we make a change of variables  $w = \frac{-1}{z}$ . We obtain

$$\begin{aligned} \widehat{a}(y) &= \int_1^i \phi_0\left(1 - \frac{1}{w-1}\right) \left(\frac{-1}{w} + 1\right)^2 w^2 e^{\pi i \|y\|^2 w} dw \\ &\quad + \int_{-1}^i \phi_0\left(1 - \frac{1}{w+1}\right) \left(\frac{-1}{w} - 1\right)^2 w^2 e^{\pi i \|y\|^2 w} dw \\ &\quad - 2 \int_{i\infty}^i \phi_0(w) e^{\pi i \|y\|^2 w} dw + 2 \int_i^0 \phi_0\left(\frac{-1}{w}\right) w^2 e^{\pi i \|y\|^2 w} dw. \end{aligned}$$

Since  $\phi_0$  is 1-periodic we have

$$\begin{aligned} \widehat{a}(y) &= \int_1^i \phi_0\left(\frac{-1}{z-1}\right) (z-1)^2 e^{\pi i \|y\|^2 z} dz + \int_{-1}^i \phi_0\left(\frac{-1}{z+1}\right) (z+1)^2 e^{\pi i \|y\|^2 z} dz \\ &\quad + 2 \int_i^{i\infty} \phi_0(z) e^{\pi i \|y\|^2 z} dz - 2 \int_0^i \phi_0\left(\frac{-1}{z}\right) z^2 e^{\pi i \|y\|^2 z} dz \\ &= a(y). \end{aligned}$$

This finishes the proof of the proposition. □

Next, we check that  $a$  has double zeroes at all  $\Lambda_8$ -lattice points of length greater than  $\sqrt{2}$ .

**Proposition 6.6.** *For  $r > \sqrt{2}$  we can express  $a(r)$  in the following form*

$$a(r) = -4 \sin(\pi r^2/2)^2 \int_0^{i\infty} \phi_0\left(\frac{-1}{z}\right) z^2 e^{\pi i r^2 z} dz. \quad (41)$$

*Proof.* We denote the right hand side of (41) by  $d(r)$ . It is easy to see that  $d(r)$  is well-defined. Indeed,

from the transformation formula (38) and the expansions (37)–(36) we obtain

$$\begin{aligned}\phi_0\left(\frac{-1}{it}\right) &= O(e^{-2\pi/t}) \quad \text{as } t \rightarrow 0 \\ \phi_0\left(\frac{-1}{it}\right) &= O(t^{-2} e^{2\pi t}) \quad \text{as } t \rightarrow \infty\end{aligned}$$

Hence, the integral (41) converges absolutely for  $r > \sqrt{2}$ . We can write

$$\begin{aligned}d(r) &= \int_{-1}^{i\infty-1} \phi_0\left(\frac{-1}{z+1}\right) (z+1)^2 e^{\pi i r^2 z} dz - 2 \int_0^{i\infty} \phi_0\left(\frac{-1}{z}\right) z^2 e^{\pi i r^2 z} dz \\ &\quad + \int_1^{i\infty+1} \phi_0\left(\frac{-1}{z-1}\right) (z-1)^2 e^{\pi i r^2 z} dz.\end{aligned}$$

From (38) we deduce that if  $r > \sqrt{2}$  then  $\phi_0\left(\frac{-1}{z}\right) z^2 e^{\pi i r^2 z} \rightarrow 0$  as  $\Im(z) \rightarrow \infty$ . Therefore, we can deform the paths of integration and rewrite

$$\begin{aligned}d(r) &= \int_{-1}^i \phi_0\left(\frac{-1}{z+1}\right) (z+1)^2 e^{\pi i r^2 z} dz + \int_i^{i\infty} \phi_0\left(\frac{-1}{z+1}\right) (z+1)^2 e^{\pi i r^2 z} dz \\ &\quad - 2 \int_0^i \phi_0\left(\frac{-1}{z}\right) z^2 e^{\pi i r^2 z} dz - 2 \int_i^{i\infty} \phi_0\left(\frac{-1}{z}\right) z^2 e^{\pi i r^2 z} dz \\ &\quad + \int_1^i \phi_0\left(\frac{-1}{z-1}\right) (z-1)^2 e^{\pi i r^2 z} dz + \int_i^{i\infty} \phi_0\left(\frac{-1}{z-1}\right) (z-1)^2 e^{\pi i r^2 z} dz.\end{aligned}$$

Now from (38) we find

$$\begin{aligned}&\phi_0\left(\frac{-1}{z+1}\right) (z+1)^2 - 2\phi_0\left(\frac{-1}{z}\right) z^2 + \phi_0\left(\frac{-1}{z-1}\right) (z-1)^2 = \\ &\phi_0(z+1)(z+1)^2 - 2\phi_0(z)z^2 + \phi_0(z-1)(z-1)^2 \\ &- \frac{12i}{\pi} \left( \phi_{-2}(z+1)(z+1) - 2\phi_{-2}(z)z + \phi_{-2}(z-1)(z-1) \right) \\ &- \frac{36}{\pi^2} \left( \phi_{-4}(z+1) - 2\phi_{-4}(z) + \phi_{-4}(z-1) \right) = \\ &2\phi_0(z).\end{aligned}$$

Thus, we obtain

$$\begin{aligned}d(r) &= \int_{-1}^i \phi_0\left(\frac{-1}{z+1}\right) (z+1)^2 e^{\pi i r^2 z} dz - 2 \int_0^i \phi_0\left(\frac{-1}{z}\right) z^2 e^{\pi i r^2 z} dz \\ &\quad + \int_1^i \phi_0\left(\frac{-1}{z-1}\right) (z-1)^2 e^{\pi i r^2 z} dz + 2 \int_i^{i\infty} \phi_0(z) e^{\pi i r^2 z} dz = a(r).\end{aligned}$$

This finishes the proof.  $\square$

Finally, we find another convenient integral representation for  $a$  and compute values of  $a(r)$  at  $r = 0$  and  $r = \sqrt{2}$ .

**Proposition 6.7.** *For  $r \geq 0$  we have*

$$\begin{aligned} a(r) = & 4i \sin(\pi r^2/2)^2 \left( \frac{36}{\pi^3 (r^2 - 2)} - \frac{8640}{\pi^3 r^4} + \frac{18144}{\pi^3 r^2} \right. \\ & \left. + \int_0^\infty \left( t^2 \phi_0\left(\frac{i}{t}\right) - \frac{36}{\pi^2} e^{2\pi t} + \frac{8640}{\pi} t - \frac{18144}{\pi^2} \right) e^{-\pi r^2 t} dt \right). \end{aligned} \quad (42)$$

The integral converges absolutely for all  $r \in \mathbb{R}_{\geq 0}$ .

*Proof.* Suppose that  $r > \sqrt{2}$ . Then by Proposition 6.6

$$a(r) = 4i \sin(\pi r^2/2)^2 \int_0^\infty \phi_0(i/t) t^2 e^{-\pi r^2 t} dt.$$

From (37)–(38) we obtain

$$\phi_0(i/t) t^2 = \frac{36}{\pi^2} e^{2\pi t} - \frac{8640}{\pi} t + \frac{18144}{\pi^2} + O(t^2 e^{-2\pi t}) \quad \text{as } t \rightarrow \infty. \quad (43)$$

For  $r > \sqrt{2}$  we have

$$\int_0^\infty \left( \frac{36}{\pi^2} e^{2\pi t} + \frac{8640}{\pi} t + \frac{18144}{\pi^2} \right) e^{-\pi r^2 t} dt = \frac{36}{\pi^3 (r^2 - 2)} - \frac{8640}{\pi^3 r^4} + \frac{18144}{\pi^3 r^2}.$$

Therefore, the identity (42) holds for  $r > \sqrt{2}$ .

On the other hand, from the definition (39) we see that  $a(r)$  is analytic in some neighborhood of  $[0, \infty)$ . The asymptotic expansion (43) implies that the right hand side of (42) is also analytic in some neighborhood of  $[0, \infty)$ . Hence, the identity (42) holds on the whole interval  $[0, \infty)$ . This finishes the proof of the proposition.  $\square$

From the identity (42) we see that the values  $a(r)$  are in  $i\mathbb{R}$  for all  $r \in \mathbb{R}_{\geq 0}$ . In particular, we have

**Proposition 6.8.** *We have*

$$a(0) = \frac{-i 8640}{\pi} \quad a(\sqrt{2}) = 0 \quad a'(\sqrt{2}) = \frac{i 72\sqrt{2}}{\pi}.$$

*Proof.* These identities follow immediately from the previous proposition.  $\square$

Now we construct function  $b$ . To this end we consider the function

**Definition 6.9.**

$$h(z) := 128 \frac{H_3(z) + H_4(z)}{H_2(z)^2}. \quad (44)$$

It is easy to see that  $h \in M_{-2}^1(\Gamma_0(2))$ . Indeed, first we check that  $h|_{-2}\gamma = h$  for all  $\gamma \in \Gamma_0(2)$ . Since the group  $\Gamma_0(2)$  is generated by elements  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  and  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  it suffices to check that  $h$  is invariant under their action. This follows immediately from (14)–(16) and (44). Next we analyze the poles of  $h$ . It is known [8, Chapter I Lemma 4.1] that  $\theta_{10}$  has no zeros in the upper-half plane and hence  $h$  has poles only at the cusps. At the cusp  $i\infty$  this modular form has the Fourier expansion

$$h(z) = q^{-1} + 16 - 132q + 640q^2 - 2550q^3 + O(q^4).$$

Let  $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ ,  $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ , and  $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$  be elements of  $\Gamma_1$ .

**Definition 6.10.** We define the followig three functions

$$\psi_I := h - h|_{-2}ST \quad (45)$$

$$\psi_T := \psi_I|_{-2}T$$

$$\psi_S := \psi_I|_{-2}S. \quad (46)$$

**Lemma 6.11.** More explicitly, we have

$$\begin{aligned} \psi_I(z) &= 128 \frac{H_3(z) + H_4(z)}{H_2(z)^2} + 128 \frac{H_4(z) - H_2(z)}{H_3(z)^2} \\ \psi_T(z) &= 128 \frac{H_3(z) + H_4(z)}{H_2(z)^2} + 128 \frac{H_2(z) + H_3(z)}{H_4(z)^2} \\ \psi_S(z) &= 128 \frac{H_2(z) + H_3(z)}{H_4(z)^2} - 128 \frac{H_2(z) - H_4(z)}{H_3(z)^2} \end{aligned} \quad (47)$$

**Lemma 6.12.** The Fourier expansions of these functions are

$$\begin{aligned} \psi_I(z) &= q^{-1} + 144 - 5120q^{1/2} + 70524q - 626688q^{3/2} + 4265600q^2 + O(q^{5/2}) \\ \psi_T(z) &= q^{-1} + 144 + 5120q^{1/2} + 70524q + 626688q^{3/2} + 4265600q^2 + O(q^{5/2}) \\ \psi_S(z) &= -10240q^{1/2} - 1253376q^{3/2} - 48328704q^{5/2} - 1059078144q^{7/2} + O(q^{9/2}). \end{aligned} \quad (48)$$

**Definition 6.13.** For  $x \in \mathbb{R}^8$  define

$$\begin{aligned} b(x) &:= \int_{-1}^i \psi_T(z) e^{\pi i \|x\|^2 z} dz + \int_1^i \psi_T(z) e^{\pi i \|x\|^2 z} dz \\ &\quad - 2 \int_0^i \psi_I(z) e^{\pi i \|x\|^2 z} dz - 2 \int_i^{i\infty} \psi_S(z) e^{\pi i \|x\|^2 z} dz. \end{aligned} \quad (49)$$

Now we prove that  $b$  satisfies condition (35).

**Proposition 6.14.** *The function  $b$  defined by (49) belongs to the Schwartz space and satisfies*

$$\widehat{b}(x) = -b(x).$$

*Proof.* Here, we repeat the arguments used in the proof of Proposition 6.5. First we show that  $b$  is a Schwartz function. We have

$$\begin{aligned} \int_{-1}^i \psi_T(z) e^{\pi i r^2 z} dz &= \int_0^{i+1} \psi_I(z) e^{\pi i r^2 (z-1)} dz = \\ &= \int_{i\infty}^{-1/(i+1)} \psi_I\left(\frac{-1}{z}\right) e^{\pi i r^2 (-1/z-1)} z^{-2} dz = \int_{i\infty}^{-1/(i+1)} \psi_S(z) z^{-4} e^{\pi i r^2 (-1/z-1)} dz. \end{aligned}$$

There exists a positive constant  $C$  such that

$$|\psi_S(z)| \leq C e^{-\pi \Im z} \quad \text{for } \Im z > \frac{1}{2}.$$

Thus, as in the proof of Proposition 6.5 we estimate the first summand in the left-hand side of (49)

$$\left| \int_{-1}^i \psi_T(z) e^{\pi i r^2 z} dz \right| \leq C_1 r K_1(2\pi r).$$

We combine this inequality with analogous estimates for the other three summands and obtain

$$|b(r)| \leq C_2 r K_1(2\pi r) + C_3 \frac{e^{-\pi(r^2+1)}}{r^2+1}.$$

Here  $C_1$ ,  $C_2$ , and  $C_3$  are some positive constants. Similar estimates hold for all derivatives  $\frac{d^k}{dr} b(r)$ .

Now we prove that  $b$  is an eigenfunction of the Fourier transform. We use identity (40) and change contour integration in  $z$  and Fourier transform in  $x$ . Thus we obtain

$$\begin{aligned} \mathcal{F}(b)(x) &= \int_{-1}^i \psi_T(z) z^{-4} e^{\pi i \|x\|^2 (\frac{-1}{z})} dz + \int_1^i \psi_T(z) z^{-4} e^{\pi i \|x\|^2 (\frac{-1}{z})} dz \\ &\quad - 2 \int_0^i \psi_I(z) z^{-4} e^{\pi i \|x\|^2 (\frac{-1}{z})} dz - 2 \int_i^{i\infty} \psi_S(z) z^{-4} e^{\pi i \|x\|^2 (\frac{-1}{z})} dz. \end{aligned}$$

We make the change of variables  $w = \frac{-1}{z}$  and arrive at

$$\mathcal{F}(b)(x) = \int_1^i \psi_T\left(\frac{-1}{w}\right) w^2 e^{\pi i \|x\|^2 w} dw + \int_{-1}^i \psi_T\left(\frac{-1}{w}\right) w^2 e^{\pi i \|x\|^2 w} dw$$

$$-2 \int_{i\infty}^i \psi_I\left(\frac{-1}{w}\right) w^2 e^{\pi i \|x\|^2 w} dw - 2 \int_i^0 \psi_S\left(\frac{-1}{w}\right) w^2 e^{\pi i \|x\|^2 w} dw.$$

Now we observe that the definitions (45)–(46) imply

$$\psi_T|_{-2S} = -\psi_T$$

$$\psi_I|_{-2S} = \psi_S$$

$$\psi_S|_{-2S} = \psi_I.$$

Therefore, we arrive at

$$\begin{aligned} \mathcal{F}(b)(x) &= \int_1^i -\psi_T(z) e^{\pi i \|x\|^2 z} dz + \int_{-1}^i -\psi_T(z) e^{\pi i \|x\|^2 z} dz \\ &\quad + 2 \int_i^{i\infty} \psi_S(z) e^{\pi i \|x\|^2 z} dz + 2 \int_0^i \psi_I(z) e^{\pi i \|x\|^2 w} dw. \end{aligned}$$

Now from (49) we see that

$$\mathcal{F}(b)(x) = -b(x).$$

□

Now we regard the radial function  $b$  as a function on  $\mathbb{R}_{\geq 0}$ . We check that  $b$  has double roots at  $\Lambda_8$ -points.

**Proposition 6.15.** *For  $r > \sqrt{2}$  function  $b(r)$  can be expressed as*

$$b(r) = -4 \sin(\pi r^2/2)^2 \int_0^{i\infty} \psi_I(z) e^{\pi i r^2 z} dz. \quad (50)$$

*Proof.* We denote the right hand side of (50) by  $c(r)$ . First, we check that  $c(r)$  is well-defined. We have

$$\psi_I(it) = O(t^2 e^{\pi/t}) \quad \text{as } t \rightarrow 0$$

$$\psi_I(it) = O(e^{2\pi t}) \quad \text{as } t \rightarrow \infty.$$

Therefore, the integral (50) converges for  $r > \sqrt{2}$ . Then we rewrite it in the following way:

$$c(r) = \int_{-1}^{i\infty-1} \psi_I(z+1) e^{\pi i r^2 z} dz - 2 \int_0^{i\infty} \psi_I(z) e^{\pi i r^2 z} dz + \int_1^{i\infty+1} \psi_I(z-1) e^{\pi i r^2 z} dz.$$

From the Fourier expansion (48) we know that  $\psi_I(z) = e^{-2\pi i z} + O(1)$  as  $\Im(z) \rightarrow \infty$ . By assumption

$r^2 > 2$ , hence we can deform the path of integration and write

$$\begin{aligned} \int_{-1}^{i\infty-1} \psi_I(z+1) e^{\pi i r^2 z} dz &= \int_{-1}^i \psi_T(z) e^{\pi i r^2 z} dz + \int_i^{i\infty} \psi_T(z) e^{\pi i r^2 z} dz \\ \int_1^{i\infty+1} \psi_I(z-1) e^{\pi i r^2 z} dz &= \int_{-1}^i \psi_T(z) e^{\pi i r^2 z} dz + \int_i^{i\infty} \psi_T(z) e^{\pi i r^2 z} dz. \end{aligned}$$

We have

$$\begin{aligned} c(r) &= \int_{-1}^i \psi_T(z) e^{\pi i r^2 z} dz + \int_1^i \psi_T(z) e^{\pi i r^2 z} dz - 2 \int_0^i \psi_I(z) e^{\pi i r^2 z} dz \\ &\quad + 2 \int_i^{i\infty} (\psi_T(z) - \psi_I(z)) e^{\pi i r^2 z} dz. \end{aligned} \tag{51}$$

Next, we check that the functions  $\psi_I, \psi_T$ , and  $\psi_S$  satisfy the following identity:

$$\psi_T + \psi_S = \psi_I. \tag{52}$$

Indeed, from definitions (45)-(46) we get

$$\begin{aligned} \psi_T + \psi_S &= (h - h|_{-2}ST)|_{-2}T + (h - h|_{-2}ST)|_{-2}S \\ &= h|_{-2}T - h|_{-2}ST^2 + h|_{-2}S - h|_{-2}STS. \end{aligned}$$

Note that  $ST^2S$  belongs to  $\Gamma_0(2)$ . Thus, since  $h \in M_{-2}^1\Gamma_0(2)$  we get

$$\psi_T + \psi_S = h|_{-2}T - h|_{-2}STS.$$

Now we observe that  $T$  and  $STS(ST)^{-1}$  are also in  $\Gamma_0(2)$ . Therefore,

$$\psi_T + \psi_S = h|_{-2}T - h|_{-2}STS = h|_{-2} - h|ST = \psi_I.$$

Combining (51) and (52) we find

$$\begin{aligned} c(r) &= \int_{-1}^i \psi_T(z) e^{\pi i r^2 z} dz + \int_1^i \psi_T(z) e^{\pi i r^2 z} dz - 2 \int_0^i \psi_I(z) e^{\pi i r^2 z} dz \\ &\quad - 2 \int_i^{i\infty} \psi_S(z) e^{\pi i r^2 z} dz \\ &= b(r). \end{aligned}$$

□

At the end of this section we find another integral representation of  $b(r)$  for  $r \in \mathbb{R}_{\geq 0}$  and compute special values of  $b$ .

**Proposition 6.16.** *For  $r \geq 0$  we have*

$$b(r) = 4i \sin(\pi r^2/2)^2 \left( \frac{144}{\pi r^2} + \frac{1}{\pi(r^2-2)} + \int_0^\infty (\psi_I(it) - 144 - e^{2\pi t}) e^{-\pi r^2 t} dt \right). \quad (53)$$

*The integral converges absolutely for all  $r \in \mathbb{R}_{\geq 0}$ .*

*Proof.* The proof is analogous to the proof of Proposition 6.7. First, suppose that  $r > \sqrt{2}$ . Then by Proposition 6.15

$$b(r) = 4i \sin(\pi r^2/2)^2 \int_0^\infty \psi_I(it) e^{-\pi r^2 t} dt.$$

From (48) we obtain

$$\psi_I(it) = e^{2\pi t} + 144 + O(e^{-\pi t}) \quad \text{as } t \rightarrow \infty. \quad (54)$$

For  $r > \sqrt{2}$  we have

$$\int_0^\infty (e^{2\pi t} + 144) e^{-\pi r^2 t} dt = \frac{1}{\pi(r^2-2)} + \frac{144}{\pi r^2}.$$

Therefore, the identity (53) holds for  $r > \sqrt{2}$ .

On the other hand, from the definition (49) we see that  $b(r)$  is analytic in some neighborhood of  $[0, \infty)$ . The asymptotic expansion (54) implies that the right hand side of (53) is also analytic in some neighborhood of  $[0, \infty)$ . Hence, the identity (53) holds on the whole interval  $[0, \infty)$ . This finishes the proof of the proposition. □

We see from (53) that  $b(r) \in i\mathbb{R}$  for all  $r \in \mathbb{R}_{\geq 0}$ . Another immediate corollary of this proposition is

**Proposition 6.17.** *We have*

$$b(0) = 0 \quad b(\sqrt{2}) = 0 \quad b'(\sqrt{2}) = \frac{i}{2\sqrt{2}\pi}.$$

## 7 Proof of Theorem 4.2

Our proof of the Theorem 4.2 relies on the following two inequalities for modular objects.



**Proposition 7.1.** Consider the function  $A : (0, \infty) \rightarrow \mathbb{C}$  defined as

$$A(t) := -t^2 \phi_0(i/t) - \frac{36}{\pi^2} \psi_I(it).$$

Then

$$A(t) < 0 \tag{55}$$

for all  $t > 0$ .

**Proposition 7.2.** Consider the function  $B : (0, \infty) \rightarrow \mathbb{C}$  defined as

$$B(t) := -t^2 \phi_0(i/t) + \frac{36}{\pi^2} \psi_I(it)$$

Then

$$B(t) > 0 \tag{56}$$

for all  $t > 0$ .

Here we formalize the proof of the inequalities by Lee [7]. First, we can rewrite the inequality in 7.1 as follows.

**Definition 7.3.** Define two (quasi) modular forms as

$$\begin{aligned} F(z) &= (E_2(z)E_4(z) - E_6(z))^2 \\ G(z) &= H_2(z)^3(2H_2(z)^2 + 5H_2(z)H_4(z) + 5H_4(z)^2). \end{aligned}$$

**Lemma 7.4.** We have

$$\phi_0 = \frac{F}{\Delta} \tag{57}$$

$$\psi_S = -\frac{1}{2} \frac{G}{\Delta} \tag{58}$$

*Proof.* (57) is clear. For (58), using (47) and (18) gives

$$\begin{aligned} \psi_S &= -128 \frac{H_3 + H_2}{H_4^2} - 128 \frac{H_2 - H_4}{H_3^2} \\ &= -128 \frac{H_3^2(H_2 - H_4) + H_4^2(H_2 - H_4)}{H_3^2 H_4^2} \\ &= -128 \frac{(H_2 + H_4)^2(2H_2 + H_4) + H_4^2(H_2 + H_4)}{H_3^2 H_4^2} \\ &= -128 \frac{H_2(2H_2^2 + 5H_2H_4 + 5H_4^2)}{H_3^2 H_4^2} \\ &= -128 \frac{H_2^3(2H_2^2 + 5H_2H_4 + 5H_4^2)}{H_2^2 H_3^2 H_4^2} \\ &= -\frac{1}{2} \frac{G}{\Delta}. \end{aligned}$$

□

**Lemma 7.5.** *Inequality (55) and (56) are equivalent to*

$$F(it) + \frac{18}{\pi^2} G(it) > 0 \quad (59)$$

$$F(it) - \frac{18}{\pi^2} G(it) > 0 \quad (60)$$

respectively.

*Proof.* By (46),

$$\psi_I(it) = (\psi_S|_{-2}S)(it) = (it)^2 \psi_S\left(-\frac{1}{it}\right) = -t^2 \psi_S\left(\frac{i}{t}\right).$$

Combined with Lemma 7.4 we can rewrite (55) as

$$A(t) = -t^2 \phi_0\left(\frac{i}{t}\right) + \frac{36}{\pi^2} \psi_S\left(\frac{i}{t}\right) < 0 \Leftrightarrow \frac{F(it)}{\Delta(it)} + \frac{18}{\pi^2} \frac{G(it)}{\Delta(it)} > 0$$

for  $t > 0$ , which is equivalent to (59) by Corollary 5.21. Equivalences of (56) and (60) follows similarly; just change the sign. □

Now, the first inequality (59) follows from the positivity of each  $F(it)$  and  $G(it)$ .

**Lemma 7.6.** *For all  $t > 0$ , we have  $F(it) > 0$  and  $G(it) > 0$ .*

*Proof.* By Ramanujan's identity (27), we have  $F(z) = 9E_4'(z)^2$  and

$$F(it) = 9E_4'(it)^2 = 9 \left( 240 \sum_{n \geq 1} n \sigma_3(n) e^{-2\pi n t} \right) > 0.$$

$G(it) > 0$  follows from positivity of  $H_2(it)$  and  $H_4(it)$  (Lemma 5.29). □

**Corollary 7.7.** (59) holds.

*Proof.* This directly follows from Lemma 7.6. □

To prove the second inequality (60), we need some identities satisfied by  $F$  and  $G$ .

**Lemma 7.8.**  *$F$  and  $G$  satisfy the following differential equations:*

$$\partial_{12} \partial_{10} F - \frac{5}{6} E_4 F = 7200 \Delta(-E_2') \quad (61)$$

$$\partial_{12} \partial_{10} G - \frac{5}{6} E_4 G = -640 \Delta H_2 \quad (62)$$

*Proof.* Both can be shown by direct computations. By Ramanujan's identities (Theorem 5.39) and the product rule of Serre derivatives (Theorem 5.42), we have

$$\begin{aligned}
\partial_5(E_2E_4 - E_6) &= (E_2E_4 - E_6)' - \frac{5}{12}E_2(E_2E_4 - E_6) \\
&= \frac{E_2^2 - E_4}{12} \cdot E_4 + E_2 \cdot \frac{E_2E_4 - E_6}{3} - \frac{E_2E_6 - E_4^2}{2} - \frac{5}{12}E_2(E_2E_4 - E_6) \\
&= -\frac{5}{12}(E_2E_6 - E_4^2) \\
\partial_7(E_2E_6 - E_4^2) &= (E_2E_6 - E_4^2)' - \frac{7}{12}E_2(E_2E_6 - E_4^2) \\
&= \frac{E_2^2 - E_4}{12} \cdot E_6 + E_2 \cdot \frac{E_2E_6 - E_4^2}{2} - 2E_4 \cdot \frac{E_2E_4 - E_6}{3} - \frac{7}{12}E_2(E_2E_6 - E_4^2) \\
&= -\frac{7}{12}E_4(E_2E_4 - E_6)
\end{aligned}$$

and using these we can compute

$$\begin{aligned}
\partial_{10}F &= \partial_{10}(E_2E_4 - E_6)^2 \\
&= 2(E_2E_4 - E_6)\partial_5(E_2E_4 - E_6) \\
&= -\frac{6}{5}(E_2E_4 - E_6)(E_2E_6 - E_4^2), \\
\partial_{12}\partial_{10}F &= -\frac{5}{6}\partial_{12}((E_2E_4 - E_6)(E_2E_6 - E_4^2)) \\
&= -\frac{5}{6}(\partial_5(E_2E_4 - E_6))(E_2E_6 - E_4^2) - \frac{5}{6}(E_2E_4 - E_6)(\partial_7(E_2E_6 - E_4^2)) \\
&= \frac{25}{72}(E_2E_6 - E_4^2)^2 + \frac{35}{72}E_4(E_2E_4 - E_6)^2, \\
\partial_{12}\partial_{10}F - \frac{5}{6}E_4F &= \frac{25}{72}(E_2E_6 - E_4^2)^2 + \frac{35}{72}E_4(E_2E_4 - E_6)^2 - \frac{5}{6}E_4(E_2E_4 - E_6)^2 \\
&= \frac{25}{72}((E_2E_6 - E_4^2)^2 - E_4(E_2E_4 - E_6)^2) \\
&= \frac{25}{72}(-E_2^2E_4^3 + E_2^2E_6^2 + E_4^4 - E_4E_6^3) \\
&= -\frac{25}{72}(E_4^3 - E_6^2)(E_2^2 - E_4) \\
&= 7200 \cdot \frac{E_4^3 - E_6^2}{1728} \cdot \frac{-E_2^2 + E_4}{12} \\
&= 7200\Delta(-E_2')
\end{aligned}$$

which proves (61). Similarly, (62) can be proved using Proposition 5.41 and Lemma 5.28.  $\square$

**Corollary 7.9.** (61) (resp. (62)) is positive (resp. negative) on the (positive) imaginary axis.

*Proof.*  $\square$

The second inequality (60) follows from the following two observations. Since  $G(it) > 0$  for all  $t > 0$ ,

we can define the quotient

$$Q(t) := \frac{F(it)}{G(it)}$$

as a function on  $(0, \infty)$ .

**Lemma 7.10.** *We have*

$$\lim_{t \rightarrow 0^+} Q(t) = \frac{18}{\pi^2}.$$

*Proof.* We have

$$\lim_{t \rightarrow 0^+} Q(t) = \lim_{t \rightarrow 0^+} \frac{F(it)}{G(it)} = \lim_{t \rightarrow \infty} \frac{F(i/t)}{G(i/t)}.$$

By using the transformation laws of Eisenstein series (24), (9) (for  $k = 4, 6$ ) and the thetanull functions, (14), (16), we get

$$\begin{aligned} F\left(\frac{i}{t}\right) &= t^{12}F(it) - \frac{12t^{11}}{\pi}(E_2(it)E_4(it) - E_6(it))E_4(it) + \frac{36t^{10}}{\pi^2}E_4(it)^2, \\ G\left(\frac{i}{t}\right) &= t^{10}H_4(it)^3(2H_4(it)^2 + 5H_4(it)H_2(it) + 5H_2(it)^2). \end{aligned}$$

Since  $F$ ,  $E_2E_4 - E_6$  and  $H_2$  are cusp forms, we have  $\lim_{t \rightarrow \infty} t^k A(it) = 0$  when  $A(z)$  is one of these forms and  $k \geq 0$ . From  $\lim_{t \rightarrow \infty} E_4(it) = 1 = \lim_{t \rightarrow \infty} H_4(it)$ , we get

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{F(i/t)}{G(i/t)} &= \lim_{t \rightarrow \infty} \frac{t^{12}F(it) - \frac{12t^{11}}{\pi}(E_2(it)E_4(it) - E_6(it))E_4(it) + \frac{36t^{10}}{\pi^2}E_4(it)^2}{t^{10}H_4(it)^3(2H_4(it)^2 + 5H_4(it)H_2(it) + 5H_2(it)^2)} \\ &= \lim_{t \rightarrow \infty} \frac{t^2F(it) - \frac{12t}{\pi}(E_2(it)E_4(it) - E_6(it))E_4(it) + \frac{36}{\pi^2}E_4(it)^2}{H_4(it)^3(2H_4(it)^2 + 5H_4(it)H_2(it) + 5H_2(it)^2)} \\ &= \frac{18}{\pi^2}. \end{aligned}$$

□

**Proposition 7.11.** *The function  $t \mapsto Q(t)$  is monotone decreasing.*

*Proof.* It is enough to show that

$$\begin{aligned} \frac{d}{dt} \left( \frac{F(it)}{G(it)} \right) < 0 &\Leftrightarrow (-2\pi) \frac{F'(it)G(it) - F(it)G'(it)}{G(it)^2} < 0 \\ &\Leftrightarrow F'(it)G(it) - F(it)G'(it) > 0 \\ &\Leftrightarrow (\partial_{10}F)(it)G(it) - F(it)(\partial_{10}G)(it) > 0. \end{aligned}$$

Let  $\mathcal{L}_{1,0} := (\partial_{10}F)G - F(\partial_{10}G)$ . Then its Fourier expansion starts with

$$\mathcal{L}_{1,0} = 5308416000q^{\frac{7}{2}} + O(q^{\frac{9}{2}})$$

and its Serre derivative  $\partial_{22}\mathcal{L}_{1,0}$  is positive by Corollary 7.9:

$$\partial_{22}\mathcal{L}_{1,0} = (\partial_{12}\partial_{10}F)G - F(\partial_{12}\partial_{10}G) = \Delta(7200(-E_2')G + 640H_2F) > 0.$$

Hence  $\mathcal{L}_{1,0}(it) > 0$  by Theorem 5.43, and the monotonicity follows.  $\square$

**Corollary 7.12.** (60) holds.

*Proof.*

$$\frac{F(it)}{G(it)} = Q(t) < \lim_{u \rightarrow 0^+} Q(u) = \frac{18}{\pi^2}$$

and by Lemma 7.6, (60) follows.  $\square$

Finally, we are ready to prove Theorem 4.2.

**Theorem 7.13.** The function

$$g(x) := \frac{\pi i}{8640} a(x) + \frac{i}{240\pi} b(x)$$

satisfies conditions (5)–(7).

*Proof.* First, we prove that (5) holds. By Propositions 6.6 and 6.15 we know that for  $r > \sqrt{2}$

$$g(r) = \frac{\pi}{2160} \sin(\pi r^2/2)^2 \int_0^\infty A(t) e^{-\pi r^2 t} dt \quad (63)$$

where

$$A(t) = -t^2 \phi_0(i/t) - \frac{36}{\pi^2} \psi_I(it).$$

from the Proposition 7.1 we know that  $A(t) < 0$  for  $t \in (0, \infty)$ . Therefore identity (63) implies (5).

Next, we prove (6). By Propositions 6.7 and 6.16 we know that for  $r > 0$

$$\widehat{g}(r) = \frac{\pi}{2160} \sin(\pi r^2/2)^2 \int_0^\infty B(t) e^{-\pi r^2 t} dt$$

where

$$B(t) = -t^2 \phi_0(i/t) + \frac{36}{\pi^2} \psi_I(it).$$

Finally, the property (7) readily follows from Proposition 6.8 and Proposition 6.17. This finishes the proof of Theorems 7.13 and 4.2.  $\square$

## References

- [1] M. ABRAMOWITZ, I. STEGUN, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Applied Mathematics Series 55 (10 ed.), New York, USA: United States Department of Commerce, National Bureau of Standards; Dover Publications, 1964.
- [2] J. BRUINIER, *Borcherds products on  $O(2,1)$  and Chern classes of Heegner divisors*, Springer Lecture Notes in Mathematics 1780 (2002)
- [3] H. COHN, N. ELKIES, *New upper bounds on sphere packings I*, Annals of Math. 157 (2003) pp. 689–714.
- [4] F. DIAMOND, J. SHURMAN, *A First Course in Modular Forms*, Springer New York, 2005.
- [5] D. HEJHAL, *The Selberg trace formula for  $\mathrm{PSL}(2, \mathbb{R})$* , Springer Lecture Notes in Mathematics 1001 (1983)
- [6] W. KOHNEN, *A Very Simple Proof of the  $q$ -Product Expansion of the  $\Delta$ -Function*, The Ramanujan Journal 10 (2005): 71–73.
- [7] S. LEE, *Algebraic proof of modular form inequalities for optimal sphere packings*, arXiv preprint arXiv:2406.14659 (2024).
- [8] D. MUMFORD, *Tata Lectures on Theta I*, Birkhäuser, 1983.
- [9] H. PETERSSON, *Ueber die Entwicklungskoeffizienten der automorphen Formen*, Acta Mathematica, Bd. 58 (1932), pp. 169–215.
- [10] H. RADEMACHER AND H. S. ZUCKERMAN, *On the Fourier coefficients of certain modular forms of positive dimension*, Annals of Math. (2) 39 (1938), pp. 433–462.
- [11] J. SERRE, *A Course in Arithmetic*, Springer New York, 1973.
- [12] MARYNA S. VIAZOVSKA, *The sphere packing problem in dimension 8*, Pages 991–1015 from Volume 185 (2017), Issue 3.
- [13] D. ZAGIER, *Elliptic Modular Forms and Their Applications*, In: The 1-2-3 of Modular Forms, (K. Ranestad, ed.) Norway, Springer Universitext, 2008.

Ecole Polytechnique Federale de Lausanne

1015 Lausanne

Switzerland

*Email address: [maryna.viazovska@epfl.ch](mailto:maryna.viazovska@epfl.ch)*