


# Endura®

Fungicide

## Lettuce Use in Arizona


### Control of *Sclerotinia* minor and *S. sclerotiorum* in Lettuce



2011-2012. Mike Matheron, University of Arizona, Yuma Ag Center.



Treatments were applied on Dec 6 and Dec 14, 2011. Disease ratings were taken Mar 12-13, 2012.

### Endura applied at seeding followed by application at thinning



2011-2012. Mike Matheron, University of Arizona, Yuma Ag Center.

Treatments were applied on Nov 7, 2011 and Dec 6, 2011. Disease ratings were taken Mar 12-13, 2012.



### Endura Use Directions and Restrictions

- For control of lettuce drop (*Sclerotinia* minor and *S. sclerotiorum*) apply at 8 to 11 oz/A
- Applications can be made at seeding or pre emergence for *Sclerotinia* control
- Maximum of 2 applications per season and 22 oz/A
- Use the higher rate when disease pressure is high
- To limit the potential for development of resistance, do not make more than 2 applications of Endura per season
- Pre Harvest interval is 14 days
- At Planting Use Not Registered in CA

Always follow FRAC guidelines when using fungicides.

**Always read and follow label directions.** Endura is a registered trademark of BASF.

Fontelis is a registered trademark of Dupont Crop Protection.

©2012 BASF Corporation. All Rights Reserved. APN 2012-19-201-KAW-1

## Technical Information Bulletin

 **BASF**  
The Chemical Company