

Merivon®


Xemium® Brand Fungicide

Disease Control in Pistachios*

Benefits of Merivon® Fungicide

- Longer-lasting disease protection
- More consistent performance for maximum crop potential
- Advanced Plant Health benefits

Long-Term Effects of Merivon Fungicide for Control of Botryosphaeria in Pistachio



2013, Butte Co., Dr. Themis Michalaidis, UC Davis, Kearney Agricultural Research and Extension Center. Buds sampled on 3/6/14, 7 months after 2013 fungicide sprays. 50 buds per replicate. Buds plated on APDA according to BUDMON technique. 2013 application timings: 4/2/13, 6/7/13, 7/10/13, & 8/1/13. Treatments sponsored by BASF and other product makers.

Merivon fungicide delivers longer-lasting disease control, along with Advanced Plant Health benefits, that will help to maximize pistachio orchard potential.




150 years

 **BASF**

We create chemistry

Technical Information Bulletin

Merivon® Fungicide Control of Alternaria Late Blight in Pistachio



2013 Dr. Themis Michalaidis – University of CA (Parlier, CA). Three applications (6/11, 7/3, 7/24/2013). BASF treatments with 0.06% Induce. Treatments are a subset of those evaluated. Treatments sponsored by BASF and other product makers.

Best Use Recommendations

- Use Rate: 5.5–6.5 oz/A
 - 5.5 oz/A for Botryosphaeria
 - 6.5 oz/A for Alternaria
- Do not make more than two sequential applications before rotation to an alternative mode of action fungicide
- Do not make more than three applications per season
- Adjuvant flexible
- Merivon fungicide may be mixed with other products, such as insecticides

Target Diseases

- Alternaria (Alternaria late blight)
- Botrytis (Botrytis blossom and shoot blight)
- Botryosphaeria (Panicle and shoot blight)

Merivon Fungicide Use Recommendation

Application Timings	Bloom (April)	Succulent Shoot Growth (May)	Early Fruit Development (June)	Fruit/Kernel Development (July)
Suggested Merivon Fungicide Rate	5.5 oz/A		6.5 oz/A	
Target Pathogen	Botryosphaeria, Botrytis		Alternaria, Botryosphaeria	

For Alternaria control:

- In areas with a history of Alternaria, three fungicide applications per season are recommended
- If only one application is made, a late June/early July timing is best



*This is a new use that may not be registered in all states. Consult your BASF representative or your state regulatory authority regarding the approval of this use in your state.

Always read and follow label directions.

Merivon, Pristine, and Xemium are registered trademarks of BASF.

Fontelis is a registered trademark of DuPont. Luna is a registered trademark of Bayer.

Inspire Super and Quadris Top are registered trademarks of Syngenta.

Quash is a registered trademark of Valent.

©2015 BASF Corporation. All Rights Reserved. APN# 1504002 Merivon-Pistachio