


Endura[®]

Fungicide

Control Disease in Broccoli

Endura[®] fungicide is a broad-spectrum multi-disease fungicide with a high level of activity on major diseases that threaten yield and quality in Broccoli like *Alternaria* spp.

For Control of Pin Rot in Broccoli



2012 Jake Zaccaria. Trial Location: Guadalupe, CA. Variety: Heritage. Application dates: Feb 6 and Feb 15 (dime size head and 1/2 dollar head). All treatments include NIS at 0.25%. Randomized complete block, 4 reps. 65 gpa. Plots were rated on Mar 13. Evaluations: % Disease severity.

Always follow FRAC guidelines when using fungicides.

Technical Information Bulletin

 **BASF**
The Chemical Company

Labeled Crops

- **Brassica Head and Stem:** Broccoli, Brussels sprouts, Chinese broccoli, Cabbage, Chinese Cabbage (napa), Chinese mustard, Cauliflower.
- **Leafy Brassica:** Broccoli raab, Chinese cabbage (bok choy), Collards: Kale, Mustard Greens, Mustard spinach.

**See label for full list of crops. **

Diseases Controlled

- Alternaria blight
- Gray mold
- Sclerotinia stem rot

Suppression

- Powdery mildew
- Rhizoctonia bottom rot



Rates and Recommendations

Use Rate: 6-9 oz*

Maximum Applications: 2

Apply on a 7-14 day spray interval

PHI: Head and Stem – 0 days, Brassica Leafy Greens – 14 days

*Use the higher rate and shorter interval when disease pressure is higher or conditions are favorable for disease.

General Guidelines

- Make your first Endura application prior to disease development or when conditions become favorable for disease.
- This is not a label.
- Refer to label for additional information.

Resistance Management

- To limit the potential for development of resistance, DO NOT make more than two (2) applications of Endura per season.
- Do not make more than 2 sequential applications of Endura before alternating to another labeled fungicide with a different mode of action for at least 1 application.