

Headline®

Fungicide

Headline® Fungicide In-Furrow

Best Use Recommendations

- Use Rate: 3 to 9 fl oz/A broadcast equivalent, based on row spacing (see rate table below)
- Maximum Rate/Season: 12 fl oz/A
- Minimum Volume of Water: 2.5 gal/A

Target Diseases

- *Rhizoctonia solani*

General Information

- Use Headline EC fungicide and follow BASF Best Management Practices when mixing with liquid fertilizer

Headline Fungicide Rate (fl oz/A)					
Rate per 1000 row feet (fl oz)	15" rows	20" rows	22" rows	30" rows	36" rows
0.1	3.5	—	—	—	—
0.2	7.0	5.2	4.7	3.5	3.0
0.3	10.5	7.8	7.1	5.2	4.5
0.4	—	10.4	9.5	6.9	6.0

DO NOT apply more than 12 fl oz/A. For 30" rows, use a maximum of 0.6 fl oz/1000 row feet.

Always read and follow label directions.

Headline is a registered trademark of BASF Corporation

©2015 BASF Corporation. All Rights Reserved.

APN# 1501005 Headline-In-Furrow-Dry Shelled Beans

Headline fungicide provides convenient and effective disease control that helps deliver more rapid and uniform emergence.

*Adzuki bean, black-eyed pea, broad bean, catjang, chickpea, crowder pea, field bean, grain lupin, guar, kidney bean, lablab bean, lima bean, moth bean, mung bean, navy bean, pink bean, pinto bean, rice bean, southern pea, sweet lupin, tepary bean, urd bean, and white lupin.

Recommendations for Use
on Dry Shelled Beans*

150 years

 **BASF**

We create chemistry