


BASF Potato Offer: A Planned Approach to Success

**Grow Smart™
with BASF**


Endura® Fungicide White Mold Control



2013 Mike Hubbard. Kootenai Valley Research, Bonners Ferry, ID. All fungicides applied twice, at row closure on July 25 and 11 days after row closure on August 5. Severity ratings taken on October 4.

Potato Fungicide Products

	Endura Fungicide	Priaxor® Fungicide
TIMING	Prior to infection for control of <i>Sclerotinia</i> white mold. Prior to onset of disease for control of <i>Alternaria</i> early blight or suppression of <i>Botrytis</i> gray mold.	Prior to disease development, continue 7- to 14-day interval
RATES	3.5-10 fl oz/A	4-8 fl oz/A
APPLICATION	Aerial, Ground	Aerial, Ground
PHI	10 days	7 days
REI	12 hr	12 hr
COMMENTS	Application rate dependent on disease	Application rate varies with disease. Adjuvant precautions, refer to label for details.



Always read and follow label directions.

Grow Smart is a trademark and Endura, Priaxor and Xemium are registered trademarks of BASF Corporation. Luna is a registered trademark of Bayer CropScience. Omega is a registered trademark of Syngenta ©2016 BASF Corporation. All Rights Reserved. APN# 1602004-Potatoes-Fungicides. For more information on BASF Crop Protection products, visit www.agproducts.basf.us

Technical Information Bulletin

BASF
We create chemistry