

**FRONT LABEL**



## **MONSANTO AMENITY GLYPHOSATE**

A foliar applied translocated herbicide for the control of emerged weeds in industrial and amenity situations, in forestry and on hard surfaces.

Degraded by micro-organisms / microbes in the soil.

A soluble concentrate containing 360 g/l glyphosate, present as 441g/l (35.3% w/w) of the potassium salt of glyphosate

<b>GROUP</b>	<b>9</b>	<b>HERBICIDES</b>
--------------	----------	-------------------

UFI: KPF1-E0Y7-K009-RH2Q

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work

### **MONSANTO AMENITY GLYPHOSATE**

Contains 360 g/l glyphosate, present as 441g/l (35.3% w/w) of the potassium salt of glyphosate.



### **Warning**

Causes serious eye irritation.

Toxic to aquatic life with long lasting effects.

Wash hands thoroughly after handling.

Wear protective gloves/ protective clothing/eye protection/face protection.

Keep only in original container.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

If eye irritation persists: Get medical advice/attention.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

MAPP Number 16382  
PROTECT FROM FROST  
Batch/lot number:  
Trade marks:

**BOOKLET AND BASE LABEL**



## MONSANTO AMENITY GLYPHOSATE

A soluble concentrate containing 360 g/l glyphosate present as 441 g/l (35.3% w/w) of the potassium salt of glyphosate

**MONSANTO (UK) LIMITED**

230 Cambridge Science Park, Milton Road, Cambridge, CB4 0WB, UK.  
Telephone: 01223 226500 ; <https://cropscience.bayer.co.uk/>

**For 24-hour emergency information contact Bayer CropScience Ltd.  
Tel: 00800 1020 3333**

### IMPORTANT INFORMATION

FOR USE ONLY AS AN INDUSTRIAL/AMENITY/FORESTRY HERBICIDE

**Crops/situations:**

Natural surfaces not intended to bear vegetation, permeable surfaces overlying soil, hard surfaces;  
Amenity vegetation;  
Forest nursery, forest (weed control, stump application and chemical thinning).

**Maximum individual dose:**

}

**Maximum number of treatments:**

}

**Latest time of application:**

} Full details are given in  
} the attached leaflet

**Other specific restrictions:**

} (Crop Specific Information – marked #)

READ ALL OTHER SAFETY PRECAUTIONS AND DIRECTIONS BEFORE USE.

### SAFETY PRECAUTIONS

#### Operator protection

*\* Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:*

*\* WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate and when handling contaminated surfaces.*

*\*WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND RUBBER BOOTS when applying by hand-held equipment, hand-held weedwiper, spot gun, hand-held controlled droplet applicator (CDA) equipment or making cut stump treatment.*

*\*WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES, RUBBER BOOTS AND FACE PROTECTION (FACESHIELD) when using stem injection equipment.*

*\* However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.*

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

### **Environmental protection**

Do not contaminate water with the product or its container except when used as directed. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

### **Storage and disposal**

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

EMPTY CONTAINER COMPLETELY and dispose of safely.

RINSE CONTAINER THOROUGHLY by using an integrated pressure-rinsing device or manually rinse three times. Add washings to sprayer at time of filling and dispose of safely. Triple rinsed containers may be disposed of as non-hazardous waste.

### **Medical advice**

For 24-hour emergency information contact Bayer CropScience Ltd. Tel: 00800 1020 3333

### **DIRECTIONS FOR USE**

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

### **Warnings**

EXTREME CARE SHOULD BE TAKEN TO AVOID SPRAY DRIFT AS THIS CAN SEVERELY DAMAGE NEIGHBOURING CROPS OR PLANTS.

DO NOT MIX, STORE OR APPLY MONSANTO AMENITY GLYPHOSATE IN GALVANISED OR UNLINED STEEL CONTAINERS OR SPRAY TANKS.

DO NOT leave spray mixtures in tank for long periods and make sure tanks are WELL VENTED.

### **Restrictions**

A period of at least 6 hours and preferably 24 hours rain-free must follow application of Monsanto Amenity Glyphosate.

Do not spray onto weeds which are naturally senescent, or where growth is impaired by drought, high temperatures, a covering of dust, flooding or frost at, or immediately after application, otherwise poor control may result.

Do not spray in windy conditions as drift onto desired crops or vegetation could severely damage or destroy them.

Do not tank mix Monsanto Amenity Glyphosate with adjuvants, pesticides or fertilisers except as specified under Directions for Use – Compatibility.

After application, large concentrations of decaying foliage, stolons, roots or rhizomes should be dispersed or buried by thorough cultivation before crop drilling.

Applications of lime, fertilizer, farmyard manure and pesticides should be delayed until 5 days after application of Monsanto Amenity Glyphosate unless cultivation and drilling of the next crop has taken place.

### **Weeds controlled**

*Monsanto Amenity Glyphosate* is a foliar acting herbicide which controls annual and perennial grasses and most broad-leaved weeds when used as directed. It is important that all weeds are at the correct growth stage when treated, otherwise some re-growth may occur and this will need re-treatment.

Apply *Monsanto Amenity Glyphosate* herbicide once grasses and broad-leaved weeds have emerged and they have ACTIVELY GROWING green leaves.

- PERENNIAL GRASSES must have a full emergence of healthy, green leaf. (Common Couch, for example, becomes susceptible at the onset of tillering and new rhizome growth commences which usually occurs when plants have 4-5 leaves, each with 10-15cm of new growth).
- PERENNIAL BROAD-LEAVED WEEDS are most susceptible around the flowering stage.
- BRACKEN should be treated after frond tips are unfurled, but pre-senescence.
- OTHER SPECIES - recommendations for specific Areas of Use are given in the Recommendation Tables,
- This product will not give an acceptable level of control of Horsetails (*Equisetum arvense*) – repeat treatment will be necessary.

### **Following crops**

Upon soil adsorption the herbicidal properties of *Monsanto Amenity Glyphosate* are lost permitting the drilling of crops 48 hrs after application.

## Crop specific information

<b>COMPLIANCE WITH THE FOLLOWING CONDITIONS OF USE AND ALL SAFETY PRECAUTIONS MARKED* IS A LEGAL REQUIREMENT</b>	
<b>Crops/situations:</b>	<b>Maximum individual dose (litres product/ hectare):</b>
Natural surfaces not intended to bear vegetation, permeable surfaces overlaying soil, hard surfaces	5.0
Amenity vegetation	5.0
Forestry, forest nursery: <ul style="list-style-type: none"> <li>Weed control</li> <li>Stump application</li> <li>Chemical thinning (by injection)</li> </ul>	10.0  200ml/litre of water (20% solution of product in water)  2 ml per 10cm diameter (or less) of tree
<b>Other specific restrictions:</b>  When applying through rotary atomisers the spray droplet spectra produced must be of a minimum Volume Median Diameter (VMD) of 200 microns.  Weed wipers may be used in any crop where the wiper or chemical does not touch the growing crop.  For weed wiper applications, the maximum concentrations must not exceed the following:  Weed wiper Mini                      1:2 dilution with water } Refer to weed wiper guidance under Other wipers                            1:1 dilution with water } 'Mixing & Spraying' section  For stump application, the maximum concentration must not exceed 200 ml of product per litre of water (i.e., a 20% solution).	

**RECOMMENDATION TABLES**

AREA OF USE	TARGET WEEDS/USAGE	CROP/SITUATION	WEED INFESTATION	APPLICATION RATE l/ha	WATER VOLUME	APPLICATION TIMING AND GUIDANCE
NATURAL SURFACES NOT INTENDED TO BEAR VEGETATION, PERMEABLE SURFACES OVERLYING SOIL, RAILWAY BALLAST	Vegetation management	Including roadsides, paths, and along fences & total weed control on industrial sites	Annual weeds	1.5	Hydraulic sprayers (boom and knapsack) at water volumes 80-400 l/ha or rotary atomisers*at water volumes 40 l/ha or hand-held equipment. See Mixing & Spraying section.	Use areas include: Cleaning up weedy ground prior to planting or sowing and as a directed spray in ornamental plantings. Hydraulic sprayers, rotary atomisers or weed wipers may be used. DO NOT USE IN OR ALONGSIDE HEDGEROWS. DO NOT USE UNDER POLYTHENE OR GLASS
			Perennial grasses and broad-leaved weeds	4.0 - 5.0		
Hard surfaces (excluding railway ballast)	Vegetation management	including roadsides, paths, concrete and alongside walls	Annual weeds	1.5	Hydraulic sprayers (boom and knapsack) at water volumes 80-400 l/ha or rotary atomisers*at water volumes 40 l/ha or hand-held equipment. See Mixing & Spraying section.	Apply this product carefully. Ensure spraying takes place only when weeds are actively growing (normally March to October) and is confined only to visible weeds including those in the 30cm swath covering the kerb edge and road gully – do not overspray drains
			Perennial grasses and broad-leaved weeds	4.0 - 5.0		
AMENITY VEGETATION	Vegetation management	Areas of semi-natural or ornamental vegetation including trees. Areas of bare soil around ornamental plants or areas intended for ornamental planting or clearance of allotments	Annual weeds	1.5	80-400 l/ha*	Hydraulic sprayers, rotary atomisers or weed wipers may be used  DO NOT USE IN OR ALONGSIDE HEDGEROWS.  DO NOT USE UNDER POLYTHENE OR GLASS
			Perennial grasses and broad-leaved weeds	4.0 - 5.0		
	Root suckers	-	All species	5.0		

Forestry weed control

Monsanto Amenity Glyphosate can be used for site preparation and for weed control in planted out trees.

Monsanto Amenity Glyphosate A2 December 2023

AREA OF USE	TARGET WEEDS/USAGE	WEED INFESTATION	APPLICATION RATE L/HA.	WATER VOLUME	APPLICATION TIMING & GUIDANCE
Forestry: - Pre-planting	Arable land, planting, replanting, & grassland areas	Arable weeds  Grassland weeds	4.0  5.0	Hydraulic sprayers: 80-250 l/ha or rotary atomisers: 40 l/ha*	All tree species may be planted 7 days or more after treatment.  *Where rotary atomisers are used their droplet diameter must fall within the range 200-300µm.
Forestry: - Post-planting (directed) in conifers & broad-leaved trees	Clean-up around trees with knapsack applications.	Annual/perennial grasses and broad-leaves	4.0	Hand held equipment.  Knapsack: Apply as a 2% concentration or Weed wiper mini: apply as a concentration of 1 part Monsanto Amenity Glyphosate to 2 parts water (see Mixing & Spraying section)	It is ESSENTIAL to use a TREE GUARD for all applications made in the growing season. Treat bracken after frond tips are unfurled but before senescence. Treat heather late August to end September. All other woody weeds are treated June-August, before leaf senescence (but after new growth of crop has hardened).
		Woody weeds:  Bracken/Beech Brush/Brambles Sycamore/Oak Hazel/Willow/Ash	3.0		
		Heather (peat soils) Heather (mineral soils)	4.0 6.0		
		Rhododendron (*)	10.0		Cut back and treat re-growth when at least 1 metre in height throughout the entire coppice. Spray to just before point of run-off.  (*) For improved control of Rhododendron add Mixture B (ADJ AO161) at a concentration of 2% final water volume to 8.0 l/ha of Monsanto Amenity Glyphosate. Application using the weed wiper is not suitable.

Monsanto Amenity Glyphosate A2 December 2023

AREA OF USE	TARGET WEEDS/USAGE	WEED INFESTATION	APPLICATION RATE L/HA.	WATER VOLUME	APPLICATION TIMING & GUIDANCE
Forestry: - Post-planting (overall application in dormant season in certain conifers – conifer release)	Grass weeds  - Lowland areas - Upland areas          Bracken Beech & Birch Brambles	Black Bent, Cock's-foot, Common Couch, Creeping Soft-grass, False Oat-grass, Fescues, Meadow-grasses, Other Bent species, Purple Moor-grass, Sweet Vernal-grass, Tufted Hair-grass, Wavy Hair-grass, Wood Small-reed (Bush grass)  All infestation levels All infestation levels All infestation levels	1.5 2.0         2.0 2.0 3.0	Hydraulic sprayers: 200-250 l/ha  or  Hand-held equipment - see 'Mixing and Spraying' section	DO NOT OVERALL SPRAY trees being grown for ORNAMENTAL PURPOSES, including CHRISTMAS TREES.  Species safe to spray when fully dormant and leader growth has hardened:  Corsican, Lodgepole and Scots Pines, Norway Spruce, Sitka Spruce, Lawson Cypress, Western Red Cedar.  Douglas Fir and Noble Fir - safe to spray when fully dormant and leader growth has hardened but NOT in spring.  If overall application takes place after the optimum timing weed control may be reduced. It is advisable to spray a limited area of forest to test crop safety under local conditions before widespread overall application in subsequent years.  These recommended application rates refer to forestry usage only.  Inadequate control may result if used in other areas.  <u>Caution:</u> The timing of hardening of leader growth varies considerably between locations and between seasons. It may occur as early as the end of July or be delayed to October or later. To avoid damage to Lammas growth, sprays should be directed away from leaders.



Monsanto Amenity Glyphosate A2 December 2023

AREA OF USE	TARGET WEEDS/USAGE	WEED INFESTATION	APPLICATION RATE L/HA.	WATER VOLUME	APPLICATION TIMING & GUIDANCE
Forestry: - Stump application for chemical thinning	Deciduous trees  Coniferous trees	All species  All species	10% solution of Monsanto Amenity Glyphosate in water  20% solution of Monsanto Amenity Glyphosate in water		Apply the solution to saturate the rim of the newly cut surface, with a suitable adapted clearing saw, spot gun or paintbrush. Treat as soon as possible (within a week) after felling, in the period November to March. Do not apply in the period of active sap flow in the spring/early summer. Do not cut trenches or drill holes and fill with the solution or use undiluted product.  Note: for ease of identification of treated areas a suitable, commercially available, water-soluble dye may be added to the prepared spray solution.
Forestry: - Chemical thinning by injection of tree stems	Coniferous and deciduous species	-	2 ml neat Monsanto Amenity Glyphosate per cut per 10 cm diameter (or less) tree		Use a hatchet to cut one notch in trees up to 10cm diameter and apply 2 ml of the solution to each cut. Use two or three notches in trees over 10cm diameter. Do not treat in the period of active sap flow in the spring/early summer.

## **Mixing and spraying**

*Monsanto Amenity Glyphosate* mixes readily with water and can be applied in spray volumes ranging from 80-400 l/ha using tractor mounted, knapsack, rotary atomisers and hand-held sprayers. Specialised application equipment such as weed wipers, stem injection and spot gun applicators may be used where indicated.

Correctly calibrate all sprayers under field or use conditions prior to application.

### a) Tractor mounted and powered sprayers

These should be capable of applying accurately 80-400 l/ha within a pressure range of 1.5-2.5 bars (20-35 psi).

Half fill the spray tank with clean water, start gentle agitation, and then add the correct amount of *Monsanto Amenity Glyphosate*. Top up the tank with water to the required level. To avoid foaming do not use top tank agitation. Use of a defoamer may be necessary.

All applications using hydraulic sprayers (including knapsack sprayers) to be as 'MEDIUM' or 'COARSE' spray quality (BCPC definition).

#### *Medium Volume application (150-300 l/ha)*

Avoid high water volumes (>300 l/ha) which may lead to run-off from the treated vegetation, resulting in reduced control. Low drift nozzles such as air induction and pre-orifice types producing a medium or coarse spray (BCPC definition) should be used to minimise the risk of drift.

#### *Low Volume Application (minimum 80 l/ha)*

Low volume application can be achieved by reducing pressure and the appropriate nozzle selection. Low drift nozzles which produce a medium spray quality (BCPC definition) should be used to minimise the risk of drift.

### b) Knapsack sprayers

Recommended delivery range is 80 - 300 l/ha. Half fill the spray tank with clean water, add the correct amount of *Monsanto Amenity Glyphosate* and top up with water.

When used at a walking speed of 1 m/sec to apply a swath of 1 m width, most knapsack sprayers fitted with a Hypro AN 0.6-AN2.4 or similar nozzle deliver approximately 200 l/ha spray volume (or 10 l per 500 m<sup>2</sup>). To apply 5.0 l/ha of MONSANTO AMENITY GLYPHOSATE, therefore, use 50ml of product for each 2 litres of spray liquid required. Similarly, knapsack sprayers fitted with low volume nozzles such as D/0.23/1- D/0.68/1 typically deliver approximately 100 l/ha spray volume. To apply 5.0 l/ha *Monsanto Amenity Glyphosate* in this case, use 100ml of product for each 2 litres of spray liquid required.

### c) Rotary Atomisers

Tractor-mounted boom sprayers and hand-held machines are suitable for use in some situations to apply a minimum spray volume of 10 l/ha.

When rotary atomisers are used to apply *Monsanto Amenity Glyphosate* ensure that the droplet diameter falls within the range 200-300 microns for all uses.

Stir the correct amount of *Monsanto Amenity Glyphosate* to control the particular target species into the sprayer bottle half filled with clean water. Top up with water, close the top and shake gently to ensure good mixing.

Do not tank mix *Monsanto Amenity Glyphosate* when using rotary atomiser sprayers.

#### d) Weed Wipers

For ropewick applicators use a concentration of 1 part *Monsanto Amenity Glyphosate* to 2 parts of water and add a water-soluble dye if required. Care should be taken to avoid dripping onto wanted vegetation.

For new generation weed wipers, use 1 part *Monsanto Amenity Glyphosate* to 10 or 20 parts of water or as directed by manufacturer's instructions. A list of machines is included in the Company Advisory section at the end of this label.

#### e) Cut Stump Applicators

For cut stump treatment an Enso attachment to rotary saws may be used.

This technique is specific to scrub clearance in forestry. A suitable water soluble dye may be used.

#### f) Stem injection

Use a hatchet to cut one notch in trees up to 10cm diameter and two to three notches in trees above 10cm diameter. Use 2.0 ml of undiluted *Monsanto Amenity Glyphosate* per notch.

#### g) Spot Gun Applicators

Spot gun applicators are for the treatment of individual weeds. Apply 5 ml of spray to target weed, using a narrow cone TG-3 or TG-5 nozzle.

Spot Diameter (metres)	Amount of Monsanto Amenity Glyphosate (ml) per 5 litres spray solution for targeted dosages of:		
	3.0 l/ha	4.0 l/ha	5.0 l/ha
0.3	20	28	35
0.6	85	110	140

## **Compatibility**

Do not tank mix *Monsanto Amenity Glyphosate* with adjuvants, pesticides or fertilisers except as advised by Monsanto. For up to date information on compatible products contact Monsanto UK Limited (Telephone: 01223 226500

*Monsanto Amenity Glyphosate* is compatible with Mixture B NF(ADJ 0161). Where conventional hydraulic sprayers are being used Mixture B NF may be added to the spray tank solution, at a rate of 2% of the final water volume.

DO NOT APPLY WITH MIXTURE B NF TO EDIBLE CROPS, OR GRASSLAND WEEDS.

Do not tank-mix *Monsanto Amenity Glyphosate* when using rotary atomiser sprayers.

For hydraulic sprayers: maintain continuous agitation when using *Monsanto Amenity Glyphosate* in tank mixture.

For knapsack sprayers: mix thoroughly and use immediately when using *Monsanto Amenity Glyphosate* in tank mixture.

## COMPANY ADVISORY INFORMATION

This section is not part of the Product Label under the Plant Protection Products Regulations 1995 and provides additional advice on the product.

### General Information

*Monsanto Amenity Glyphosate* herbicide is a foliar-acting herbicide with broad-spectrum activity. It is taken up by foliage and translocated to underground roots, rhizomes and stolons, providing control of both annual and perennial grasses and broad-leaved weeds. *Monsanto Amenity Glyphosate* is rapidly adsorbed onto particulate matter in soils and water and is quickly degraded by the micro-organisms present in soil and aquatic bottom sediments. Upon adsorption, the herbicidal properties of *Monsanto Amenity Glyphosate* are lost, permitting drilling of crops within 48 hours of application. When used as directed, any water subjected to *Monsanto Amenity Glyphosate* spray drift may be used immediately for irrigation purposes. Until degraded, the active ingredient in *Monsanto Amenity Glyphosate*, glyphosate, is practically immobile in soils and is, therefore, unlikely to contaminate groundwater.

To maximise the intrinsic safety of *Monsanto Amenity Glyphosate* to operator, consumer and environment, the label recommendations and the DEFRA/HSC/NAW publication "Code of Practice for Using Plant Protection Products" of January 2006, should be adhered to.

### Symptoms on the weeds

Symptoms of treatment are generally first seen 7-10 days, or longer (if growth is slow), after spraying. These take the form of leaf reddening followed by yellowing and are usually quicker to appear on grasses than on broad-leaved weeds. Reaction of nettles is slow.

### **Effects of weather**

See Directions for Use (Restrictions).

*Monsanto Amenity Glyphosate* will remain efficacious at low but not freezing temperatures however the onset of symptoms will be delayed.

A covering of dew may reduce efficacy where run-off occurs.

Reduced control is likely where weed growth is impaired by natural senescence, drought, high temperature, a covering of dust, flooding or severe/prolonged frost at, or immediately after, application.

### **Weed resistance strategy**

There is low risk for the development of weed resistance to *Monsanto Amenity Glyphosate*.

There are no known cases of weed resistance to glyphosate in UK.

Strains of some annual weeds (e.g. Black-grass, Wild oats and Italian Ryegrass) have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. This should include integrating herbicides with a programme of cultural control measures. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer (Monsanto).

Growers are encouraged to implement a weed resistance strategy based on (a) Good Agricultural Practices and (b) Good Plant Protection Practices by:

Following label recommendations

The adoption of complimentary weed control practices

Minimising the risk of spreading weed infestations

The implementation of good spraying practice to maintain effective weed control

Using the correct nozzles to maximise coverage

Application only under appropriate weather conditions

Monitoring performance and reporting any unexpected results to Monsanto UK Ltd (Telephone: 01223 226500)

### **General Cautions**

Take extreme care to avoid drift, particularly when using near or alongside hedgerows. The use of low drift nozzles such as 'air induction' and 'pre-orifice' nozzles are recommended.

After application, large concentrations of decaying foliage, stolons, roots or rhizomes should be dispersed or buried by thorough cultivation before crop drilling.

### **Sprayer Maintenance**

Ensure the sprayer is in good working order and replace damaged, worn or malfunctioning parts before use. Carry out maintenance according to the instructions of the sprayer manufacturer.

### **Sprayer Hygiene**

It is essential to thoroughly clean-out spray tanks, pumps and pipelines and nozzle or disc assemblies, with a recommended detergent cleaner, between applying this product and other pesticides to avoid contamination from pesticide residues. Traces of *Monsanto Amenity Glyphosate* left in the equipment may seriously damage or destroy crops sprayed later.

### **Calibration**

All sprayers should always be calibrated before use. This is essential when nozzles are changed or if a different dose of product is to be applied.

### **Unused Spray Mixture**

Once *Monsanto Amenity Glyphosate* has been diluted in the spray tank, it should be used as soon as possible. However, if unexpected delays occur the diluted spray can be safely stored. Agitate well before use. Storage for longer than 3 days may result in reduced efficacy.

### **Disposal**

Follow the guidance on the disposal of surplus spray solution, tank washings, concentrate and containers as given in Section 5 of the DEFRA/HSC/NAW publication "Code of Practice for Using Plant Protection Products", January 2006.

### **Material Safety Data Sheet**

A material safety data sheet for this product is available on request.

<https://cropscience.bayer.co.uk/> for MSDS and larger label.

### **Trade Mark References**

Monsanto® and the Vine symbol are registered trade marks of Monsanto Technology LLC. All other brand names referred to are trade marks of other manufacturers in which proprietary rights may exist.

Monsanto does not warrant that the purchase or use of equipment mentioned in this document will not infringe any patent or trade mark registration.