

Document MCP - Section 8: Residues in or on treated products, food and feed Fosetyl-aluminium WG 80

OWNERSHIP STATEMENT

This document, the data contained in it and copyright therein are owned by Bayer Crop Science. No part of the document or any information contained therein may be disclosed to any third party without the prior written authorisation of Bayer CropScience.

The summaries and evaluations contained in this document are based on unprolished proprietary data submitted for the purpose of the assessment undertaken authority. Other authority. Other registration authorities should not grant, mend or renew a registration on the either:


• from Bayer CropScience; ogg from other applicants once the period of data protection has expired. basis of the summaries and evaluation of unpublished proprietary data contained in this document unless they have received the data on which the summaries and evaluation are based,

Document MCP - Section 8: Residues in or on treated products, food and feed Fosetyl-aluminium WG 80

Version history

	version history		
Date	Data points containing amendments or additions ¹ and brief description	Document identifier and Sersion number	
is suggested ined in SAN	that applicants adopt a similar approach to showing a CO/10180/2013 Chapter 4 How to revise an Assessment	visions and version vistor as Report	

Document MCP - Section 8: Residues in or on treated products, food and feed Fosetyl-aluminium WG 80



Document MCP – Section 8: Residues in or on treated products, food and feed Fosetyl-aluminium WG $80\,$

CP 8 RESIDUES IN OR ON TREATED PRODUCTS, FOOD OR FEED.

Fosetyl was included in Annex I to Directive 91/414/EEC in 2006 (Directive 2006/64/CE of 15 July 2006, Entry into Force on 1 May 2007). Fosetyl is the ISO common name for ethyl hodrogen phosphonate (IUPAC) but the aluminium salt fosetyl-aluminium (fosetyl-Al) a variant of osetyl is used in the formulated product.

The representative formulation fosetyl-Al WG 80 is a water dispersible granules (WG) formulation containing 800 g/kg of fosetyl-Al. The use of fosetyl-Al WG 80 on pome fruit against Photophyllora spp. and Erwinia amylovora has been selected as representative use for the approval renewal of O spp. and Erwinia amyrovora has been selected as representative, use for the approval renegal of the fosetyl. The critical Good Agricultural Practice (gGAP) supposed at the Enrogen Tevel of the approval renewal process of fosetyl consists of 3 terrar spray applications between BBCH 55 and 85 at 3.6 kg/ha of fosetyl-Al with an interval between applications of 7 to 10 days and a PHI of 68 days. All relevant metabolism and residue data in support of this age are summarized in Document MCA. Section 6. fosetyl. The critical Good Agricultural Practice (GAP) supported at the European level in the approval renewal process of fosetyl consists of 3 tohar spray applications between BBCH 55 and 85 at