

Cell Type Plant, suspension

Molecules Electroporated DNA: pCH (pUC vector with hygromycin resistant gene)

Species Used *Oryza sativa*, cv. Yamahouci or cv. Nihonbare

Before the Pulse

Cell Growth Medium AA medium (Amino Acid Medium)

Growth Phase at Harvest Log phase

Pre-pulse Incubation 0.5 mM MES, 70 mM KCl, 4 mM CaCl₂, 0.36 M Mannitol

Wash Solution 0.5 mM MES, 70 mM KCl, 4 mM CaCl₂, 0.36 M Mannitol

The Pulse

Instruments Used Gene Pulser® apparatus , Capacitance

Electroporation Temperature Room temperature

Electroporation Medium 0.5 mM MES, 70 mM KCl, 4 mM CaCl₂, 0.36 M Mannitol

Cuvette Gap 0.4 cm

Cell Density 10 (6) / ml

Voltage 0.25 kV

Volume of Cells 500 µl

Field Strength 0.625 kV/cm

DNA Concentration Not given

Capacitor 250 µF

DNA Resuspension Buffer Not given

Resistor (Pulse Controller) 200 Ω

Volume of DNA 50 µl

Time Constant 10 to 20 msec

After the Pulse

Outgrowth Medium Not given

Relevant Publications and/or Comments

Note: exponential values designated in parentheses.

Outgrowth Temperature Not given

Length of Incubation Not given

Selection Method or Assay Used Not given

Electroporation Efficiency Not given

Per Cent Survival Not given

Name of Submitter Dr. Seiichi Toki

Institution Address Hokkaido Univ College of Science
Biology, Plant structural studies
Sapporo-shi, kita-ku, 10 jo nishi 8 chome
JAPAN 7060

Survey Number

185