

Cell Type Other Cell Types

Molecules Electroporated DNA: supercoiled DNA used for transient transfections.

Species Used Chicken, TS34 a6 L1, [LSCC HD2], erythroblast

Before the Pulse

Cell Growth Medium Not given

Growth Phase at Harvest 50 to 70% confluency

Pre-pulse Incubation 4° C, 10 min.

Wash Solution Wash two times in electroporation buffer

The Pulse

Instruments Used Gene Pulser® apparatus & Capacitance

Electroporation Temperature Room temperature

Electroporation Medium Phosphate Buffered Saline

Cuvette Gap 0.4 cm

Cell Density 5 x 10 (6) cells / pulse

Voltage 0.270 kV

Volume of Cells 0.5 ml

Field Strength 0.675 kV/cm

DNA Concentration 10 µg / pulse

DNA Resuspension Buffer Not given; pulse volume: 0.8 ml

Capacitor 960 µF

Volume of DNA Not given; pulse volume: 0.8 ml

Resistor (Pulse Controller) Ω not used

After the Pulse

Time Constant 24 msec

Outgrowth Medium Not given

Relevant Publications and/or Comments

Note: exponential values designated in parentheses.
PBS: 1x = 8g NaCl, 0.2g KCl, 0.2g KH₂PO₄, 1.15g Na₂HPO₄

Outgrowth Temperature 37 °C

Length of Incubation 48 to 72 hrs.

Selection Method or Assay Used Transient assays

Electroporation Efficiency Not given

Per Cent Survival about 50%

Name of Submitter Jackie Beall

Institution Address University of Adelaide
Department of Biochemistry
P.O. Box 498
Adelaide 5001
Australia

Survey Number

187