

Cell Type Other Cell Types

Molecules Electroporated Analogue cytoskeletal proteins (like tubulin).

Species Used Hydra cells, *Cnidaria*

Before the Pulse

Cell Growth Medium (Dissociation buffer) : 1.2 mM MgSO₄, 6 mM CaCl₂, 3.6 mM KCl, 6 mM pyruvate; 6 mM Na-Citrate, 12.5 mM TES buffer, 6 mM glucose, pH 6.9.

Growth Phase at Harvest Not given

Pre-pulse Incubation Not given

Wash Solution Not given

The Pulse

Instruments Used Gene Pulser® apparatus

Electroporation Temperature Room temperature

Electroporation Medium The same as cell growth medium

Cuvette Gap 0.4 cm

Cell Density 10 (5) cells / ml

Voltage 0.230 kV

Volume of Cells 10 (4) cells / pulse

Field Strength 0.575 kV/cm

DNA Concentration Not given

DNA Resuspension Buffer Not given

Capacitor 25 µF

Volume of DNA Not given

Resistor (Pulse Controller) none Ω

After the Pulse

Time Constant 4.3 msec

Outgrowth Medium The same

Relevant Publications and/or Comments

Note: exponential values designated in parentheses.

Outgrowth Temperature Room temperature

Length of Incubation Not given

Selection Method or Assay Used These cells are "primary cell cultures"

Electroporation Efficiency Not given

Per Cent Survival Not given

Name of Submitter Charo Gonzalez Agosti

Institution Address Zoologisches Inst.
The Univ. Zurich
Winterthurerstr. 190
8057 Zurich, Switzerland

Survey Number

189