

Cell Type Bacterial, gram negative

Molecules Electroporated DNA: pUC 18, including cDNA

Species Used *E. coli*, unspecified strain

Before the Pulse

Cell Growth Medium LB

Growth Phase at Harvest O.D. (600) =0.55

Pre-pulse Incubation 5 seconds on ice

Wash Solution 10 mM HEPES, pH 7.0

The Pulse

Instruments Used Gene Pulser® apparatus

Electroporation Temperature Not given

Electroporation Medium 10% glycerol

Cuvette Gap 0.2 cm

Cell Density 1 x 10 (a11) cells / ml

Voltage 2.5 kV

Volume of Cells 40 µl

Field Strength 12.5 kV/cm

DNA Concentration 20 ng

DNA Resuspension Buffer 2 µl water

Capacitor 25 µF

Volume of DNA 2 µl

Resistor 200 Ω (Pulse Controller)

After the Pulse

Time Constant 4.6 to 4.7 msec

Outgrowth Medium Not given

Relevant Publications and/or Comments

Note: exponential values designated in parentheses.

LB: 1% Bacto tryptone, 0.5% Bacto yeast extract, 0.5% NaCl.

Outgrowth Temperature 37 °C

Length of Incubation 1.5 hours

Selection Method or Assay Used ampicillin

Electroporation Efficiency 2-5 x 10(9)/µg pUC18 or 2-5 x 10(6)/µg double stranded blunt-end ligated cDNA

Per Cent Survival Not given

Name of Submitter Rainer Fiodas

Institution Address Beckman Research Institute of City of Hope
Molecular Biochemistry
1450 East Duarte Road
Duarte, CA 91010

Survey Number

044