

Cell Type Bacterial, gram negative
Species Used *E. coli*, unspecified strain

Molecules Electroporated DNA

Before the Pulse

Cell Growth Medium LB

Growth Phase at Harvest O.D. (600) =0.25

Pre-pulse Incubation 1 min.

Wash Solution water

The Pulse

Instruments Used Gene Pulser® apparatus

Electroporation Temperature 4 °C

Electroporation Medium water

Cell Density 10 (10) cells / pulse

Volume of Cells 40 µl

DNA Concentration 1 ng / µl

DNA Resuspension Buffer TE buffer

Volume of DNA 1 µl

Cuvette Gap 0.2 cm

Voltage 2.5 kV

Field Strength 12.5 kV/cm

Capacitor 25 µF

Resistor 300 Ω (Pulse Controller)

Time Constant 4.7 msec

After the Pulse

Outgrowth Medium Not given

Relevant Publications and/or Comments

Note: exponential values designated in parentheses.

LB: 1% Bacto tryptone, 0.5% Bacto yeast extract, 0.5% NaCl.

Outgrowth Temperature 37 °C

Length of Incubation 1 hour

Selection Method or Assay Used Ampicillin

Electroporation Efficiency Not given

Per Cent Survival Not given

Name of Submitter Dr. D. Wilson

Institution Address Cornell University, Biochemistry Dept.
Biotechnology Bld.
Ithaca, NY 14853

Survey Number
045