31"

1/16" White | 4 1/2" Back Handle | 5/16" Live Area | 10 9/16" Side | 1/4" Live Area | 4 1/2" Front Spout | 5/16" Live Area | 10 7/16" Side | 1/16" White

2 1/2" Fold Line | 15 11/16" Fold Line | 13 3/16" Fold Line

One-Step Disinfectant – Cleaner – Fungicide – Mildewstat – Virucide* – Deodorizer for Restaurants, Hospitals, Institutional and Industrial Use

This product will disinfect, clean and deodorize hard, non-porous surfaces in rest rooms and toilet areas, behind and under sinks and counters, garbage cans and garbage storage areas, and other places where bacterial growth can cause malodors.

This product is a concentrated one-step hospital use disinfectant cleaner that is effective against a broad spectrum of bacteria, is virucide* (including HIV-1, the AIDS Virus, Hepatitis B Virus and Hepatitis C Virus), fungicidal, and inhibits the growth of mold and mildew, and their odors, when used as directed.

Cross-contamination is of major housekeeping concern. This product has been formulated to aid in the reduction of cross-contamination on hard non-porous surfaces not only in hospitals, but in schools, institutions and industry.

This product leaves a clean fresh scent.

This product is a one-step hospital use disinfectant, bactericidal according to the current AOAC Use-Dilution Test Method and Virucide* according to the virucidal qualification modified in the presence of 200 ppm hard water plus 5% organic serum against:

Bacteria:
Pseudomonas aeruginosa (Pseudomonas)
Staphylococcus aureus (Staph)
Salmonella enterica (Salmonella)
Bronchiseptica bronchiseptica
Klebsiella pneumoniae
Campylobacter jejuni
Corynebacterium ammoniagenes
Enterobacter aerogenes
Enterobacter cloacae NDM-1 positive
Enterococcus faecalis (Enterococcus)
Enterococcus faecalis - Vancomycin Resistant (VRE)
Escherichia coli (E. coli)
Escherichia coli, **New Delhi** Metallo-Beta Lactamase (NDM-1)
Escherichia coli - Shiga toxin producing Shiga toxin lactamase producing E. coli)
Klebsiella pneumoniae (Klebsiella)
Klebsiella pneumoniae - NDM-1 positive
Legionella pneumophila
Listeria monocytogenes (Listeria)
Salmonella schottmuelleri (Salmonella)
Salmonella typhi (Salmonella)
Serratia marcescens (Serratia)
Shigella dysenteriae (Shigella)
Staphylococcus aureus – Community Associated Methicillin-Resistant (CA-MRSA) (ABCSAR) (USA300)
Staphylococcus aureus – Community Associated Methicillin-Resistant (CA-MRSA) (ARC172) (USA400)
Staphylococcus aureus – Methicillin-Resistant (MRSA)
Staphylococcus aureus – Multi-Drug Resistant (resistant to tetracycline (Tc), penicillin (Pc), streptomycin (Sm) and erythromycin (Em) and susceptible to chloramphenicol (Cm) in vitro)
Staphylococcus aureus – Vancomycin Intermediate Resistant – (VISA)
Streptococcus pyogenes (Strep) (a cause of scarlet fever)
Vibrio cholerae

Viruses:
*Hepatitis B Virus (HBV)
*Hepatitis C Virus (HCV)
*Herpes Simplex Virus Type 1
*Herpes Simplex Virus Type 2
*HIV-1 (AIDS Virus)
*Human Coronavirus
*Influenza Virus Type A / Brazil (Influenza)
*Influenza Virus Type A / Hong Kong (Influenza)
*2010 Influenza A Virus (H1N1)
*Respiratory Syncytial Virus (RSV)
*SARS Associated Coronavirus (SARS) (cause of Severe Acute Respiratory Syndrome)
*Vaccinia Virus (Pox Virus)
*Adenovirus Type 5 (2 sec. per gallon)
*Adenovirus Type 7 (2 sec. per gallon)
*Norwalk Virus (Feline Calicivirus) (Norovirus)
*Rotavirus

Kills Pandemic 2009 H1N1 influenza A virus.

Animal Viruses:
Avian Influenza Virus (H7N2)
Avian Influenza Virus (H5N1)
Avian Infectious Bronchitis Virus
Canine Distemper Virus
Feline Calicivirus
Feline Panleukopenia Virus (8 sec. per gallon)
Marine Reovirus*
Newcastle's Disease Virus
Pseudorabies Virus

Fungi:
Aspergillus fumigatus
Aspergillus niger (mildew)
Trichophyton mentagrophytes (The Athlete's Foot Fungus) (A cause of Ringworm) (A cause of Ringworm of the foot)

*Tested in laboratory

Dilution:
Disinfection (1:256) ½ oz. per gallon of water ..... .844 ppm active
Disinfection (1:128) 1 oz. per gallon of water ..... 1655 ppm active
Disinfection (1:64) 2 oz. per gallon of water ..... 3280 ppm active
Disinfection (1:16) 8 oz. per gallon of water ..... 12582 ppm active

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
If you do not understand the label, find someone to explain it to you in detail.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected.

Preparation of Use-Solution: Mix ½ - 1 oz. per gallon of water. For heavy duty use, mix 2 oz. per gallon of water. Use 2 oz. per gallon of water to kill Adenovirus Type 5 and Type 7. Use 8 oz. per gallon of water to kill Feline Panleukopenia Virus.

DISINFECTION / VIRUCIDAL* / FUNGICIDAL / MOLD AND MILDEW CONTROL DIRECTIONS: Apply use-solution to hard, non-porous surfaces, thoroughly wetting surfaces with a cloth, mop, sponge, sprayer or by immersion. Treated surfaces must remain wet for 10 minutes.

For Influenza Virus Type A and Human Coronavirus, treated surfaces must remain wet for 1 minute. Wipe dry with a cloth, sponge or mop or allow to air dry. For heavily soiled areas, a preliminary cleaning is required. For sprayer applications, use a coarse spray device. Spray 6 - 8 inches from the surface; rub with a brush, sponge or cloth. Do not breathe spray. Rinse all surfaces that come in contact with food such as countertops, exteriors of appliances, tables and stowetops with potable water before reuse. Do not use on utensils, glassware and dishes.

For fungicidal activity, this product is an effective fungicide against *Trichophyton mentagrophytes* (the athlete's foot fungus) in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities. Dilute at ½ - 1 oz. per gallon of water in 200 ppm hard water and allow surface to remain wet for 10 minutes.

For mold and mildew, at ½ - 1 oz. per gallon of water, this product will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, non-porous surfaces. Follow disinfection instructions. Repeat treatment every seven days, or more often if new growth appears.

BACTERICIDAL STABILITY OF USE-DILUTION: Tests confirm that the use-solution of this product, when stored in a sealed container such as a spray bottle, remains effective for up to 1 year. If product becomes visibly dirty or contaminated, the use-dilution must be discarded and fresh product prepared. Always use clean, properly labeled containers when diluting this product. Bactericidal stability of the use-dilution does not apply to open containers such as buckets or pails. This product is not for use on medical device surfaces.

***KILLS HIV AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids, and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OR HBV OR HCV ON SURFACES / OBJECTS SOILED WITH BLOOD / BODY FLUIDS.

Personal Protection: Clean-up must always be done wearing protective gloves, gowns, masks and eye protection.

Cleaning Procedure: Blood and other body fluids containing HIV or HBV or HCV must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: Leave surfaces wet for 1 minute (60 seconds) for HIV-1 and 10 minutes for HBV and HCV with ½ - 1 oz. per gallon use-solution. Use 2 oz. per gallon of water to kill Adenovirus Type 5 and type 7. Use a 10-minute contact time for disinfection against all other viruses, bacteria and fungi claimed.

Disposal of Infectious Material: Blood, body fluids, cleaning materials and clothing must be autoclaved and disposed of according to local regulations for infectious waste disposal.

NON-ACID TOILET BOWL AND URINAL DISINFECTANT / CLEANER DIRECTIONS: Remove gross filth prior to disinfection.

From Concentrate: Add ½ - 1 oz. to the bowl and mix. Brush over exposed surfaces and under the rim, allow to stand for 10 minutes and flush.

From Use Solution: Empty toilet bowl or urinal and apply ½ - 1 oz. per gallon use-solution to exposed surfaces and under the rim with a cloth, mop, sponge or mechanical spray. Brush or swab thoroughly; allow to stand for 10 minutes and flush.

For Heavy Duty Use: Empty toilet bowl or urinal and apply 2 oz. per gallon use-solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer, brush or swab thoroughly, allow to stand for 10 minutes and flush.

VETERINARY PRACTICE / ANIMAL CARE / ANIMAL LABORATORY / ZOOS / PET SHOPS / KENNELS DISINFECTION / VIRUCIDAL* DIRECTIONS: For cleaning and disinfecting hard non-porous surfaces: equipment used for feeding or watering animals, utensils, instruments, cages, kennels, stables, catteries, pens, stalls and etc. Remove all animals and feeds from premises, animal transportation vehicles, crates, pens, stalls and etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of ½ - 1 oz. of this product per gallon of water for a period of 10 minutes. Use 2 oz. per gallon of water to kill Adenovirus Type 5 and Type 7. Use 8 oz. per gallon of water to kill Feline Panleukopenia Virus. For marine reoviruses use dilution must be made with deionized water. Wipe or allow to air dry.
Immerse all halters, ropes, and other types of equipment used in handling and restraining animals as well as forks, shovels, and scrapers used to remove litter and manure. Thoroughly scrub all treated surfaces, then rinse all surfaces that come in contact with food, including equipment used for feeding or watering, with potable water before reuse. Ventilate buildings, animal enclosures, vehicles and other closed spaces. Do not house animals or empty equipment until treatment has been absorbed, set or dried.

POULTRY AND SWINE PREMISE DISINFECTION / VIRUCIDAL DIRECTIONS: Remove all animals and feeds from premises, vehicles and enclosures such as coops and crates. Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use ½ - 1 oz. of this product per gallon of water. Use 2 oz. per gallon of water to kill Adenovirus Type 5 and type 7. Saturate surfaces with the disinfecting solution for 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, coops and other closed spaces. Do not house animals or empty equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, managers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

TO CLEAN AND DEODORIZE: Mix ½ - 1 oz. of this product per gallon of water to clean and deodorize surfaces. Apply using a cloth, mop, sponge or sprayer. Wipe or allow to air dry.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE STORAGE: Open dumping is prohibited. Store in original container in areas inaccessible to children.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Fill container ¼ full with water and reclose the container. Agitate vigorously, and dispose of rinsate consistent with pesticide disposal instructions. Repeat two more times. Then offer for recycling if available or puncture and dispose in sanitary landfill or by other procedures approved by state and local authorities. Follow pesticide disposal instructions for rinsate. If not triple rinsed, these containers are acute hazardous wastes and must be disposed in accordance with local, state, and federal regulations.

ECOLAB® 6100281

Oasis™ 499 HBV Disinfectant

NET CONTENTS: 2.5 US GAL (9.46 L) Oasis™

ACTIVE INGREDIENTS:	
Octyl decyl dimethyl ammonium chloride	6.510%
Dioctyl dimethyl ammonium chloride	2.604%
Didecyl dimethyl ammonium chloride	3.906%
Alkyl (C12, 50%; C12, 40%; C14, 10%) dimethyl benzyl ammonium chloride	8.680%
OTHER INGREDIENTS:	78.300%
TOTAL:	100.000%

KEEP OUT OF REACH OF CHILDREN

DANGER

See side panel for precaudionary statements and first aid.

ECOLAB® 6100281

Oasis™ 499 HBV Disinfectant

NET CONTENTS: 2.5 US GAL (9.46 L) Oasis™

ACTIVE INGREDIENTS:
Octyl decyl dimethyl ammonium chloride 6.510%
Dioctyl dimethyl ammonium chloride 2.604%
Didecyl dimethyl ammonium chloride 3.906%
Alkyl (C12, 50%; C12, 40%; C14, 10%) dimethyl benzyl ammonium chloride 8.680%
OTHER INGREDIENTS: 78.300%
TOTAL: 100.000%

KEEP OUT OF REACH OF CHILDREN

DANGER

See side panel for precaudionary statements and first aid.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin, or on clothing. Wear protective eyewear (goggles, face shield or safety glasses), protective clothing and protective rubber or chemical resistant gloves. Harmful if swallowed and/or if absorbed through the skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.
FOR EMERGENCY MEDICAL INFORMATION, CALL TOLL-FREE 1-800-328-0026
OUTSIDE NORTH AMERICA, CALL 1-651-222-5352
Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor or going for treatment.

EPA Reg. No. 6836-78-9677
EPA Est. No.: 9677-IL-2 (L), 303-IN-1 (L), 9677-MV-1 (V), 9677-CA-2 (R), 9677-TX-1 (D), 5389-NC-1 (D), 9677-GA-1 (R), 9677-MN-1 (P)
Superscript refers to first letter of date code.

Distributed by:
Ecolab Inc., 1 Ecolab Place - St. Paul, MN 55102 USA
© 2019 Ecolab USA Inc. All rights reserved
Made in USA - 7404403/5409/0819

0 25469 57368 1

5 1/2"

5 1/8"

31 3/8"

DIELINE #
57129

REV #	DATE	DESCRIPTION	PROOF BY
Orig.	02/02/94	Created from Original Container	dfb/jbd
Rev A	10/02/09	Revised Container	dfb/rb
Rev B	10/06/09	Finalized Container	dfb/rb

SleeveCo
195 LUMPKIN CAMPGROUND RD., N
MOUNTAIN VIEW, GA 30052
(770) 279-0121 FAX: (770)279-0119
www.sleeveco.com

Ecolab:2 1/2 Gallon F-Style
Layflat: 15 11/16"
Height: 5 1/2"
Container#: BO4417& 4418