

ICBpharma

NAZWA PLIKU: ID13 Ensystex AUSTRALIA Attrathor 500ml BOOKLET 80x75mm REV-13-202305-1-3-4



MATERIAŁ: folia

WYMIARY: 80 x 75 mm

KOLORYSTYKA: CMYK

CUT

SIZE



BOOKLET AWERS

DIRECTIONS FOR USE: RESTRAINT. DO NOT apply to excessively wet soils, immediately after or during heavy rain, to avoid runoff of the chemical.			
SITUATION	PEST	RATE	CRITICAL COMMENTS
Internal & external areas & surrounds of Domestic, Commercial, Public and Industrial buildings, facilities & structures (including but not limited to food processing establishments, shops, factories, farm buildings, ships, trains, aeroplanes, offices, schools, storerooms, hospitals, barracks, private houses, and external areas associated with such establishments.)	Cockroaches	50 mL per 5 L water	Inspect premises thoroughly and locate all insect harbourages. Apply directly into harbourages such as cracks and crevices away from where it is likely to be removed by cleaning programs. Numerous smaller placements will provide faster control than fewer larger spots, especially for German cockroach control. Apply behind sinks, stoves, and pipes paying particular attention to all cracks and crevices which might contain insects. Repeat treatments when pests begin to reappear.
	Ants		Apply prepared spray to indoor and outdoor surfaces where ants are tracking. Internal harbourage sites include (but are not restricted to) areas such as walls, behind and under sinks, under furniture and indoor plants. External harbourage sites include (but are not restricted to) areas such as building exterior, eaves, walls, fences, garages, sheds, gazebos, ornamental plants, bushes, shrubs, hedges, shady or damp areas around buildings. Reapply as necessary. For indoor use, pay particular attention to dark protected areas such as cracks and crevices.
	Crawling insects and pests including: Earwigs (<i>Forficula auricularia</i>) House crickets (<i>Acheta domestica</i>) Silverfish (<i>Lepismatidae</i>) Slaters (<i>Porcellionidae</i>) Springtails (<i>Collembola</i>)		Apply directly onto pests as a spot treatment.
	Houseflies (<i>Musca domestica</i>)		Apply to surfaces where flies are likely to rest. On non-porous surfaces apply as a coarse spray at the rate of 1 L of emulsion per 20 m². On porous surfaces or use through power equipment, spray at the rate of 750 mL to 1 L of emulsion per 10 m² depending on porosity of surface. When treating non-porous and porous surfaces do not exceed the point of run-off.
	Millipedes (<i>Ommatolulus moreleti</i>)	115 mL per 5 L of water	Outdoor use only. Treat external surfaces 300 mm up and 300 mm out from where the building or structure touches the ground. Apply at the rate of 1 L of prepared suspension per 25 linear metres. Pay particular attention to potential entry points, such as weep holes, cracks and crevices. Also apply to where pests may be active away from the building including retaining walls, fences, garden beds, sheds etc. Apply at the rate of 1 L of prepared suspension per 16 m². May be applied directly to pests as a spot treatment.
NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.			