

WARNING

KEEP OUT OF REACH OF CHILDREN
READ LABEL COMPLETELY BEFORE OPENING OR USING

FOR USE ONLY AS A RODENTICIDE




18/10/2017

Storm® Secure

The fast single feed block bait
for rat and mouse control

Fixable Bait Blocks | Especially valuable for rodent control
scenarios where a fixable block is needed

- ✓ Proven single feed flocoumafen efficacy in fixable bait blocks
- ✓ Excellent palatability and durability
- ✓ Quick and easy to use
- ✓ Maximum value, least cost pulse baiting

3 kg

 **BASF**
We create chemistry

Contains 0.05 g/kg flocoumafen in the form of a ready-to-use bait.

® Registered trademark of BASF

Hazard Classification

6.1D, 6.9B, 9.3C



WARNING

HARMFUL. May be harmful if swallowed. May cause organ damage through prolonged or repeated oral exposure at high doses. **HARMFUL TO TERRESTRIAL VERTEBRATES.** May be harmful to pets and domestic animals if swallowed.

18/10/2017

PRECAUTIONS

STORAGE: Store in original container, tightly closed, in a cool, well-ventilated place, locked up, out of reach of children and away from foods and animal feeds.

TRANSPORT: DO NOT carry more than 3 kg of this product on a passenger service vehicle.

SPILLAGE: Prevent further spillage and product from entering waterways. Wear gloves. Collect spilled baits and place in sealable container for disposal. Wash affected area with water and detergent. Absorb and collect washings and place in the same sealable container as the spilled material. Dispose of the waste safely in a suitable landfill.

PERSONAL PROTECTION: When handling baits, wear suitable impervious gloves. Gloves are also recommended to help protect against rodent-borne diseases.

GENERAL SAFETY AND HYGIENE MEASURES: Avoid contact with eyes, skin and clothing. DO NOT eat, drink or smoke while using this product. Wash hands and exposed skin thoroughly after handling and before meals. After each day's work wash gloves.

USE: Place baits inside bait stations, or attach with wire or nails, at each bait location. PREVENT ACCESS TO THE BAIT by children, birds, livestock and non-target animals (particularly cats, dogs, pigs and poultry). DO NOT allow movement of the baits outside the bait location to occur. DO NOT place baits in areas liable to flooding. Look for and burn or bury all rodent carcasses. Remove all remains of the bait blocks and bait containers after treatment and dispose of them in a responsible manner.

DISPOSAL: DO NOT contaminate any water supply with bait or used container. DO NOT reuse this container for any purpose. Ensure container is completely empty, and make it unsuitable for further use. Scrub and triple rinse empty container and recycle through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise crush and bury in a suitable landfill. For information on disposal of unused, unwanted product, contact the local council.

FIRST AID AND MEDICAL ADVICE

The active ingredient flocoumafen is an indirect blood anticoagulant. Ingestion of a relatively large quantity of bait would be required before a toxic effect is expected. In general, an increased tendency to bleed could indicate poisoning. If you feel unwell, seek medical advice. **IF SWALLOWED:** DO NOT induce vomiting. In case of ingestion or an increased tendency to bleed, contact the National Poisons Centre 0800 POISONS (0800 764 766) or a doctor immediately. Show the product label to the doctor. **IF ON SKIN:** Wash with plenty of soap and water.

NOTE FOR PHYSICIANS ONLY: Vitamin K1 (phytomenadione) is antidotal. In cases of suspected poisoning, determine prothrombin time not less than 18 hours after consumption. If elevated, administer Vitamin K1 in divided doses, 40 mg/day for adults, 20 mg/day for children. Continue until prothrombin time normalises. Treatment may be required daily for several weeks. Withdraw antidote with care and monitor prothrombin times for two weeks, and resume treatment if elevation occurs in that time. Note: Vitamin K3 is not effective.

DOMESTIC ANIMAL POISONING

In cases of accidental poisoning by STORM SECURE consult a veterinarian who will advise of suitable Vitamin K1 antidote therapy.

GENERAL INFORMATION

STORM SECURE can be used both indoors and outdoors. Rats and mice that are resistant to warfarin are controlled, and no pre-baiting is required. A lethal dose can be readily consumed in a single feed but death may not occur for up to seven days afterwards. STORM SECURE contains Bitrex[®], a human taste deterrent, to help prevent accidental human consumption.

- **Exposure (when used on stocked land):** By law, all practicable measures must be taken to minimise all access to baits by livestock, pets and other non-target animals.
- **Carcasses:** By law, access by livestock, pets and other non-target animals must be prevented by collecting and burying or burning poisoned carcasses where practical, or necessary.
- In all instances when using this product, appropriate best practice guidelines must be followed where these exist.
- Failure to comply with these label directions is an offence under the ACVM Act.

DIRECTIONS FOR USE

Baits must only be placed in areas frequented by the target pests. DO NOT place baits in areas accessible to children and non-target animals, or in areas liable to flooding. Alternative food sources should be removed or reduced as much as possible. The baits should preferably be placed in bait stations or covered to prevent access by domestic pets, birds and non-target animals, and to prevent them becoming unpalatable. Suitable covers would include drain pipes, tiles or boards leant against, especially inside, walls. Rodents feel more secure when feeding under cover. Use gloves to avoid touching the bait with hands as this may deter rats and mice from eating it.

Rats: Select suitable bait locations, such as in active rat holes, along runs and under rubbish. Try to establish a barrier of bait locations between living and feeding areas. Bait locations may be as close as 5 metres apart in heavy infestations. Attach 3 STORM SECURE blocks at each bait location on wire or nails* or inside rat bait stations. Also place 1 bait block in each rat hole. Inspect the bait locations after 3 days and select new locations where there has been no take. Inspect again 4 days later and replace any blocks that have been eaten. Place a fresh block in each active rat hole. Repeat the inspection of all bait locations at 7 day intervals; placing fresh blocks on each occasion only when blocks have been eaten. Normally only 3 to 4 baiting rounds (pulses) are required.

Mice: Mice have very limited home ranges and do not need to drink. They are more difficult to control than rats because their feeding patterns are more erratic. Selection of bait locations is therefore even more important than for rats. Attach single STORM SECURE blocks on wire or nails*, or inside mouse bait stations, up to 2 metres apart in areas where activity is obvious, particularly where droppings are seen. In domestic premises, secure single blocks on wire or nails, or inside mouse bait stations, in sheltered positions, inaccessible to children and pets, where mice are active such as under flooring, inside ceiling cavities, along pipe runs and conduits and behind skirting boards, furniture and appliances. Inspect bait locations and replace blocks in the same sequence as for rats.

* Do not use a wire, nail or screw thickness greater than 2.5 mm.

CONDITIONS OF SUPPLY

Read conditions of supply before buying and using. If these conditions are unacceptable, return at once unopened. All goods supplied by BASF New Zealand Limited are of a high grade and conform to the chemical description of the label. However, to the extent permitted by law, neither BASF New Zealand Limited, nor any of its Distributors, accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential, howsoever arising, in connection with the supply of those goods including without limitation their storage, application, handling or use. BASF New Zealand Limited, or any of its Distributors, makes no warranties of any kind, express or implied, with respect to fitness for purpose (including crop injury, ineffectiveness or any other unintended consequence), merchantable quality or any other matter with respect to these goods. When you are a consumer as defined in the Consumer Guarantees Act 1993 (the "Act") these conditions will be subject to the provisions of the Act. However, the Act will not apply to goods supplied to you for business purposes.

Approved pursuant to the HSNO Act 1996, Code HSR001601.

See www.epa.govt.nz for approval controls.

Registered pursuant to the ACVM Act 1997, No. V9219.

See www.foodsafety.govt.nz for registration conditions.

MPI Approved Type C (All animal product except dairy).

MPI Approved for use in farm dairies and recognized for use in dairy processing.

Expiry Date: 3 years from DOM

Date of Manufacture:

Batch No.:



18/10/2017



Registered to and distributed by:

BASF New Zealand Limited
P.O. Box 407, Auckland 1140
Phone: 0800 932 273
Fax: 0800 940 000

EMERGENCY PHONELINK
24 Hour Advice in an Emergency Only
0800 944 955

® Registered Trademark of BASF

®1 Registered trademark of Johnson Matthey Plc., UK.

© Copyright 2017, BASF New Zealand Limited

IIIIIIII BAR CODE IIIIIIII

