

Main Panel

ECOTOXIC
READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN

 Maxforce[®] Fusion
COCKROACH GEL
ACTIVE INGREDIENT: 21.5 g/kg IMIDACLOPRID in the form of a ready to use gel

30 g NET

For the control of cockroaches in the situations specified in the
Directions for Use.

APPROVED PURSUANT TO THE HSNO ACT 1996, No. HSR000676
See www.epa.govt.nz for approval controls.

MPI Approved Type D (All animal products except dairy).



Side Panels

HSNO CLASSIFICATIONS: 6.9B, 9.1B, 9.2C, 9.4A**WARNING**

May cause organ damage from repeated oral exposure.
 Toxic to aquatic life with long lasting effects.
 Harmful to the soil environment.
 Very toxic to terrestrial invertebrates.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. Wash hands after use.

STORAGE AND DISPOSAL

The syringe must not be sold separately. Do not destroy carton while product still remains. Keep out of reach of children. Do not store in direct sunlight. Store in the closed, original carton, in a cool dry place. Completely empty syringe in normal use. Dispose of empty syringe and carton by wrapping in paper, placing in plastic bag and putting in garbage.

When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product which is older than this.

ENVIRONMENTAL PROTECTION

Do not apply this product onto or into water. Do not contaminate streams, rivers or waterways with this product or used container. Do not use in a manner contrary to the label which may contaminate the soil and cause harm to terrestrial invertebrates. Collect spillage.

EMERGENCY RESPONSE

For specialist advice in an emergency call 0800 734 607 (24 hours).

FIRST AID

For advice contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately.

It is an offence to use this product on animals.

DIRECTIONS FOR USE

Situations	Pest	Rate	Critical Comments
Barracks, farm buildings, food processing establishments, factories, homes, hospitals, meatworks (non- product areas), offices, restaurants, schools, shops, storerooms, transport vehicles (including aircrafts) & facilities	<u>Small cockroaches</u> e.g. German Cockroach (<i>Blattella germanica</i>)	<u>Light infestations</u> 1 placement per m ² <u>Heavy infestations</u> 2 placements per m ² 1 placement = a 5 mm ball (about the size of a small pea or approx. 0.1 g) or a line about 2.5 cm long	Apply gel in out of the way locations where cockroaches occur e.g., under fridges, near water heaters, behind microwave ovens, garbage bins etc. Re-assessment of treated areas one month after initial baiting is recommended to evaluate if additional baiting is required. Bait placements should be replenished when consumed or at least every three months as long as pest populations persist.
	<u>Large cockroaches</u> e.g. American Cockroach and Australian Cockroach (<i>Periplaneta</i> spp.)	<u>Light infestations</u> 2 placements per m ² <u>Heavy infestations</u> 3 placements per m ²	<i>Periplaneta</i> spp. require more bait because of the quantity which can be consumed by a single cockroach. Reassess baiting effectiveness as above.

GENERAL INSTRUCTIONS

Apply this product using a suitable application device. Gel should be applied out of reach of children, pets and other animals. Do not place gel on surfaces which are regularly washed.

PRECAUTIONS

Do not allow to contact food, livestock feed or their water. Do not place gel on surfaces where food is prepared or on utensils that are used for food preparation, on food packing materials or on surfaces where food packing materials are stored.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet.

RECORD KEEPING

Records of use, as described in the current version of NZS 8409 Management of Agrichemicals, must be kept if more than 3 kg is applied or discharged within 24 hours in a place where the substance is likely to enter air or water and leave the place.

CONDITIONS OF SUPPLY

READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING. IF THESE CONDITIONS ARE UNACCEPTABLE RETURN AT ONCE UNOPENED.

To the extent permitted by law Bayer does not accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with these goods, including without limitation their storage, application, handling or use. Bayer does not make any warranties (express or implied) with respect to fitness for purpose, merchantable quality or any other matter. To the extent only that Bayer is prevented by law from contracting out of the Consumer Guarantees Act then these conditions apply subject to that Act. The Act shall not apply where these goods are acquired for business purposes.

Maxforce® is a Registered Trademark of the Bayer Group.

Bayer CropScience Pty Ltd
Level 1, 8 Redfern Rd,
Hawthorn East, Vic, 3123
Australia
+61 3 9248 6888
www.es.bayer.com.au

Distributed by Garrards (NZ) Ltd.
Unit 4/27B Cain Road,
Penrose,
New Zealand
Telephone (09) 526 5232
www.garrards.com.nz

Batch No.
Date of Manufacture

Syringe label

READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN



Maxforce[®] Fusion

COCKROACH GEL

ACTIVE INGREDIENT: 21.5 g/kg IMIDACLOPRID
in the form of a ready to use bait

30 g NET

For the control of cockroaches in the situations specified in the Directions for Use.

APPROVED PURSUANT TO THE HSNO ACT 1996, No. HSR000676

See www.epa.govt.nz for approval controls.

MPI Approved Type D (All animal products except dairy).

Before using product read directions on the outer pack. Sale of this syringe except in outer carton is illegal. DO NOT destroy carton while product still remains.

HSNO CLASSIFICATIONS: 6.9B, 9.1B, 9.2C, 9.4A

WARNING

May cause organ damage from repeated oral exposure.

Toxic to aquatic life with long lasting effects.

Harmful to the soil environment.

Very toxic to terrestrial invertebrates.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. Wash hands after use.

STORAGE AND DISPOSAL

The syringe must not be sold separately. Do not destroy carton while product still remains. Keep out of reach of children. Do not store in direct sunlight. Store in the closed, original carton, in a cool dry place. Completely empty syringe in normal use. Dispose of empty syringe and carton by wrapping in paper, placing in plastic bag and putting in garbage.

When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product which is older than this.

ENVIRONMENTAL PROTECTION

Do not contaminate streams, rivers or waterways with this product or used container. Do not use in a manner contrary to the label which may contaminate the soil and cause harm to terrestrial invertebrates. Collect spillage.

EMERGENCY RESPONSE

For specialist advice in an emergency call 0800 734 607(24 hours).

Distributed by Garrards (NZ) Ltd., Telephone (09) 526 5232.