



International

WARNING

KEEP OUT OF REACH OF CHILDREN
READ LABEL BEFORE USE



ROBAN

RODENTICIDE PASTE

Contains difenacoum 0.005% w/w and denatonium benzoate 0.001% w/w.

Specially developed for New Zealand

Ready for use rodenticide bait for the control of brown or Norway rats (*Rattus norvegicus*) and mice (*Mus musculus*) in and around industrial, commercial, public services, agricultural and domestic buildings.

6kg



ROBAN RODENTICIDE PASTE

Contains difenacoum 0.005% w/w and denatonium benzoate 0.001% w/w.

6kg

GENERAL INFORMATION

The ROBAN RODENTICIDE PASTE bait is formulated for use indoors and outdoors. It will control rats and mice. It is approved for use as a ground-based application in a bait station. Use only with an appropriate bait station. Most rodents die within 10 days.

Baiting strategy.

- Eliminate as much as practical, all alternative food sources.
- Locate baits under cover, along walls or rafters and in dark sheltered areas where there are signs of rodent activity.
- For mice - use up to 45gm (1-3 bait sachets) in a bait station placed 5m apart, or 2m apart for high infestations.
- For rats - use up to 195gm (2-13 bait sachets) in a bait station placed 10m apart reducing to 5m apart for high infestations.
- Record location and number of bait stations to assist in checking and later removal.
- Inspect baits regularly.
- Replace baits every 7 days until no more is taken. If baits are ingested more quickly then replace baits more frequently.
- Baiting for at least 2 weeks will be necessary to reduce rodent numbers.
- Ensure untaken baits are removed and safely disposed of when rodent activity has ceased.

By law this product must only be used in a bait station and as specified on the label.

Under regulation 13.19 of the Health and Safety at Work (Hazardous Substances) Regulations 2017, WorkSafe has specified a period of time for which a PCBU with management or control of work involving a vertebrate toxic agent (VTA) being applied or laid will must ensure that compliant signage is displayed. Please refer to www.worksafe.govt.nz for more details. Further information can also be obtained from the Pest Management Association of New Zealand (PMANZ) www.pmanz.nz

DIRECTIONS FOR USE

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it. Where possible, prior to the treatment inform any bystanders (e.g., users of the treated areas and their surroundings) about the rodent control campaign. Consider preventative control measures (e.g. plug holes, remove potential food and drink as far as possible) to improve product intake and reduce the likelihood of re invasion. Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation. The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve. The product should be placed in the immediate vicinity of places where rodent activity has previously occurred (e.g. rodent runs, nesting sites, holes, burrows etc.).

FOR AMATEURS

Rats: The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

Mice: The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

Baits must be placed to minimise the exposure to non-target species and children. If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation. The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control. For outdoor use, baiting points must be covered and placed in strategic sites to minimise exposure to non-target species. Do not use this product in pulsed baiting treatments. Protect bait from atmospheric conditions. Place the baiting points in areas not liable to flooding.

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided. Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened. Where possible, bait stations must be fixed to the ground or other structures. Bait should be secured so that it cannot be dragged away from the bait station. Do not wash the bait stations or utensils used in covered and protected bait points with water between applications. Place the product out of the reach of children, birds, pets and farm animals and other non-target animals. Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these. When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product. Do not use in areas where resistance to the active substance is suspected. Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment. Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.

FOR AMATEURS

Using this product should eliminate rodents within 35 days. In case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), advice should be sought from the product supplier or a professional pest control service. To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals.

PERMANENT BAITING

Permanent baiting is strictly limited to sites with a high potential for re invasion when other methods of control have proven insufficient. The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation. Sites under a permanent baiting regime should be inspected regularly in accordance with product label directions. The period between visits should be determined by the technician in charge but will not be longer than every four weeks when permanent baiting is conducted outdoors. If after a treatment period of 35 days baits continue to be consumed and no decline in rodent activity is observed, the likely cause must be determined. Where other elements have been excluded, it is likely that there are resistant rodents so consider the use of a non anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

WARNING

May be harmful if swallowed. May cause damage to the blood system with long-lasting effects.

PRECAUTIONS

Avoid skin contact. Wear gloves. Wash hands and exposed skin after handling. Avoid contamination of any water supply with product or empty container. Do NOT lay baits in places accessible to children, pets, domestic animals, livestock or birds. Do NOT lay baits in bush area where stock or feral animals have access to baits. Dispose of any carcasses by burial or by wrapping in paper, placing in a plastic bag and putting in the garbage.

FIRST AID

If swallowed, do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor, immediately. Treatment of domestic animals: Depending on the amount eaten, this bait may kill or injure dogs, cats, pigs, poultry and wildlife. Vitamin K1 is an effective antidote against accidental poisoning of domestic animals.

Note to Doctors and Veterinarians: Difenacoum is a long-lasting anticoagulant compound which if ingested by humans, domestic animals or pets, can reduce the clotting ability of the blood and haemorrhage may result. Vitamin K1 (Phytomenadione) should be administered urgently by medical, appropriate para-medical personnel or veterinarians. If administered intravenously the injection must be given slowly.

STORAGE

Store in original container tightly closed and in a locked, dry, cool area away from direct sunlight and foodstuffs.

DISPOSAL

Ensure container is completely empty. Do NOT use container for any other purpose. Burn empty container if circumstances such as wind direction permit, otherwise crush and bury in a suitable landfill. Wrap untaken or unused baits, place in plastic bag and put into the garbage.

Registered pursuant to the ACVM Act 1997, No.V9689 See www.foodsafety.govt.nz for registration conditions.

EPA Approval Number: HSR101550 **GHS Classifications:** Reproductive Tox. Cat 1, Specific Target Organ Tox. – RE Cat 2, Aquatic Chronic Cat 4

In a transport emergency dial 111, Police or Fire Brigade.



International

Registered by:

PelGar International Ltd.
Unit 13 Newman Lane, Alton,
Hampshire, GU34 2QR. UK.
Tel. +44 (0)1420 80744
www.PelGar.co.uk

ROBAN is a registered trade mark of PelGar International Ltd.

New Zealand Agent:

PelGar International (NZ) Ltd.
PO Box 11825
Ellerslie, Auckland 1542.
Tel. + 64 21 670 105

New Zealand Distributor :

PelGar International (NZ) Ltd.
PO Box 11825
Ellerslie, Auckland 1542.
Tel. + 64 21 670 105



TOXIC



ECOTOXIC

BATCH:

DOM:

DOE: