Components	Species	Calculated/Test Results
2,2',2''-Nitrilotriethanol (CAS 102-71-6)		
Acute		
Dermal		
LD50	Rabbit	> 20000 mg/kg
Oral		
LD50	Guinea pig	5300 mg/kg
	Rat	8 g/kg
Other		
LD50	Mouse	1450 mg/kg
POTASSIUM HYDROXIDE (CAS 1310-58-3)		
Acute		
Oral		
LD50	Rat	273 mg/kg
		1.23 g/kg

Skin corrosion/irritation Prolonged skin contact may cause temporary irritation.

Serious eye damage/eye irritation Direct contact with eyes may cause temporary irritation.

Respiratory or skin sensitization

Respiratory sensitization Not a respiratory sensitizer.

Skin sensitization This product is not expected to cause skin sensitization.

Germ cell mutagenicity No data available to indicate product or any components present at greater than 0.1% are mutagenic or genotoxic.

Carcinogenicity Not classifiable as to carcinogenicity to humans.

IARC Monographs. Overall Evaluation of Carcinogenicity

Not listed.

OSHA Specifically Regulated Substances (29 CFR 1910.1001-1053)

Not listed.

Reproductive toxicity This product is not expected to cause reproductive or developmental effects.

Specific target organ toxicity - single exposure Not classified.

Specific target organ toxicity - repeated exposure Not classified.

Aspiration hazard Not an aspiration hazard.

Chronic effects Prolonged inhalation may be harmful. May be harmful if absorbed through skin.

Prolonged or repeated exposure may cause liver and kidney damage. These effects have not been observed in humans.

12. Ecological information

Ecotoxicity The product is not classified as environmentally hazardous. However, this does not exclude the possibility that large or frequent spills can have a harmful or damaging effect on the environment.

Ecotoxicity

Components		Species	Calculated/Test Results
2,2',2''-Nitrilotriethanol (CAS 102-71-6)			
Aquatic			
Crustacea	EC50	Water flea (Ceriodaphnia dubia)	565.2 - 658.3 mg/l, 48 hours
Fish	LC50	Fathead minnow (Pimephales promelas)	10610 - 13010 mg/l, 96 hours
POTASSIUM HYDROXIDE (CAS 1310-58-3)			
Aquatic			
Fish	LC50	Western mosquitofish (Gambusia affinis)	80 mg/l, 96 hours

Persistence and degradability No data is available on the degradability of any ingredients in the mixture.

Bioaccumulative potential