

Decomposition temperature Not available.

Viscosity 16.27 cSt

10. Stability and reactivity

Reactivity The product is stable and non-reactive under normal conditions of use, storage and transport.

Chemical stability Material is stable under normal conditions.

Possibility of hazardous reactions No dangerous reaction known under conditions of normal use.

Conditions to avoid Avoid temperatures exceeding the flash point. Contact with incompatible materials.

Incompatible materials Strong oxidizing agents.

Hazardous decomposition products Upon decomposition, this product emits carbon monoxide, carbon dioxide and/or low molecular weight hydrocarbons.

11. Toxicological information

Information on likely routes of exposure

Inhalation May cause irritation to the respiratory system. Prolonged inhalation may be harmful.

Skin contact Causes skin irritation.

2-Butoxy ethanol may be absorbed through the skin in toxic amounts if contact is repeated and prolonged. These effects have not been observed in humans.

Eye contact Causes serious eye irritation.

Ingestion Droplets of the product aspirated into the lungs through ingestion or vomiting may cause a serious chemical pneumonia.

Symptoms related to the physical, chemical and toxicological characteristics Aspiration may cause pulmonary edema and pneumonitis. Headache. Nausea, vomiting. Diarrhea. Severe eye irritation. Symptoms may include stinging, tearing, redness, swelling, and blurred vision. May cause respiratory irritation. Skin irritation. May cause redness and pain.

Information on toxicological effects

Acute toxicity May be fatal if swallowed and enters airways.

Components	Species	Calculated/Test Results
------------	---------	-------------------------

2-BUTOXYETHANOL (CAS 111-76-2)

Acute

Dermal

LD50	Rabbit	400 mg/kg
------	--------	-----------

Inhalation

LC50	Mouse	700 ppm, 7 Hours
	Rat	486 ppm, 4 Hours
		450 ppm, 4 Hours

Oral

LD50	Guinea pig	1.2 g/kg
	Mouse	1519 mg/kg
		1.2 g/kg
	Rabbit	0.32 g/kg
	Rat	560 mg/kg
		1.48 g/kg

Other

LD50	Mouse	1130 mg/kg
	Rabbit	280 mg/kg
	Rat	550 mg/kg
		340 mg/kg