



US 20250257360A1

(19) United States

(12) Patent Application Publication

Ruiz Echevarria et al.

(10) Pub. No.: US 2025/0257360 A1

(43) Pub. Date: Aug. 14, 2025

(54) OLIGONUCLEOTIDE INTERFERENCE  
TREATMENTS OF PROSTATE CANCER(71) Applicant: The Board of Regents of the  
University of Oklahoma, Norman, OK  
(US)(72) Inventors: Maria Jesus Ruiz Echevarria,  
Edmond, OK (US); Joshua Moses  
Corbin, Oklahoma City, OK (US)

(21) Appl. No.: 18/855,017

(22) PCT Filed: Apr. 7, 2023

(86) PCT No.: PCT/US2023/065490

§ 371 (c)(1),

(2) Date: Oct. 8, 2024

## Related U.S. Application Data

(60) Provisional application No. 63/329,132, filed on Apr.  
8, 2022.

## (52) U.S. Cl.

CPC ..... C12N 15/1138 (2013.01); A61P 35/00  
(2018.01); C12N 2310/141 (2013.01); C12N  
2310/321 (2013.01); C12N 2310/322  
(2013.01); C12N 2310/351 (2013.01); C12N  
2310/531 (2013.01)

(57)

## ABSTRACT

Interfering nucleic acids and methods of their use in treat prostate cancers, such as aggressive prostate cancers. The nucleic acids may be, for example, short interfering RNA (siRNA), short hairpin RNA (shRNA), antisense RNA, DNA, antisense DNA, Chimeric Antisense DNA/RNA, and microRNA (miRNA) oligonucleotides. The oligonucleotide has a seed sequence that is complementary to a sequence of either a gene or an mRNA encoding an androgen receptor (AR) coregulator or a fragment having AR coregulator activity. The nucleic acid compound may have a non-natural modification in the oligonucleotide, and/or an organic moiety conjugated to the oligonucleotide. The oligonucleotide has inhibitory activity against the expression or biological activity of the AR coregulator.

Specification includes a Sequence Listing.

## Publication Classification


## (51) Int. Cl.

C12N 15/113

(2010.01)

A61P 35/00

(2006.01)






FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 3




FIG. 4A




FIG. 4B





FIG. 4C




FIG. 4D




FIG. 5




FIG. 6

## OLIGONUCLEOTIDE INTERFERENCE TREATMENTS OF PROSTATE CANCER

### CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Patent Application Ser. No. 63/329,132, filed Apr. 8, 2022, the content of which is expressly incorporated herein by reference in its entirety.

### SEQUENCE LISTING STATEMENT

[0002] The ASCII file, entitled "820233-02546\_seq-listing", created on Apr. 6, 2023, comprising 617,171 bytes, submitted concurrently with the filing of this application is incorporated herein by reference.

### BACKGROUND

[0003] Ribonucleic acid interference (RNAi) via small interfering ribonucleic acids (siRNA) and short hairpin RNAs (shRNA) is a useful tool for knocking-down messenger RNA (mRNA) expression and studying gene function. Both siRNA and shRNA utilize the evolutionarily conserved microRNA (miRNA) gene regulatory process to decrease targeted mRNA expression. In short, after shRNA processing by the endogenous miRNA processing enzymes, one strand of the resulting mature double stranded RNA is loaded onto the RNA-induced silencing complex (RISC). Similar to endogenous mature miRNAs, the loaded RNA guides RISC to target transcripts via sequence complementarity, resulting in subsequent transcript degradation and/or translation inhibition. The RNA strand that is intended to guide RISC to the desired target transcript is called the guide strand, while the complementary strand is called the passenger. The seed sequence, nucleotides 2-7, on the 5' end of the mature guide strand is the most important sequence for determining RISC transcript targeting. It has been found that, like miRNAs, numerous transcripts can be targeted by a single sh/siRNA via seed sequence complementarity, especially to the three-prime untranslated region (3' UTR) of target transcripts, often resulting in undesired off target effects (OTEs). For this reason, numerous strategies have been developed to determine whether RNAi-induced phenotypes are generated by knocking-down the target gene of interest or off target transcripts. These strategies include: the use multiple independent RNAi target sequences, and demonstration of phenotypic recovery using recombinant over-expression of the target gene. The discovery of alternate knockdown/knockout methods, such as the potent CRISPR Cas9 system, has also aided researchers in the study of gene function.

[0004] Prostate cancer (PCa) is a major public health concern, as it remains the most commonly diagnosed non-skin malignancy and the second leading cause of cancer related deaths in men. The androgen signaling is essential for cell growth, differentiation and survival during prostate development, homeostasis and cancer progression. This dependency is exploited in PCa therapy with androgen deprivation therapy and targeted androgen receptor (AR) inhibitors. While these therapies are initially beneficial, the majority of patients eventually relapse with a resistant and lethal form of the disease called castration-resistant PCa (CRPC). In most cases, CRPC cells retain AR expression and remain dependent on its activity for growth and survival,

and adaptations, such as, increased AR coregulator expression, AR amplification and/or mutation and constitutively AR splice variants that lack ligand binding, such as AR variant 7 (AR-V7), allow PCa cells to tolerate and grow in the androgen depleted environment. However, in an increasing subset of patients, CRPC cells lose AR expression and gain complete independence from AR transcriptional activity for growth and survival. Currently, the poor prognosis for patients with AR positive and AR negative CRPC underscores the need for the development of novel therapeutic agents and strategies. It is to such a need that the present disclosure is directed.

### BRIEF DESCRIPTION OF THE DRAWINGS

[0005] FIG. 1A shows Western blot analysis of nine independent shRNAs targeting 8 of 10 exons and 3'UTR of the TMEFF2 transcript. The schematic bar in the lower panel indicates targeted exon were exons 2, 3, and 5-10 and that untargeted exons were exons 1 and 4. Numbers above the bar indicate the corresponding shRNA name. shScramble control shRNA were transduced into LNCaP cells. Western blot analysis shows the levels of TMEFF2 (target gene), AR, and two androgen responsive proteins, PSA and FKBP5. Cells were grown in the presence and absence of 10 nM DHT for 24 hours to test androgen response. Calnexin was used as a loading control. Multiple shRNAs targeting TMEFF2 reduce AR and androgen responsive protein levels in LNCaP and 22Rv1 PCa cell lines.

[0006] FIG. 1B shows a Western blot analysis of PSA and TMEFF2 protein levels in response to increasing dose of shTMEFF2-3 and shScramble control shRNA in LNCaP cells four days after transduction (upper panel). Dose is presented as percentage of lentiviral supernatant in transduction media. Lower panel is a graphical representation of Calnexin normalized PSA and TMEFF2 levels relative to shScramble control at each dose. Band intensity was quantified using Biorad Image Lab. Multiple shRNAs targeting TMEFF2 reduce AR and androgen responsive protein levels in LNCaP and 22Rv1 PCa cell lines.

[0007] FIG. 1C shows a Western blot analysis in 22Rv1 cells showing protein levels of TMEFF2, AR (including both full length and the constitutively active AR V7 isoform) and FKBP5 in response to two TMEFF2 targeted shRNA (-2 and -3) in the presence and absence of 10 nM DHT. Calnexin was used as a loading control.

[0008] FIG. 1D shows a Western blot analysis of cell fraction lysates shows cytoplasmic and nuclear levels of both AR isoforms in response to shRNAs targeting TMEFF2 in 22Rv1 cells. Calnexin, an ER protein, was used as a loading and fractionation control for cytoplasmic/membrane lysates, and H3 was used as a loading and fractionation control for nuclear lysates.

[0009] FIG. 2A shows a Western blot analysis of TMEFF2, PSA and AR levels in C4-2B cells transfected with an empty vector (EV) or an overexpressing doxycycline-inducible recombinant TMEFF2 vector (TMEFF2o/e) and transduced with either shTMEFF2-3 or shScramble shRNA. 250 ng/ml doxycycline was used to induce TMEFF2 overexpression for 2 days prior and three days after shRNA transductions. Calnexin was used as a loading control.

[0010] FIG. 2B shows a Western blot analysis showing TMEFF2 protein levels in LNCaP cells expressing doxycycline inducible sgRNA (sgTMEFF2-1 and sgGFP) after 6

days in the presence or absence of 500 ng/ml doxycycline. Calnexin was used as a loading control.

[0011] FIG. 2C shows Western blot and RT qPCR analyses. TMEFF2 and PSA protein levels (upper panel), and TMEFF2 and KLK3 (PSA transcript) mRNA levels (lower panel), were measured by western blot and RT qPCR analysis in sgTMEFF2-1 knockdown and sgGFP control LNCaP Cas9 cells grown in the presence and absence of DHT. Calnexin was used as a loading control in western blot analysis. CANX and HPRT1 mRNA levels were used for normalization in RT qPCR analysis, and relative gene expression was normalized to the sgGFP control grown in the absence of DHT (-DHT) sample. N=3, error bars $\pm$ SD, \*p<0.05.

[0012] FIG. 2D shows Western blot analysis showing TMEFF2, PSA, FKBP5 and AR protein levels in LNCaP Cas9 sgGFP control and sgTMEFF2-1 knockdown cell lines, which were transduced with shScramble or shTMEFF2-3 shRNA and grown in the presence or absence of 10 nM DHT. Calnexin was used as a loading control. LNCaP Cas9 sgGFP and sgTMEFF2-1 cells were incubated in 500 ng/ml doxycycline for 2 weeks to induce TMEFF2 knockdown. Doxycycline was then removed from culture media for 2 weeks, at which time cells were transduced with shScramble or shTMEFF2 shRNA. FIG. 2A-D results demonstrate that TMEFF2-targeted shRNA mediated low androgen signaling phenotype is disconnected from TMEFF2 expression.

[0013] FIG. 3 is a flow chart illustrating workflow for determining whether shTMEFF2-3 works in a DISE-like mechanism, downregulating essential and AR coregulatory genes through shRNA seed matches to the 3'UTR.

[0014] FIG. 4A is a bar graph showing percent growth inhibition of targeted shRNAs relative to shScramble control in HEK293T, Panc-1, DU145, 22Rv1 and LNCaP cell lines. N=3, error bars $\pm$ SD, \*p<0.05, as calculated by two tailed T-test. TMEFF-targeted shRNA are toxic to transformed cells, and toxicity correlates with reduced expression of androgen responsive genes in LNCaP cells.

[0015] FIG. 4B is a bar graph showing percent viability of targeted shRNA relative to shScramble control in HEK293T, Panc-1, DU145, 22Rv1 and LNCaP cell lines. N=3, error bars $\pm$ SD, \*p<0.05, as calculated by two tailed T-test. TMEFF2-targeted shRNA are toxic to transformed cells, and toxicity correlates with reduced expression of androgen responsive genes in LNCaP cells.

[0016] FIG. 4C depicts a Western blot analysis showing procaspase 3 and cleaved caspase 3 levels in response to shRNA 4 days after transductions in LNCaP cells (upper panel), and a graph showing that relative % Caspase 3 cleavage (cleaved caspase 3/procaspase 3), as determined using Biorad Image Lab, correlates with relative viability detected for shRNA expressing LNCaP cell lines (lower panel).

[0017] FIG. 4D shows representative micrograph images demonstrating cell morphology in four LNCaP shRNA cell lines 4 days after transductions.

[0018] FIG. 5 is a schematic showing a strategy for a negative selection screen with pooled shRNAs to validate and optimize identification of toxic seed sequences.

[0019] FIG. 6 is a schematic showing a comparison of the conventional AR-inhibitory therapeutic strategy with the Androgen Network-Death Induced by Survival gene Elimination (AN-DISE) strategy disclosed herein. Left panel:

Current FDA approved therapeutic approaches in prostate cancer target androgen signaling either by inhibiting global androgen production (androgen deprivation therapy, ADT) or by direct AR inhibition (i.e. enzalutamide, Enz), resulting in the depletion of androgen regulated essential survival genes in PCa cells, and initial beneficial clinical response. However, most patients eventually develop resistance, and present with either AR+ or AR- CRPC. Right Panel: Using shTMEFF2-3 as an example, AN-DISE, directly reduces the expression of non-androgen regulated essential survival genes, as well as, androgen signaling regulatory genes, such as, AR, TAF1 and USP12, through shRNA seed complementarity to the 3'UTR. Resultant inhibition of androgen signaling leads to the reduction of androgen regulated essential survival genes, and increased PCa cell death. The global nature of the DISE mechanism, and the sensitivity of both AR+ and AR- CRPC cells, indicate resistance may be less probable compared to standard androgen signaling targeted therapies.

#### DETAILED DESCRIPTION

[0020] The present disclosure is directed to a novel therapeutic strategy for treating prostate cancer (PCa) that relies on large-scale RNA interference (RNAi) of a network of androgen receptor (AR) coregulators. RNA interference (RNAi) is a biological mechanism to modulate gene expression which is mediated by small RNA molecules that recognize and bind to target mRNA(s) of specific genes thereby inhibiting their expression or promoting their degradation. The present work shows that certain small RNAs can inhibit networks of androgen receptor (AR) co-regulators, promoting growth inhibition and death of PCa cells, which depend on the activity of the AR and its coregulators to grow. This death mechanism is specific to cancer cells and does not affect viability of normal cells, and can kill PCa cells that have become resistant to conventional therapies. As further shown below, this disclosure relates to the therapeutic use of toxic small RNAs to kill prostate cancer cells via RNAi of networks of AR-coregulators.

[0021] Death Induced by Survival gene Elimination (DISE) is a process in which certain small interfering ribonucleic acids (siRNAs) or short hairpin ribonucleic acids (shRNAs) induce toxicity specifically in transformed cells through off-target seed region complementarity to the three-prime untranslated region (3' UTR) of essential survival genes, resulting in reduced expression. Paradoxically, DISE was identified using shRNA derived from the CD95/CD95L tumor suppressor and subsequently observed using shRNAs from several other tumor suppressors. We hypothesized that a DISE mechanism targeting genes that are regulated by androgen signaling is relevant to PCa. Consequently we have demonstrated that RNAi to the transmembrane protein with EGF like and two follistatin like domains 2 (TMEFF2), an androgen regulated tumor suppressor gene with restricted expression to adult brain and prostate, triggers a DISE-like mechanism in PCa cells. The present disclosure shows that certain shRNAs targeted to TMEFF2 inhibit androgen signaling and are toxic to prostate cancer (PCa) cells through a similar off-target seed mediated mechanism that also targets androgen signaling regulatory genes, resulting in potent inhibition of androgen mediated transcriptional response and cell death. We call this PCa-specific DISE-like mechanism the Androgen Network-DISE (AN-DISE). Importantly, AN-DISE effectively kills castration resistant PCa (CRPC)

cells that arise as a consequence of conventional therapies targeting androgen signaling. Given the sensitivity of CRPC cells, and the general toxicity of essential gene depletion in cancer cells, AN-DISE provides a useful strategy for treating prostate cancer, particularly CRPC.

[0022] Before further describing various embodiments of the compounds, compositions, and methods of the present disclosure in more detail by way of exemplary description, examples, and results, it is to be understood that the compounds, compositions, and methods of present disclosure are not limited in application to the details of specific embodiments and examples as set forth in the following description. The description provided herein is intended for purposes of illustration only and is not intended to be construed in a limiting sense. As such, the language used herein is intended to be given the broadest possible scope and meaning; and the embodiments and examples are meant to be exemplary, not exhaustive. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting unless otherwise indicated as so. Moreover, in the following detailed description, numerous specific details are set forth in order to provide a more thorough understanding of the present disclosure. However, it will be apparent to a person having ordinary skill in the art that the present disclosure may be practiced without these specific details. In other instances, features which are well known to persons of ordinary skill in the art have not been described in detail to avoid unnecessary complication of the description. It is intended that all alternatives, substitutions, modifications and equivalents apparent to those having ordinary skill in the art are included within the scope of the present disclosure. While the compounds, compositions, and methods of the present disclosure have been described in terms of particular embodiments, it will be apparent to those of skill in the art that variations may be applied to the compounds, compositions, and methods and in the steps or in the sequence of steps of the methods described herein without departing from the concept, spirit, and scope of the inventive concepts.

[0023] All patents, published patent applications, and non-patent publications mentioned in the specification or referenced in any portion of this application, including U.S. Provisional Patent Application Ser. No. 63/329,132, filed Apr. 8, 2022, U.S. application Ser. No. 17/441,895 filed Sep. 22, 2021, PCT Application PCT/US2020/060753 filed Nov. 16, 2020, and U.S. Provisional Patent Application Ser. No. 62/935,521, filed Nov. 14, 2019, are herein expressly incorporated by reference in their entirety to the same extent as if each individual patent or publication was specifically and individually indicated to be incorporated by reference.

[0024] Unless otherwise defined herein, scientific and technical terms used in connection with the present disclosure shall have the meanings that are commonly understood by those having ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Where used herein, the specific term "single" is limited to only "one".

[0025] As utilized in accordance with the methods, compounds, and compositions of the present disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

[0026] The use of the word "a" or "an" when used in conjunction with the term "comprising" in the claims and/or

the specification may mean "one," but it is also consistent with the meaning of "one or more," "at least one," and "one or more than one." The use of the term "or" in the claims is used to mean "and/or" unless explicitly indicated to refer to alternatives only or when the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and "and/or." The use of the term "at least one" will be understood to include one as well as any quantity more than one, including but not limited to, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 100, or any integer inclusive therein. The term "at least one" may extend up to 100 or 1000 or more, depending on the term to which it is attached; in addition, the quantities of 100/1000 are not to be considered limiting, as higher limits may also produce satisfactory results. In addition, the use of the term "at least one of X, Y and Z" will be understood to include X alone, Y alone, and Z alone, as well as any combination of X, Y and Z.

[0027] As used herein, all numerical values or ranges include fractions of the values and integers within such ranges and fractions of the integers within such ranges unless the context clearly indicates otherwise. Thus, to illustrate, reference to a numerical range, such as 1-10 includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, as well as 1.1, 1.2, 1.3, 1.4, 1.5, etc., and so forth. Reference to a range of 1-50 therefore includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, etc., up to and including 50, as well as 1.1, 1.2, 1.3, 1.4, 1.5, etc., 2.1, 2.2, 2.3, 2.4, 2.5, etc., and so forth. Reference to a series of ranges includes ranges which combine the values of the boundaries of different ranges within the series. Thus, to illustrate reference to a series of ranges, for example, of 1-10, 10-20, 20-30, 30-40, 40-50, 50-60, 60-75, 75-100, 100-150, 150-200, 200-250, 250-300, 300-400, 400-500, 500-750, 750-1,000, includes ranges of 1-20, 10-50, 50-100, 100-500, and 500-1,000, for example. Reference to an integer with more (greater) or less than includes any number greater or less than the reference number, respectively. Thus, for example, reference to less than 100 includes 99, 98, 97, etc. all the way down to the number one (1); and less than 10 includes 9, 8, 7, etc. all the way down to the number one (1). A strand length in a range of 12 to 50 nucleotides refers to an oligonucleotide having at least 12 nucleotides and less than 51 nucleotides, and includes any range bounded by two different integers in said range of 12, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, and 50 nucleotides, including for example 18 to 25, 20 to 24, or 20-22.

[0028] As used in this specification and claims, the words "comprising" (and any form of comprising, such as "comprise" and "comprises"), "having" (and any form of having, such as "have" and "has"), "including" (and any form of including, such as "includes" and "include") or "containing" (and any form of containing, such as "contains" and "contain") are inclusive or open-ended and do not exclude additional, unrecited elements or method steps.

[0029] The term "or combinations thereof" as used herein refers to all permutations and combinations of the listed items preceding the term. For example, "A, B, C, or combinations thereof" is intended to include at least one of: A, B, C, AB, AC, BC, or ABC, and if order is important in a particular context, also BA, CA, CB, CBA, BCA, ACB, BAC, or CAB. Continuing with this example, expressly included are combinations that contain repeats of one or

more item or term, such as BB, AAA, AAB, BBC, AAABCCCC, CBBAAA, CABABB, and so forth. The skilled artisan will understand that typically there is no limit on the number of items or terms in any combination, unless otherwise apparent from the context.

[0030] Throughout this application, the terms “about” or “approximately” are used to indicate that a value includes the inherent variation of error in the disclosed embodiments, or the variation that exists among the study subjects. As used herein the qualifiers “about” or “approximately” are intended to include not only the exact value, amount, degree, orientation, or other qualified characteristic or value, but are intended to include some slight variations due to measuring error, manufacturing tolerances, stress exerted on various parts or components, observer error, wear and tear, and combinations thereof, for example. The term “about” or “approximately”, where used herein when referring to a measurable value such as an amount, a temporal duration, and the like, is meant to encompass, for example, variations of or  $\pm 25\%$ ,  $\pm 20\%$ , or  $\pm 10\%$ , or  $\pm 5\%$ , or  $\pm 1\%$ , or  $\pm 0.1\%$  from the specified value, as such variations are appropriate to perform the disclosed methods and as understood by persons having ordinary skill in the art. As used herein, the term “substantially” means that the subsequently described event or circumstance completely occurs or that the subsequently described event or circumstance occurs to a great extent or degree. For example, the term “substantially” means that the subsequently described event or circumstance occurs at least 90% of the time, or at least 95% of the time, or at least 98% of the time.

[0031] As used herein any reference to “one embodiment” or “an embodiment” means that a particular element, feature, structure, or characteristic described in connection with the embodiment is included in at least one embodiment and may be included in other embodiments. The appearances of the phrase “in one embodiment” in various places in the specification are not necessarily all referring to the same embodiment and are not necessarily limited to a single or particular embodiment.

[0032] Where used herein the term “active agent” refers to an oligonucleotide comprising a seed-sequence as described herein, or to a nucleic acid compound comprising an oligonucleotide comprising a seed sequence as described herein. The oligonucleotide may be an siRNA, shRNA, DNA or RNA antisense oligonucleotide, chimeric antisense DNA/RNA, or microRNA, or other biochemical or molecule that is biologically active as described herein. By “biologically active” is meant the ability to modify the molecular, biochemical, or physiological system of a cell, organ, or organism, without reference to how the active agent has its physiological effects. In one non-limiting embodiment, biologically active refers to the ability of an active agent to target and interfere with the normal function of one or more AR coregulators.

[0033] The term “pharmaceutically acceptable” refers to active agents, compounds and compositions which are suitable for administration to humans and/or animals without undue adverse side effects such as toxicity, irritation and/or allergic response commensurate with a reasonable benefit/risk ratio. The compounds or conjugates of the present disclosure may be combined with one or more pharmaceutically-acceptable excipients, including carriers, vehicles, and diluents which may improve solubility, deliverability,

dispersion, stability, and/or conformational integrity of the compounds or conjugates thereof.

[0034] As used herein, “pure,” or “substantially pure” means an object species is the predominant species present (i.e., on a molar basis it is more abundant than any other object species in the composition thereof), and particularly a substantially purified fraction is a composition wherein the object species comprises at least about 50 percent (on a molar basis) of all macromolecular species present. Generally, a substantially pure composition will comprise more than about 80% of all macromolecular species present in the composition, more particularly more than about 85%, more than about 90%, more than about 95%, or more than about 99%. The term “pure” or “substantially pure” also refers to preparations where the object species is at least 60% (w/w) pure, or at least 70% (w/w) pure, or at least 75% (w/w) pure, or at least 80% (w/w) pure, or at least 85% (w/w) pure, or at least 90% (w/w) pure, or at least 92% (w/w) pure, or at least 95% (w/w) pure, or at least 96% (w/w) pure, or at least 97% (w/w) pure, or at least 98% (w/w) pure, or at least 99% (w/w) pure, or 100% (w/w) pure.

[0035] Non-limiting examples of animals within the scope and meaning of the term “mammal” include dogs, cats, rats, mice, guinea pigs, chinchillas, horses, goats, cattle, sheep, zoo animals, Old and New World monkeys, non-human primates, and humans.

[0036] “Treatment” refers to therapeutic treatments. “Prevention” refers to prophylactic or preventative treatment measures or reducing the onset of a condition or disease. The term “treating” refers to administering a composition to a subject for therapeutic purposes and/or for prevention, or to a procedure conducted on or to the subject, e.g., a surgical procedure.

[0037] The terms “therapeutic composition” and “pharmaceutical composition” refer to an active agent-containing composition that may be administered to a subject by any method known in the art or otherwise contemplated herein, wherein administration of the composition brings about a therapeutic effect as described elsewhere herein. In addition, the compositions of the present disclosure may be designed to provide delayed, controlled, extended, and/or sustained release using formulation techniques which are well known in the art.

[0038] The term “effective amount” refers to an amount of an active agent which is sufficient to exhibit a detectable therapeutic or treatment effect in a subject without excessive adverse side effects (such as substantial toxicity, irritation and allergic response) commensurate with a reasonable benefit/risk ratio when used in the manner of the present disclosure. The effective amount for a subject will depend upon the subject’s type, size and health, the nature and severity of the condition to be treated, the method of administration, the duration of treatment, the nature of concurrent therapy (if any), the specific formulations employed, and the like. Thus, it is not possible to specify an exact effective amount in advance. However, the effective amount for a given situation can be determined by one of ordinary skill in the art using routine experimentation based on the information provided herein.

[0039] The term “ameliorate” means a detectable or measurable improvement in a subject’s condition, disease or symptom thereof. A detectable or measurable improvement includes a subjective or objective decrease, reduction, inhibition, suppression, limit or control in the occurrence, fre-

quency, severity, progression, or duration of the condition or disease, or an improvement in a symptom or an underlying cause or a consequence of the disease, or a reversal of the disease. A successful treatment outcome can lead to a "therapeutic effect," or "benefit" of ameliorating, decreasing, reducing, inhibiting, suppressing, limiting, controlling or preventing the occurrence, frequency, severity, progression, or duration of a disease or condition, or consequences of the disease or condition in a subject.

[0040] A decrease or reduction in worsening, such as stabilizing the condition or disease, is also a successful treatment outcome. A therapeutic benefit therefore need not be complete ablation or reversal of the disease or condition, or any one, most or all adverse symptoms, complications, consequences or underlying causes associated with the disease or condition. Thus, a satisfactory endpoint may be achieved when there is an incremental improvement such as a partial decrease, reduction, inhibition, suppression, limit, control or prevention in the occurrence, frequency, severity, progression, or duration, or inhibition or reversal of the condition or disease (e.g., stabilizing), over a short or long duration of time (hours, days, weeks, months, etc.). Effectiveness of a method or use, such as a treatment that provides a potential therapeutic benefit or improvement of a condition or disease, can be ascertained by various methods and testing assays.

[0041] Where used herein, the pronoun "we" is intended to refer to all persons involved in a particular aspect of the investigation disclosed herein and as such may include non-inventor laboratory assistants and non-inventor collaborators working under the supervision of the inventor(s).

[0042] In at least certain embodiments, the active agents of the present disclosure are oligonucleotides (generally single stranded), and may be referred to as antisense oligonucleotides ("ASO"). As used herein, the terms "antisense oligonucleotide" and "ASO" refer to an oligomeric nucleic acid that is capable of hybridizing with its complementary target sequence, generally resulting in the modulation of the normal function of the nucleic acid (e.g., mRNA or gene) having the target sequence. "Antisense" further refers to an oligomer having a sequence of nucleotide bases and a subunit-to-subunit backbone that allows the antisense oligomer to hybridize to a target sequence in an RNA by Watson-Crick base pairing, to form an RNA:oligomer heteroduplex within the target sequence, e.g., with an mRNA. The antisense oligomer (oligonucleotide) may have exact sequence complementarity to the target sequence or near complementarity, and may include modified (non-natural) nucleobases in place of naturally complementary nucleobases. Examples of such modified nucleobases are shown below.

[0043] As used herein, the term "shRNA" or "short hairpin RNA" refers to a sequence of ribonucleotides comprising a single-stranded RNA polymer that makes a tight hairpin turn on itself to provide a "double-stranded" or duplexed region. shRNA can be used to silence gene expression via RNA interference. shRNA hairpin is cleaved into short interfering RNAs (siRNA) by the cellular machinery and then bound to the RNA-induced silencing complex (RISC). It is believed that the complex inhibits RNA as a consequence of the complexed siRNA hybridizing to and cleaving RNAs that match the siRNA that is bound thereto.

[0044] As used herein, the term "RNA interference" or "RNAi" refers to the silencing or decreasing of gene expres-

sion by siRNAs. It is the process of sequence-specific, post-transcriptional gene silencing in animals and plants, initiated by siRNA that is homologous in its duplex region to the sequence of the silenced gene. The gene may be endogenous or exogenous to the organism, present integrated into a chromosome or present in a transfection vector that is not integrated into the genome. The expression of the gene is either completely or partially inhibited. RNAi inhibits the gene by compromising the function of a target RNA, completely or partially. Both plants and animals mediate RNAi by the RNA-induced silencing complex (RISC); a sequence-specific, multicomponent nuclease that destroys messenger RNAs homologous to the silencing trigger. RISC is known to contain short RNAs (approximately 22 nucleotides) derived from the double-stranded RNA trigger, although the protein components of this activity are unknown. However, the 22-nucleotide RNA sequences are homologous to the target gene that is being suppressed. Thus, the 22-nucleotide sequences appear to serve as guide sequences to instruct a multicomponent nuclease, RISC, to destroy the specific mRNAs. Biochemical reactions that recapitulate this phenomenon generate RNA fragments of 21 to 23 nucleotides from the double-stranded RNA. These stably associate with an RNA endonuclease, and probably serve as a discriminator to select mRNAs. Once selected, mRNAs are cleaved at sites 21 to 23 nucleotides apart.

[0045] As used herein, the term "siRNA" refers to a short interfering RNA. In some embodiments, siRNAs comprise a duplex, or double-stranded region, of about 18-25 nucleotides long; often siRNAs contain from about two to four unpaired nucleotides at the 3' end of each strand. At least one strand of the duplex or double-stranded region of a siRNA is substantially homologous to or substantially complementary to a target RNA molecule. The strand complementary to a target RNA molecule is the "antisense strand"; the strand homologous to the target RNA molecule is the "sense strand", and is also complementary to the siRNA antisense strand. siRNAs may also contain additional sequences; non-limiting examples of such sequences include linking sequences, or loops, as well as stem and other folded structures. siRNAs appear to function as key intermediaries in triggering RNA interference in invertebrates and in vertebrates, and in triggering sequence-specific RNA degradation during posttranscriptional gene silencing in plants.

[0046] In at least certain embodiments, the active agents (e.g., oligonucleotides) may have strand lengths comprising, for example, approximately 12 to 50, or 18 to 40, or 20 to 30 nucleotides, including a targeting sequence (i.e., a seed sequence) that is complementary to a target sequence of a nucleic acid which comprises a portion of an AR coregulator, such as an AR coregulator as listed elsewhere herein, a pre-mRNA transcribed from an AR coregulator, and/or (2) a mature mRNA processed from said pre-mRNA. For example, when an oligonucleotide binds to the target sequence of a preprocessed mRNA, it effectively inhibits splicing at the normal splice acceptor site and thus produces a splice variant mRNA, leading to truncated or otherwise aberrant versions of the encoded protein upon translation, or when the oligonucleotide binds to the target region of a mature mRNA, it effectively inhibits proper translation of the mRNA into an encoded protein.

[0047] The term "nucleic acid" is well known in the art. A "nucleic acid" as used herein will generally refer to a molecule (i.e., a strand) of DNA, RNA or a derivative or

analog thereof, comprising a nucleobase. A nucleobase includes, for example, a naturally-occurring purine or pyrimidine base found in DNA (e.g., an adenine “A,” a guanine “G,” a thymine “T” or a cytosine “C”) or RNA (e.g., an “A,” a “G,” a uracil “U” or a “C”). The term nucleobase also includes non-natural bases as described below. The term “nucleic acid” encompasses the terms “oligonucleotide” and “polynucleotide,” each as a subgenus of the term “nucleic acid.” The term “oligonucleotide” generally refers to a molecule of between about 3 and about 100 nucleobases in length. The term “polynucleotide” generally refers to at least one molecule of greater than about 100 nucleobases in length. These definitions generally refer to a single-stranded molecule, but in specific embodiments will also encompass an additional strand that is partially, substantially or fully complementary to the single-stranded molecule. Thus, a nucleic acid may encompass a double-stranded molecule that comprises a complementary strand or “complement” of a particular sequence comprising a molecule. As used herein, a single-stranded nucleic acid may be denoted by the prefix “ss,” and a double-stranded nucleic acid by the prefix “ds. The terms “polynucleotide sequence” or “nucleic acid,” as used herein, include any polynucleotide sequence which encodes a peptide or fusion protein (or polypeptide) including polynucleotides in the form of RNA, such as mRNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The RNA or DNA may be double-stranded or single-stranded. Single-stranded DNA may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

[0048] Where a strand is designated herein as RNA, and thus comprises uracil (U) nucleobases, the present disclosure is also directed to an equivalent DNA sequence where the U nucleobase is replaced with a thymine (T) nucleobase. For example, where an RNA active agent described herein comprises the seed sequence GUCUGA, the equivalent DNA active agent comprises the seed sequence GTCTGA.

[0049] As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

[0050] Therefore, in the context of the present disclosure, the term “oligonucleotide” refers to an oligomer or polymer of RNA or DNA or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-oc-

curing nucleobases, sugars and synthetic heterocycles and covalent internucleoside (backbone) linkages which function similarly. Such modified or substituted non-natural oligonucleotides, as compared to native (natural) forms may have desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

[0051] Where used herein, the term “oligonucleotide,” is also intended to include linked nucleobase sequences containing modified backbones comprising non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. Further, for the purposes of this specification, the term “nucleoside” is intended to refer to a nucleobase linked to a ribose or deoxyribose sugar (a natural nucleoside), and to a nucleobase linked to a non-ribose or non-deoxyribose heterocycle, e.g., a morpholine structure (a non-natural, or modified, nucleoside or other structures described elsewhere herein). Thus, a series of such modified, non-natural, nucleosides linked together via an internucleoside backbone can also be considered to be an oligonucleotide (a non-natural, or modified, oligonucleotide). Further, the term “sugar” where used herein in the context of a nucleoside, is intended to include “non-sugar” heterocyclic compounds, such as morpholines, as the portion of the internucleoside backbone which is linked to the nucleobase.

[0052] Oligonucleotides useful in the compounds and methods disclosed herein also include those comprising entirely or partially of naturally occurring nucleobases. Naturally occurring nucleobases as defined herein, include adenine, guanine, thymine, cytosine, and uracil. Although 5-methylcytosine (5-me-C) is technically a naturally occurring nucleobase, for the purposes of the present disclosure it will be included in the list of non-natural (a.k.a., modified) nucleobases.

[0053] As noted above, oligonucleotides of the present disclosure may further include those comprised entirely or partially of modified nucleobases and their corresponding nucleosides. Specific modified nucleobases include, but are not limited to, 5-uracil (pseudouridine), dihydrouracil, 2-thiouracil, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, dihydrouracil, 5-trifluoromethyl uracil, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, 5-carboxymethylaminomethyl-2-thiouracil, 5-(carboxyhydroxymethyl) uracil, 5-methoxycarboxymethyluracil, 5-methoxyuracil, 5-methyl-2-thiouracil, 5-propynyl uracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, 5-carboxymethylaminomethyluracil, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, alkynyl uracil, 4-thiouracil, 6-azo uracil, 2-aminoadenine, 2-methyladenine, 6-methyladenine, 7-methyladenine, 2-F-adenine, 2-amino-adenine, 2-propyladenine, N6-adenine, N6-isopentenyladenine, 2-methylthio-N6-isopentenyladenine, alkynyl adenine, 8-fluoro-adenine, 8-chloro-adenine, 8-bromo-adenine, 8-ido-adenine, 8-amino adenine, 8-thiol adenine, 8-thioalkyl adenine, 8-hydroxyl adenine, 3-deazaadenine, 7-deazaadenine, 8-azaadenine, 3-methylcytosine, 5-methylcytosine, 5-trifluoromethyl cytosine, 2-thiocytosine, 5-hydroxymethyl cytosine, 5-propynyl cytosine, 6-azo cytosine, 4-acetylcytosine, 2-methylguanine, 6-methylguanine, 2-propylguanine, 1-methylguanine, 7-methylguanine, 2,2-dim-

ethylguanine, 8-azaguanine, 7-deazaguanine, 8-fluoro-guanine, 8-chloro-guanine, 8-bromo-guanine, 8-iodo-guanine, 8-amino guanine, 8-thiol guanine, 8-thioalkyl guanine, 8-hydroxyl guanine, 3-deazaguanine, 2-thiothymine, 6-azo thymine, alkynyl thymine, ribothymine, inosine, hypoxanthine, xanthine, beta-D-galactosylqueosine, beta-D-manno-sylqueosine, 1-methylinosine, 2,6-diaminopurine, queosine, tricyclic pyrimidines, phenoxazine cytidine(1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), and phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one).

[0054] The present disclosure also encompasses oligonucleotides which comprise targeting sequences (base sequences) that are complementary to particular nucleic acid target sequences taught herein. A nucleic acid is a “complement” or is “complementary” to another nucleic acid when it is capable of base-pairing with the other nucleic acid according to the standard Watson-Crick, Hoogsteen or reverse Hoogsteen binding complementarity rules. Polynucleotides (nucleic acids) are described as “complementary” to one another when hybridization occurs in an anti-parallel configuration between two single-stranded polynucleotides.

[0055] More particularly, “complementary,” as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, “specifically hybridizable” and “complementary” are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target, and as such, as is understood in the art, the targeting sequence of an antisense oligonucleotide of the present disclosure need not be 100% complementary to that of its target sequence to be specifically hybridizable. An oligonucleotide is specifically hybridizable when binding of the oligonucleotide to the target sequence of the DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the oligonucleotide to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed. An oligonucleotide and a target sequence are thus complementary to each other when a sufficient number of nucleobases of the oligonucleotide can hydrogen bond with the corresponding nucleobases of the target sequence, such that a desired effect will occur (e.g., antisense inhibition of a target nucleic acid, such as an AR coregulator).

[0056] For example, an oligonucleotide in which 18 of 20 nucleobases of the oligonucleotide are complementary to a target sequence, and would therefore specifically hybridize, would represent 90 percent complementarity. In this example, the remaining noncomplementary nucleobases may be clustered or interspersed with complementary nucleobases and need not be contiguous to each other or to

complementary nucleobases. As such, an oligonucleotide which is 18 nucleobases in length having three noncomplementary nucleobases which are flanked by two regions of complete complementarity with the target nucleic acid, or are distributed in non-contiguous positions, would have 83% overall complementarity with the target sequence.

[0057] In other embodiments, the seed sequence of the antisense oligonucleotide provided herein is fully complementary (i.e. 100% complementary) to a target sequence of a nucleic acid, e.g., of an AR coregulator. As used herein, “fully complementary” means each nucleobase of the referenced portion of an oligonucleotide (e.g., the seed sequence) is capable of precise base pairing with the corresponding nucleobases of a target nucleic acid.

[0058] The term “target sequence” where used herein refers to a contiguous series of nucleobases in a specific nucleotide sequence (target region), for example of an mRNA. The term “target sequence” refers to a sequence that is a subsequence (portion or segment) of the target region, or to the entire sequence of the target region. A target sequence may include the 5' terminal nucleobase of a nucleic acid sequence plus adjacent internal nucleobases of the sequence, or the 3' terminal nucleobase plus adjacent internal nucleobases of the sequence, or only internal nucleobases within the sequence, or the target sequence may be 100% identical to the target region. In certain embodiments, a nucleic acid compound of the present disclosure comprises an oligonucleotide having a nucleobase sequence that, when written in the 5' to 3' direction, comprises the reverse complement of a target sequence of a nucleic acid target region to which it is targeted.

[0059] The terms “complementary” and “antisense” can be used interchangeably. Complementary also refers to polynucleotide sequences that are substantially complementary (antisense) over their entire length and have very few base mismatches. For example, sequences of fifteen bases in length may be termed complementary when they have complementary nucleotides at thirteen or fourteen positions. Naturally, sequences which are completely complementary will be sequences which are entirely complementary throughout their entire length and have no base mismatches.

[0060] In certain embodiments, oligonucleotides of the present disclosure are synthesized using one or more modified nucleotides. As used herein, the terms “modified” and “modification” when used in the context of the constituents of a nucleotide monomer, i.e., sugar, nucleobase and internucleoside linkage (backbone), refer to non-natural changes to the chemical structure of these naturally occurring constituents or the substitutions of these constituents with non-naturally occurring ones, i.e., mimetics. For example, the “unmodified” or “naturally occurring” sugar ribose (of RNA) can be modified by replacing the hydrogen at the 2'-position of ribose with a methyl group. Similarly, the naturally occurring internucleoside linkage of nucleic acids is a 3' to 5' phosphodiester linkage that can be modified, in one embodiment, by replacing one of the non-bridging oxygen atoms of the phosphate linker with a sulfur atom to create a phosphorothioate linkage. Modified oligonucleotides are structurally distinguishable, but functionally interchangeable with naturally occurring or synthetic unmodified oligonucleotides and usually have enhanced properties such as increased resistance to degradation by exonucleases and endonucleases, or increased binding affinity.

**[0061]** As noted above, in certain embodiments, modifications to the oligonucleotides of the present disclosure encompass substitutions or changes in internucleoside linkages, sugar moieties, or nucleobases. Where used herein in reference to an oligonucleotide, the term “non-natural” or “unnatural” refers to an oligonucleotide which comprises at least one modification in an internucleoside linkage, a sugar, and/or a nucleobase thereof, wherein such modified internucleoside linkage, modified sugar, and/or modified nucleobase is not found naturally in DNA or RNA (unless specifically defined otherwise herein).

**[0062]** Non-naturally occurring internucleoside linkages of the oligonucleotides of the present disclosure include those that contain a phosphorus atom and also those that do not contain a phosphorus atom. Numerous phosphorus-containing modified oligonucleotide backbones are known in the art and may be used in the oligonucleotides of the present disclosure. Examples of phosphorus-containing internucleoside linkages of non-natural (modified) oligonucleotide backbones which may occur in the presently disclosed oligonucleotides include, but are not limited to, phosphorothioate, phosphorodithioate, phosphoramidite, phosphorodiamidate, morpholino, phosphotriester, aminoalkylphosphotriester, phosphonate, chiral phosphorothioates, methyl and other alkyl phosphonates including 3'-alkylene phosphonate, 5'-alkylene phosphonate and chiral phosphonate, phosphinate, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidate, thionophosphoramidate, thionoalkylphosphonate, thionoalkylphosphotriester, selenophosphates and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', 5' to 5' or 2' to 2' linkage, and oligonucleotides having inverted polarity comprise a single 3' to 3' linkage at the 3'-most internucleotide linkage i.e., a single inverted nucleoside residue which may be abasic (the nucleobase is missing or has a hydroxyl group in place thereof) linkages. Examples of U.S. patents that teach the preparation of such phosphorus-containing linkages include, but are not limited to, U.S. Pat. Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,194,599; 5,565,555; 5,527,899; 5,721,218; 5,672,697 and 5,625,050.

**[0063]** As noted above, in some embodiments, the internucleoside linkages are without phosphorus atoms and may instead comprise short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. In further embodiments, the non-naturally occurring internucleoside linkages are uncharged and in others, the linkages are achiral. In some embodiments, the non-naturally occurring internucleoside linkages are uncharged and achiral, such as peptide nucleic acids (PNAs).

**[0064]** It is understood that the sequence set forth in each sequence or SEQ ID NO contained herein is independent of any modification to sugar moieties, internucleoside linkages, or nucleobases of the sequence, unless otherwise specified. As such, antisense oligonucleotides of the present disclosure may be defined by a complementary correspondence to a sequence or SEQ ID NO disclosed herein, or segment

thereof, and may comprise, independently, one or more modifications to a sugar moiety, an internucleoside linkage, or a nucleobase. Other embodiments of oligonucleotide backbones include siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; riboacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH<sub>2</sub> component parts. Examples of U.S. patents that teach the preparation of such non-phosphorus containing oligonucleotides include, but are not limited to, U.S. Pat. Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; 5,792,608; 5,646,269 and 5,677,439.

**[0065]** In certain oligonucleotide mimetics of the present disclosure, both the sugar moiety and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with non-natural groups. One such oligomeric compound is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Examples of U.S. patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262.

**[0066]** The oligonucleotides described herein stabilized against nucleolytic degradation such as by the incorporation of a modification, e.g., a nucleotide modification. The oligonucleotides can include a non-natural nucleoside linkage such as a phosphorothioate linkage as the first, second, and/or third internucleotide linkage at the 5' or 3' end of the oligonucleotide sequence. In certain embodiments, the oligonucleotides can include a 2'-modified nucleotide, e.g., a 2'-deoxy, 2'-fluoro, 2'-deoxy-2'-fluoro, 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), 2'-O-guanidinopropyl-, or 2'-O—N-methylacetamido (2'-O-NMA) nucleotide. In a particular embodiment, the oligonucleotides include at least one 2'-O-methyl-modified nucleotide, and in some embodiments, all of the nucleotides include a 2'-O-methyl modification.

**[0067]** As noted elsewhere herein, the oligonucleotide can be further modified so as to be conjugated to an organic moiety such as a biogenic molecule that is selected to improve stability, distribution and/or cellular uptake of the oligonucleotide, e.g., cholesterol, forming the nucleic acid compound of the present disclosure. Such an organic moiety can be attached, e.g., to the 3' or 5' end of the oligonucleotide, and/or at the 2' position of the sugar moiety of a nucleotide of the oligonucleotide, such as the 2' ribose position.

**[0068]** The nucleic acid compound can further be in isolated form or can be part of a pharmaceutical composition, such as a pharmaceutical composition formulated for parental administration. The pharmaceutical compositions can contain one or more nucleic acid compounds, and in

some embodiments will contain two or more inhibitory nucleic acid compounds, each one directed to a different target gene.

[0069] The oligonucleotides can be delivered in any of a variety of forms, including in liposomes as described above, and via expression vectors. The oligonucleotide can be endogenously expressed from transcription units inserted into DNA or RNA vectors. The recombinant vectors can be DNA plasmids or viral vectors for example. Viral vectors suitable for producing the presently disclosed oligonucleotides capable of reducing expression or activity of an AR coregulator can be constructed based on, but not limited to, adeno-associated virus, retrovirus, lentivirus, adenovirus, or alphavirus. The recombinant vectors which contain a nucleic acid for expressing the oligonucleotides disclosed herein can be delivered as described above and can persist in target cells. Alternatively, viral vectors can be used that provide for transient expression of the oligonucleotides. Such vectors can be repeatedly administered as necessary. Once expressed, in one embodiment, the oligonucleotides may interact with the target RNA and inhibit mRNA activity for example. The delivery vehicles (vectors) for the oligonucleotides optionally comprise an expression construct which includes an enhancer sequence, a promoter sequence, and other sequences necessary for expression of the products of the oligonucleotide sequence desired to be produced. In one embodiment, the promoter is cell-specific. The term "cell-specific" means that the particular promoter selected for the recombinant vector can direct expression of the selected transgene only in a particular cell type. As one example, the promoter is specific for expression in prostate cells. A number of viruses can be used in connection with the methods described herein, including papovaviruses, e.g., SV40, adenovirus, vaccinia virus, adeno-associated virus, herpesviruses including HSV and EBV, and retroviruses of avian, murine, and human origin. In certain embodiments, lentiviral vectors can be used in connection with the methods described herein. In certain embodiments, the lentiviral vector can be a doxycycline-inducible lentiviral vector engineered to express one or more shRNAs or siRNAs.

[0070] Specific vectors which may be used include, but are not limited to, adeno-associated virus vectors (e.g., as disclosed in U.S. Pat. Nos. 5,139,941, 5,436,146, and 5,622, 856), an attenuated or gutless adenoviral vectors, (e.g., as disclosed in U.S. Pat. No. 5,935,935), lentiviral vectors (such as are disclosed in U.S. Pat. Nos. 5,665,577; 5,994, 136; and 6,013,516), plasmids or synthetic (non-viral) vectors (such as disclosed in U.S. Pat. Nos. 4,394,448 and 5,676,954), and/or nanoparticles (such as disclosed, for example, in U.S. Pat. Nos. 6,217,912; 7,514,098; and 8,323, 618), retroviral vectors (such as are disclosed in U.S. Pat. Nos. 5,672,510; 5,707,865; and 5,817,491), herpes virus vectors (such as are disclosed in U.S. Pat. No. 5,288,641), and sindbis virus vectors and papilloma virus vectors (such as are disclosed in EP 820 773). The vectors may be either monocistronic, bicistronic, or multicistronic. A recombinant vector (e.g., lenti-, parvo-, AAV) sequence can be packaged as a "particle" for subsequent infection (transduction) of a cell, ex vivo, in vitro or in vivo. Where a recombinant vector sequence is encapsulated or packaged into an AAV particle, the particle can also be referred to as a "rAAV." Such particles include proteins that encapsulate or package the vector genome. Particular examples include viral envelope proteins, and in the case of AAV, capsid proteins.

[0071] Thus, the oligonucleotides of the present disclosure may be used as a form of gene therapy. The term "gene therapy" as used herein means genetic modification of cells by the introduction of exogenous DNA or RNA into these cells, such as via an expression vector containing the oligonucleotide, for the purpose of expressing or replicating one or more peptides, polypeptides, proteins, oligonucleotides, or polynucleotides in vivo for the treatment or prevention of disease or deficiencies in humans or animals. Examples of gene therapy are disclosed for example in U.S. Pat. No. 5,399,346. Any suitable route of administration of the oligonucleotide-containing vector may be employed. For example, parenteral (subcutaneous, subretinal, suprachoroidal, intramuscular, intravenous, transdermal) and like forms of administration may be employed. Dosage formulations include injections, implants, or other known and effective gene therapy delivery methods.

[0072] Delivery of the oligonucleotide-expressing vectors can be systemic, such as by intravenous or intra-muscular administration, direct administration to a tumor site, such as a prostate tumor, by administration to target cells ex-planted from a subject followed by reintroduction into the subject, or by any other means that would allow for introduction into the desired target cell. The therapeutic and/or pharmaceutical compositions, in non-limiting embodiments, contain viral particles per dose in a range of, for example, from about  $10^4$  to about  $10^{11}$  particles, from about  $10^5$  to about  $10^{10}$  particles, or from about  $10^6$  to about  $10^9$  particles. In the context of AAV vectors, vector genomes are provided in a range of, for example, from about  $10^4$  to about  $10^{14}$  vector genomes, from about  $10^5$  to about  $10^{13}$  vector genomes, from about  $10^6$  to about  $10^{13}$  vector genomes, from about  $10^7$  to about  $10^{13}$  vector genomes, from about  $10^8$  to about  $10^{13}$  vector genomes, or from about  $10^9$  to about  $10^{13}$  vector genomes. Such doses/quantities of AAV vector are useful in the methods set forth herein.

[0073] Because nucleases that cleave the phosphodiester linkages are expressed in almost every cell, unmodified nucleic acid molecules such as the inhibitory oligonucleotides of the present disclosure may be modified to resist degradation, as described above for example. Other biogenic molecules may be conjugated to the oligonucleotides to improve their ability to resist degradation, target certain cells, or to cross barriers like cell membranes or the blood brain barrier. Examples of biogenic molecules that can be conjugated to the oligonucleotides include lipids such as, but not limited to, stearic acid, palmitic acid, docosanoic acid, docosahexanoic acid, docosahexaenoic acid, cholesterol, tocopherol, and other C12-C22 saturated or unsaturated fatty acids; peptides such as but not limited to, cell-penetrating peptides (CPPs) such as penetratin, HIV-1 Tat peptides, pVEC-Cadherin 615-634, polyarginines (6-12), and transportan, linear and cyclic RGD-containing peptides, and SPACE peptide; receptor-specific ligands; aptamers (synthetic oligoribonucleotides); antibodies or antibody fragments; CpG-containing oligonucleotides; polyamines, such as spermine and spermidine; polymers such as dendrimers and polyethylene glycols (e.g., PEG 0.6 kDa-5,000 kDa); and saccharides such as N-acetylgalactosamine (GalNAc) and cyclodextrins. The biogenic molecule may be conjugated to the oligonucleotide by any suitable means, such as via linker or a cleavable bond such as but not limited to disulfide, thioether, pH sensitive (e.g., hydrazone or carboxymethylmaleic anhydride), or ethylene glycol.

**[0074]** The oligonucleotides or nucleic acid compounds of the present disclosure may be delivered in the form of nanoparticles and microparticles which encapsulate the nucleic acid compounds within liposomes of cationic lipids or within PEG, for example. These delivery systems can enhance intracellular delivery either by protecting the nucleic acid compound from nuclease degradation and/or by promoting absorptive endocytosis. Further, the addition of dioleylphosphatidylethanolamine to liposome delivery systems results in the destabilization of endosomal membranes and promotion of release of the oligonucleotide after endocytosis. The nucleic acid compounds can be administered to cells by a variety of other methods known to those of skill in the art, including, but not limited to, ionophoresis, or by incorporation into other vehicles, such as hydrogels, cyclodextrins, biodegradable nanocapsules, and bioadhesive microspheres, or by proteinaceous vectors. In one example, the nucleic acid compounds can be delivered via the nanoparticle system shown in U.S. Patent Application Publication 2019/0255088. The liposomes may comprise amphiphatic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Pat. Nos. 4,235,871; 4,501,728; 4,837,028; and 4,737,323.

**[0075]** In certain embodiments, the nanoparticles which contain the nucleic acid compounds of the present disclosure may comprise a pharmaceutically acceptable carrier such as, but not limited to, poly(ethylene-co-vinyl acetate), PVA, partially hydrolyzed poly(ethylene-co-vinyl acetate), poly(ethylene-co-vinyl acetate-co-vinyl alcohol), a cross-linked poly(ethylene-co-vinyl acetate), a cross-linked partially hydrolyzed poly(ethylene-co-vinyl acetate), a cross-linked poly(ethylene-co-vinyl acetate-co-vinyl alcohol), poly-D,L-lactic acid, poly-L-lactic acid, polyglycolic acid, PGA, copolymers of lactic acid and glycolic acid, polycaprolactone, polyvalerolactone, poly(anhydrides), copolymers of polycaprolactone with polyethylene glycol, copolymers of polylactic acid with polyethylene glycol, polyethylene glycol; fibrin, Gelfoam<sup>TM</sup> (which is a water-insoluble, off-white, nonelastic, porous, pliable gel foam prepared from purified gelatin and water for injection), and combinations and blends thereof. Copolymers can comprise from about 1% to about 99% by weight of a first monomer unit such as ethylene oxide and from 99% to about 1% by weight of a second monomer unit such as propylene oxide. Blends of a first polymer such as gelatin and a second polymer such as poly-L-lactic acid or polyglycolic acid can comprise from about 1% to about 99% by weight of the first polymer and from about 99% to about 1% of the second polymer.

**[0076]** The oligonucleotides or nucleic acid compounds can be delivered directly by systemic administration such as using oral formulations or stereotactic injection into prostate or prostate tumor, typically in saline with chemical modifications to enable uptake, or other methods described elsewhere herein. In certain embodiments, such as when the oligonucleotide of the nucleic acid compound has a phosphorothioate backbone, the oligonucleotide binds to serum proteins, slowing excretion by the kidney. The aromatic nucleobases also interact with other hydrophobic molecules

in serum and on cell surfaces. In certain embodiments, siRNA delivery systems involve complexing the RNA with cationic and neutral lipids, although encouraging results have also been obtained using peptide transduction domains and cationic polymers. Including PEGylated lipids in the formulation prolongs the circulating half-life of the particles.

**[0077]** As noted, one type of optimization of single-stranded DNA or RNA oligonucleotides is the use of chemical modifications to increase the nuclease resistance such as the introduction of phosphorothioate ("PS") linkages in place of the phosphodiester bond. This modification improves protection from digestion by nucleases. PS linkages also improved binding to serum proteins *in vivo*, increasing half-life and permitting greater delivery of active compound to tissues. Chemical modifications to subunits of the nucleotides can also improve potency and selectivity by increasing binding affinity of oligonucleotides for their complementary sequences. Examples of such modifications to the nucleoside sugars include 2'-O-methyl (2'-O-Me), 2'-fluoro (2'-F), and 2'-O-methoxyethyl (2'-MOE) RNA, and others as discussed elsewhere herein. Even more affinity can be gained using oligonucleotides modified with locked nucleic acid (LNA), which contains a methylene bridge between the 2' and 4' position of the ribose. This bridge "locks" the ribose ring in a conformation that is ideal for binding, leading to high affinity for complementary sequences. Related bridged nucleic acid (BNA) compounds have been developed and share these favorable properties. Their high affinity has permitted the development of far shorter oligonucleotides than previously thought possible which nonetheless retain high potency. The chemistry for introducing 2'-O-Me, 2'-MOE, 2'-F, or LNA into oligonucleotides is compatible with DNA or RNA synthesis, allowing chimeras with DNA or RNA bases to be easily obtained. This compatibility allows the properties of chemically modified oligonucleotides to be fine-tuned for specific applications, which is a major advantage for development that makes LNAs and other BNAs convenient tools for many applications.

**[0078]** Therapeutic administration of the active agents described herein include any method by which a nucleic acid (e.g., DNA or RNA), as known to one of ordinary skill in the art. For treatment of aggressive prostate cancer, delivery may be via, for example, oral administration and/or injection into the prostate gland or tumor or both.

**[0079]** In certain embodiments, the active agents can be delivered to an organelle, a cell, a tissue, a tumor or an organism via one or more injections (i.e., a needle injection), such as, for example, orally, subcutaneously, intradermally, intramuscularly, intravenously, or intraperitoneally.

**[0080]** A described inhibitory nucleic acid or other active agent can be incorporated into pharmaceutical compositions suitable for administration. For example, pharmaceutical compositions can comprise one or more the active agents and a pharmaceutically acceptable carrier.

**[0081]** Then active agent may be provided in a sustained release composition. The use of immediate or sustained release compositions depends on the nature of the condition being treated. If the condition consists of an acute or over-acute disorder, treatment with an immediate release form can be conducted over a prolonged release composi-

tion. Alternatively, for certain preventative or long-term treatments, a sustained release composition may be appropriate.

[0082] The active agent can be administered in a single dose or in multiple doses. Where the administration of the active agent is by infusion, the infusion can be a single sustained dose or can be delivered by multiple infusions. Injection of the active agent can be directly into the tissue at or near the site of aberrant or unwanted target gene expression. Multiple injections of the active agent can be made into the tissue, for example, into the prostate gland, into the prostate tumor, or near the tumor.

[0083] In addition to treating pre-existing aggressive or non-aggressive prostate cancers, active agents of the disclosure can be administered prophylactically in order to prevent or slow the conversion of a non-aggressive prostate cancer to an aggressive form. The active agent can be employed in combination therapies, meaning that the present compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutic agents or medical procedures. The combination of therapies (therapeutic agents or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutic agents and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies employed can achieve a desired effect for the same disorder (for example, a compound described herein can be administered concurrently with another therapeutic agent used to treat the same disorder), or they can achieve different effects (e.g., control of any adverse effects).

[0084] Known agents useful for treating cancers can be combined with the presently described active agents to treat a prostate cancer. For example, the active can be administered with another therapeutic agent simultaneously or sequentially in separate unit dosage forms or together in a single unit dosage form. The active agents are useful for individuals who have received prior medication for prostate cancer, as well as individuals who have received no prior medication for a cancer. Individuals of any age can be treated by the methods compositions of the disclosure.

[0085] It is understood that the appropriate dose of an active agent depends upon a number of factors within the knowledge of the ordinarily skilled physician, veterinarian, or researcher. The dose(s) vary, for example, depending upon the identity, size, and condition of the subject or sample being treated, further depending upon the route by which the active agents and pharmaceutical compositions are to be administered, and the effect which the practitioner desires the an active agent to have. It is furthermore understood that appropriate doses of an active agent depend upon the potency with respect to the expression or activity to be modulated. Such appropriate doses may be determined using the assays described herein. When one or more of these active agents are to be administered to an animal (e.g., a human), a relatively low dose may be prescribed at first, with the dose subsequently increased until an appropriate response is obtained. In addition, it is understood that the specific dose level for any particular subject will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, gender, and diet of the subject, the time of administration, the route of administration, the rate of excretion, any drug combination, and the degree of expression or activity to be modulated.

[0086] In certain embodiments, an effective amount of the nucleic acid compound of the present disclosure will generally contain sufficient active agent to deliver from about 0.1 µg/kg to about 100 mg/kg (weight of active agent/body weight of patient). Particularly, the composition will deliver at least 0.5 µg/kg to 50 mg/kg, and more particularly at least 1 µg/kg to 25 mg/kg. Without wishing to be held to a specific dosage, it is contemplated that the various pharmaceutical compositions used to practice the method of the present disclosure may contain, but are not limited to, about 0.01 mg to about 10 mg of the peptide per kg body weight per dose.

[0087] Practice of the method of the present disclosure may include administering to a subject an effective amount of the nucleic acid compound in any suitable systemic or local formulation, in an amount effective to deliver the dosages listed above, including but not limited to an implantable material. In one embodiment, an effective, particular therapeutic dosage is 1 µg/kg to 10 mg/kg. The dosage can be administered on a one-time basis, or (for example) from one to five times per day or once or twice per week, or continuously via a venous drip, depending on the desired therapeutic effect. In one therapeutic method of the present disclosure, the composition is provided in an IV infusion in the range of from 1 mg/kg-10 mg/kg of body weight once a day. The duration of an intravenous therapy using the pharmaceutical composition of the present disclosure will vary, depending on the condition being treated and the condition and potential idiosyncratic response of each individual patient. In at least one embodiment, it is contemplated that the duration of each application of the nucleic acid compound may be in the range of 1 to 4 hours and given once every 12 or 24 hours by continuous intravenous administration. Other therapeutic drugs, intravenous fluids, cardiovascular and respiratory support could also be provided if requested by the attending physician in a manner known to one of ordinary skill in the art.

[0088] As used herein, the terms "complementary" or "complement" also refer to a nucleic acid comprising a sequence of consecutive nucleobases or semiconsecutive nucleobases (e.g., one or more nucleobase moieties are not present in the molecule) capable of hybridizing to another nucleic acid strand or duplex even if less than all the nucleobases do not base pair with a counterpart nucleobase. In certain embodiments, a "complementary" nucleic acid comprises a sequence in which about 70%, about 71%, about 72%, about 73%, about 74%, about 75%, about 76%, about 77%, about 77%, about 78%, about 79%, about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100%, and any range derivable therein, of the nucleobase sequence is capable of base-pairing with a single or double stranded nucleic acid molecule during hybridization. In certain embodiments, the term "complementary" refers to a nucleic acid that may hybridize to another nucleic acid strand or duplex in stringent conditions, as would be understood by one of ordinary skill in the art.

[0089] The term "homologous" or "% identity" as used herein means a nucleic acid (or fragment thereof) or a protein (or a fragment thereof) having a degree of homology to the corresponding natural reference nucleic acid or protein that may be in excess of 70%, or in excess of 80%, or in excess of 85%, or in excess of 90%, or in excess of 91%,

or in excess of 92%, or in excess of 93%, or in excess of 94%, or in excess of 95%, or in excess of 96%, or in excess of 97%, or in excess of 98%, or in excess of 99%. For example, in regard to peptides or polypeptides, the percentage of homology or identity as described herein is typically calculated as the percentage of amino acid residues found in the smaller of the two sequences which align with identical amino acid residues in the sequence being compared, when four gaps in a length of 100 amino acids may be introduced to assist in that alignment (as set forth by Dayhoff, in *Atlas of Protein Sequence and Structure*, Vol. 5, p. 124, National Biochemical Research Foundation, Washington, D.C. (1972)). In one embodiment, the percentage homology as described above is calculated as the percentage of the components found in the smaller of the two sequences that may also be found in the larger of the two sequences (with the introduction of gaps), with a component being defined as a sequence of four, contiguous amino acids. Also included as substantially homologous is any protein product which may be isolated by virtue of cross-reactivity with antibodies to the native protein product. Sequence identity or homology can be determined by comparing the sequences when aligned so as to maximize overlap and identity while minimizing sequence gaps. In particular, sequence identity may be determined using any of a number of mathematical algorithms. A non-limiting example of a mathematical algorithm used for comparison of two sequences is the algorithm of Karlin & Altschul, *Proc. Natl. Acad. Sci. USA* 1990, 87, 2264-2268, modified as in Karlin & Altschul, *Proc. Natl. Acad. Sci. USA* 1993, 90, 5873-5877.

[0090] In one embodiment "% identity" represents the number of amino acids or nucleotides which are identical at corresponding positions in two sequences of a protein or nucleic acids, respectively. For example, two amino acid sequences each having 100 residues will have 95% identity when 95 of the amino acids at corresponding positions are the same. Similarly, two nucleic acid sequences each having 100 bases will have 95% identity when 95 of the bases at corresponding positions are the same.

[0091] Another example of a mathematical algorithm used for comparison of sequences is the algorithm of Myers & Miller, CABIOS 1988, 4, 11-17. Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used. Yet another useful algorithm for identifying regions of local sequence similarity and alignment is the FASTA algorithm as described in Pearson & Lipman, *Proc. Natl. Acad. Sci. USA* 1988, 85, 2444-2448. Another algorithm is the WU-BLAST (Washington University BLAST) version 2.0 software (WU-BLAST version 2.0 executable programs for several UNIX platforms). This program is based on WU-BLAST version 1.4, which in turn is based on the public domain NCBI-BLAST version 1.4 (Altschul & Gish, 1996, Local alignment statistics, Doolittle ed., *Methods in Enzymology* 266, 460-480; Altschul et al., *Journal of Molecular Biology* 1990, 215, 403-410; Gish & States, *Nature Genetics*, 1993, 3: 266-272; Karlin & Altschul, 1993, *Proc. Natl. Acad. Sci. USA* 90, 5873-5877; all of which are incorporated by reference herein).

[0092] In addition to those otherwise mentioned herein, mention is made also of the programs BLAST, gapped

BLAST, BLASTN, BLASTP, and PSI-BLAST, provided by the National Center for Biotechnology Information. These programs are widely used in the art for this purpose and can align homologous regions of two amino acid sequences. In all search programs in the suite, the gapped alignment routines are integral to the database search itself. Gapping can be turned off if desired. The default penalty (Q) for a gap of length one is Q=9 for proteins and BLASTP, and Q=10 for BLASTN, but may be changed to any integer. The default per-residue penalty for extending a gap (R) is R=2 for proteins and BLASTP, and R=10 for BLASTN, but may be changed to any integer. Any combination of values for Q and R can be used in order to align sequences so as to maximize overlap and identity while minimizing sequence gaps. The default amino acid comparison matrix is BLOSUM62, but other amino acid comparison matrices such as PAM can be utilized.

[0093] Specific amino acids may be referred to herein by the following designations: alanine: ala or A; arginine: arg or R; asparagine: asn or N; aspartic acid: asp or D; cysteine: cys or C; glutamic acid: glu or E; glutamine: gln or Q; glycine: gly or G; histidine: his or H; isoleucine: ile or I; leucine: leu or L; lysine: lys or K; methionine: met or M; phenylalanine: phe or F; proline: pro or P; serine: ser or S; threonine: thr or T; tryptophan: trp or W; tyrosine: tyr or Y; and valine: val or V.

[0094] The terms "oligonucleotide", "polynucleotide sequence" or "nucleic acid," as used herein, may include polynucleotide sequences which have particular activity or encode peptides or polypeptides, e.g., proteins, and include polynucleotides in the form of RNA, such as mRNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof, or a non-coding sequence such as siRNA, shRNA, miRNA, or other such short RNA sequences. The DNA may be double-stranded or single-stranded. Single-stranded DNA may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

[0095] As noted, the term "antisense" refers to a polynucleotide or oligonucleotide molecule that is substantially complementary or 100% complementary to a particular polynucleotide or oligonucleotide molecule (RNA or DNA), i.e., a "sense" strand, or portion thereof. For example, the antisense molecule may be complementary in whole or in part to a molecule of messenger RNA, miRNA, pRNA, tRNA, rRNA or hnRNA, or a sequence of DNA that is either coding or non-coding.

[0096] The term "operably linked" where used herein refers to an association of two chemical moieties linked in such a way so that the function of one is not affected by the other, e.g., an arrangement of elements wherein the components so described are configured so as to perform their usual function. The two moieties may be linked directly, or may be linked indirectly via a linker sequence of molecule.

[0097] The term "primer" refers to an oligonucleotide sequence which serves as a starting point for DNA synthesis in the polymerase chain reaction (PCR). A primer generally comprises from about 12 to about 30 nucleotides and hybridizes with a complementary region of a target sequence, for example a microRNA molecule.

[0098] The term "probe" refers to an oligonucleotide which is bound to or configured to bind to a target sequence,

and includes for example, an antisense nucleic acid sequence which is designed to hybridize by a sequence-specific method with a complementary region of a specific nucleic acid sequence such as a target nucleic acid, such as an miRNA as disclosed herein. An oligonucleotide probe can comprise any number of nucleotides, such as 10 to 25, as long as the oligonucleotide probe comprises a sufficient number of nucleotides to bind to the target nucleic acid with the necessary specificity for the particular use of the probe. For purposes of quantification of the probe-target sequence complex, the probe may further optionally comprise a tag or label operably linked thereto, wherein the tag or label comprises, for example, a fluorescent (e.g., fluorophore), luminescent, or chemiluminescent label or reporter group. Oligonucleotides with binding specificity to the RNAs and cDNAs expressed from the biomarker panel disclosed herein may be referred to as "capture molecules" in the assay systems and methods disclosed herein.

**[0099]** The term "fluorophore" or "fluorochrome" or "fluorescent species" or "fluorescent label" or "fluorescent tag," as used herein indicates a substance which itself fluoresces or can be made to fluoresce. Each term is interchangeable. Fluorophores can be used alone or covalently attached ("operably-linked") or non-covalently linked to another molecule, such as an oligonucleotide primer, probe, or miRNA, such as described herein. The process of covalently attaching a fluorophore to another molecule or compound is referred to as "fluorescent labeling" and may be conducted by, for example, an enzyme effective in forming the covalent bond therebetween.

**[0100]** Examples of fluorophores which may be used in various embodiments of the present disclosure include but are not limited to: hydroxycoumarin, methoxycoumarin, Alexa fluor 345, aminocoumarin, 7-diethylaminocoumarin-3-carboxylic acid, Cy2 (cyanine 2), FAM, Alexa fluor 350, Alexa fluor 405, Alexa fluor 488, Fluorescein (FITC), Alexa fluor 430, Alexa fluor 532, HEX 535, Cy3, Alexa fluor 546, Alexa fluor 555, R-phycerythrin (PE), tetramethyl rhodamine (TRITC), Rhodamine Red-X, Tamara, Cy3.5, Rox, Alexa fluor 568, Red 613 480, Texas Red 615, Alexa fluor 594, Alexa fluor 633, Allophycocyanin, Alexa fluor 647, Cy5, Alexa fluor 660, Cy5.5, TruRed 490, Alexa fluor 680, Alexa fluor 750, Cy7, DAPI, QSY 7, QSY 33, dabsyl, BODIPY FL, BODIPY630/650, BODIPY 650/665, BODIPY TMR-X, BODIPY TR-X, Hoechst 33258, SYTOX blue, Hoechst 33342, YOYO-1 509, SYTOX green, TOTO1, TO-PRO-1, SYTOX orange, Chromomycin A3, Mithramycin, propidium iodide, ethidium bromide, Pacific Orange, Pacific Green, Pacific Blue, Oregon Green 488, Oregon Green 514, red fluorescent protein (RFP), green fluorescent protein (GFP), and cyan fluorescent protein (CFP).

**[0101]** Certain novel embodiments of the present disclosure, having now been generally described, will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present disclosure, and are not intended to be limiting. The following examples are to be construed, as noted above, only as illustrative, and not as limiting of the present disclosure in any way whatsoever. Those skilled in the art will promptly recognize appropriate variations from the various compositions, structures, components, procedures and methods.

## EXPERIMENTAL

### Material and Methods

#### Cell Culture and Plasmid Constructs

**[0102]** LNCaP, 22Rv1, DU145, RWPE1, PC3 and HEK 293T cell lines were obtained from American Type Culture Collection (ATCC), VCaP cells were obtained from Sigma-Aldrich, Panc-1 cells were obtained from Dr. Priyabrata Mukherjee (Oklahoma University Health Sciences Center), and LentiX-293T cells were obtained from Clonetech. LNCaP and 22Rv1 cells were maintained in RPMI Glutamax growth media (Gibco), DU145, HEK293T, VCaP, PC3 and Panc-1 and LentiX-293T cells were maintained in DMEM growth media (Gibco), and RWPE1 were maintained in KSF media (Gibco). Both RPMI and DMEM media were supplemented with 10% fetal bovine serum (FBS), 100 units/mL penicillin, 100 g/mL streptomycin, Amphotericin B and 2 mM L-Glutamine. pLKO.1 vector was used for shRNA expression, and plasmids were obtained from Open Biosystems or cloned using pLKO.1-TRC (pLKO.1-TRC cloning vector was a gift from Dr. David Root (Addgene plasmid #10878; <http://n2t.net/addgene:10878>; RRID: Addgene\_10878)) and the pLKO.1-TRC cloning vector protocol obtained from Addgene. shRNA target sequences were as follows (Note: Most shRNAs used in this work (Table 1) come from the RNAi consortium shRNA library, and TRC numbers are provided for these sequences. shTMEFF2-6 and shTMEFF2-9 sequences were generated using the siRNA Wizard online tool by Invitrogen: <https://www.invivogen.com/simawizard/>).

TABLE 1

shRNA sequences and TRC numbers		
shRNA	Sequence	TRC No.
		SEQ ID NO
shTMEFF2-1	CTGGTTATGATGACAGAGAAA	TRCN0000073522 1
shTMEFF2-2	CGTCTGTCAGTTCAAGTGCAA	TRCN0000222559 2
shTMEFF2-3	GCGCTTCTGATGGAAATCTT	TRCN0000073521 3
shTMEFF2-4	GCAGGTGTATGCTGGTTATA	TRCN0000073520 4
shTMEFF2-5	CCTTGCATTGTTGTTAATCTA	TRCN0000073518 5
shTMEFF2-6	GGCTCTGGAGAAACTAGTCAA	
shTMEFF2-7	ATGCAGAGAACATGCTAACAAAT	TRCN0000373776 7
shTMEFF2-8	CATACCTTGTCCGGAACATTA	TRCN0000373700 8
shTMEFF2-9	GGCACTACAGTTCAGACAATA	
shScramble	CCTAAGGTTAACGCCCCCTCG	
shGFP	CGACGTAAACGGGCCAACAGTT	
shL3	ACTGGGCTGTACTTTGTATAT	TRCN0000059000 12
shR6	CCTGAAACAGTGGCAATAAT	TRCN0000038696 13

**[0103]** CRISPR Cas9 (pRCCH-CMV-Cas9-2A-Hygro) and doxycycline inducible sgRNA (pRSGT16-U6Tet-(sg)-CMV-TetRep-2A-TagRFP-2A-Puro) plasmids were obtained from Cellecta. Constitutive sgRNA expressing

vector, pLX-sgRNA, was a gift from Dr. Eric Lander & Dr. David Sabatini (Addgene plasmid #50662; <http://n2t.net/addgene:50662>; RRID: Addgene\_50662), and sgRNA were cloned using an Addgene protocol. Cloned sgRNA target sequences were: sgGFP (AAGATCGAGTGCCGCATCAC) (SEQ ID NO:14) and sgTMEFF2-3 (CCAGCTGCTGCACTGCCGCG) (SEQ ID NO: 15).

[0104] RetroX-ON and RetroX-Tight-Pur vectors transduced sequentially into C4-2B cells, were used for doxycycline inducible TMEFF2 overexpression. LentiX 293T cells (Clonetech) and CalPhos transfection reagents (Clonetech) were used for viral particle packaging. psPAX2 and VSV-G plasmids were used for lentiviral packaging. psPAX2 was a gift from Dr. Didier Trono (Addgene plasmid #12260; <http://n2t.net/addgene:12260>; RRID: Addgene\_12260), and pCMV-VSV-G was a gift from Dr. Robert Weinberg (Addgene plasmid #8454; <http://n2t.net/addgene:8454>; RRID: Addgene\_8454). pCL-10A1 plasmid was used for retroviral packaging. Viral concentrations necessary for approximately 90% or greater survival (for non-toxic constructs) in selection antibiotic were used for transductions. For retroviral and lentiviral transductions, cells were seeded in 6 cm plates at 50% confluence, and viral particle containing supernatant was diluted in 1.5 ml of 8 µg/ml polybrene serum free media and added to cells. After 5 hours, 1.5 ml of 10% FBS growth media was added to transduction media. Growth media refreshed 24 hours after initial viral particle exposure. Transduced cells were stably selected for an average of 10 days using the following selection antibiotic concentrations: Puromycin 1 µg/ml, hygromycin 750 µg/ml, G418 800 µg/ml. Selection antibiotic was not used for short time course experiments (1 week or less) with shRNA constructs when the generation of stable cell lines was not possible.

#### Antisense Oligo Transfections

[0105] FANA antisense oligos (ASOs) were obtained from AUM Biotech. LNCaP cells were transfected with 250 nM ASO (Non-target or pool of 4 targeting TMEFF2) using 0.19% Dharmafect #3 transfection reagent. 48 hours after transfection, cells were treated with 10% CSS RPMI for 24 hours, followed by 24 hours in 10 nM dihydrotestosterone (DHT) or 0.0001% ethanol (EtOH).

#### RNA Extractions and Lysate Preparations

[0106] For experiments measuring gene expression response to DHT, cells were washed with PBS and treated with 10% charcoal stripped serum (CSS) containing growth media for 24 hours for hormone depletion, followed by 24 hours in 10 nM DHT (Sigma Aldrich) or 0.0001% EtOH vehicle in 10% CSS growth media, at which time samples were collected. Cells were maintained in 10% FBS growth media for all experiments not using DHT. RNeasy (Qiagen) with on-column DNase treatment (Qiagen) was used for RNA extractions. Cell Signaling lysis buffer 9803 was used for whole cell lysates, and NE-PER kit (Thermo) was used for Nuclear/Cytoplasmic protein extractions.

#### Western Blot Analysis and Antibodies

[0107] Proteins were separated via 10% SDS/PAGE using mini-PROTEAN TGX stain free gels (Biorad) and transferred onto 0.2 µm nitrocellulose membranes using the semi-dry turbo transfer system (Biorad). One-hour incubation in 5% NFDM in tris-buffered saline 0.1% tween 20

(TBST) was used for blocking, followed by overnight incubation at 4 degrees Celsius with primary antibody diluted in 5% NFDM TBST. anti-TMEFF2 (Sigma HPA015587) 1:1000, anti-PSA (abcam 76113) 1:1000, anti-FKBP5 (abcam 2901) 1:1000, anti-AR (Cell Signaling D6F11) 1:1000, anti-ERG (abcam 92513), anti-Calnexin (abcam 22595) 1:4000. Goat anti-rabbit HRP conjugated secondary antibody (ThermoFisher 31460) was diluted in 5% NFDM at a concentration of 80 ng/ml. Clarity Western ECL (Biorad) or SuperSignal West Femto (Thermo) were used for chemiluminescent detection

#### Quantitative RT PCR

[0108] RNA samples were reverse transcribed using iScript reverse transcription supermix (500 ng/rxn) (Biorad). RT qPCR reactions were carried out in 96 well plates (25 ng cDNA/well and 200 nM per primer/well) using Sso Advanced Universal SYBR Green supermix (Biorad) and the Biorad CFX96 touch RTPCR detection system. Each reaction (sample/primer set combination) was run in duplicate and triplicate to ensure accurate loading. Forward and reverse primers used are shown in para. [0104] of U.S. Provisional application Ser. No. 62/935,521. Relative gene expression was calculated via  $\Delta\Delta CT$  or pfaffl method, when appropriate, and LinRegPCR software was used to calculate primer amplification efficiency. At least 2-4 housekeeping genes were used for normalization per run, with the geometric mean of CT values being used for normalization of gene expression. Housekeeping genes used for normalization included, CANX, HPRT1, PSMA1, RPL8, RPL38, and PPP2CA. Statistics: Two-tailed T tests were used to calculate significant differential gene expression. P<0.05 was considered to be statistically significant.

#### RNA Sequencing

[0109] Deep paired-end RNA sequencing analysis targeted to the TMEFF2 locus for de novo isoform detection was carried out by Oklahoma Medical Research Foundation's Sequencing Facility. One sample of LNCaP RNA was used for analysis, and over 317 million reads were obtained after decontamination. The genomic region of focus was the TMEFF2 locus -/+1Mbp upstream and downstream (chr2: 190949046-193194933). StringTie (v.1.2.3) was used for transcript reconstruction.

[0110] For differential gene expression analyses with LNCaP shScramble/shTMEFF2-3-/+DHT, 36 hours post-transduction cells were grown in CSS RPMI for 24 hours, followed by 24 hours in 10 nM DHT or 0.0001% EtOH vehicle control. In both cases RNA samples were prepared as described in the RNA extractions and lysate preparations section of the materials and methods. Three repeats were analyzed for each RNA sequencing analysis. RNA sequencing and initial statistical analyses were carried out by Novogene. RNA integrity was analyzed by Agilent 2100 to ensure sample quality. mRNAs were isolated using polyT magnetic beads, which was followed by fragmentation. Two cDNA libraries were synthesized using random hexamer primers; one with dTTP to dUTP substitutions in the second strand to allow for strand specificity, and one library without substitution. cDNA fragments from both libraries were ligated to NEBNext Adaptor and purified using AMPure XP beads after PCR amplification. A read depth of 20 million reads was used for each sample using an Illumina Next-

Generation sequencer. For quality control purposes, error rate distributions and G/C content were analyzed in reads, and low-quality reads containing adapter sequences, >10% unknown nucleotides or low Qscore values were eliminated. STAR was used for mapping clean reads to the human transcriptome and genome, and differential gene expression was determined using the DESeq2 R package. P-values were adjusted using the Benjamini and Hochberg method. Transcripts with average log 2 fold change >0.5 (n=3) and adjusted p-value <0.05 were considered significantly differentially expressed.

#### Enrichment Analysis

**[0111]** Essential gene lists were derived from the following publications: Wang et al., 2015 (Wang T, Birsoy K, Hughes N W, Krupczak K M, Post Y, Wei J J, Lander E S, Sabatini D M. *Identification and characterization of essential genes in the human genome*. Science. 2015 Nov. 27; 350(6264):1096-101) and Fei et al., 2017 (Teng Fei, Yiwen Chen, Tengfei Xiao, Wei Li, Laura Cato, Peng Zhang, Maura B. Cotter, Michaela Bowden, Rosina T. Lis, Shuang G. Zhao, Qiu Wu, Felix Y. Feng, Massimo Loda, Housheng Hansen He, X. Shirley Liu, Myles Brown. *Genome-wide CRISPR screen identifies HNRNPL as a prostate cancer dependency regulating RNA splicing*. Proceedings of the National Academy of Sciences Jun 2017, 114 (26) E5207-E5215). For the essential gene list derived from Wang et al., 2015, genes with under-represented reads were identified from the sequencing data from the following cell lines: KBM7, K562, Raji, Jiyoye. Only genes that were essential in two or more of the four cell lines were considered essential in our study (1739 total genes, with 1705 genes having consistent RNA seq reads from our RNA seq analysis with LNCaP shScramble/shTMEFF2-3-/+DHT samples). For the Fei et al., 2017 list, all of the top 999 essential LNCaP genes were used for our study. In addition, all 274 AR co-regulators from DePriest, et al., 2016 ((DePriest, A. D., Fiandalo, M. V., Schlanger, S., Heemers, F., Mohler, J. L., Liu, S., & Heemers, H. V. *Regulators of Androgen Action Resource: a one-stop shop for the comprehensive study of androgen receptor action*. Database: the journal of biological databases and curation, 2016, bav125) were used for AR coregulatory gene list. Gene ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) lists were also used in enrichment analyses. Where used herein in regard to gene lists, the term “essential” does not imply that it is essential for a particular gene or gene product to be targeted in order for the presently disclosed method be effective.

**[0112]** P-values for enrichment analyses were calculated using the following formula:

$$p = 1 - \sum_{i=0}^{m-1} \frac{\binom{M}{i} \binom{N-M}{n-i}}{\binom{N}{n}}$$

**[0113]** where p=p value, N=total genes, M=genes of pathway/gene list, n=differentially expressed genes, and i=overlap of M and n.

#### Sylamer Analysis

**[0114]** The following online resource was used for sylamer analyses: <https://www.genomique.info/sylamer/>.

Genes from a given comparison were rank ordered based on fold differential gene expression from most downregulated to most up regulated and submitted to sylamer analysis. Sylamer analysis compares enrichment p values of all possible words of a given length (6,7,8) from sequences of transcript 5'UTR, coding region or 3'UTR, and provides the most enriched words and corresponding enrichment log 10 p values.

#### shRNA Seed Match Analysis

**[0115]** The following online resource was used to generate strand specific shRNA seed match gene lists: <https://dharmacon.horizondiscovery.com/resources/tools-and-calculators/sirna-seed-locator/>. Since the online resource is intended for siRNA with a length of 19 nucleotides, while TRC-shRNAs contain 21 nucleotide target sequences, probable Drosha/Dicer cuts were taken into account when submitting 19 nucleotide sequences. A 2-nucleotide trim was used for the passenger strand and a 3 nucleotide trim was used for the guide strand. As an additional control, a shTMEFF2-3 guide strand sequence with 2 nucleotides trimmed was also used (shTMEFF2-3 Guide +1, see Table S4. Appendix 1). After seed match lists were generated, genes with multiple seed matches were eliminated from single seed match lists. Single and multiple seed match lists were compared to essential and AR coregulatory gene lists. Statistics: Pearson’s Chi-square test of independence was used to identify significant associations between seed match and gene downregulation or upregulation. Yates correction was used for comparisons that contained values of 10 genes or fewer to limit type 1 error rate associated with small sample size. P<0.05 was considered statistically significant.

#### Cell Growth and Viability Assays

**[0116]** For growth and viability analyses, LNCaP, 22Rv1, DU145, Panc-1 and HEK293T cell lines were transduced with pLKO.1 shRNA constructs as described in the cell culture and plasmid constructs section of the materials and methods. Cells were trypsinized and seeded in 6 well plates 24 hours post-transduction at the following concentrations: 1×10<sup>5</sup> cells/well (HEK293T, DU145, Panc-1), 2×10<sup>5</sup> cells/well (LNCaP, 22Rv1). Cells were counted using Nexcelom Auto T4 Cellometer, and trypan blue was used to selectively count live cells and assess viability. Cells were trypsinized and counted using the same method 24 hours after seeding (36 hours after seeding for LNCaP, which require longer time for attachment) for initial cell count. Growth was analyzed by counting cells again at 48 hours after initial cell count, and lastly 72 hours (DU145, Panc-1, HEK293T) or 96 hours after initial cell count (LNCaP, 22Rv1 due to slower growth rate of these cells). Viability was assessed at each time point via trypan blue, and percent viability was averaged for 48 and 72 hour time points (HEK293T, DU145, Panc-1) or 48 and 96 hour time points for slower growing cell lines (22Rv1, LNCaP). Viability of RWPE1 cells transduced with shRNA was determined at 96 hours and 120 hours after transduction. Relative percent viability was then calculated by dividing percent viability of knockdown cell lines by percent viability in shScramble control. Percent growth inhibition was calculated using the following formula:

$$\% GI = 1 - \left[ \frac{\frac{(K_f - K_i)}{K_i}}{\left[ \frac{(S_f - S_i)}{S_i} \right]} \right]$$

[0117] where % GI=% Growth Inhibition,  $K_f$ =Final cell count of knockdown cell line,  $K_i$ =Initial cell count of knockdown cell line,  $S_f$ =Final cell count of shScramble cell line,  $S_i$ =Initial cell count of shScramble cell line. Growth and viability analyses were repeated 3 times for each cell line.

[0118] Statistics Significance was calculated using two-tailed T-test. P<0.05 was considered to be statistically significant.

#### Identification of Toxic Seed Sequences by Large-Scale Screening

[0119] A large-scale viability screen was used to identify 6-mer seed RNAi sequences that were most efficient at promoting toxicity in prostate cancer at different stages of the disease. The general strategy, schematically represented in FIG. 5, conducted a viability screen using a pooled shRNA library that represents all possible 6-mer seed sequences during the course of 0-28 days, followed by next generation sequencing (NGS) to quantify shRNA representation. shRNAs that decrease viability (i.e., were toxic) are depleted in the final population. Comparison across cell lines/conditions define the shRNAs that preferentially target the different stages of the disease. shRNAs were synthesized as oligonucleotides differing in a 7-nt seed sequence. Adding a nucleotide to the conventional 6-mer seed accounts for potential Dicer “shifts” resulting in noncanonical cuts. The 16,384 ( $4^7$ ) different 7-nt seed sequences were inserted in a shRNA backbone and cloned into a doxycycline (Dox) inducible PLKO.1 TRC lentiviral cloning vector modified to include primer target sequences for PCR shRNA recovery, and unique barcode sequences to identify the presence of each of the different shRNAs (Celllecta). Only 15,572 were used in the final screen after removal of redundant sequences that could compromise the expression (i.e. TTTTTTT) or that contained restriction sites incompatible with the cloning. The resulting plasmid pool was sequenced to confirm shRNA representation, and then packaged into lentiviral particles, titrated and transduced into the selected cell lines (22Rv1 grown in the presence or absence of dihydrotestosterone, DHT) at a MOI of 0.3-0.5 to avoid multiple infection. After puromycin selection, a portion of cells was collected as day 0 sample. The rest was continuously cultured in the presence of Dox and maintained at an average of 1000-fold shRNA representation for reproducibility. To compute the relative effects of the shRNAs, we determined the average barcode representation from samples collected at 7 and 28 days of growth (corresponding to 2 and 10 population doublings) and compared that with the average initial barcode representations at day 0. Two biological replicates will be used per cell line/condition/time point. shRNAs that are toxic to the cell and are thus depleted at the end of the screen, were considered hits. shRNA hits are selected based on comparison across the cell lines/conditions to identify those that are most toxic to each specific stage of the disease. Selected cell lines/conditions used in this screen represent distinct disease stages: 22Rv1 is a

CRPC cell line that constitutively expresses an active AR splice variant, AR-V7, as well as the full-length AR (ARfl), and, as a result, it displays hormone-independent growth representing advanced disease (in -DHT condition), but can also respond to hormonal stimulation (+DHT condition).

#### Results

##### Multiple shRNAs Targeting TMEFF2 Inhibit Androgen Signaling

[0120] Western blot analysis was used to determine relative TMEFF2 and AR protein expression in three benign prostate epithelial, three ADPC and four CRPC cell lines. We found a strong coincidence of TMEFF2, PSA and AR expression, and noticed a trend toward higher TMEFF2 levels in ADPC cell lines (see FIG. S1A of U.S. Provisional application Ser. No. 62/935,521), correlating with a previous finding that TMEFF2 mRNA levels are higher in primary PCa compared to benign prostate and CRPC. In order to test our hypothesis that TMEFF2 functions as a regulator of androgen signaling in PCa, we used shRNA to knockdown TMEFF2 in four TMEFF2+/AR+PCa cell lines, including two ADPC cell lines (LNCaP and VCaP), and two CRPC cell lines, (22Rv1 and C4-2B), and measured androgen responsive protein levels in presence and absence of DHT via western blot analyses. In LNCaP cells, transductions with nine independent shRNAs (shTMEFF2-1 thru shTMEFF2-9), targeting eight of the ten exons and 3'UTR of the TMEFF2 transcript, resulted in lower protein expression of two androgen responsive genes (PSA and FKBP5) and AR compared to cells transduced with scramble control shRNA (shScramble) (FIG. 1A). Furthermore, lentiviral dose response experiments with two TMEFF2-targeted shRNAs demonstrated that TMEFF2 and PSA reductions tightly correlate with one another, even at low lentiviral dose (FIG. 1B herein, FIG. S1B of U.S. Provisional application Ser. No. 62/935,521). TMEFF2 was knocked-down with two shRNAs in 22Rv1, C4-2B and VCaP cell lines, resulting in reductions in AR targets in all cases (FIG. 1C herein, FIG. S1C of U.S. Provisional application Ser. No. 62/935,521). TMEFF2-targeted shRNAs resulted in reductions in total and nuclear levels of full length AR isoform 1 and the constitutively active AR-V7 in 22Rv1 cells (FIGS. 1C-D), suggesting that the effect on androgen signaling is not dependent on the AR ligand binding domain. RT qPCR analysis of 53 genes commonly upregulated by the AR showed down regulation in 88% of genes (47 of 53; median log 2 fold change: -0.7212) in shTMEFF2-2 vs shScramble control expressing 22Rv1 cells (FIG. S1D of U.S. Provisional application Ser. No. 62/935,521), further supporting androgen signaling inhibition in response TMEFF2-targeted shRNA. In correlation with reduced androgen signaling, we observed a high amount of toxicity and difficulty generating cell lines with stable TMEFF2 knockdown. In LNCaP shTMEFF2-3 cells, TMEFF2 mRNA levels recovered after approximately 6 days in culture, while reductions in KLK3 (PSA transcript) remained for more than two weeks after transduction (FIG. S1E of U.S. Provisional application Ser. No. 62/935,521).

The Inhibition of Androgen Signaling by TMEFF2 shRNA is not a Consequence of Low TMEFF2

[0121] To further confirm that the androgen signaling phenotype was on-target, we overexpressed recombinant TMEFF2 isoform 1 in C4-2B cells using a doxycycline (Dox) inducible vector (FIG. S2A of U.S. Provisional appli-

cation Ser. No. 62/935,521), and determined TMEFF2 over-expression resulted in negligible changes in PSA expression (FIG. S2A of U.S. Provisional application Ser. No. 62/935, 521), and was also insufficient to rescue the reduction of PSA or AR protein levels by TMEFF2 targeted shRNA (FIG. 2A). One potential explanation for this discrepancy is that TMEFF2 has multiple isoforms; therefore, targeting of other isoforms could be responsible for the effect on androgen signaling seen with shRNA-mediated knockdown. To determine whether this is the case, we used a Dox inducible CRISPR Cas9 system in LNCaP cells to knockdown TMEFF2 protein expression via induced frameshift mutations in exon 1 of TMEFF2. Of note, all known TMEFF2 isoforms contain exon 1 in their open reading frame (ORF). Using this method, TMEFF2 protein expression was knocked-down >90% with the addition of Dox in LNCaP cells expressing the TMEFF2 targeted sgRNA (sgTMEFF2-1) compared to the control sgRNA targeting GFP (sgGFP) (FIG. 2B). Significant reductions in TMEFF2 mRNA were also noticed, presumably due to nonsense mediated decay of the transcript; however, PSA levels were not affected by CRISPR Cas9 mediated TMEFF2 knockdown (FIG. 2C), nor was toxicity evident in these cells (data not shown). In order to detect potentially unknown isoforms lacking the canonical AUG or exon 1 in ORFs, or long non-coding RNAs (lncRNAs) potentially targeted by TMEFF2 shRNAs, we conducted a deep paired-end RNA-seq analysis of the TMEFF2 locus in LNCaP cells for de novo isoform detection. Only transcript isoform 1 (ENST00000272771.9) contains all nine of the TMEFF2 shRNA target sequences, while isoform 2 (ENST00000392314.5) contains eight of nine target sequences (Table S1 and Appendix 1 of U.S. Provisional application Ser. No. 62/935,521), and both contain exon 1 and the canonical AUG. All other transcripts detected contain two or fewer TMEFF2 shRNA target sequences. Therefore, sequencing analyses suggest that it is highly improbable for the nine TMEFF2 shRNAs to be targeting an isoform from the TMEFF2 locus not targeted by CRISPR Cas9.

[0122] The divergent androgen signaling phenotypes in shRNA and CRISPR Cas9 mediated TMEFF2 knockdown could have been the result of OTEs, CRISPR Cas9 mediated concealment of low androgen signaling phenotype, or low TMEFF2 levels could have been interacting with shRNA-induced cell stress to elicit the low androgen signaling phenotype. To test these possibilities, CRISPR Cas9 sgGFP and sgTMEFF2 LNCaP cells were Dox treated to induce knockdown, followed by transductions with either shTMEFF2-3 or shScramble shRNA. In both CRISPR Cas9 LNCaP cell lines (sgGFP and sgTMEFF2), TMEFF2 targeted shRNA led to a reduction in PSA, FKPB5 and AR protein levels, while the levels of these proteins remained unaffected in shScramble control cells (FIG. 2D). Similar results were also obtained using a different sgRNA targeting TMEFF2 (sgTMEFF2-3) expressed from a constitutive expression vector (pLX-sgRNA) in LNCaP cells (FIG. S2B of U.S. Provisional application Ser. No. 62/935,521).

[0123] In addition to CRISPR Cas9, pooled FANA anti-sense oligos (ASOs), which work via nuclear RNaseH mediated transcript degradation, were also used to knockdown TMEFF2 in LNCaP cells. Similar to CRISPR Cas9 mediated knockdown, no reductions in PSA or AR protein levels were evident using ASO mediated TMEFF2 knockdown (FIG. S2C of U.S. Provisional application Ser. No.

62/935,521). Together, these data suggest that while the low androgen signaling phenotype is consistent using multiple independent TMEFF2-targeting shRNAs and several PCa cell lines, it was not the result of low TMEFF2 expression, but likely the result of OTEs.

#### shRNA Targeting TMEFF2 Downregulates Essential and AR Coregulator Gene Expression and Inhibits Global Androgen Transcriptional Response

[0124] In order to delineate target genes responsible for the low androgen signaling phenotype induced by TMEFF2 targeted shRNA, we conducted RNA seq analyses from LNCaP shTMEFF2-3 and shScramble cells grown in the presence and absence of DHT. RNA was extracted three days after shRNA transductions in order to ensure adequate expression of TMEFF2 targeted shRNA at the time of analysis. Log 2 fold change >0.5 and adj p-value <0.05 were used to define significant differential gene expression. Differential gene expression was determined between TMEFF2 knockdown and control cells and in response to DHT treatment in both cell lines. A near total inhibition of androgen transcriptional response was evident in LNCaP shTMEFF2-3 cells, in which 13 genes were differentially expressed with DHT treatment, compared to 1,423 genes in shScramble control LNCaP cells (data shown in FIG. 3A of U.S. Provisional application Ser. No. 62/935,521). Furthermore, of the genes differentially expressed with DHT treatment in LNCaP shScramble cells, a majority (867 of 1423 genes) were also differentially expressed between LNCaP shTMEFF2-3 and shScramble cells in the presence or absence of DHT (data shown in FIG. 3A of U.S. Provisional application Ser. No. 62/935,521). Gene ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analyses indicate that many of the gene sets and pathways enriched among DHT induced genes in LNCaP shScramble control cells, such as, cell division and DNA replication, are also enriched among genes downregulated by shTMEFF2-3 in the presence of DHT (data shown in FIG. 3B of U.S. Provisional application Ser. No. 62/935, 521), suggesting inhibition of androgen signaling is a strong contributor to the transcriptomic landscape and phenotype of shTMEFF2-3 expressing LNCaP cells. Genes within the p53 signaling pathway were enriched among genes elevated in LNCaP shTMEFF2-3 cells compared to shScramble in the presence of DHT (data shown in FIG. 3B of U.S. Provisional application Ser. No. 62/935,521), consistent with toxicity noticed in LNCaP shTMEFF2-3 cells. In effort to delineate the mechanism through which shTMEFF2-3 inhibits androgen signaling, we utilized Qiagen's Ingenuity Pathway Analysis (IPA) to identify potential causal networks and master regulators affected by shTMEFF2-3. AR and multiple AR coregulators were among the master regulators identified (data shown in FIG. 3C of U.S. Provisional application Ser. No. 62/935,521), and of those, AR, TAF1 and USP12 were also significantly downregulated by shTMEFF2-3 (data shown in FIGS. 3C and S3A-B of U.S. Provisional application Ser. No. 62/935,521). In order to determine whether the downregulation of AR coregulators is a broader effect in LNCaP shTMEFF2-3 cells, we utilized a published list of AR coregulators (DePriest et al., 2016) and found that a significant number (38 out of 274,  $p<7\times 10^{-6}$ ) were downregulated by shTMEFF2-3 in the presence or absence of DHT, while similar significance levels were not found among differentially expressed genes in other comparisons (data shown in FIGS. 3C and S2A of U.S. Provisional

application Ser. No. 62/935,521). These data indicate that shTMEFF2-3 inhibits androgen signaling in LNCaP cells, at least partly, through the depletion of AR and AR coregulators.

**[0125]** To confirm the effect of shRNA to TMEFF2 on AR and AR-coregulators, we conducted RNA-Seq analysis in LNCaP cells expressing 3 different shRNAs to TMEFF2, shTMEFF2-3, -4, and -9, and with shL3, a shRNA that targets CD45L and is known to promote DISE. For these experiments, cells were grown in complete media and RNA was extracted 55 hours after transfection with the shRNAs to circumvent potential changes in AR targets secondary to loss of viability, which we observe only after 72 hrs after transfection. Differential gene expression (DGE) was determined comparing cells expressing the different shRNAs to those expressing the shScramble control, followed by Metascape Enrichment Analysis of the down- and up-regulated gene lists. Enriched terms for downregulated genes common to all shRNAs were related to androgen response and endocrine therapy resistance (top 3 categories); these were not represented in the top GO terms enriched for upregulated genes (data shown in FIGS. 3D-F of U.S. Provisional application Ser. No. 62/935,521). These results indicate that shRNA to TMEFF2 promotes global downregulation of the AR transcriptional response in PCa cells. Importantly, shL3 also promoted downregulation of the AR transcriptional response in LNCaP cells, suggesting that DISE in PCa cells involves AR signaling, perhaps anticipated based on the essential role of AR signaling in survival of PCa cells. In addition, using the published list of AR coregulators (DePriest et al., 2016), we observed that the downregulated genes in the LNCaP cells transfected with for each of the shRNAs were significantly enriched in AR coregulators (data shown in FIG. 3B of U.S. Provisional application Ser. No. 62/935,521).

**[0126]** We hypothesized that the toxicity induced by TMEFF2 targeted shRNA was similar to the previously described RNAi-mediated cell death mechanism, DISE, which is the result of OTEs that involve si/shRNA seed interactions with the 3' UTR of multiple essential survival genes, and is characterized, in part, by transcriptomic effects that include the downregulation of essential survival and histone transcripts. Using essential gene lists generated from two independent studies of CRISPR Cas9 lethality screens, which used leukemia and lymphoma cell lines (Wang et al., 2015), or LNCaP cell line (Fei et al., 2017), we found significant enrichments for both essential gene lists in the set of genes downregulated in LNCaP shTMEFF2-3 cells ( $p<5\times 10^{-15}$ ,  $p<2\times 10^{-10}$ ) (data shown in FIG. 3G, Table S2B and Appendix 1 of U.S. Provisional application Ser. No. 62/935, 521). We also detected an enrichment of essential genes induced by DHT in LNCaP shScramble cells ( $p<4\times 10^{-65}$ ,  $p<4\times 10^{-10}$ ) as expected, due to the androgen dependent nature of LNCaP cells. Given the impact of shTMEFF2-3 on androgen signaling, it seemed plausible that androgen signaling inhibition was a major contribution of essential gene downregulation by shTMEFF2-3. Indeed, a significant overlap of common targets was evident between essential genes induced by DHT in LNCaP shScramble cells and those downregulated by shTMEFF2-3 for both essential gene lists ( $p<4\times 10^{-59}$ ,  $p<2\times 10^{-13}$ ) (data shown in FIG. 3G, Table S2C and Appendix 1 of U.S. Provisional application Ser. No. 62/935,521). In addition to essential survival genes, histone transcripts, while not significantly modulated by DHT in

LNCaP shScramble cells, were also significantly enriched among genes downregulated by shTMEFF2-3 (23 of 87 histone genes,  $p<3.8\times 10^{-9}$ ) (data shown in FIG. 3G, Table S2D and Appendix 1 of U.S. Provisional application Ser. No. 62/935,521), resembling the described DISE transcriptomic phenotype.

**[0127]** The shTMEFF2-3 guide strand seed region targets the 3'UTR of essential and AR coregulatory genes, leading to downregulation

**[0128]** In order to determine whether shTMEFF2-3 may downregulate essential and AR coregulatory genes in LNCaP cells through an off-target DISE-like mechanism (see FIG. 3 for workflow), we utilized Sylamer analysis (we ee) to determine significantly enriched 6mer and 8mer sequences associated with LNCaP shTMEFF2-3 downregulated genes in rank ordered gene expression lists. Sylamer analysis is a useful tool for miRNA discovery and delineating RNAi OTEs. The 6mer 3'UTR sequence (GGGAAA), complementary to a seed sequence on the shTMEFF2-3 guide strand (UUUCCC), was found to be the most significantly enriched 6mer sequence associated with total, as well as essential and AR coregulator, gene downregulation in LNCaP shTMEFF2-3 cells in the presence and absence of DHT (data shown in FIGS. 5A, S4A, S4B, Table S3 and Appendix 1 of U.S. Provisional application Ser. No. 62/935, 521). shTMEFF2-3 seed matched sequences were significantly enriched in coding region of shTMEFF2-3 total downregulated genes in LNCaP cells (data shown in FIG. S4C and Table S3 of U.S. Provisional application Ser. No. 62/935,521), although to a lesser extent, but not in the coding region of downregulated essential and AR coregulatory genes (data shown in Table S3 of U.S. Provisional application Ser. No. 62/935,521). shTMEFF2-3 seed matched sequences were also significantly enriched among 8mer 3'UTR sequences (data shown in Table S3 of U.S. Provisional application Ser. No. 62/935,521). These analyses indicate that shTMEFF2-3 guide strand seed-mediated OTEs significantly influence gene expression mainly through binding to complementary 3'UTR sequences in LNCaP cells expressing shTMEFF2-3.

**[0129]** To confirm the association between 3'UTR seed matches and shTMEFF2-3 downregulated essential and AR coregulatory genes, we used Dharmacon seed locator software (<https://dharmacon.horizondiscovery.com/resources/tools-and-calculators/sima-seed-locator/>) to find genes with single and multiple 3'UTR seed matches to the shTMEFF2-3 and shScramble shRNA passenger and guide strands. We then tested for associations between essential and AR coregulatory genes containing seed matches and shTMEFF2-3 downregulated genes in LNCaP cells; shTMEFF2-3 upregulated and genes induced by DHT in LNCaP shScramble cells served as control comparisons. The shTMEFF2-3 guide strand seed was significantly associated with LNCaP shTMEFF2-3 downregulated essential (Wang et al., 2015,  $p=5.1\times 10^{-7}$ ; Fei et al., 2017,  $p=8.4\times 10^{-5}$ ) and AR coregulatory genes ( $p=0.006$ ) (data shown in FIG. 5B, Table S4 of U.S. Provisional application Ser. No. 62/935, 521), further supporting the results obtained by Sylamer analyses. While associations are much weaker compared to the shTMEFF2-3 guide strand, some off target activity from the passenger strand cannot be ruled out. Together these data suggest that shTMEFF2-3 shRNA leads to the downregulation of essential and AR coregulatory genes through guide strand complementarity to the 3'UTR of said genes.

**[0130]** Next, because a significant number of androgen-induced essential genes are downregulated by shTMEFF2-3 (data shown in FIG. 3G of U.S. Provisional application Ser. No. 62/935,521), we repeated the association analyses after separating essential genes into androgen induced and non-androgen induced categories. The significance of the association between shTMEFF2-3 guide strand seed match and LNCaP shTMEFF2-3 downregulated genes increased for non-androgen induced essential genes (Wang et al., 2015, essential genes:  $p=5.2\times10^{-17}$ , Fei et al., 2017, essential genes:  $p=5.2\times10^{-7}$ , FIG. 5C of U.S. Provisional application Ser. No. 62/935,521), compared to all essential genes (Wang et al., 2015, essential genes:  $p=5.1\times10^{-7}$ , Fei et al., 2017, essential genes:  $p=8.4\times10^{-5}$ , FIG. 5B of U.S. Provisional application Ser. No. 62/935,521); however, associations with downregulated androgen induced essential genes were not significant (data shown in FIG. 5C of U.S. Provisional application Ser. No. 62/935,521). 86% and 74% of essential genes downregulated by shTMEFF2-3 in LNCaP cells either contained seed matches, were androgen responsive or both in Wang et al., 2015 and Fei et al., 2017 lists, respectively (data shown in FIG. S5 of U.S. Provisional application Ser. No. 62/935,521). These data indicate that androgen induced essential gene downregulation in shTMEFF2-3 cells is a consequence of androgen signaling inhibition, and likely the result of the seed-mediated downregulation of androgen signaling regulatory genes. Among those genes, the 3'UTR of AR was found to have four seed matches to the shTMEFF2-3 guide strand. A significant negative correlation was evident between AR mRNA expression and the number of guide strand seed matches to the AR 3'UTR when multiple TMEFF2-targeted shRNAs were compared to shScramble in LNCaP cells (data shown in FIG. 5D of U.S. Provisional application Ser. No. 62/935,521). One other TMEFF2 shRNA, shTMEFF2-9, was found to have four seed matches to the AR 3'UTR and significantly lower AR mRNA expression in LNCaP cells as well. Therefore, AR mRNA, in addition to numerous AR coreulatory genes, is a likely target of the shTMEFF2-3 seed, resulting in a strong repression of androgen signaling and the downregulation of androgen-induced essential genes.

**[0131]** Furthermore, Sylamer analysis identified miR-4446-5p as having an identical seed sequence to the shTMEFF2-3 guide strand. Using an online miRNA database (<http://www.mirbase.org/>), we found that miR-4446-5p predicted targets were significantly enriched with LNCaP shTMEFF2-3 downregulated genes, as well as, essential genes in particular (data shown in FIG. 5E of U.S. Provisional application Ser. No. 62/935,521). Additionally, shTMEFF2-3 downregulated genes in LNCaP cells had a significantly higher miR-4446-5p miRDB target score compared to other predicted targets (data shown in FIG. 5E of U.S. Provisional application Ser. No. 62/935,521), and miRDB target score negatively correlated with fold change gene expression in LNCaP shTMEFF2-3 cells, with the absence of DHT being more significant than in the presence of DHT (data shown in FIGS. S6A-B of U.S. Provisional application Ser. No. 62/935,521). Genes involved in cell proliferation were also found to be enriched among the 1148 predicted miR-4446-5p targets (FIG. 4C); these proliferative gene included genes downregulated by shTMEFF2-3 in LNCaP cells, such as RAD21, CENPA and CDK7. Since miR-4446-5p has an identical 6mer seed to shTMEFF2-3, and little sequence similarity outside the seed region (data shown in

FIG. 5E of U.S. Provisional application Ser. No. 62/935,521), this correlative relationship between predicted targets and shTMEFF2-3 downregulated genes in LNCaP cells suggests a tumor suppressive function for miR-4446-5p.

**[0132]** Sylamer followed by seed match analyses were also conducted with downregulated genes from LNCaP cells expressing shTMEFF2-3, -4, -9, or shL3 and grown in complete media. Confirming the data presented above, we observed that for these shRNAs, downregulation of AR-coregulators is also significantly associated with shRNA seed matches (data shown in FIGS. 5F,G,H of U.S. Provisional application Ser. No. 62/935,521). Importantly, using the miRNA database (<https://www.mirbase.org/>), we found that miR-634 predicted targets were significantly enriched with LNCaP shTMEFF2-4 downregulated genes, and miRDB target score negatively correlated with fold change gene expression in LNCaP shTMEFF2-4 cells (data shown in FIG. S1 of U.S. Provisional application Ser. No. 62/935,521).

#### TMEFF2 shRNAs are Toxic to Transformed Cells and Sensitivity in LNCaP Cells Correlates with Lower Expression of AR Regulated Genes

**[0133]** To further characterize the toxicity elicited by TMEFF2-targeted shRNA, we analyzed the growth and viability of multiple transformed cell lines in response to four TMEFF2-targeted shRNAs (shTMEFF2-2, shTMEFF2-3, shTMEFF2-8 and shTMEFF2-9) and two shRNAs previously shown to induce DISE (shL3 targeting CD95L and shR6 targeting CD95), in comparison to shScramble control. For these experiments, we selected two AR positive PCa cell lines, 22Rv1 (CRPC) and LNCaP (ADPC), and three transformed cell lines of different origin, HEK293T (transformed embryonic kidney), Panc-1 (pancreatic carcinoma) and DU145 (AR negative CRPC). Of note, DU145 and Panc-1 cell lines express little to no TMEFF2 (data shown in FIG. S1A of U.S. Provisional application Ser. No. 62/935,521). All of the targeted shRNAs tested significantly reduced the growth and viability of transformed cells, with few exceptions, when compared to shScramble control cells (data shown in FIGS. 4A-B herein, and in S7A of U.S. Provisional application Ser. No. 62/935,521), which maintained high viability percentage in all cell lines tested (70%-86%) (data shown in FIG. S7B of U.S. Provisional application Ser. No. 62/935,521); however, two of the most toxic TMEFF2-targeted shRNAs in LNCaP cells did not significantly affect the viability of RWPE1 cells, a benign prostate epithelial cell line, while shL3 resulted in only a 15% reduction in viability compared to RWPE1 shScramble cells (data shown in FIG. S7C of U.S. Provisional application Ser. No. 62/935,521), in agreement to previous reports that DISE inducing shRNAs are non-toxic to normal cells. Using LNCaP cells expressing shL3, shTMEFF2-3 and shTMEFF2-9 as examples, toxic shRNAs induced a similar rounded cell morphology with long thin extensions (FIG. 4D), as well as, Caspase-3 cleavage as measured by western blot, indicative of cell stress and apoptosis activation; furthermore, the level of Caspase-3 cleavage correlated with reduction in measured viability in LNCaP cells (FIG. 4C).

**[0134]** Since multiple shRNAs targeting TMEFF2 inhibited androgen signaling and induced similar toxicity to DISE-inducing shRNAs, we investigated whether all nine TMEFF2-targeted shRNAs contain seed sequences with predicted toxicity using an in silico scoring system generated by Putzbach et al., 2017 (Putzbach, W., Gao, Q.Q., Pate,

1 M., van Dongen, S., Haluck-Kangas, A., Sarshad, A. A., Bartom, E. T., Kim, K. A., Scholten,s D. M., Hafner, M., Zhao, J. C., Murmann, A E., Peter, M. E. *Many si/shRNAs can kill cancer cells by targeting multiple survival genes through an off-target mechanism.* Elife. 2017; 6. Epub 2017/10/25) based on seed complementarity to the 3' UTR of essential genes vs nonessential genes, and using an in vitro seed toxicity screen data from Gao et al., 2018 (Gao Q Q, Putzbach W E, Murmann A E, Chen S, Sarshad A A, Peter J M, Bartom E T, Hafner M, Peter M E. *6mer seed toxicity in tumor suppressive microRNAs.* Nat Commun. 2018 Oct. 29; 9(1):4504). All possible 6mer seed sequences were analyzed in both sequence-based and viability analyses, and all possible 8mer seed sequences were also analyzed in sequence-based analyses. All seeds were rank ordered for the number of complementary sequences in essential survival gene 3' UTRs, toxicity index (normalized #of sequences in essential genes divided normalized #of sequences in a list of non-essential genes) and percent viability. TMEFF2-targeted shRNA seed sequences within the passenger and guide strands were consistently high percentile for the number of complementary sequences to essential genes (median percentile rank: 77.5, 3<sup>rd</sup> quartile rank: 91) (data shown in Table S5A and Appendix 1 of U.S. Provisional application Ser. No. 62/935,521), while 6mer toxic seeds, as predicted from the viability screen, were mostly located in the guide strand (median percentile rank: 69, 3<sup>rd</sup> quartile rank: 90) (data shown in Table S5B and Appendix 1 of U.S. Provisional application Ser. No. 62/935,521). While the viability screens did not include PCa cells, these data support our findings that TMEFF2 targeted shRNAs contain toxic seed sequences that induce DISE in transformed cells.

**[0135]** When comparing viability effects of the six toxic shRNAs across cell lines, we noticed a trend of higher toxicity in the two AR+PCa compared to the three other transformed cell lines, with shTMEFF2-9 demonstrating the most dramatic difference (data shown in FIG. S7D of U.S. Provisional application Ser. No. 62/935,521). Since shRNAs targeting TMEFF2 inhibit androgen signaling, a critical survival pathway in these cells, we next determined the effect of the four shTMEFF2 and shL3/shR6 shRNAs on androgen responsive gene expression in LNCaP cells. All six shRNAs, with few exceptions, significantly reduced the expression of four androgen responsive transcripts often used as surrogates for AR activity, KLK3, KLK2, NKX3-1 and TMPRSS2, when compared to shScramble control LNCaP cells 48 hours to 72 hours after shRNA transductions (data shown in FIG. 6D of U.S. Provisional application Ser. No. 62/935,521). Both shTMEFF2-3 and shTMEFF2-9, two of the more toxic shRNAs in LNCaP cells, demonstrated significantly more potent reductions in androgen responsive gene expression when compared to shTMEFF2-2, shTMEFF2-8 and shL3 (data shown in FIG. 6D of U.S. Provisional application Ser. No. 62/935,521). Additionally, the level of androgen responsive gene reduction for each shRNA significantly correlated with relative viability reduction in LNCaP cells compared to the three AR-independent transformed cell lines tested (HEK293T, Panc-1 and DU145) (data shown in FIG. 6E of U.S. Provisional application Ser. No. 62/935,521). These data demonstrate a relationship between the effect of toxic shRNA-mediated androgen signaling inhibition, likely through the depletion

of AR and/or AR coregulators as demonstrated with shTMEFF2-3, and toxicity in AR+PCa cells.

Identification of AR Coregulators by Complementarity to the shRNA

**[0136]** The following procedure can be carried out to identify AR coregulators whose expression is affected by shRNAs or have one or more sequences in the 3'-UTR that can be target by the shRNA: (1) Cells are transfected with each one of the shRNAs to be tested and with a control; (2) RNA-Seq is carried out to determine which genes are affected by each of the tested shRNAs (mRNA level increase or decrease with respect to the control); (3) The genes that are affected by each specific shRNA are identified and compared to a list of AR-coregulators; (4) Also, the 3'-UTR of all of the AR-coregulator mRNAs are searched (independently of whether their expression is changed or not based on the RNA-Seq data) for sequences that are complementary to the seed sequence of the tested shRNAs. If those sequences are present in any or more of the AR-coregulator's 3'-UTR, that would suggest that those specific AR-coregulators are targets for that particular shRNA; (5) Genes that are downregulated and contain one or more sequences complementary to the seed sequence of the specific shRNA would be bona fide targets.

**[0137]** In each of Tables 2, 3 and 4, (shTMEFF2-3, shTMEFF2-4 and shTMEFF2-9), there are 3 columns. The righthand column lists AR-coregulators that are definitely targets of the specific shRNA. The lefthand column lists AR-coregulators whose expression is not affected by the shRNA as seen in the RNA-Seq results, but which have one sequence in its 3'-UTR that can be target by the tested shRNA. The center column lists AR-coregulators whose expression is not affected by the shRNA as seen in the RNA-Seq results, but which have more than one sequence in their 3'-UTR that can be targeted by the tested shRNA.

TABLE 2

AR coregulators downregulated by or having binding sites (single or multiple) to shTMEFF2-3		
Single	Multiple	Downregulated
AKT1	ACTN2	ADAM10
ARID5A	ARHGDA	APPBP2
ATXN7L3	ARID1A	CCND3
BRCA2	ARRB2	CDK7
CDC25A	BRCA1	CTDSP2
CDC25B	CBX1	FKBP5
CDK6	CCND1	HELZ2
DDC	CREBBP	MED1
DYRK1A	DDX17	PER1
EGFR	FKBP4	PIAS1
ENY2	HDAC3	PIK3CB
FHL2	HDAC4	RANBP10
GSK3B	IL6ST	TAF1
GSN	KAT2B	TPD52
HEY1	MAGEA11	YWHAH
HIP1	NCOA3	
HSP90AA1	NCOA6	
HSPA4	NRIP1	
HTATIP2	NSD1	
IDE	PATZ1	
IRS1	PIAS2	
JMJD1C	PPARGC1A	
KAT7	PSPC1	
KDM4C	RBAK	
KDM5B	RBM14	
KHDRBS1	RCHY1	
KIF1A	RNF4	

TABLE 2-continued

AR coregulators downregulated by or having binding sites (single or multiple) to shTMEFF2-3		
Single	Multiple	Downregulated
MAK	SENP1	
MAPK1	SMAD3	
MED24	SMAD4	
MYO6	SMARCC1	
NCOA1	STAT5B	
NCOA2	TCF4	
NCOA4	WDR77	
NCOR1		
NONO		
PRKD1		
PRMT2		
PSMC3IP		
PXN		
RPS6KA3		
SFN		
SFPQ		
SMAD1		
SOD2		
SORBS3		
SRA1		
SRC		
SRCAP		
SUMO2		
SUMO3		
SVIL		
TDG		
TOB2		
TRIM68		
UBE2I		
UBE2L3		
UBE3A		
VAV3		
ZMIZ1		
ZNF318		
ZNF451		

TABLE 3

AR coregulators downregulated by or having binding sites (single or multiple) to shTMEFF2-4		
Single	Multiple	Downregulated
RCHY1	CREBBP	FKBP5
TPD52	PXN	CTDSP2
HEY1	ATXN7L3	RANBP10
HAP1	MAK	BRCA1
VAV3	KIF1A	CALM1
CASP8	PMEPA1	CALR
TCF4	PIK3CB	CCND1
DDX17	ZNF451	CDC25A
CCND3	DYRK1A	CTNNB1
SMAD3	MDM2	EHMT2
GSK3B	STAT3	FKBP4
CHD8	SMAD4	IDE
BAG1	IRS1	KAT2B
KAT7	AATF	MAPK1
KAT5	USP22	PIK3R1
PRMT2	ZMIZ1	PSMC3IP
HIP1	CCAR2	RPS6KA1
UBE2L3	ARRB2	RPS6KA3
CDK9		SMAD1
EP300		
PPP1CC		
UBE2I		
KMT2D		
NRIP1		
TRIM24		
MED17		

TABLE 3-continued

AR coregulators downregulated by or having binding sites (single or multiple) to shTMEFF2-4		
Single	Multiple	Downregulated
MED1		
SFPQ		
USP10		
EGFR		
STUB1		
HDAC4		
DNAJB1		
POU4F1		
ANP32A		
CDC37		
LATS2		
KDM4A		
TRIM68		
PAK6		
KAT2A		
NSD1		
CALCOCO1		
PRMT1		
PPP2R1A		
PSPC1		
SENP1		

TABLE 4

AR coregulators downregulated by or having binding sites (single or multiple) to shTMEFF2-9		
Single site	Multiple sites	Downregulated
HSP90AA1	SMAD1	ADAM10
SRA1	HDAC7	APPBP2
HAP1	SMARCA4	BAG1
TCF4	SMAD3	CALM1
DDX17	ENY2	CDK2AP1
ATXN7L3	KMT2D	FKBP5
SIRT1	NRIP1	GSK3B
KDM4C	SMAD4	HIP1
NCOA3	CTDSP2	HIPK3
KMT2A	HDAC4	IL6ST
RBM14	PIAS2	KAT2B
KAT5	PSPC1	MAPK1
PIK3R1	BRCA1	MED1
PPP1CC	SENP1	MED21
ZNF451		MKRN1
CDK9		NCOA2
DYRK1A		PMEP1
FKBP4		PRMT2
MDM2		RANBP9
NPM1		RPS6KA3
SFRP1		SMARCD1
TDG		UBE2L3
TGIF1		ZMIZ1
UBE2I		
NCOA1		
ARRB2		
SMAD1		
KHDRBS1		
TMF1		
STAT5B		
PATZ1		
PSIP1		
CCAR2		

Identification of AR Coregulators by Analysis of Copy Number Alterations (CNA)

**[0138]** Androgen receptor (AR)-mediated transcription is dependent on the recruitment of AR coregulators, which modulate its transcriptional response. Expression of AR coregulators is usually de-regulated in prostate cancer (PCa), and generally increased expression correlate with aggressive disease and poor clinical outcome. Studies on AR signaling during PCa progression have indicated that: 1) aberrant expression of coregulators is one of the mechanisms that leads to therapeutic resistance; 2) specific coregulators control the expression of subsets of AR target genes, and 3) the AR drives a distinct transcriptional program in advanced CRPC. Accordingly, some coregulators promote AR signaling in low androgen conditions, such as under androgen-deprivation therapy (ADT), or in CRPC. We subjected a list of 274 AR coregulators (reported to modulate AR transcriptional output) to analysis of copy number alterations (CNA) in multiple clinical cohorts of metastatic prostate cancer (PCa) extracted from cBioportal for cancer genomics. The analysis was carried out in three different cohorts, and specific coregulators were selected if their copy number was increased in at least 10% of the patients. We observed 21, 14 and 8 genes with high copy number in the Fred Hutchinson CRC dataset (Kumar, A., Coleman, I., Morrissey, C., Zhang, X., True, L. D., Gulati, R., Etzioni, R., Boulari, H., Montgomery, B., White, T., et al., 2016, Substantial interindividual and limited intraindividual genomic diversity among tumors from men with metastatic prostate cancer. *Nat. Med.* 22, 369-378), the SU2/PCF Dream Team dataset (Abida, W., Cyrtà, J., Heller, G., Prandi, D., Armenia, J., Coleman, I., . . . Sawyers, C. L., 2019, Genomic correlates of clinical outcome in advanced prostate cancer. *Proceedings of the National Academy of Sciences of the United States of America*, 116(23), 11428-11436. doi:10.1073/pnas.1902651116); and the University of Michigan dataset (Grasso, C. S., Wu, Y. M., Robinson, D. R., Cao, X., Dhanasekaran, S. M., Khan, A. P., Tomlins, S. A., 2012, The mutational landscape of lethal castration-resistant prostate cancer. *Nature*, 487(7406), 239-243. doi:10.1038/nature11125). Twelve of the genes were in common for the first two datasets and those included the 8 amplified in the University of Michigan dataset. These results indicate that these 12 genes have a role in advanced disease: ATAD2, CCND1, COPS5, ENY2, HEY1, MAPK15, NCOA2, PQBP1, PRKDC, RPS6KA3, TPD52, UXT.

Seed Sequences of Interfering RNAs

**[0139]** Examples of RNA oligomers that can be used in accordance with the present disclosure to treat prostate cancer include, but are not limited to, RNAs such as the 5'-3' sequences in Tables 5 and 8 below, which comprise 6-mer "seed" sequences that bind to target AR coregulator nucleic acids. The RNAs may be single-stranded or may be double stranded and include the complementary strand. The 6-mer seed sequence portions of the sequences that are underlined in Table 5 are specifically shown in Table 6. DNA equivalents of the sequences in Table 6 below (i.e., where U is replaced with T) may also be used in certain embodiments of the present disclosure, for example as shown in Table 9.

TABLE 5

Interfering RNA sequences*		
siTMEFF2-1	<u>CUCUGUCAUCAUAACCAGA</u>	SEQ ID NO: 16
siTMEFF2-2	<u>CACUUGAACUGACAGACGC</u>	SEQ ID NO: 17
siTMEFF2-3	<u>AUUUCCCAUCAGAACGCGA</u>	SEQ ID NO: 18
siTMEFF2-4	<u>AACCAGCAUCACACCUGCA</u>	SEQ ID NO: 19
siTMEFF2-5	<u>AUUAACCAACAAUGCAAGGC</u>	SEQ ID NO: 20
siTMEFF2-6	<u>ACUAGUUUCUCCAGAGCCU</u>	SEQ ID NO: 21
siTMEFF2-7	<u>UGUUAGCAUUCUCUGCAUA</u>	SEQ ID NO: 22
siTMEFF2-8	<u>UGUUCGGACAAGGUAGU</u>	SEQ ID NO: 23
siTMEFF2-9	<u>UGUCUGAACUGUAGUGCCC</u>	SEQ ID NO: 24
miR-634	<u>AACCAGCACCCCAACUUUGGAC</u>	SEQ ID NO: 25
miR-4446-5p	<u>AUUUCCUGCCAUCUCCUUGGC</u>	SEQ ID NO: 26
miR-6888-3p	<u>AUCUGUCUCGAUUGUUUCCAG</u>	SEQ ID NO: 27

\*6-mer seed sequence portions of the sequences are underlined

TABLE 6

6-mer seed sequences of source siRNA sequences of Table 5	
siTMEFF2-1	UCUGUC
siTMEFF2-2	ACUUGA
siTMEFF2-3	UUUCCC
siTMEFF2-4	ACCAGC
siTMEFF2-5	UUACCA
siTMEFF2-6	CUAGUU
siTMEFF2-7	GUUAGC
siTMEFF2-8	GUUCCG
siTMEFF2-9	GUCUGA
miR-634	ACCAGC
miR-4446-5p	UUUCCC
miR-6888-3p	UCUGUC

**[0140]** Table 7 shows generic 10-mer sequences which contain, after a first generic nucleotide, a 6-mer seed sequences for RNA or DNA oligonucleotides, where X=U or T or other pyrimidine nucleobases as described elsewhere herein, followed by 3 additional generic nucleotides. The 10-mer sequence can comprise portions of larger oligonucleotides (e.g., 12-50 nucleotides) in the diagnostic and therapeutic nucleic acid compounds of the present disclosure. Also included are complementary sequences. Generic, RNA, and DNA versions of toxic seed sequences identified through the large-scale viability screening procedure shown in FIG. 5 are included in Tables 7, 8 and 9, respectively.

TABLE 7

Generic oligonucleotides with 6-mer seed sequences *		
Sequence	SEQ ID NO:	Nucleobase Alternatives
X <sub>1</sub> X <sub>2</sub> CX <sub>3</sub> GX <sub>4</sub> CX <sub>5</sub> X <sub>6</sub> X <sub>7</sub>	SEQ ID NO: 28	X <sub>1</sub> , X <sub>5</sub> , X <sub>6</sub> , X <sub>7</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>4</sub> = U or T***
X <sub>1</sub> ACX <sub>2</sub> X <sub>3</sub> GAX <sub>4</sub> X <sub>5</sub> X <sub>6</sub>	SEQ ID NO: 29	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> X <sub>2</sub> X <sub>3</sub> X <sub>4</sub> CCCCX <sub>5</sub> X <sub>6</sub> X <sub>7</sub>	SEQ ID NO: 30	X <sub>1</sub> , X <sub>5</sub> , X <sub>6</sub> , X <sub>7</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>4</sub> = U or T***
X <sub>1</sub> ACCAGCX <sub>2</sub> X <sub>3</sub> X <sub>4</sub>	SEQ ID NO: 31	X <sub>1</sub> -X <sub>4</sub> = C, G, U, A or T **
X <sub>1</sub> X <sub>2</sub> X <sub>3</sub> ACCA <sub>4</sub> X <sub>5</sub> X <sub>6</sub>	SEQ ID NO: 32	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> CX <sub>2</sub> AGX <sub>3</sub> X <sub>4</sub> X <sub>5</sub> X <sub>6</sub> X <sub>7</sub>	SEQ ID NO: 33	X <sub>1</sub> , X <sub>5</sub> , X <sub>6</sub> , X <sub>7</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>4</sub> = U or T***
X <sub>1</sub> GX <sub>2</sub> X <sub>3</sub> AGCX <sub>4</sub> X <sub>5</sub> X <sub>6</sub>	SEQ ID NO: 34	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> GX <sub>2</sub> X <sub>3</sub> CCGX <sub>4</sub> X <sub>5</sub> X <sub>6</sub>	SEQ ID NO: 35	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> GX <sub>2</sub> CX <sub>3</sub> GAX <sub>4</sub> X <sub>5</sub> X <sub>6</sub>	SEQ ID NO: 36	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> X <sub>2</sub> CCX <sub>3</sub> GX <sub>4</sub> X <sub>5</sub> X <sub>6</sub> X <sub>7</sub>	SEQ ID NO: 37	X <sub>1</sub> , X <sub>5</sub> , X <sub>6</sub> , X <sub>7</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>4</sub> = U or T***
X <sub>1</sub> X <sub>2</sub> ACX <sub>3</sub> GGX <sub>4</sub> X <sub>5</sub> X <sub>6</sub>	SEQ ID NO: 38	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> AX <sub>2</sub> CX <sub>3</sub> GX <sub>4</sub> X <sub>5</sub> X <sub>6</sub> X <sub>7</sub>	SEQ ID NO: 39	X <sub>1</sub> , X <sub>5</sub> , X <sub>6</sub> , X <sub>7</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>4</sub> = U or T***
X <sub>1</sub> X <sub>2</sub> X <sub>3</sub> GCAGX <sub>4</sub> X <sub>5</sub> X <sub>6</sub>	SEQ ID NO: 40	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> X <sub>2</sub> CCX <sub>3</sub> GGX <sub>4</sub> X <sub>5</sub> X <sub>6</sub>	SEQ ID NO: 41	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> ACX <sub>2</sub> GACX <sub>3</sub> X <sub>4</sub> X <sub>5</sub>	SEQ ID NO: 42	X <sub>1</sub> , X <sub>3</sub> X <sub>4</sub> , X <sub>5</sub> = C, G, U, A or T ** X <sub>2</sub> = U or T***
X <sub>1</sub> CCAGCCX <sub>2</sub> X <sub>3</sub> X <sub>4</sub>	SEQ ID NO: 43	X <sub>1</sub> -X <sub>4</sub> = C, G, U, A or T
X <sub>1</sub> X <sub>2</sub> ACACCX <sub>3</sub> X <sub>4</sub> X <sub>5</sub>	SEQ ID NO: 44	X <sub>1</sub> , X <sub>3</sub> X <sub>4</sub> , X <sub>5</sub> = C, G, U, A or T ** X <sub>2</sub> = U or T***
X <sub>1</sub> AAGX <sub>2</sub> ACX <sub>3</sub> X <sub>4</sub> X <sub>5</sub>	SEQ ID NO: 45	X <sub>1</sub> , X <sub>3</sub> X <sub>4</sub> , X <sub>5</sub> = C, G, U, A or T ** X <sub>2</sub> = U or T***
X <sub>1</sub> X <sub>2</sub> AAGX <sub>3</sub> AX <sub>4</sub> X <sub>5</sub> X <sub>6</sub>	SEQ ID NO: 46	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> CCCX <sub>2</sub> GGCX <sub>3</sub> X <sub>4</sub> X <sub>5</sub>	SEQ ID NO: 47	X <sub>1</sub> , X <sub>3</sub> X <sub>4</sub> , X <sub>5</sub> = C, G, U, A or T X <sub>2</sub> = U or T***
X <sub>1</sub> ACCX <sub>2</sub> GGX <sub>3</sub> X <sub>4</sub> X <sub>5</sub>	SEQ ID NO: 48	X <sub>1</sub> , X <sub>3</sub> X <sub>4</sub> , X <sub>5</sub> = C, G, U, A or T ** X <sub>2</sub> = U or T***
X <sub>1</sub> GCAGGX <sub>2</sub> X <sub>3</sub> X <sub>4</sub> X <sub>5</sub>	SEQ ID NO: 49	X <sub>1</sub> , X <sub>3</sub> X <sub>4</sub> , X <sub>5</sub> = C, G, U, A or T ** X <sub>2</sub> = U or T***
X <sub>1</sub> X <sub>2</sub> GCAGGX <sub>3</sub> X <sub>4</sub> X <sub>5</sub>	SEQ ID NO: 50	X <sub>1</sub> , X <sub>3</sub> X <sub>4</sub> , X <sub>5</sub> = C, G, U, A or T ** X <sub>2</sub> = U or T***
X <sub>1</sub> GACX <sub>2</sub> X <sub>3</sub> GX <sub>4</sub> X <sub>5</sub> X <sub>6</sub>	SEQ ID NO: 51	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> GAX <sub>2</sub> GCCX <sub>3</sub> X <sub>4</sub> X <sub>5</sub>	SEQ ID NO: 52	X <sub>1</sub> , X <sub>3</sub> X <sub>4</sub> , X <sub>5</sub> = C, G, U, A or T ** X <sub>2</sub> = U or T***

TABLE 7-continued

Generic oligonucleotides with 6-mer seed sequences *		
Sequence	SEQ ID NO:	Nucleobase Alternatives
X <sub>1</sub> <u>X<sub>2</sub>AAGGCX<sub>3</sub>X<sub>4</sub>X<sub>5</sub></u>	SEQ ID NO: 53	X <sub>1</sub> , X <sub>3</sub> X <sub>4</sub> , X <sub>5</sub> = C, G, U, A or T ** X <sub>2</sub> = U or T***
X <sub>1</sub> X <sub>2</sub> <u>GAX<sub>3</sub>AGX<sub>4</sub>X<sub>5</sub>X<sub>6</sub></u>	SEQ ID NO: 54	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> X <sub>2</sub> X <sub>3</sub> <u>AGCCX<sub>4</sub>X<sub>5</sub>X<sub>6</sub></u>	SEQ ID NO: 55	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> X <sub>2</sub> <u>GX<sub>3</sub>X<sub>4</sub>X<sub>5</sub>AX<sub>6</sub>X<sub>7</sub>X<sub>8</sub></u>	SEQ ID NO: 56	X <sub>1</sub> , X <sub>6</sub> X <sub>7</sub> , X <sub>8</sub> = C, G, U, A or T * X <sub>2</sub> -X <sub>5</sub> = U or T***
X <sub>1</sub> X <sub>2</sub> X <sub>3</sub> <u>CGGCX<sub>4</sub>X<sub>5</sub>X<sub>6</sub></u>	SEQ ID NO: 57	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> X <sub>2</sub> <u>ACX<sub>3</sub>GX<sub>4</sub>X<sub>5</sub>X<sub>6</sub>X<sub>7</sub></u>	SEQ ID NO: 58	X <sub>1</sub> , X <sub>5</sub> , X <sub>6</sub> , X <sub>7</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>4</sub> = U or T***
X <sub>1</sub> X <sub>2</sub> <u>ACX<sub>3</sub>GAX<sub>4</sub>X<sub>5</sub>X<sub>6</sub></u>	SEQ ID NO: 59	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> <u>GX<sub>2</sub>GX<sub>3</sub>GX<sub>4</sub>X<sub>5</sub>X<sub>6</sub>X<sub>7</sub></u>	SEQ ID NO: 60	X <sub>1</sub> , X <sub>5</sub> , X <sub>6</sub> , X <sub>7</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>4</sub> = U or T***
X <sub>1</sub> <u>AX<sub>2</sub>GX<sub>3</sub>ACX<sub>4</sub>X<sub>5</sub>X<sub>6</sub></u>	SEQ ID NO: 61	X <sub>1</sub> , X <sub>4</sub> , X <sub>5</sub> , X <sub>6</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>3</sub> = U or T***
X <sub>1</sub> <u>X<sub>2</sub>AX<sub>3</sub>CAX<sub>4</sub>X<sub>5</sub>X<sub>6</sub>X<sub>7</sub></u>	SEQ ID NO: 62	X <sub>1</sub> , X <sub>5</sub> , X <sub>6</sub> , X <sub>7</sub> = C, G, U, A or T ** X <sub>2</sub> -X <sub>4</sub> = U or T***
X <sub>1</sub> <u>GAX<sub>2</sub>AGAX<sub>3</sub>X<sub>4</sub>X<sub>5</sub></u>	SEQ ID NO: 63	X <sub>1</sub> , X <sub>3</sub> X <sub>4</sub> , X <sub>5</sub> = C, G, U, A or T ** X <sub>2</sub> = U or T***

\*6-mer seed sequences are underlined

\*\*or other modified nucleobases as described elsewhere herein

\*\*\*or other pyrimidine nucleobases as described elsewhere herein.

[0141] Shown in Table 8 are examples of generic RNA oligonucleotides, comprising RNA versions of the generic 6-mer seed sequences of Table 7, for use in diagnostic or therapeutic treatments of the present disclosure. The RNA oligonucleotides represented may be single-stranded or may be a single strand of a double-stranded oligonucleotide. Also included are complementary sequences thereof and sequences in which U is replaced with T.

TABLE 8

Generic RNA oligonucleotide sequences with 6-mer seed sequences	
xUCUGUxxxxxxxxxxxxxx	SEQ ID NO: 64
xACUUGAxxxxxxxxxxxxx	SEQ ID NO: 65
xUUUCCCxxxxxxxxxxxxx	SEQ ID NO: 66
xACCAGCxxxxxxxxxxxxx	SEQ ID NO: 67
xUUACCAxxxxxxxxxxxxx	SEQ ID NO: 68
xCUAGGUxxxxxxxxxxxxx	SEQ ID NO: 69
xGUUAGCxxxxxxxxxxxxx	SEQ ID NO: 70
xGUUCCGxxxxxxxxxxxxx	SEQ ID NO: 71
xGUCUGAxxxxxxxxxxxxx	SEQ ID NO: 72
xUCCUGUxxxxxxxxxxxxx	SEQ ID NO: 73

TABLE 8-continued

Generic RNA oligonucleotide sequences with 6-mer seed sequences	
xUACUGGxxxxxxxxxxxxx	SEQ ID NO: 74
xAUCUGUxxxxxxxxxxxxx	SEQ ID NO: 75
xUUGCAGxxxxxxxxxxxxx	SEQ ID NO: 76
xUCCUGGxxxxxxxxxxxxx	SEQ ID NO: 77
xACUGACxxxxxxxxxxxxx	SEQ ID NO: 78
xCCAGGCCxxxxxxxxxxxx	SEQ ID NO: 79
xUACACCxxxxxxxxxxxxx	SEQ ID NO: 80
xAAGUACxxxxxxxxxxxxx	SEQ ID NO: 81
xUAAGUAxxxxxxxxxxxxx	SEQ ID NO: 82
xCCUGGCxxxxxxxxxxxxx	SEQ ID NO: 83
xACCUGGxxxxxxxxxxxxx	SEQ ID NO: 84
xGCAGGUxxxxxxxxxxxxx	SEQ ID NO: 85
xUGCAGGxxxxxxxxxxxxx	SEQ ID NO: 86
xGACUUGxxxxxxxxxxxxx	SEQ ID NO: 87

TABLE 8-continued

Generic RNA oligonucleotide sequences with 6-mer seed sequences	
xGAUGCCxxxxxxxxxxxxxx	SEQ ID NO: 88
xUAAGGCxxxxxxxxxxxxxx	SEQ ID NO: 89
xUGAUAGxxxxxxxxxxxxxx	SEQ ID NO: 90
xUUAGCCxxxxxxxxxxxxxx	SEQ ID NO: 91
xUGUUUAxxxxxxxxxxxxxx	SEQ ID NO: 92
xUUCGGCxxxxxxxxxxxxxx	SEQ ID NO: 93
xUACUGUxxxxxxxxxxxxxx	SEQ ID NO: 94
xUACUGAxxxxxxxxxxxxxx	SEQ ID NO: 95
xGUGUGUxxxxxxxxxxxxxx	SEQ ID NO: 96
xAUGUACxxxxxxxxxxxxxx	SEQ ID NO: 97
xUAUCAUxxxxxxxxxxxxxx	SEQ ID NO: 98
xGAUAGAxxxxxxxxxxxxxx	SEQ ID NO: 99

x = C, G, U, or A, or other modified nucleobases as described elsewhere herein.

[0142] Shown in Table 9 are examples of generic DNA oligonucleotides, comprising DNA versions of the generic 6-mer seed sequences of Table 7, for use in diagnostic or therapeutic treatments of the present disclosure. The sequences comprise a strand of a double-stranded DNA oligonucleotide. Also included are complementary sequences thereof and sequences in which T is replaced with U.

TABLE 9

Generic DNA oligonucleotides with 6-mer seed sequences	
xTCTGTCxxxxxxxxxxxxxx	SEQ ID NO: 100
xACTTGAXxxxxxxxxxxxxx	SEQ ID NO: 101
xTTTCCCxxxxxxxxxxxxxx	SEQ ID NO: 102
xACCAGCxxxxxxxxxxxxxx	SEQ ID NO: 67
xTTACCAxxxxxxxxxxxxxx	SEQ ID NO: 103
xCTAGTTxxxxxxxxxxxxxx	SEQ ID NO: 104
xGTTAGCxxxxxxxxxxxxxx	SEQ ID NO: 105
xGTTCCGxxxxxxxxxxxxxx	SEQ ID NO: 106
xGTCTGAxxxxxxxxxxxxxx	SEQ ID NO: 107
xTCCTGTxxxxxxxxxxxxxx	SEQ ID NO: 108
xTACTGGxxxxxxxxxxxxxx	SEQ ID NO: 109
xATCTGTxxxxxxxxxxxxxx	SEQ ID NO: 110
xTTGCAGxxxxxxxxxxxxxx	SEQ ID NO: 111
xTCCTGGxxxxxxxxxxxxxx	SEQ ID NO: 112
xACTGACxxxxxxxxxxxxxx	SEQ ID NO: 113

TABLE 9-continued

Generic DNA oligonucleotides with 6-mer seed sequences	
xCCAGCCxxxxxxxxxxxxxx	SEQ ID NO: 114
xAACACCxxxxxxxxxxxxxx	SEQ ID NO: 115
xAAGTACxxxxxxxxxxxxxx	SEQ ID NO: 116
xAAGTAxxxxxxxxxxxxxx	SEQ ID NO: 117
xCCTGGCxxxxxxxxxxxxxx	SEQ ID NO: 118
xACCTGGxxxxxxxxxxxxxx	SEQ ID NO: 119
xCAGGGTxxxxxxxxxxxxxx	SEQ ID NO: 120
xtGCAGGxxxxxxxxxxxxxx	SEQ ID NO: 121
xACTTGxxxxxxxxxxxxxx	SEQ ID NO: 122
xAATGCCxxxxxxxxxxxxxx	SEQ ID NO: 123
xtAAGGCxxxxxxxxxxxxxx	SEQ ID NO: 124
xtGATAGxxxxxxxxxxxxxx	SEQ ID NO: 125
xtTAGCCxxxxxxxxxxxxxx	SEQ ID NO: 126
xtGTTTAxxxxxxxxxxxxxx	SEQ ID NO: 127
xtTCGGCxxxxxxxxxxxxxx	SEQ ID NO: 128
xtACTGTxxxxxxxxxxxxxx	SEQ ID NO: 129
xtACTGAXxxxxxxxxxxxxx	SEQ ID NO: 130
xtGTGTGTxxxxxxxxxxxxx	SEQ ID NO: 131
xtATGTACxxxxxxxxxxxxx	SEQ ID NO: 132
xtATCATxxxxxxxxxxxxxx	SEQ ID NO: 133
xtGATAGAxxxxxxxxxxxxx	SEQ ID NO: 134

x = C, G, T, or A, or other modified nucleobases as described elsewhere herein.

[0143] Table 10 shows 7-mer DNA oligonucleotides, where X=C, G, T, or A, or other modified nucleobases as described elsewhere herein. The 7-mer sequence can comprise portions (“seed sequences”) of larger DNA oligonucleotides (e.g., 12-50 nucleotides), for example as shown in Table 11, wherein x or X=C, G, T, or A, or other modified cytosine-type, guanine-type, thymine-type, or adenine-type nucleobases as described elsewhere herein. For example, an oligonucleotide based on the generic nucleotide seed sequence TGXTXXX (SEQ ID NO:136), can have the sequence TGTTTGC, TGTTTGG, TGTTTGA, or TGTTTGT. Further, for example, in an oligonucleotide based on the generic nucleotide seed sequence xTGXTXXXxxxxxxxxxxxx (SEQ ID NO:173), the TGXTXXX portion can have the sequence TGTTTGC, TGTTTGG, TGTTTGA, or TGTTTGT. The 7-mer sequences may be used in diagnostic or therapeutic treatments of the present disclosure. The sequences may comprise a strand of a double-stranded DNA oligonucleotide. Also included are complementary sequences of Tables 10 and 11 and sequences thereof in which T is replaced with U. The letters A, T, G, and C can also represent adenine-type, thymine-type, guanine-type, and cytosine-type modified nucleobases such as those described above.

TABLE 10

7-mer seed sequences for DNA oligonucleotides	
TGXXXCX	SEQ ID NO: 135
TGXXXX	SEQ ID NO: 136
XXGAAXX	SEQ ID NO: 137
AGXXCXX	SEQ ID NO: 138
XXTGTXX	SEQ ID NO: 139
TAXXXGX	SEQ ID NO: 140
XXXTGXG	SEQ ID NO: 141
TATAXXX	SEQ ID NO: 142
TATGXXX	SEQ ID NO: 143
TAXGTXX	SEQ ID NO: 144
TAXGXAX	SEQ ID NO: 145
TAXXXAC	SEQ ID NO: 146
TAXXAGX	SEQ ID NO: 147
TAXXXGC	SEQ ID NO: 148
TXTGTXX	SEQ ID NO: 149
TXTXTAX	SEQ ID NO: 150
TXTXTXC	SEQ ID NO: 151
TXTAXXC	SEQ ID NO: 152
TXXXAGT	SEQ ID NO: 153
TXXXGGC	SEQ ID NO: 154
XATGTXX	SEQ ID NO: 155
XXATGGX	SEQ ID NO: 156
XXXTGGC	SEQ ID NO: 157
XXAXGGC	SEQ ID NO: 158
GCAAXXX	SEQ ID NO: 159
ATGXXXC	SEQ ID NO: 160
ACXXXGG	SEQ ID NO: 161
GAXXXTC	SEQ ID NO: 162
XXTXCTC	SEQ ID NO: 163
AGTAAGX	SEQ ID NO: 164
AGTGGGX	SEQ ID NO: 165
CATGGCX	SEQ ID NO: 166
GTGTATX	SEQ ID NO: 167
TACTCAX	SEQ ID NO: 168
TCGGTCTX	SEQ ID NO: 169

TABLE 10-continued

7-mer seed sequences for DNA oligonucleotides	
ACAGTAX	SEQ ID NO: 170
TCCTCTX	SEQ ID NO: 171
X = C, G, T, or A, or other modified nucleobases as described elsewhere herein.	
xTGXXXCXxxxxxxxxxxxxxx	SEQ ID NO: 172
xTGXTXXXxxxxxxxxxxxxxx	SEQ ID NO: 173
xxXGAAXXxxxxxxxxxxxxxx	SEQ ID NO: 174
xAGXXCXXxxxxxxxxxxxxxx	SEQ ID NO: 175
xxXTGTXXxxxxxxxxxxxxxx	SEQ ID NO: 176
xTAXXXGXxxxxxxxxxxxxxx	SEQ ID NO: 177
xxXTTGXGxxxxxxxxxxxxxx	SEQ ID NO: 178
xTATAXXXxxxxxxxxxxxxxx	SEQ ID NO: 179
xTATGXXXxxxxxxxxxxxxxx	SEQ ID NO: 180
xTAXGTXXxxxxxxxxxxxxxx	SEQ ID NO: 181
xTAXGXAXxxxxxxxxxxxxxx	SEQ ID NO: 182
xTAXXXACxxxxxxxxxxxxxx	SEQ ID NO: 183
xTAXXAGXxxxxxxxxxxxxxx	SEQ ID NO: 184
xTAXXGCxxxxxxxxxxxxxx	SEQ ID NO: 185
xtXTGTXXxxxxxxxxxxxxxx	SEQ ID NO: 186
xtTXTXTAXxxxxxxxxxxxxxx	SEQ ID NO: 187
xtTXTXTXCxxxxxxxxxxxxxx	SEQ ID NO: 188
xtXTTAXXCxxxxxxxxxxxxxx	SEQ ID NO: 189
xTXXXAGTxxxxxxxxxxxxxx	SEQ ID NO: 190
xTXXXGGCxxxxxxxxxxxxxx	SEQ ID NO: 191
xxATGTXXxxxxxxxxxxxxxx	SEQ ID NO: 192
xxXTATGGXxxxxxxxxxxxxxx	SEQ ID NO: 193
xxXTTGGCxxxxxxxxxxxxxx	SEQ ID NO: 194
xxAXAGGCxxxxxxxxxxxxxx	SEQ ID NO: 195
xGCAAXXXxxxxxxxxxxxxxx	SEQ ID NO: 196
xATGXXXCxxxxxxxxxxxxxx	SEQ ID NO: 197
xACXXXGGxxxxxxxxxxxxxx	SEQ ID NO: 198
xGAXXXTCxxxxxxxxxxxxxx	SEQ ID NO: 199
xxXTXCTCxxxxxxxxxxxxxx	SEQ ID NO: 200
xAGTAAGXxxxxxxxxxxxxxx	SEQ ID NO: 201

TABLE 11

Generic DNA oligonucleotide sequences with 7-mer seed sequences

TABLE 11-continued

Generic DNA oligonucleotide sequences with 7-mer seed sequences	
xAGTGGGXXXXXXXXXXXXXX	SEQ ID NO: 202
xCATGGCXXXXXXXXXXXXXX	SEQ ID NO: 203
xGTGTATXXXXXXXXXXXXXX	SEQ ID NO: 204
xACTCAXXXXXXXXXXXXXX	SEQ ID NO: 205
xCCTCTXXXXXXXXXXXXXX	SEQ ID NO: 206
xAACAGTAXXXXXXXXXXXXXX	SEQ ID NO: 207
xCCTCTXXXXXXXXXXXXXX	SEQ ID NO: 208

x or X = C, G, T, or A, or other modified nucleobases as described elsewhere herein.

[0144] Table 12 shows 7-mer RNA oligonucleotides, where X=C, G, U, or A, or other modified nucleobases as described elsewhere herein. The 7-mer sequence can comprise portions (“seed sequences”) of larger RNA oligonucleotides (e.g., 12-50 nucleotides), for example as shown in Table 13, wherein x or X=C, G, U, or A, or other modified cytosine-type, guanine-type, thymine-type, or adenine-type nucleobases as described elsewhere herein. For example, an oligonucleotide based on the generic nucleotide seed sequence UGXUXXX (SEQ ID NO:210) or based on the sequence UGUUUGX, can have the sequence UGUUUGC, UGUUUGG, UGUUUGA, or UGUUUGU. Further, for example, in an oligonucleotide based on the generic nucleotide seed sequence xUGXUXXXxxxxxxxxxxxx (SEQ ID NO:254), the UGXUXXX portion can, in non-limiting embodiments, have the sequence UGUUUGC, UGUUUGG, UGUUUGA, or UGUUUGU. The 7-mer sequences may be used in diagnostic or therapeutic treatments of the present disclosure. The sequences may comprise a strand of a double-stranded RNA oligonucleotide. Also included are complementary sequences of Tables 12 and 13 and sequences thereof in which U is replaced with T. The letters A, U, G, and C can also represent adenine-type, uracil-type, guanine-type, and cytosine-type modified nucleobases such as those described above.

TABLE 12  
7-mer seed sequences for RNA oligonucleotides

UGXXXCX	SEQ ID NO: 209
UGXUXXX	SEQ ID NO: 210
XXGAAXX	SEQ ID NO: 211
AGXXCXX	SEQ ID NO: 212
XXUGUXX	SEQ ID NO: 213
UAXXXGX	SEQ ID NO: 214
UAUAXXX	SEQ ID NO: 215
UAUGXXX	SEQ ID NO: 216
UAXGUXX	SEQ ID NO: 217
UAXGXAX	SEQ ID NO: 218

TABLE 12-continued

7-mer seed sequences for RNA oligonucleotides	
UAXXXAC	SEQ ID NO: 219
UAXXAGX	SEQ ID NO: 220
UAXXXGC	SEQ ID NO: 221
UXUGUXX	SEQ ID NO: 222
UXUXUAX	SEQ ID NO: 223
UXUXUXC	SEQ ID NO: 224
UXUAXXC	SEQ ID NO: 225
UXXXAGU	SEQ ID NO: 226
UXXXGGC	SEQ ID NO: 227
XAUGUXX	SEQ ID NO: 228
XXAUGGX	SEQ ID NO: 229
XXXUGGC	SEQ ID NO: 230
XXAXGGC	SEQ ID NO: 231
GCAAXXX	SEQ ID NO: 232
AUGXXXC	SEQ ID NO: 233
ACXXXGG	SEQ ID NO: 234
GAXXXUC	SEQ ID NO: 235
XXUXCUC	SEQ ID NO: 236
AGUAAGX	SEQ ID NO: 237
AGUGGGX	SEQ ID NO: 238
CAUGGCX	SEQ ID NO: 239
GUGUAUX	SEQ ID NO: 240
UACUCAX	SEQ ID NO: 241
UCGUCUX	SEQ ID NO: 242
ACAGUAX	SEQ ID NO: 243
UCCUCUX	SEQ ID NO: 244
GUGUGUA	SEQ ID NO: 245
GUGUGUU	SEQ ID NO: 246
GUGUGUG	SEQ ID NO: 247
GUGUGUC	SEQ ID NO: 248
UGUUUGC	SEQ ID NO: 249
UGUUUGG	SEQ ID NO: 250
UGUUUGA	SEQ ID NO: 251
UGUUUGU	SEQ ID NO: 252

X = C, G, U, or A, or other modified nucleobases as described elsewhere herein.

TABLE 13

Generic RNA oligonucleotide sequences with 7-mer seed sequences	
xUGXXXCXXXXXXXXXXXXXX	SEQ ID NO: 253
xUGXUXXXXXXXXXXXXXXX	SEQ ID NO: 254
xXXGAAXXXXXXXXXXXXXXX	SEQ ID NO: 255
xAGXXCXXXXXXXXXXXXXX	SEQ ID NO: 256
xXXUGUXXXXXXXXXXXXXXX	SEQ ID NO: 257
xUAXXXGXXXXXXXXXXXXXX	SEQ ID NO: 258
xUAUAXXXXXXXXXXXXXXX	SEQ ID NO: 259
xUAUGXXXXXXXXXXXXXX	SEQ ID NO: 260
xUAXGUXXXXXXXXXXXXXX	SEQ ID NO: 261
xUAXGXAXXXXXXXXXXXXX	SEQ ID NO: 262
xUAXXXACXXXXXXXXXXXX	SEQ ID NO: 263
xUAXXAGXXXXXXXXXXXX	SEQ ID NO: 264
xUAXXXGCXXXXXXXXXXXX	SEQ ID NO: 265
xUXUGUXXXXXXXXXXXXXXX	SEQ ID NO: 266
xUXUXUAXXXXXXXXXXXXXX	SEQ ID NO: 267
xUXUXUXCXXXXXXXXXXXX	SEQ ID NO: 268
xUXUAXXCXXXXXXXXXXXX	SEQ ID NO: 269
xUXXXAGUXXXXXXXXXXXX	SEQ ID NO: 270
xUXXXGGCXXXXXXXXXXXX	SEQ ID NO: 271
xXAUGUXXXXXXXXXXXXXX	SEQ ID NO: 272
xxxAUGGXXXXXXXXXXXXX	SEQ ID NO: 273
xxxXUGGCXXXXXXXXXXXX	SEQ ID NO: 274
xxxAXGGCXXXXXXXXXXXX	SEQ ID NO: 275
xGCAAXXXXXXXXXXXXXXX	SEQ ID NO: 276
xAUGXXXCXXXXXXXXXXXX	SEQ ID NO: 277
xACXXXGGXXXXXXXXXXXX	SEQ ID NO: 278
xGAXXXUCXXXXXXXXXXXX	SEQ ID NO: 279
xXXUXCUCXXXXXXXXXXXX	SEQ ID NO: 280
xAGUAAGXXXXXXXXXXXXXX	SEQ ID NO: 281
xAGUGGGXXXXXXXXXXXXX	SEQ ID NO: 282
xCAUGGCXXXXXXXXXXXXXX	SEQ ID NO: 283
xGUGUAUXXXXXXXXXXXXX	SEQ ID NO: 284
xUACUCAXXXXXXXXXXXXXX	SEQ ID NO: 285
xUCGUCUCXXXXXXXXXXXXXX	SEQ ID NO: 286
xACAGUAAXXXXXXXXXXXXX	SEQ ID NO: 287
xUCCUCUCXXXXXXXXXXXXXX	SEQ ID NO: 288
xGUGUGUAXXXXXXXXXXXXXX	SEQ ID NO: 289

TABLE 13 -continued

Generic RNA oligonucleotide sequences with 7-mer seed sequences	
xGUGUGUUXXXXXXXXXXXXXX	SEQ ID NO: 290
xGUGUGUGXXXXXXXXXXXXXX	SEQ ID NO: 291
xGUGUGUCXXXXXXXXXXXXXX	SEQ ID NO: 292
xUGUUUGCXXXXXXXXXXXXXX	SEQ ID NO: 293
xUGUUUGGXXXXXXXXXXXXXX	SEQ ID NO: 294
xUGUUUGAXXXXXXXXXXXXXX	SEQ ID NO: 295
xUGUUUGUXXXXXXXXXXXXXX	SEQ ID NO: 296

x or X = C, G, U, or A, or other modified nucleobases as described elsewhere herein.

### Discussion

[0145] As is the case for many malignancies, therapy resistance is a major cause of PCa-related morbidities and mortalities. Androgen deprivation therapy often gives rise to AR positive or AR negative CRPC, which represent two subclasses of resistant disease in which genetic, epigenetic, RNA splicing and other molecular aberrations contribute to either sustained androgen signaling or complete independence from the AR. These adaptive resistance mechanisms pose grave challenges to the effective treatment and management of this disease, underscoring the importance for the development of novel therapeutic agents and strategies.

[0146] Herein it is demonstrated that shRNAs targeting TMEFF2 inhibited androgen signaling in AR+PCa cell lines (both ADPC and CRPC) and elicited toxicity in transformed cells. RNA seq analysis with representative TMEFF2-targeted shRNAs suggests these phenotypes are induced by reduced essential gene expression via off target shRNA seed/3' UTR interactions, resembling a previously described process, Death Induced by Survival gene Elimination (DISE). Unique to this work, we showed that the 3' UTR of AR and AR coregulatory genes are also targets of TMEFF2 targeted shRNA seed interactions, leading to androgen signaling inhibition and further essential gene downregulation. Consequently, the inhibition of androgen signaling by DISE-inducing shRNAs correlated with toxicity in LNCaP cells compared to cells that do not depend on AR activity for growth and survival. We term this DISE-subtype, in which essential genes are eliminated in AR+PCa cells through direct off-target seed interactions and the depletion of the AR and androgen signaling coregulatory genes, androgen network DISE (AN-DISE) (FIG. 6). Unlike androgen deprivation therapy or other AR targeted therapies, AN-DISE, in addition to inhibiting androgen signaling, directly downregulates numerous non-androgen regulated essential genes, making developed resistance less likely. In support of this concept, DU145, an AR negative CRPC cell line, like other transformed cells, are also sensitive to DISE-inducing shRNAs. In addition to AN-DISE in PCa cells, other cancer cell types that are largely dependent on a single pathway for the expression of essential survival genes, such as estrogen receptor positive breast cancer, may also show increased sensitivity to the DISE mechanism if the central survival pathway is also compromised. Previous reports and our viability experiments with normal prostate epithelial cell

line, RWPE1, suggest that DISE preferentially occurs in transformed cells, further supporting therapeutic potential. Importantly, DISE-inducing siRNAs have already been shown to be effective in reducing tumor growth *in vivo* using an ovarian cancer mouse model, with little to no evident systemic toxicity.

**[0147]** Herein, also identified were at least three miRNAs, miR-4446-5p, miR-634 and miR6888-3p, that have identical seed sequence and many common predicted targets to three of the TMEFF2 shRNAs described herein. These data indicate that miR-4446-5p, miR-634, and miR6888-3p, have an effect on tumor suppressive miRNAs, especially in PCa.

**[0148]** In summary, in addition to corroborating previous studies describing DISE induction through shRNA seed-mediated off target effects, data from this work indicate that DISE-inducing shRNAs that inhibit AR transcriptional activity through the depletion of androgen signaling regulatory genes are particularly toxic to AR positive PCa cells via subsequent elimination of additional survival genes, demonstrating therapeutic use of these RNAi seeds in treatment of PCa.

**[0149]** Thus, in accordance with the present disclosure, there have been provided compositions, as well as methods of producing and using same, which fully satisfy the objectives and advantages set forth hereinabove. Although the present disclosure has been described in conjunction with specific formulas, compositions, and methods set forth hereinabove, it is evident that many alternatives, modifications, and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications, and variations that fall within the spirit and broad scope of the present disclosure.

**[0150]** In at least certain embodiments, the present disclosure is directed to the following non-limiting clauses.

**[0151]** Clause 1. A nucleic acid compound, comprising an oligonucleotide selected from the group consisting of short interfering RNA (siRNA), short hairpin RNA (shRNA), antisense RNA, DNA, antisense DNA, Chimeric Antisense DNA/RNA, and microRNA (miRNA), wherein the oligonucleotide has a 6-mer seed sequence that is complementary to a sequence of either a gene or an mRNA encoding an androgen receptor (AR) coregulator or a fragment thereof having AR coregulator activity; wherein the oligonucleotide comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO:36, SEQ ID NO:52, SEQ ID NO:35, SEQ ID NO:60, SEQ ID NO:51, SEQ ID NO:54, SEQ ID NO:44, SEQ ID NO:38, SEQ ID NO:37, SEQ ID NO:31, SEQ ID NO:50, SEQ ID NO:43, SEQ ID NO:49, SEQ ID NO:42, SEQ ID NO:40, SEQ ID NO:47, SEQ ID NO:39, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NOS:28-30, SEQ ID NOS:32-34, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:53, SEQ ID NOS:57-59, SEQ ID NOS:61-63, SEQ ID NOS:135-171, SEQ ID NOS:209-252, particularly including SEQ ID NO:249, SEQ ID NO:210, SEQ ID NOS:250-252, SEQ ID NO:254, SEQ ID NOS:293-296, SEQ ID NO:136, SEQ ID NO:173, SEQ ID NO:135, SEQ ID NOS:137-171, SEQ ID NO:209, and SEQ ID NOS:211-244, and nucleic acid sequences complementary to each of said sequences; and wherein the nucleic acid compound comprises at least one of (1) a non-natural modification in the oligonucleotide, and (2) an organic moiety conjugated to the oligonucleotide; and wherein the nucleic acid compound has inhibitory activity

against the expression or biological activity of the AR coregulator or biologically active fragment thereof.

**[0152]** Clause 2. The nucleic acid compound of clause 1, wherein the oligonucleotide has a strand length in a range of 12 to 50 nucleotides.

**[0153]** Clause 3. The nucleic acid compound of clause 1 or 2, wherein the at least one non-natural modification in the oligonucleotide is selected from the group consisting of a modified nucleoside sugar, a modified internucleoside linkage, and a modified nucleobase.

**[0154]** Clause 4. The nucleic acid compound of clause 3, wherein the modified nucleoside sugar comprises a 2'-modified sugar.

**[0155]** Clause 5. The nucleic acid compound of clause 4, wherein the 2'-modified sugar is selected from the group consisting of 2'-O-methyl-, 2'-deoxy-, 2'-fluoro-, 2'-deoxy-2'-fluoro-, 2'-O-methoxyethyl-(2'-O-MOE)-, 2'-O-amino-propyl-(2'-O-AP)-, 2'-O-dimethylaminoethyl-(2'-O-DMAOE)-, 2'-O-dimethylaminopropyl-(2'-O-DMAP)-, 2'-O-dimethylaminoethoxyethyl-(2'-O-DMAEOE)-, 2'-O-guanidinopropyl-, and 2'-O—N-methylacetamido-(2'-O-NMA)-modified sugars.

**[0156]** Clause 6. The nucleic acid compound of any one of clauses 3-5, wherein the modified internucleoside linkage is selected from the group consisting of phosphorothioate, phosphorodithioate, phosphoramidite, phosphorodiamidate, morpholino, phosphotriester, aminoalkylphosphotriester, phosphonate, alkyl phosphonate, 3'-alkylene phosphonate, 5'-alkylene phosphonate, chiral phosphonate, and phosphinate linkages.

**[0157]** Clause 7. The nucleic acid compound of any one of clauses 3-6, wherein the modified internucleoside linkage comprises at least the first, second, or third internucleoside linkage at the 5' and/or 3' end of the oligonucleotide.

**[0158]** Clause 8. The nucleic acid compound of any one of clauses 3-7, wherein the modified nucleobase is selected from the group consisting of 5-uracil (pseudouridine), dihydrouracil, 2-thiouracil, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, dihydrouracil, 5-trifluoromethyl uracil, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, 5-carboxymethylaminomethyl-2-thiouracil, 5-(carboxyhydroxymethyl) uracil, 5-methoxycarboxymethyluracil, 5-methoxyuracil, 5-methyl-2-thiouracil, 5-propynyl uracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, 5-carboxymethylaminomethyluracil, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, alkynyl uracil, 4-thiouracil, 6-azo uracil, 2-aminoadenine, 2-methyladenine, 6-methyladenine, 7-methyladenine, 2-F-adenine, 2-amino-adenine, 2-propyladenine, N6-adenine, N6-isopentenyladenine, 2-methylthio-N6-isopentenyladenine, alkynyl adenine, 8-fluoro-adenine, 8-chloro-adenine, 8-bromo-adenine, 8-ido-adenine, 8-amino adenine, 8-thiol adenine, 8-thioalkyl adenine, 8-hydroxyl adenine, 3-deazaadenine, 7-deazaadenine, 8-azaadenine, 3-methylcytosine, 5-methylcytosine, 5-trifluoromethyl cytosine, 2-thiocytosine, 5-hydroxymethyl cytosine, 5-propynyl cytosine, 6-azo cytosine, 4-acetylcytosine, 2-methylguanine, 6-methylguanine, 2-propylguanine, 1-methylguanine, 7-methylguanine, 2,2-dimethylguanine, 8-azaguanine, 7-deazaguanine, 8-fluoro-guanine, 8-chloro-guanine, 8-bromo-guanine, 8-ido-guanine, 8-amino guanine, 8-thiol guanine, 8-thioalkyl guanine, 8-hydroxyl guanine, 3-deazaguanine, 2-thiothymine, 6-azo thymine, alkynyl thymine,

ribothymine, inosine, hypoxanthine, xanthine, beta-D-galactosylqueosine, beta-D-mannosylqueosine, 1-methylinosine, 2,6-diaminopurine, queosine, tricyclic pyrimidines, phenoxazine cytidine(1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), and phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one).

[0159] Clause 9. The nucleic acid compound of any one of clauses 3-8, wherein the organic moiety is selected from the group consisting of lipids, peptides, receptor-specific ligands, aptamers, antibodies or antibody fragments, CpG-containing oligonucleotides, polyamines, polymers, dendrimers, saccharides, polysaccharides, and cyclodextrins.

[0160] Clause 10. The nucleic acid compound of any one of clauses 3-9, wherein the AR coregulator is selected from the group consisting of CCND1, ADAM10, APPBP2, ATAD2, BAG1, BRCA1, CALM1, CALR, CCND3, CDC25A, CDK2AP1, CTNNB1, CDK7, CTDSP2, COPSS5, CTDSP2, ENY2, EHMT2, FKBP4, FKBP5, GSK3B, HELZ2, HEY1, HIP1, HIPK3, IDE, IL6ST, KAT2B, MAPK1, MAPK15, MED1, MED21, MKRN1, NCOA2, PER1, PIAS1, PIK3CB, PIK3R1, PMEPA1, PRMT2, PSMC3IP, PQBP1, PRKDC, RANBP9, RANBP10, RPS6KA1, RPS6KA3, SMAD1, SMARCD1, TAF1, TPD52, UBE2L3, UXT, YWHAH, and ZMIZ1.

[0161] Clause 11. The nucleic acid compound of any one of clauses 3-10, wherein the AR coregulator is an AR coregulator of an aggressive prostate cancer.

[0162] Clause 12. An expression vector comprising the oligonucleotide of any one of clauses 3-11.

[0163] Clause 13. A composition, comprising at least one nucleic acid compound of any one of clauses 3-12, and a pharmaceutically-acceptable carrier.

[0164] Clause 14. The composition of clause 13, comprising a nanoparticle.

[0165] Clause 15. A method for treating prostate cancer in a subject in need of such treatment, comprising: administering to the subject a therapeutically effective amount of an oligonucleotide selected from the group consisting of short interfering RNA (siRNA), short hairpin RNA (shRNA), antisense RNA, DNA, antisense DNA, Chimeric Antisense DNA/RNA, and microRNA (miRNA), wherein the oligonucleotide has a 6-mer seed sequence that is complementary to a sequence of either a gene or an mRNA encoding an androgen receptor (AR) coregulator or a fragment thereof having AR coregulator activity; wherein the oligonucleotide comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO:36, SEQ ID NO:31, SEQ ID NO:35, SEQ ID NO:60, SEQ ID NO:51, SEQ ID NO:54, SEQ ID NO:44, SEQ ID NO:38, SEQ ID NO:37, SEQ ID NO:52, SEQ ID NO:50, SEQ ID NO:43, SEQ ID NO:49, SEQ ID NO:42, SEQ ID NO:40, SEQ ID NO:47, SEQ ID NO:39, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NOS:28-30, SEQ ID NOS:32-34, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:53, SEQ ID NOS:57-59, SEQ ID NOS:61-63, SEQ ID NOS:135-171, SEQ ID NOS:209-252, particularly including SEQ ID NO:249, SEQ ID NO:210, SEQ ID NOS:250-252, SEQ ID NO:254, SEQ ID NOS:293-296, SEQ ID NO:136, SEQ ID NO:173, SEQ ID NO:135, SEQ ID NOS:137-171, SEQ ID NO:209, and SEQ ID NOS:211-244, and nucleic acid sequences complementary to each of said sequences; and wherein the oligonucleotide has inhibitory activity against the expression or biological activity of the AR coregulator or biologically active fragment thereof.

[0166] Clause 16. The method of clause 15, wherein the prostate cancer is aggressive prostate cancer.

[0167] Clause 17. The method of clause 15 or 16, wherein the oligonucleotide comprises a portion of a nucleic acid compound or nucleic acid composition.

[0168] Clause 18. The method of any one of clauses 15-17, wherein the oligonucleotide comprises at least one of (1) a non-natural modification in the oligonucleotide, and (2) an organic moiety conjugated to the oligonucleotide.

[0169] Clause 19. The method of clause 18, wherein the at least one non-natural modification in the oligonucleotide is selected from the group consisting of a modified nucleoside sugar, a modified internucleoside linkage, and a modified nucleobase.

[0170] Clause 20. The method of clause 19, wherein the modified nucleoside sugar comprises a 2'-modified sugar.

[0171] Clause 21. The method of clause 20, wherein the 2'-modified sugar is selected from the group consisting of 2'-O-methyl-, 2'-deoxy-, 2'-fluoro-, 2'-deoxy-2'-fluoro-, 2'-O-methoxyethyl-(2'-O-MOE)-, 2'-O-aminopropyl-(2'-O-AP)-, 2'-O-dimethylaminooethyl-(2'-O-DMAOE)-, 2'-O-dimethylaminopropyl-(2'-O-DMAP)-, 2'-O-dimethylaminooethoxyethyl-(2'-O-DMAEOE)-, 2'-O-guanidinopropyl-, and 2'-O—N-methylacetamido-(2'-O-NMA)-modified sugars.

[0172] Clause 22. The method of any one of clauses 19-21, wherein the modified internucleoside linkage is selected from the group consisting of phosphorothioate, phosphordithioate, phosphoramidite, phosphorodiamidate, morpholino, phosphotriester, aminoalkylphosphotriester, phosphonate, alkyl phosphonate, 3'-alkylene phosphonate, 5'-alkylene phosphonate, chiral phosphonate, and phosphinate linkages.

[0173] Clause 23. The method of any one of clauses 19-22, wherein the modified internucleoside linkage comprises at least the first, second, or third internucleoside linkage at the 5' and/or 3' end of the oligonucleotide.

[0174] Clause 24. The method of any one of clauses 19-23, wherein the modified nucleobase is selected from the group consisting of 5-uracil (pseudouridine), dihydrouracil, 2-thiouracil, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, dihydrouracil, 5-trifluoromethyl uracil, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, 5-carboxymethylaminomethyl-2-thiouracil, 5-(carboxyhydroxymethyl) uracil, 5-methoxycarboxymethyluracil, 5-methoxyuracil, 5-methyl-2-thiouracil, 5-propynyl uracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, 5-carboxymethylaminomethyluracil, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, alkynyl uracil, 4-thiouracil, 6-azo uracil, 2-aminoadenine, 2-methyladenine, 6-methyladenine, 7-methyladenine, 2-F-adenine, 2-amino-adenine, 2-propyladenine, N6-adenine, N6-isopentenyladenine, 2-methylthio-N6-isopentenyladenine, alkynyl adenine, 8-fluoro-adenine, 8-chloro-adenine, 8-bromo-adenine, 8-iodo-adenine, 8-amino adenine, 8-thiol adenine, 8-thioalkyl adenine, 8-hydroxyl adenine, 3-deazaadenine, 7-deazaadenine, 8-azaadenine, 3-methylcytosine, 5-methylcytosine, 5-trifluoromethyl cytosine, 5-propynyl cytosine, 6-azo cytosine, 4-acetylcytosine, 2-methylguanine, 6-methylguanine, 2-propylguanine, 1-methylguanine, 7-methylguanine, 2,2-dimethylguanine, 8-azaguanine, 7-deazaguanine, 8-fluoro-gua-

nine, 8-chloro-guanine, 8-bromo-guanine, 8-ido-guanine, 8-amino guanine, 8-thiol guanine, 8-thioalkyl guanine, 8-hydroxyl guanine, 3-deazaguanine, 2-thiothymine, 6-azo thymine, alkynyl thymine, ribothymine, inosine, hypoxanthine, xanthine, beta-D-galactosylqueosine, beta-D-mannosylqueosine, 1-methylinosine, 2,6-diaminopurine, queosine, tricyclic pyrimidines, phenoxazine cytidine(1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), and phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one).

[0175] Clause 25. The method of any one of clauses 18-24, wherein the organic moiety is selected from the group consisting of lipids, peptides, receptor-specific ligands, aptamers, antibodies or antibody fragments, CpG-containing oligonucleotides, polyamines, polymers, dendrimers, saccharides, polysaccharides, and cyclodextrins.

[0176] Clause 26. The method of any one of clauses 19-25, wherein the AR coregulator is selected from the group consisting of CCND1, ADAM10, APPBP2, ATAD2, BAG1, BRCA1, CALM1, CALR, CCND3, CDC25A, CDK2AP1, CTNNB1, CDK7, CTDSP2, COPS5, CTDSP2, ENY2, EHMT2, FKBP4, FKBP5, GSK3B, HELZ2, HEY1, HIP1, HIPK3, IDE, IL6ST, KAT2B, MAPK1, MAPK15, MED1, MED21, MKRN1, NCOA2, PER1, PIAS1, PIK3CB, PIK3R1, PMEPA1, PRMT2, PSMC3IP, PQBP1, PRKDC, RANBP9, RANBP10, RPS6KA1, RPS6KA3, SMAD1, SMARCD1, TAF1, TPD52, UBE2L3, UXT, YWHAH, and ZMIZ1.

[0177] Clause 27. The method of any one of clauses 19-26, wherein the AR coregulator is an AR coregulator of an aggressive prostate cancer.

---

#### SEQUENCE LISTING

```

Sequence total quantity: 296
SEQ ID NO: 1          moltype = RNA  length = 21
FEATURE
modified_base          Location/Qualifiers
2
mod_base = OTHER
note = thymine
5..6
mod_base = OTHER
note = thymine
8
mod_base = OTHER
note = thymine
11
mod_base = OTHER
note = thymine
1..21
note = RNAi consortium shRNA library - TRCN0000073522
source
mol_type = other RNA
organism = synthetic construct

SEQUENCE: 1
ctggtatga tgacagaga a                                21

SEQ ID NO: 2          moltype = RNA  length = 21
FEATURE
modified_base          Location/Qualifiers
3
mod_base = OTHER
note = thymine
5
mod_base = OTHER
note = thymine
7
mod_base = OTHER
note = thymine
11..12
mod_base = OTHER
note = thymine
17
mod_base = OTHER
note = thymine
1..21
note = RNAi consortium shRNA library - TRCN0000222559
source
mol_type = other RNA
organism = synthetic construct

SEQUENCE: 2
cgtctgtcag ttcaagtgc a                                21

SEQ ID NO: 3          moltype = RNA  length = 21
FEATURE
modified_base          Location/Qualifiers
5..6
mod_base = OTHER
note = thymine
8
mod_base = OTHER

```

---

-continued

---

```

modified_base      note = thymine
                   11
                   mod_base = OTHER
modified_base      note = thymine
                   18
                   mod_base = OTHER
modified_base      note = thymine
                   20..21
                   mod_base = OTHER
note = thymine
1..21
note = RNAi consortium shRNA library - TRCN0000073521
source            1..21
mol_type = other RNA
organism = synthetic construct

SEQUENCE: 3
gcgcttctga tggaaatct t                                21

SEQ ID NO: 4      moltype = RNA  length = 21
FEATURE          Location/Qualifiers
modified_base     6
mod_base = OTHER
note = thymine
modified_base     8
mod_base = OTHER
note = thymine
modified_base     11
mod_base = OTHER
note = thymine
modified_base     14
mod_base = OTHER
note = thymine
modified_base     17..18
mod_base = OTHER
note = thymine
modified_base     20
mod_base = OTHER
note = thymine
note = RNAi consortium shRNA library - TRCN0000073520
source            1..21
mol_type = other RNA
organism = synthetic construct

SEQUENCE: 4
gcaggtgtga tgctggttat a                                21

SEQ ID NO: 5      moltype = RNA  length = 21
FEATURE          Location/Qualifiers
modified_base     3..4
mod_base = OTHER
note = thymine
modified_base     8..10
mod_base = OTHER
note = thymine
modified_base     12
mod_base = OTHER
note = thymine
modified_base     15
mod_base = OTHER
note = thymine
modified_base     18
mod_base = OTHER
note = thymine
modified_base     20
mod_base = OTHER
note = thymine
note = RNAi consortium shRNA library - TRCN0000073518
source            1..21
mol_type = other RNA
organism = synthetic construct

SEQUENCE: 5
ccttgcattt gtggtaatct a                                21

SEQ ID NO: 6      moltype = RNA  length = 21
FEATURE          Location/Qualifiers

```

---

-continued

---

```

modified_base      4
                  mod_base = OTHER
                  note = thymine
modified_base      6
                  mod_base = OTHER
                  note = thymine
modified_base      15
                  mod_base = OTHER
                  note = thymine
modified_base      18
                  mod_base = OTHER
                  note = thymine
misc_feature       1..21
                  note = Completely synthesized in the lab.
source            1..21
                  mol_type = other RNA
                  organism = synthetic construct
SEQUENCE: 6
ggctctggag aaactagtca a                                21

SEQ ID NO: 7      moltype = RNA length = 21
FEATURE          Location/Qualifiers
modified_base     2
                  mod_base = OTHER
                  note = thymine
modified_base     11
                  mod_base = OTHER
                  note = thymine
modified_base     14
                  mod_base = OTHER
                  note = thymine
modified_base     21
                  mod_base = OTHER
                  note = thymine
misc_feature      1..21
                  note = RNAi consortium shRNA library - TRCN0000373776
source            1..21
                  mol_type = other RNA
                  organism = synthetic construct
SEQUENCE: 7
atgcagagaa tgctaacaaa t                                21

SEQ ID NO: 8      moltype = RNA length = 21
FEATURE          Location/Qualifiers
modified_base     3
                  mod_base = OTHER
                  note = thymine
modified_base     7..8
                  mod_base = OTHER
                  note = thymine
modified_base     10
                  mod_base = OTHER
                  note = thymine
modified_base     19..20
                  mod_base = OTHER
                  note = thymine
misc_feature      1..21
                  note = RNAi consortium shRNA library - TRCN0000373700
source            1..21
                  mol_type = other RNA
                  organism = synthetic construct
SEQUENCE: 8
catacattgt ccggaacatt a                                21

SEQ ID NO: 9      moltype = RNA length = 21
FEATURE          Location/Qualifiers
modified_base     6
                  mod_base = OTHER
                  note = thymine
modified_base     11..12
                  mod_base = OTHER
                  note = thymine
modified_base     20
                  mod_base = OTHER
                  note = thymine
misc_feature      1..21

```

---

-continued

---

```

source          note = Completely synthesized in the lab.
                1..21
                mol_type = other RNA
                organism = synthetic construct

SEQUENCE: 9
ggcactacag ttcagacaat a                                21

SEQ ID NO: 10      moltype = RNA  length = 21
FEATURE          Location/Qualifiers
modified_base    3
                  mod_base = OTHER
                  note = thymine
modified_base    8..9
                  mod_base = OTHER
                  note = thymine
modified_base    13
                  mod_base = OTHER
                  note = thymine
modified_base    19
                  mod_base = OTHER
                  note = thymine
misc_feature     1..21
                  note = Completely synthesized in the lab.
source          1..21
                mol_type = other RNA
                organism = synthetic construct

SEQUENCE: 10
cctaagggtta agtcgcccctc g                                21

SEQ ID NO: 11      moltype = RNA  length = 21
FEATURE          Location/Qualifiers
modified_base    6
                  mod_base = OTHER
                  note = thymine
modified_base    20..21
                  mod_base = OTHER
                  note = thymine
misc_feature     1..21
                  note = Completely synthesized in the lab.
source          1..21
                mol_type = other RNA
                organism = synthetic construct

SEQUENCE: 11
cgacgtaaac ggccacaagt t                                21

SEQ ID NO: 12      moltype = RNA  length = 21
FEATURE          Location/Qualifiers
modified_base    3
                  mod_base = OTHER
                  note = thymine
modified_base    8
                  mod_base = OTHER
                  note = thymine
modified_base    10
                  mod_base = OTHER
                  note = thymine
modified_base    13..15
                  mod_base = OTHER
                  note = thymine
modified_base    17
                  mod_base = OTHER
                  note = thymine
modified_base    19
                  mod_base = OTHER
                  note = thymine
modified_base    21
                  mod_base = OTHER
                  note = thymine
misc_feature     1..21
                  note = RNAi consortium shRNA library - TRCN0000059000.
source          1..21
                mol_type = other RNA
                organism = synthetic construct

SEQUENCE: 12
actgggctgt actttgtata t                                21

```

---

-continued

---

```

SEQ ID NO: 13      moltype = RNA  length = 21
FEATURE          Location/Qualifiers
modified_base     3
                  mod_base = OTHER
                  note = thymine
modified_base     11
                  mod_base = OTHER
                  note = thymine
modified_base     17
                  mod_base = OTHER
                  note = thymine
modified_base     21
                  mod_base = OTHER
                  note = thymine
1..21
note = RNAi consortium shRNA library - TRCN0000038696.
source           1..21
                  mol_type = other RNA
                  organism = synthetic construct

SEQUENCE: 13
cctgaaacag tggcaataaa t                                21

SEQ ID NO: 14      moltype = RNA  length = 20
FEATURE          Location/Qualifiers
modified_base     5
                  mod_base = OTHER
                  note = thymine
modified_base     10
                  mod_base = OTHER
                  note = thymine
modified_base     17
                  mod_base = OTHER
                  note = thymine
1..20
note = Completely synthesized in the lab.
source           1..20
                  mol_type = other RNA
                  organism = synthetic construct

SEQUENCE: 14
aagatcgagt gccgcacatcac                                20

SEQ ID NO: 15      moltype = RNA  length = 20
FEATURE          Location/Qualifiers
modified_base     6
                  mod_base = OTHER
                  note = thymine
modified_base     9
                  mod_base = OTHER
                  note = thymine
modified_base    14
                  mod_base = OTHER
                  note = thymine
1..20
note = Completely synthesized in the lab.
source           1..20
                  mol_type = other RNA
                  organism = synthetic construct

SEQUENCE: 15
ccagctgctg cactgccgcg                                20

SEQ ID NO: 16      moltype = RNA  length = 19
FEATURE          Location/Qualifiers
misc_feature     1..19
                  note = Completely synthesized in the lab.
source           1..19
                  mol_type = other RNA
                  organism = synthetic construct

SEQUENCE: 16
ctctgtcatc ataaccaga                                19

SEQ ID NO: 17      moltype = RNA  length = 19
FEATURE          Location/Qualifiers
misc_feature     1..19
                  note = Completely synthesized in the lab.
source           1..19
                  mol_type = other RNA

```

---

-continued

---

SEQUENCE: 17	organism = synthetic construct
cacttgaact gacagacgc	19
SEQ ID NO: 18	moltype = RNA length = 19
FEATURE	Location/Qualifiers
misc_feature	1..19
source	note = Completely synthesized in the lab.
	1..19
	mol_type = other RNA
	organism = synthetic construct
SEQUENCE: 18	
atttccatc agaagcgca	19
SEQ ID NO: 19	moltype = RNA length = 19
FEATURE	Location/Qualifiers
misc_feature	1..19
source	note = Completely synthesized in the lab.
	1..19
	mol_type = other RNA
	organism = synthetic construct
SEQUENCE: 19	
aaccaggatc acacctgca	19
SEQ ID NO: 20	moltype = RNA length = 19
FEATURE	Location/Qualifiers
misc_feature	1..19
source	note = Completely synthesized in the lab.
	1..19
	mol_type = other RNA
	organism = synthetic construct
SEQUENCE: 20	
attaccacaa atgcaaggc	19
SEQ ID NO: 21	moltype = RNA length = 19
FEATURE	Location/Qualifiers
misc_feature	1..19
source	note = Completely synthesized in the lab.
	1..19
	mol_type = other RNA
	organism = synthetic construct
SEQUENCE: 21	
actagttct ccagagcct	19
SEQ ID NO: 22	moltype = RNA length = 19
FEATURE	Location/Qualifiers
misc_feature	1..19
source	note = Completely synthesized in the lab.
	1..19
	mol_type = other RNA
	organism = synthetic construct
SEQUENCE: 22	
tgttagatt ctctgcata	19
SEQ ID NO: 23	moltype = RNA length = 19
FEATURE	Location/Qualifiers
misc_feature	1..19
source	note = Completely synthesized in the lab.
	1..19
	mol_type = other RNA
	organism = synthetic construct
SEQUENCE: 23	
tgttccggac aaggtatgt	19
SEQ ID NO: 24	moltype = RNA length = 19
FEATURE	Location/Qualifiers
misc_feature	1..19
source	note = Completely synthesized in the lab.
	1..19
	mol_type = other RNA
	organism = synthetic construct
SEQUENCE: 24	
tgtctgaact gtagtgccc	19
SEQ ID NO: 25	moltype = RNA length = 22
FEATURE	Location/Qualifiers

---

-continued

---

```

misc_feature          1..22
                      note = Completely synthesized in the lab.
source               1..22
                      mol_type = other RNA
                      organism = synthetic construct
SEQUENCE: 25
aaccaggcacc ccaactttgg ac                                22

SEQ ID NO: 26      moltype = RNA  length = 22
FEATURE           Location/Qualifiers
misc_feature       1..22
                      note = Completely synthesized in the lab.
source             1..22
                      mol_type = other RNA
                      organism = synthetic construct
SEQUENCE: 26
atttccctgc cattcccttg gc                                22

SEQ ID NO: 27      moltype = RNA  length = 21
FEATURE           Location/Qualifiers
misc_feature       1..21
                      note = Completely synthesized in the lab.
source             1..21
                      mol_type = other RNA
                      organism = synthetic construct
SEQUENCE: 27
atctgtctcg attgtttcca g                                 21

SEQ ID NO: 28      moltype =   length =
SEQUENCE: 28
000

SEQ ID NO: 29      moltype =   length =
SEQUENCE: 29
000

SEQ ID NO: 30      moltype =   length =
SEQUENCE: 30
000

SEQ ID NO: 31      moltype =   length =
SEQUENCE: 31
000

SEQ ID NO: 32      moltype =   length =
SEQUENCE: 32
000

SEQ ID NO: 33      moltype =   length =
SEQUENCE: 33
000

SEQ ID NO: 34      moltype =   length =
SEQUENCE: 34
000

SEQ ID NO: 35      moltype =   length =
SEQUENCE: 35
000

SEQ ID NO: 36      moltype =   length =
SEQUENCE: 36
000

SEQ ID NO: 37      moltype =   length =
SEQUENCE: 37
000

SEQ ID NO: 38      moltype =   length =
SEQUENCE: 38
000

SEQ ID NO: 39      moltype =   length =
SEQUENCE: 39
000

```

-continued

---

SEQ ID NO: 40	moltype =	length =
SEQUENCE: 40		
000		
SEQ ID NO: 41	moltype =	length =
SEQUENCE: 41		
000		
SEQ ID NO: 42	moltype =	length =
SEQUENCE: 42		
000		
SEQ ID NO: 43	moltype =	length =
SEQUENCE: 43		
000		
SEQ ID NO: 44	moltype =	length =
SEQUENCE: 44		
000		
SEQ ID NO: 45	moltype =	length =
SEQUENCE: 45		
000		
SEQ ID NO: 46	moltype =	length =
SEQUENCE: 46		
000		
SEQ ID NO: 47	moltype =	length =
SEQUENCE: 47		
000		
SEQ ID NO: 48	moltype =	length =
SEQUENCE: 48		
000		
SEQ ID NO: 49	moltype =	length =
SEQUENCE: 49		
000		
SEQ ID NO: 50	moltype =	length =
SEQUENCE: 50		
000		
SEQ ID NO: 51	moltype =	length =
SEQUENCE: 51		
000		
SEQ ID NO: 52	moltype =	length =
SEQUENCE: 52		
000		
SEQ ID NO: 53	moltype =	length =
SEQUENCE: 53		
000		
SEQ ID NO: 54	moltype =	length =
SEQUENCE: 54		
000		
SEQ ID NO: 55	moltype =	length =
SEQUENCE: 55		
000		
SEQ ID NO: 56	moltype =	length =
SEQUENCE: 56		
000		
SEQ ID NO: 57	moltype =	length =
SEQUENCE: 57		
000		
SEQ ID NO: 58	moltype =	length =
SEQUENCE: 58		
000		

-continued

---

SEQ ID NO: 59	moltype =	length =
SEQUENCE: 59		
000		
SEQ ID NO: 60	moltype =	length =
SEQUENCE: 60		
000		
SEQ ID NO: 61	moltype =	length =
SEQUENCE: 61		
000		
SEQ ID NO: 62	moltype =	length =
SEQUENCE: 62		
000		
SEQ ID NO: 63	moltype =	length =
SEQUENCE: 63		
000		
SEQ ID NO: 64	moltype =	length =
SEQUENCE: 64		
000		
SEQ ID NO: 65	moltype =	length =
SEQUENCE: 65		
000		
SEQ ID NO: 66	moltype =	length =
SEQUENCE: 66		
000		
SEQ ID NO: 67	moltype =	length =
SEQUENCE: 67		
000		
SEQ ID NO: 68	moltype =	length =
SEQUENCE: 68		
000		
SEQ ID NO: 69	moltype =	length =
SEQUENCE: 69		
000		
SEQ ID NO: 70	moltype =	length =
SEQUENCE: 70		
000		
SEQ ID NO: 71	moltype =	length =
SEQUENCE: 71		
000		
SEQ ID NO: 72	moltype =	length =
SEQUENCE: 72		
000		
SEQ ID NO: 73	moltype =	length =
SEQUENCE: 73		
000		
SEQ ID NO: 74	moltype =	length =
SEQUENCE: 74		
000		
SEQ ID NO: 75	moltype =	length =
SEQUENCE: 75		
000		
SEQ ID NO: 76	moltype =	length =
SEQUENCE: 76		
000		
SEQ ID NO: 77	moltype =	length =
SEQUENCE: 77		
000		

-continued

---

SEQ ID NO: 78	moltype =	length =
SEQUENCE: 78		
000		
SEQ ID NO: 79	moltype =	length =
SEQUENCE: 79		
000		
SEQ ID NO: 80	moltype =	length =
SEQUENCE: 80		
000		
SEQ ID NO: 81	moltype =	length =
SEQUENCE: 81		
000		
SEQ ID NO: 82	moltype =	length =
SEQUENCE: 82		
000		
SEQ ID NO: 83	moltype =	length =
SEQUENCE: 83		
000		
SEQ ID NO: 84	moltype =	length =
SEQUENCE: 84		
000		
SEQ ID NO: 85	moltype =	length =
SEQUENCE: 85		
000		
SEQ ID NO: 86	moltype =	length =
SEQUENCE: 86		
000		
SEQ ID NO: 87	moltype =	length =
SEQUENCE: 87		
000		
SEQ ID NO: 88	moltype =	length =
SEQUENCE: 88		
000		
SEQ ID NO: 89	moltype =	length =
SEQUENCE: 89		
000		
SEQ ID NO: 90	moltype =	length =
SEQUENCE: 90		
000		
SEQ ID NO: 91	moltype =	length =
SEQUENCE: 91		
000		
SEQ ID NO: 92	moltype =	length =
SEQUENCE: 92		
000		
SEQ ID NO: 93	moltype =	length =
SEQUENCE: 93		
000		
SEQ ID NO: 94	moltype =	length =
SEQUENCE: 94		
000		
SEQ ID NO: 95	moltype =	length =
SEQUENCE: 95		
000		
SEQ ID NO: 96	moltype =	length =
SEQUENCE: 96		
000		

-continued

---

SEQ ID NO: 97	moltype =	length =
SEQUENCE: 97		
000		
SEQ ID NO: 98	moltype =	length =
SEQUENCE: 98		
000		
SEQ ID NO: 99	moltype =	length =
SEQUENCE: 99		
000		
SEQ ID NO: 100	moltype =	length =
SEQUENCE: 100		
000		
SEQ ID NO: 101	moltype =	length =
SEQUENCE: 101		
000		
SEQ ID NO: 102	moltype =	length =
SEQUENCE: 102		
000		
SEQ ID NO: 103	moltype =	length =
SEQUENCE: 103		
000		
SEQ ID NO: 104	moltype =	length =
SEQUENCE: 104		
000		
SEQ ID NO: 105	moltype =	length =
SEQUENCE: 105		
000		
SEQ ID NO: 106	moltype =	length =
SEQUENCE: 106		
000		
SEQ ID NO: 107	moltype =	length =
SEQUENCE: 107		
000		
SEQ ID NO: 108	moltype =	length =
SEQUENCE: 108		
000		
SEQ ID NO: 109	moltype =	length =
SEQUENCE: 109		
000		
SEQ ID NO: 110	moltype =	length =
SEQUENCE: 110		
000		
SEQ ID NO: 111	moltype =	length =
SEQUENCE: 111		
000		
SEQ ID NO: 112	moltype =	length =
SEQUENCE: 112		
000		
SEQ ID NO: 113	moltype =	length =
SEQUENCE: 113		
000		
SEQ ID NO: 114	moltype =	length =
SEQUENCE: 114		
000		
SEQ ID NO: 115	moltype =	length =
SEQUENCE: 115		
000		

-continued

---

SEQ ID NO: 116	moltype =	length =
SEQUENCE: 116		
000		
SEQ ID NO: 117	moltype =	length =
SEQUENCE: 117		
000		
SEQ ID NO: 118	moltype =	length =
SEQUENCE: 118		
000		
SEQ ID NO: 119	moltype =	length =
SEQUENCE: 119		
000		
SEQ ID NO: 120	moltype =	length =
SEQUENCE: 120		
000		
SEQ ID NO: 121	moltype =	length =
SEQUENCE: 121		
000		
SEQ ID NO: 122	moltype =	length =
SEQUENCE: 122		
000		
SEQ ID NO: 123	moltype =	length =
SEQUENCE: 123		
000		
SEQ ID NO: 124	moltype =	length =
SEQUENCE: 124		
000		
SEQ ID NO: 125	moltype =	length =
SEQUENCE: 125		
000		
SEQ ID NO: 126	moltype =	length =
SEQUENCE: 126		
000		
SEQ ID NO: 127	moltype =	length =
SEQUENCE: 127		
000		
SEQ ID NO: 128	moltype =	length =
SEQUENCE: 128		
000		
SEQ ID NO: 129	moltype =	length =
SEQUENCE: 129		
000		
SEQ ID NO: 130	moltype =	length =
SEQUENCE: 130		
000		
SEQ ID NO: 131	moltype =	length =
SEQUENCE: 131		
000		
SEQ ID NO: 132	moltype =	length =
SEQUENCE: 132		
000		
SEQ ID NO: 133	moltype =	length =
SEQUENCE: 133		
000		
SEQ ID NO: 134	moltype =	length =
SEQUENCE: 134		
000		

-continued

---

SEQ ID NO: 135	moltype =	length =
SEQUENCE: 135		
000		
SEQ ID NO: 136	moltype =	length =
SEQUENCE: 136		
000		
SEQ ID NO: 137	moltype =	length =
SEQUENCE: 137		
000		
SEQ ID NO: 138	moltype =	length =
SEQUENCE: 138		
000		
SEQ ID NO: 139	moltype =	length =
SEQUENCE: 139		
000		
SEQ ID NO: 140	moltype =	length =
SEQUENCE: 140		
000		
SEQ ID NO: 141	moltype =	length =
SEQUENCE: 141		
000		
SEQ ID NO: 142	moltype =	length =
SEQUENCE: 142		
000		
SEQ ID NO: 143	moltype =	length =
SEQUENCE: 143		
000		
SEQ ID NO: 144	moltype =	length =
SEQUENCE: 144		
000		
SEQ ID NO: 145	moltype =	length =
SEQUENCE: 145		
000		
SEQ ID NO: 146	moltype =	length =
SEQUENCE: 146		
000		
SEQ ID NO: 147	moltype =	length =
SEQUENCE: 147		
000		
SEQ ID NO: 148	moltype =	length =
SEQUENCE: 148		
000		
SEQ ID NO: 149	moltype =	length =
SEQUENCE: 149		
000		
SEQ ID NO: 150	moltype =	length =
SEQUENCE: 150		
000		
SEQ ID NO: 151	moltype =	length =
SEQUENCE: 151		
000		
SEQ ID NO: 152	moltype =	length =
SEQUENCE: 152		
000		
SEQ ID NO: 153	moltype =	length =
SEQUENCE: 153		
000		

-continued

---

SEQ ID NO: 154	moltype =	length =
SEQUENCE: 154		
000		
SEQ ID NO: 155	moltype =	length =
SEQUENCE: 155		
000		
SEQ ID NO: 156	moltype =	length =
SEQUENCE: 156		
000		
SEQ ID NO: 157	moltype =	length =
SEQUENCE: 157		
000		
SEQ ID NO: 158	moltype =	length =
SEQUENCE: 158		
000		
SEQ ID NO: 159	moltype =	length =
SEQUENCE: 159		
000		
SEQ ID NO: 160	moltype =	length =
SEQUENCE: 160		
000		
SEQ ID NO: 161	moltype =	length =
SEQUENCE: 161		
000		
SEQ ID NO: 162	moltype =	length =
SEQUENCE: 162		
000		
SEQ ID NO: 163	moltype =	length =
SEQUENCE: 163		
000		
SEQ ID NO: 164	moltype =	length =
SEQUENCE: 164		
000		
SEQ ID NO: 165	moltype =	length =
SEQUENCE: 165		
000		
SEQ ID NO: 166	moltype =	length =
SEQUENCE: 166		
000		
SEQ ID NO: 167	moltype =	length =
SEQUENCE: 167		
000		
SEQ ID NO: 168	moltype =	length =
SEQUENCE: 168		
000		
SEQ ID NO: 169	moltype =	length =
SEQUENCE: 169		
000		
SEQ ID NO: 170	moltype =	length =
SEQUENCE: 170		
000		
SEQ ID NO: 171	moltype =	length =
SEQUENCE: 171		
000		
SEQ ID NO: 172	moltype =	length =
SEQUENCE: 172		
000		

-continued

---

SEQ ID NO: 173	moltype =	length =
SEQUENCE: 173		
000		
SEQ ID NO: 174	moltype =	length =
SEQUENCE: 174		
000		
SEQ ID NO: 175	moltype =	length =
SEQUENCE: 175		
000		
SEQ ID NO: 176	moltype =	length =
SEQUENCE: 176		
000		
SEQ ID NO: 177	moltype =	length =
SEQUENCE: 177		
000		
SEQ ID NO: 178	moltype =	length =
SEQUENCE: 178		
000		
SEQ ID NO: 179	moltype =	length =
SEQUENCE: 179		
000		
SEQ ID NO: 180	moltype =	length =
SEQUENCE: 180		
000		
SEQ ID NO: 181	moltype =	length =
SEQUENCE: 181		
000		
SEQ ID NO: 182	moltype =	length =
SEQUENCE: 182		
000		
SEQ ID NO: 183	moltype =	length =
SEQUENCE: 183		
000		
SEQ ID NO: 184	moltype =	length =
SEQUENCE: 184		
000		
SEQ ID NO: 185	moltype =	length =
SEQUENCE: 185		
000		
SEQ ID NO: 186	moltype =	length =
SEQUENCE: 186		
000		
SEQ ID NO: 187	moltype =	length =
SEQUENCE: 187		
000		
SEQ ID NO: 188	moltype =	length =
SEQUENCE: 188		
000		
SEQ ID NO: 189	moltype =	length =
SEQUENCE: 189		
000		
SEQ ID NO: 190	moltype =	length =
SEQUENCE: 190		
000		
SEQ ID NO: 191	moltype =	length =
SEQUENCE: 191		
000		

-continued

---

SEQ ID NO: 192	moltype =	length =
SEQUENCE: 192		
000		
SEQ ID NO: 193	moltype =	length =
SEQUENCE: 193		
000		
SEQ ID NO: 194	moltype =	length =
SEQUENCE: 194		
000		
SEQ ID NO: 195	moltype =	length =
SEQUENCE: 195		
000		
SEQ ID NO: 196	moltype =	length =
SEQUENCE: 196		
000		
SEQ ID NO: 197	moltype =	length =
SEQUENCE: 197		
000		
SEQ ID NO: 198	moltype =	length =
SEQUENCE: 198		
000		
SEQ ID NO: 199	moltype =	length =
SEQUENCE: 199		
000		
SEQ ID NO: 200	moltype =	length =
SEQUENCE: 200		
000		
SEQ ID NO: 201	moltype =	length =
SEQUENCE: 201		
000		
SEQ ID NO: 202	moltype =	length =
SEQUENCE: 202		
000		
SEQ ID NO: 203	moltype =	length =
SEQUENCE: 203		
000		
SEQ ID NO: 204	moltype =	length =
SEQUENCE: 204		
000		
SEQ ID NO: 205	moltype =	length =
SEQUENCE: 205		
000		
SEQ ID NO: 206	moltype =	length =
SEQUENCE: 206		
000		
SEQ ID NO: 207	moltype =	length =
SEQUENCE: 207		
000		
SEQ ID NO: 208	moltype =	length =
SEQUENCE: 208		
000		
SEQ ID NO: 209	moltype =	length =
SEQUENCE: 209		
000		
SEQ ID NO: 210	moltype =	length =
SEQUENCE: 210		
000		

-continued

---

SEQ ID NO: 211	moltype =	length =
SEQUENCE: 211		
000		
SEQ ID NO: 212	moltype =	length =
SEQUENCE: 212		
000		
SEQ ID NO: 213	moltype =	length =
SEQUENCE: 213		
000		
SEQ ID NO: 214	moltype =	length =
SEQUENCE: 214		
000		
SEQ ID NO: 215	moltype =	length =
SEQUENCE: 215		
000		
SEQ ID NO: 216	moltype =	length =
SEQUENCE: 216		
000		
SEQ ID NO: 217	moltype =	length =
SEQUENCE: 217		
000		
SEQ ID NO: 218	moltype =	length =
SEQUENCE: 218		
000		
SEQ ID NO: 219	moltype =	length =
SEQUENCE: 219		
000		
SEQ ID NO: 220	moltype =	length =
SEQUENCE: 220		
000		
SEQ ID NO: 221	moltype =	length =
SEQUENCE: 221		
000		
SEQ ID NO: 222	moltype =	length =
SEQUENCE: 222		
000		
SEQ ID NO: 223	moltype =	length =
SEQUENCE: 223		
000		
SEQ ID NO: 224	moltype =	length =
SEQUENCE: 224		
000		
SEQ ID NO: 225	moltype =	length =
SEQUENCE: 225		
000		
SEQ ID NO: 226	moltype =	length =
SEQUENCE: 226		
000		
SEQ ID NO: 227	moltype =	length =
SEQUENCE: 227		
000		
SEQ ID NO: 228	moltype =	length =
SEQUENCE: 228		
000		
SEQ ID NO: 229	moltype =	length =
SEQUENCE: 229		
000		

-continued

---

SEQ ID NO: 230	moltype =	length =
SEQUENCE: 230		
000		
SEQ ID NO: 231	moltype =	length =
SEQUENCE: 231		
000		
SEQ ID NO: 232	moltype =	length =
SEQUENCE: 232		
000		
SEQ ID NO: 233	moltype =	length =
SEQUENCE: 233		
000		
SEQ ID NO: 234	moltype =	length =
SEQUENCE: 234		
000		
SEQ ID NO: 235	moltype =	length =
SEQUENCE: 235		
000		
SEQ ID NO: 236	moltype =	length =
SEQUENCE: 236		
000		
SEQ ID NO: 237	moltype =	length =
SEQUENCE: 237		
000		
SEQ ID NO: 238	moltype =	length =
SEQUENCE: 238		
000		
SEQ ID NO: 239	moltype =	length =
SEQUENCE: 239		
000		
SEQ ID NO: 240	moltype =	length =
SEQUENCE: 240		
000		
SEQ ID NO: 241	moltype =	length =
SEQUENCE: 241		
000		
SEQ ID NO: 242	moltype =	length =
SEQUENCE: 242		
000		
SEQ ID NO: 243	moltype =	length =
SEQUENCE: 243		
000		
SEQ ID NO: 244	moltype =	length =
SEQUENCE: 244		
000		
SEQ ID NO: 245	moltype =	length =
SEQUENCE: 245		
000		
SEQ ID NO: 246	moltype =	length =
SEQUENCE: 246		
000		
SEQ ID NO: 247	moltype =	length =
SEQUENCE: 247		
000		
SEQ ID NO: 248	moltype =	length =
SEQUENCE: 248		
000		

-continued

---

SEQ ID NO: 249	moltype =	length =
SEQUENCE: 249		
000		
SEQ ID NO: 250	moltype =	length =
SEQUENCE: 250		
000		
SEQ ID NO: 251	moltype =	length =
SEQUENCE: 251		
000		
SEQ ID NO: 252	moltype =	length =
SEQUENCE: 252		
000		
SEQ ID NO: 253	moltype =	length =
SEQUENCE: 253		
000		
SEQ ID NO: 254	moltype =	length =
SEQUENCE: 254		
000		
SEQ ID NO: 255	moltype =	length =
SEQUENCE: 255		
000		
SEQ ID NO: 256	moltype =	length =
SEQUENCE: 256		
000		
SEQ ID NO: 257	moltype =	length =
SEQUENCE: 257		
000		
SEQ ID NO: 258	moltype =	length =
SEQUENCE: 258		
000		
SEQ ID NO: 259	moltype =	length =
SEQUENCE: 259		
000		
SEQ ID NO: 260	moltype =	length =
SEQUENCE: 260		
000		
SEQ ID NO: 261	moltype =	length =
SEQUENCE: 261		
000		
SEQ ID NO: 262	moltype =	length =
SEQUENCE: 262		
000		
SEQ ID NO: 263	moltype =	length =
SEQUENCE: 263		
000		
SEQ ID NO: 264	moltype =	length =
SEQUENCE: 264		
000		
SEQ ID NO: 265	moltype =	length =
SEQUENCE: 265		
000		
SEQ ID NO: 266	moltype =	length =
SEQUENCE: 266		
000		
SEQ ID NO: 267	moltype =	length =
SEQUENCE: 267		
000		

-continued

---

SEQ ID NO: 268	moltype =	length =
SEQUENCE: 268		
000		
SEQ ID NO: 269	moltype =	length =
SEQUENCE: 269		
000		
SEQ ID NO: 270	moltype =	length =
SEQUENCE: 270		
000		
SEQ ID NO: 271	moltype =	length =
SEQUENCE: 271		
000		
SEQ ID NO: 272	moltype =	length =
SEQUENCE: 272		
000		
SEQ ID NO: 273	moltype =	length =
SEQUENCE: 273		
000		
SEQ ID NO: 274	moltype =	length =
SEQUENCE: 274		
000		
SEQ ID NO: 275	moltype =	length =
SEQUENCE: 275		
000		
SEQ ID NO: 276	moltype =	length =
SEQUENCE: 276		
000		
SEQ ID NO: 277	moltype =	length =
SEQUENCE: 277		
000		
SEQ ID NO: 278	moltype =	length =
SEQUENCE: 278		
000		
SEQ ID NO: 279	moltype =	length =
SEQUENCE: 279		
000		
SEQ ID NO: 280	moltype =	length =
SEQUENCE: 280		
000		
SEQ ID NO: 281	moltype =	length =
SEQUENCE: 281		
000		
SEQ ID NO: 282	moltype =	length =
SEQUENCE: 282		
000		
SEQ ID NO: 283	moltype =	length =
SEQUENCE: 283		
000		
SEQ ID NO: 284	moltype =	length =
SEQUENCE: 284		
000		
SEQ ID NO: 285	moltype =	length =
SEQUENCE: 285		
000		
SEQ ID NO: 286	moltype =	length =
SEQUENCE: 286		
000		

-continued

---

SEQ ID NO: 287	moltype =	length =
SEQUENCE: 287		
000		
SEQ ID NO: 288	moltype =	length =
SEQUENCE: 288		
000		
SEQ ID NO: 289	moltype =	length =
SEQUENCE: 289		
000		
SEQ ID NO: 290	moltype =	length =
SEQUENCE: 290		
000		
SEQ ID NO: 291	moltype =	length =
SEQUENCE: 291		
000		
SEQ ID NO: 292	moltype =	length =
SEQUENCE: 292		
000		
SEQ ID NO: 293	moltype =	length =
SEQUENCE: 293		
000		
SEQ ID NO: 294	moltype =	length =
SEQUENCE: 294		
000		
SEQ ID NO: 295	moltype =	length =
SEQUENCE: 295		
000		
SEQ ID NO: 296	moltype =	length =
SEQUENCE: 296		
000		

---

**1.** A nucleic acid compound, comprising an oligonucleotide selected from the group consisting of short interfering RNA (siRNA), short hairpin RNA (shRNA), antisense RNA, DNA, antisense DNA, Chimeric Antisense DNA/RNA, and microRNA (miRNA), wherein the oligonucleotide has a 6-mer seed sequence that is complementary to a sequence of either a gene or an mRNA encoding an androgen receptor (AR) coregulator or a fragment thereof having AR coregulator activity;

wherein the oligonucleotide comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO:249, SEQ ID NO:250, SEQ ID NO:251, and SEQ ID NO:252, and nucleic acid sequences complementary to each of said sequences; and wherein SEQ ID NO:249, SEQ ID NO:250, SEQ ID NO:251, and SEQ ID NO:252 comprise positions 2-8 of their respective oligonucleotides, and wherein the nucleic acid compound comprises at least one of (1) a non-natural modification in the oligonucleotide, and (2) an organic moiety conjugated to the oligonucleotide; and wherein the nucleic acid compound has inhibitory activity against the expression or biological activity of the AR coregulator or biologically active fragment thereof.

**2.** The nucleic acid compound of claim 1, wherein the nucleic acid sequence of the oligonucleotide is selected from the group consisting of SEQ ID NO:293, SEQ ID NO:294, SEQ ID NO:295, and SEQ ID NO:296.

**3.** The nucleic acid compound of claim 1, wherein the at least one non-natural modification in the oligonucleotide is

selected from the group consisting of a modified nucleoside sugar, a modified internucleoside linkage, and a modified nucleobase.

**4.** The nucleic acid compound of claim 3, wherein the modified nucleoside sugar comprises a 2'-modified sugar.

**5.** The nucleic acid compound of claim 4, wherein the 2'-modified sugar is selected from the group consisting of 2'-O-methyl-, 2'-deoxy-, 2'-deoxy-2'-fluoro-, 2'-O-methoxyethyl-(2'-O-MOE)-, 2'-O-aminopropyl-(2'-O-AP)-, 2'-O-dimethylaminoethyl-(2'-O-DMAOE)-, 2'-O-dimethylamino-propyl-(2'-O-DMAP)-, 2'-O-dimethylaminoethoxyethyl-(2'-O-DMAEOE)-, 2'-O-guanidinopropyl-, and 2'-O—N-methylacetamido-(2'-O-NMA)-modified sugars.

**6.** The nucleic acid compound of claim 3, wherein the modified internucleoside linkage is selected from the group consisting of phosphorothioate, phosphordithioate, phosphoramidite, phosphordiamidate, morpholino, phosphotriester, aminoalkylphosphotriester, phosphonate, alkyl phosphonate, 3'-alkylene phosphonate, 5'-alkylene phosphonate, chiral phosphonate, and phosphinate linkages.

**7.** The nucleic acid compound of claim 3, wherein the modified internucleoside linkage comprises at least the first, second, or third internucleoside linkage at the 5' and/or 3' end of the oligonucleotide.

**8.** The nucleic acid compound of claim 3, wherein the modified nucleobase is selected from the group consisting of 5-uracil (pseudouridine), dihydrouracil, 2-thiouracil, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, dihydrouracil, 5-trifluoromethyl uracil, 5-methyl-2-thiouracil,

2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, 5-carboxymethylaminomethyl-2-thiouracil, 5-(carboxyhydroxylmethyl) uracil, 5-methoxycarboxymethyluracil, 5-methoxyuracil, 5-methyl-2-thiouracil, 5-propynyl uracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, 5-carboxymethylaminomethyluracil, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, alkynyl uracil, 4-thiouracil, 6-azo uracil, 2-aminoadenine, 2-methyladenine, 6-methyladenine, 7-methyladenine, 2-F-adenine, 2-amino-adenine, 2-propyladenine, N6-adenine, N6-isopentenyladenine, 2-methylthio-N6-iso-pentenyladenine, alkynyl adenine, 8-fluoro-adenine, 8-chloro-adenine, 8-bromo-adenine, 8-iodo-adenine, 8-amino adenine, 8-thiol adenine, 8-thioalkyl adenine, 8-hydroxyl adenine, 3-deazaadenine, 7-deazaadenine, 8-azaadenine, 3-methylcytosine, 5-methylcytosine, 5-trifluoromethyl cytosine, 2-thiocytosine, 5-hydroxymethyl cytosine, 5-propynyl cytosine, 6-azo cytosine, 4-acetylcytosine, 2-methylguanine, 6-methylguanine, 2-propylguanine, 1-methylguanine, 7-methylguanine, 2,2-dimethylguanine, 8-azaguanine, 7-deazaguanine, 8-fluoro-guanine, 8-chloro-guanine, 8-bromo-guanine, 8-ido-guanine, 8-amino guanine, 8-thiol guanine, 8-thioalkyl guanine, 8-hydroxyl guanine, 3-deazaguanine, 2-thiothymine, 6-azo thymine, alkynyl thymine, ribothymine, inosine, hypoxanthine, xanthine, beta-D-galactosylqueosine, beta-D-mannosylqueosine, 1-methylinosine, 2,6-diaminopurine, queosine, tricyclic pyrimidines, phenoxazine cytidine(1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), and phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one).

9. The nucleic acid compound of claim 1, wherein the organic moiety is selected from the group consisting of lipids, peptides, receptor-specific ligands, aptamers, antibodies or antibody fragments, CpG-containing oligonucleotides, polyamines, polymers, dendrimers, saccharides, polysaccharides, and cyclodextrins.

10. The nucleic acid compound of claim 1, wherein the AR coregulator is selected from the group consisting of CCND1, ADAM10, APPBP2, ATAD2, BAG1, BRCA1, CALM1, CALR, CCND3, CDC25A, CDK2AP1, CTNNB1, CDK7, CTDSP2, COP5, CTDSP2, ENY2, EHMT2, FKBP4, FKBP5, GSK3B, HELZ2, HEY1, HIP1, HIPK3, IDE, IL6ST, KAT2B, MAPK1, MAPK15, MED1, MED21, MKRN1, NCOA2, PER1, PIAS1, PIK3CB, PIK3R1, PMEPA1, PRMT2, PSMC3IP, PQBP1, PRKDC, RANBP9, RANBP10, RPS6KA1, RPS6KA3, SMAD1, SMARCD1, TAF1, TPID52, UBE2L3, UXT, YWHAH, and ZMIZ1.

11. The nucleic acid compound of claim 1, wherein the AR coregulator is an AR coregulator of an aggressive prostate cancer.

12. An expression vector comprising the oligonucleotide of claim 1.

13. A composition, comprising at least one nucleic acid compound of claim 1, and a pharmaceutically-acceptable carrier.

14. The composition of claim 13, comprising a nanoparticle.

15. A method for treating prostate cancer in a subject in need of such treatment, comprising: administering to the subject a therapeutically effective amount of an oligonucleotide selected from the group consisting of short interfering RNA (siRNA), short hairpin RNA (shRNA), antisense RNA, DNA, antisense DNA, Chimeric Antisense DNA/RNA, and microRNA (miRNA), wherein the oligonucleotide has a

6-mer seed sequence that is complementary to a sequence of either a gene or an mRNA encoding an androgen receptor (AR) coregulator or a fragment thereof having AR coregulator activity;

wherein the oligonucleotide comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO:249, SEQ ID NO:250, SEQ ID NO:251, and SEQ ID NO:252, and nucleic acid sequences complementary to each of said sequences; and wherein SEQ ID NO:249, SEQ ID NO:250, SEQ ID NO:251, and SEQ ID NO:252 comprise positions 2-8 of their respective oligonucleotides, and wherein the oligonucleotide has inhibitory activity against the expression or biological activity of the AR coregulator or biologically active fragment thereof.

16. The method of claim 15, wherein the prostate cancer is aggressive prostate cancer.

17. The method of claim 15, wherein the oligonucleotide comprises a portion of a nucleic acid compound or nucleic acid composition.

18. The method of claim 17, wherein the nucleic acid compound comprises at least one of (1) a non-natural modification in the oligonucleotide, and (2) an organic moiety conjugated to the oligonucleotide.

19. The method of claim 18, wherein the at least one non-natural modification in the oligonucleotide is selected from the group consisting of a modified nucleoside sugar, a modified internucleoside linkage, and a modified nucleobase.

20. The method of claim 19, wherein the modified nucleoside sugar comprises a 2'-modified sugar.

21. The method of claim 20, wherein the 2'-modified sugar is selected from the group consisting of 2'-O-methyl-, 2'-deoxy-, 2'-fluoro-, 2'-deoxy-2'-fluoro-, 2'-O-methoxyethyl-(2'-O-MOE)-, 2'-O-aminopropyl-(2'-O-AP)-, 2'-O-dimethylaminoethyl-(2'-O-DMAOE)-, 2'-O-dimethylaminopropyl-(2'-O-DMAP)-, 2'-O-dimethylaminoethoxyethyl-(2'-O-DMAEOE)-, 2'-O-guanidinopropyl-, and 2'-O—N-methylacetamido-(2'-O-NMA)-modified sugars.

22. The method of claim 19, wherein the modified internucleoside linkage is selected from the group consisting of phosphorothioate, phosphorodithioate, phosphoramidite, phosphordiamidate, morpholino, phosphotriester, aminoalkylphosphotriester, phosphonate, alkyl phosphonate, 3'-alkylene phosphonate, 5'-alkylene phosphonate, chiral phosphonate, and phosphinate linkages.

23. The method of claim 19, wherein the modified internucleoside linkage comprises at least the first, second, or third internucleoside linkage at the 5' and/or 3' end of the oligonucleotide.

24. The method of claim 19, wherein the modified nucleobase is selected from the group consisting of 5-uracil (pseudouridine), dihydrouracil, 2-thiouracil, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, dihydrouracil, 5-trifluoromethyl uracil, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, 5-carboxymethylaminomethyl-2-thiouracil, 5-(carboxyhydroxylmethyl) uracil, 5-methoxycarboxymethyluracil, 5-methoxyuracil, 5-methyl-2-thiouracil, 5-propynyl uracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, 5-carboxymethylaminomethyluracil, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, alkynyl uracil, 4-thiouracil, 6-azo uracil, 2-aminoadenine, 2-methyladenine, 6-methyladenine, 7-methylad-

enine, 2-F-adenine, 2-amino-adenine, 2-propyladenine, N6-adenine, N6-isopentenyladenine, 2-methylthio-N6-isopentenyladenine, alkynyl adenine, 8-fluoro-adenine, 8-chloro-adenine, 8-bromo-adenine, 8-ido-adenine, 8-amino adenine, 8-thiol adenine, 8-thioalkyl adenine, 8-hydroxyl adenine, 3-deazaadenine, 7-deazaadenine, 8-azaadenine, 3-methylcytosine, 5-methylcytosine, 5-trifluoromethyl cytosine, 2-thiocytosine, 5-hydroxymethyl cytosine, 5-propynyl cytosine, 6-azo cytosine, 4-acetylcytosine, 2-methylguanine, 6-methylguanine, 2-propylguanine, 1-methylguanine, 7-methylguanine, 2,2-dimethylguanine, 8-azaguanine, 7-deazaguanine, 8-fluoro-guanine, 8-chloro-guanine, 8-bromo-guanine, 8-ido-guanine, 8-amino guanine, 8-thiol guanine, 8-thioalkyl guanine, 8-hydroxyl guanine, 3-deazaguanine, 2-thiothymine, 6-azo thymine, alkynyl thymine, ribothymine, inosine, hypoxanthine, xanthine, beta-D-galactosylqueosine, beta-D-mannosylqueosine, 1-methylinosine, 2,6-diaminopurine, queosine, tricyclic pyrimidines, phenoxazine cytidine(1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), and phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one.

**25.** The method of claim **18**, wherein the organic moiety is selected from the group consisting of lipids, peptides, receptor-specific ligands, aptamers, antibodies or antibody fragments, CpG-containing oligonucleotides, polyamines, polymers, dendrimers, saccharides, polysaccharides, and cyclodextrins.

**26.** The method of claim **15**, wherein the AR coregulator is selected from the group consisting of CCND1, ADAM10, APPBP2, ATAD2, BAG1, BRCA1, CALM1, CALR, CCND3, CDC25A, CDK2AP1, CTNNB1, CDK7, CTDSP2, COPS5, CTDSP2, ENY2, EHMT2, FKBP4, FKBP5, GSK3B, HELZ2, HEY1, HIP1, HIPK3, IDE, IL6ST, KAT2B, MAPK1, MAPK15, MED1, MED21, MKRN1, NCOA2, PER1, PIAS1, PIK3CB, PIK3R1, PMEPA1, PRMT2, PSMC3IP, PQBP1, PRKDC, RANBP9, RANBP10, RPS6KA1, RPS6KA3, SMAD1, SMARCD1, TAF1, TPD52, UBE2L3, UXT, YWHAH, and ZMIZ1.

**27.** The method of claim **15**, wherein the AR coregulator is an AR coregulator of an aggressive prostate cancer.

\* \* \* \* \*