

US Patent & Trademark Office

Patent Public Search | Text View

United States Reissue Patent

RE50527

Kind Code

E

Date of Reissued Patent

August 12, 2025

Inventor(s)

Ren; Pingda et al.

Covalent inhibitors of KRAS G12C

Abstract

Irreversible inhibitors of G12C mutant K-Ras protein are provided. Also disclosed are methods to modulate the activity of G12C mutant K-Ras protein and methods of treatment of disorders mediated by G12C mutant K-Ras protein.

Inventors: Ren; Pingda (San Diego, CA), Liu; Yi (San Diego, CA), Li; Liansheng (San Diego, CA), Feng; Jun (San Diego, CA), Wu; Tao (Carlsbad, CA)

Applicant: Araxes Pharma LLC (San Diego, CA)

Family ID: 1000008446269

Assignee: Araxes Pharma LLC (San Diego, CA)

Appl. No.: 18/109726

Filed: February 14, 2023

Related U.S. Application Data

division parent-doc US 15891304 20180207 US 10273207 20190430 child-doc US 18109726
division parent-doc US 14933734 20151105 US 9926267 20180327 child-doc US 15891304
division parent-doc US 14212656 20140314 US 9227978 20160105 child-doc US 14933734
reissue parent-doc US 16355258 20190315 GRANTED US 10919850 20210216 child-doc US 18109726
us-provisional-application US 61889480 20131010
us-provisional-application US 61852123 20130315

Publication Classification

Int. Cl.: C07D205/04 (20060101); C07D211/56 (20060101); C07D211/58 (20060101); C07D211/60 (20060101); C07D211/62 (20060101); C07D213/64 (20060101); C07D213/74 (20060101); C07D231/40 (20060101); C07D237/04 (20060101); C07D285/14 (20060101); C07D285/16 (20060101); C07D295/16 (20060101); C07D401/04 (20060101); C07D401/12 (20060101); C07D403/04 (20060101); C07D403/06 (20060101); C07D403/12 (20060101); C07D405/06 (20060101); C07D419/12 (20060101); C07D471/04 (20060101); C07D487/04 (20060101); C07D487/10 (20060101); G01N33/58 (20060101)

U.S. Cl.:

CPC C07D205/04 (20130101); C07D211/56 (20130101); C07D211/58 (20130101); C07D211/60 (20130101); C07D211/62 (20130101); C07D213/64 (20130101); C07D213/74 (20130101); C07D231/40 (20130101); C07D237/04 (20130101); C07D285/14 (20130101); C07D285/16 (20130101); C07D295/16 (20130101); C07D401/04 (20130101); C07D401/12 (20130101); C07D403/04 (20130101); C07D403/06 (20130101); C07D403/12 (20130101); C07D405/06 (20130101); C07D419/12 (20130101); C07D471/04 (20130101); C07D487/04 (20130101); C07D487/10 (20130101); G01N33/58 (20130101);

Field of Classification Search

CPC: C07D (205/04); C07D (211/56); C07D (211/58); C07D (211/60); C07D (211/62); C07D (213/64); C07D (213/74); C07D (231/40); C07D (237/04); C07D (285/14); C07D (285/16); C07D (295/16); C07D (401/04); C07D (401/12); C07D (403/04); C07D (403/06); C07D (403/12); C07D (405/06); C07D (419/12); C07D (471/04); C07D (487/04); C07D (487/10); G01N (33/58); A61P (35/00)

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
3702849	12/1971	Cronin et al.	N/A	N/A
3752660	12/1972	Little	N/A	N/A
4436748	12/1983	Ong et al.	N/A	N/A
4439606	12/1983	Du et al.	N/A	N/A
4649219	12/1986	Itoh et al.	N/A	N/A
4656181	12/1986	Sunkel et al.	N/A	N/A
4861760	12/1988	Mazuel et al.	N/A	N/A
4911920	12/1989	Jani et al.	N/A	N/A
5010175	12/1990	Rutter et al.	N/A	N/A
5033252	12/1990	Carter	N/A	N/A
5052558	12/1990	Carter	N/A	N/A
5212162	12/1992	Missel et al.	N/A	N/A
5288514	12/1993	Ellman	N/A	N/A
5323907	12/1993	Kalvelage	N/A	N/A
5403841	12/1994	Lang et al.	N/A	N/A
5538848	12/1995	Livak et al.	N/A	N/A
5539083	12/1995	Cook et al.	N/A	N/A

5593853	12/1996	Chen et al.	N/A	N/A
5605798	12/1996	Köster	N/A	N/A
5670505	12/1996	Matsuo et al.	N/A	N/A
5731352	12/1997	Lesieur et al.	N/A	N/A
5777324	12/1997	Hillenkamp	N/A	N/A
5858659	12/1998	Sapolsky et al.	N/A	N/A
5861510	12/1998	Piscopio et al.	N/A	N/A
5863949	12/1998	Robinson et al.	N/A	N/A
5919626	12/1998	Shi et al.	N/A	N/A
5925525	12/1998	Fodor et al.	N/A	N/A
5952174	12/1998	Nikiforov et al.	N/A	N/A
6002008	12/1998	Wissner et al.	N/A	N/A
6017696	12/1999	Heller	N/A	N/A
6043031	12/1999	Köster et al.	N/A	N/A
6045996	12/1999	Cronin et al.	N/A	N/A
6051380	12/1999	Sosnowski et al.	N/A	N/A
6068818	12/1999	Ackley et al.	N/A	N/A
6214872	12/2000	Robinson	N/A	N/A
6251639	12/2000	Kurn	N/A	N/A
6511993	12/2002	Dack et al.	N/A	N/A
6586447	12/2002	Lyssikatos et al.	N/A	N/A
6849639	12/2004	Dominguez et al.	N/A	N/A
6903118	12/2004	Biedermann et al.	N/A	N/A
7074801	12/2005	Yoshida et al.	N/A	N/A
7547691	12/2008	Dahnke et al.	N/A	N/A
7595397	12/2008	Zindell et al.	N/A	N/A
7632838	12/2008	Xiang et al.	N/A	N/A
7879863	12/2010	Tokumasu et al.	N/A	N/A
8399454	12/2012	Bian et al.	N/A	N/A
8426401	12/2012	Bian et al.	N/A	N/A
8466288	12/2012	Aronov et al.	N/A	N/A
8604017	12/2012	Bian et al.	N/A	N/A
8697684	12/2013	Bian et al.	N/A	N/A
8741887	12/2013	Bian et al.	N/A	N/A
8759333	12/2013	Connolly et al.	N/A	N/A
9227978	12/2015	Ren et al.	N/A	N/A
9273057	12/2015	Purandare et al.	N/A	N/A
9376559	12/2015	Holtcamp et al.	N/A	N/A
9745319	12/2016	Ren et al.	N/A	N/A
9810690	12/2016	Patricelli et al.	N/A	N/A
9840516	12/2016	Li et al.	N/A	N/A
9862701	12/2017	Li et al.	N/A	N/A
9926267	12/2017	Ren et al.	N/A	N/A
9988357	12/2017	Mani et al.	N/A	N/A
10111874	12/2017	Janes et al.	N/A	N/A
10118931	12/2017	Chesworth et al.	N/A	N/A
10144724	12/2017	Li et al.	N/A	N/A
10246424	12/2018	Li et al.	N/A	N/A
10273207	12/2018	Ren et al.	N/A	N/A
10280172	12/2018	Li et al.	N/A	N/A

10351550	12/2018	Li et al.	N/A	N/A
10370386	12/2018	Li et al.	N/A	N/A
10377743	12/2018	Li et al.	N/A	N/A
10414757	12/2018	Li et al.	N/A	N/A
10428064	12/2018	Li et al.	N/A	N/A
10556906	12/2019	Kuramoto et al.	N/A	N/A
10646488	12/2019	Liu et al.	N/A	N/A
10647703	12/2019	Li et al.	N/A	N/A
10689356	12/2019	Li et al.	N/A	N/A
10723738	12/2019	Li et al.	N/A	N/A
10730867	12/2019	Li et al.	N/A	N/A
10736897	12/2019	Li et al.	N/A	N/A
10745385	12/2019	Li et al.	N/A	N/A
11008334	12/2020	Ostrem et al.	N/A	N/A
2002/0169300	12/2001	Waterman et al.	N/A	N/A
2003/0022344	12/2002	Williams et al.	N/A	N/A
2003/0078277	12/2002	Hibi et al.	N/A	N/A
2003/0166620	12/2002	Lee et al.	N/A	N/A
2003/0171400	12/2002	Pikul et al.	N/A	N/A
2004/0106634	12/2003	Satoh et al.	N/A	N/A
2005/0012070	12/2004	Inoue et al.	N/A	N/A
2005/0119266	12/2004	Shi et al.	N/A	N/A
2005/0227997	12/2004	Noe et al.	N/A	N/A
2007/0185089	12/2006	Bamford et al.	N/A	N/A
2007/0249648	12/2006	Bladh et al.	N/A	N/A
2008/0004285	12/2007	De Jonghje et al.	N/A	N/A
2008/0004348	12/2007	Yous et al.	N/A	N/A
2008/0021026	12/2007	Kahraman et al.	N/A	N/A
2008/0039450	12/2007	Jensen et al.	N/A	N/A
2008/0070892	12/2007	Harris et al.	N/A	N/A
2009/0036430	12/2008	De Jonghje et al.	N/A	N/A
2009/0054402	12/2008	Wang et al.	N/A	N/A
2009/0124636	12/2008	Barber et al.	N/A	N/A
2009/0298075	12/2008	Travers et al.	N/A	N/A
2010/0331300	12/2009	Bian et al.	N/A	N/A
2011/0046370	12/2010	Sim et al.	N/A	N/A
2011/0217309	12/2010	Buck et al.	N/A	N/A
2011/0230476	12/2010	Niu et al.	N/A	N/A
2011/0269244	12/2010	Petter et al.	N/A	N/A
2011/0311447	12/2010	Tu et al.	N/A	N/A
2011/0319290	12/2010	Raymond et al.	N/A	N/A
2012/0040020	12/2011	Charrier et al.	N/A	N/A
2012/0083476	12/2011	Breitenbucher et al.	N/A	N/A
2013/0012489	12/2012	Mederski	N/A	N/A
2013/0029964	12/2012	Aoki et al.	N/A	N/A
2013/0040934	12/2012	Cmiljanovic et al.	N/A	N/A
2013/0274252	12/2012	Pandey et al.	N/A	N/A
2013/0302407	12/2012	Rao et al.	N/A	N/A
2014/0288045	12/2013	Ren et al.	N/A	N/A
2014/0315886	12/2013	Suzuki et al.	N/A	N/A

2015/0005277	12/2014	Wang et al.	N/A	N/A
2015/0087628	12/2014	Ostrem et al.	N/A	N/A
2015/0239900	12/2014	Li et al.	N/A	N/A
2015/0374678	12/2014	Chamberlain et al.	N/A	N/A
2016/0016948	12/2015	Bryan et al.	N/A	N/A
2016/0031898	12/2015	Ren et al.	N/A	N/A
2016/0108019	12/2015	Li et al.	N/A	N/A
2016/0159738	12/2015	Ren et al.	N/A	N/A
2016/0166571	12/2015	Janes et al.	N/A	N/A
2016/0368930	12/2015	Ostrem et al.	N/A	N/A
2017/0022184	12/2016	Li et al.	N/A	N/A
2017/0131278	12/2016	Patricelli et al.	N/A	N/A
2017/0190672	12/2016	Mani et al.	N/A	N/A
2017/0197945	12/2016	Li et al.	N/A	N/A
2017/0247376	12/2016	Li et al.	N/A	N/A
2018/0015087	12/2017	Liu et al.	N/A	N/A
2018/0086753	12/2017	Li et al.	N/A	N/A
2018/0141927	12/2017	Li et al.	N/A	N/A
2018/0246102	12/2017	Patricelli et al.	N/A	N/A
2018/0273515	12/2017	Li et al.	N/A	N/A
2018/0273523	12/2017	Li et al.	N/A	N/A
2018/0282307	12/2017	Li et al.	N/A	N/A
2018/0282308	12/2017	Li et al.	N/A	N/A
2018/0319775	12/2017	Li et al.	N/A	N/A
2019/0055211	12/2018	Li et al.	N/A	N/A
2019/0062313	12/2018	Li et al.	N/A	N/A
2019/0062314	12/2018	Li et al.	N/A	N/A
2019/0092767	12/2018	Li et al.	N/A	N/A
2019/0119295	12/2018	Ostrem et al.	N/A	N/A
2019/0262342	12/2018	Janes et al.	N/A	N/A
2019/0284144	12/2018	Li et al.	N/A	N/A
2019/0343838	12/2018	Allen et al.	N/A	N/A
2019/0345158	12/2018	Li et al.	N/A	N/A
2019/0367489	12/2018	Li et al.	N/A	N/A
2019/0375743	12/2018	Hudson et al.	N/A	N/A
2019/0382377	12/2018	Li et al.	N/A	N/A
2019/0389851	12/2018	Li et al.	N/A	N/A
2020/0010454	12/2019	Li et al.	N/A	N/A
2020/0102321	12/2019	Li et al.	N/A	N/A
2020/0115363	12/2019	Li et al.	N/A	N/A
2020/0181123	12/2019	Li et al.	N/A	N/A
2021/0230183	12/2020	Ostrem et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
1267291	12/1999	CN	N/A
104 418 860	12/2015	CN	N/A
0094498	12/1982	EP	N/A
0 606 046	12/1993	EP	N/A
0 780 386	12/1996	EP	N/A

0 818 442	12/1997	EP	N/A
0 931 788	12/1998	EP	N/A
1 004 578	12/1999	EP	N/A
0 655 442	12/2000	EP	N/A
1 348 434	12/2002	EP	N/A
1 736 465	12/2005	EP	N/A
2 133 334	12/2008	EP	N/A
2 270 002	12/2010	EP	N/A
2 889 291	12/2014	EP	N/A
939516	12/1962	GB	N/A
9912961.1	12/1998	GB	N/A
58-203966	12/1982	JP	N/A
S58203966	12/1982	JP	N/A
59-163372	12/1983	JP	N/A
S59163372	12/1983	JP	N/A
S6143190	12/1985	JP	N/A
2002-508356	12/2001	JP	N/A
2002-371078	12/2001	JP	N/A
2003-535860	12/2002	JP	N/A
2004250329	12/2003	JP	N/A
2005-502623	12/2004	JP	N/A
2005-179557	12/2004	JP	N/A
2007-16011	12/2006	JP	N/A
2007016011	12/2006	JP	N/A
2008-524154	12/2007	JP	N/A
4775259	12/2010	JP	N/A
2013-504325	12/2012	JP	N/A
2013-516422	12/2012	JP	N/A
2013-107855	12/2012	JP	N/A
2013-522249	12/2012	JP	N/A
2016532656	12/2015	JP	N/A
86/01207	12/1985	WO	N/A
90/05719	12/1989	WO	N/A
91/19735	12/1990	WO	N/A
92/00091	12/1991	WO	N/A
93/20242	12/1992	WO	N/A
96/05309	12/1995	WO	N/A
96/13262	12/1995	WO	N/A
96/27583	12/1995	WO	N/A
96/33172	12/1995	WO	N/A
97/00271	12/1996	WO	N/A
97/30992	12/1996	WO	N/A
98/03516	12/1997	WO	N/A
98/07697	12/1997	WO	N/A
98/30566	12/1997	WO	N/A
98/33456	12/1997	WO	N/A
98/33768	12/1997	WO	N/A
98/34915	12/1997	WO	N/A
98/35951	12/1997	WO	N/A
WO 9833496	12/1997	WO	N/A

98/43960	12/1997	WO	N/A
98/34918	12/1997	WO	N/A
98/57948	12/1997	WO	N/A
99/07675	12/1998	WO	N/A
99/29667	12/1998	WO	N/A
99/52910	12/1998	WO	N/A
99/32454	12/1998	WO	N/A
99/52889	12/1998	WO	N/A
99/67641	12/1998	WO	N/A
WO 9967641	12/1998	WO	N/A
00/39587	12/1999	WO	N/A
00/74681	12/1999	WO	N/A
WO 0168186	12/2000	WO	N/A
02/04420	12/2001	WO	N/A
02/080928	12/2001	WO	N/A
02/088107	12/2001	WO	N/A
03/004480	12/2002	WO	N/A
03/053958	12/2002	WO	N/A
2004/033427	12/2003	WO	N/A
2004/074283	12/2003	WO	N/A
2004/080976	12/2003	WO	N/A
2005/070891	12/2004	WO	N/A
2005/082892	12/2004	WO	N/A
2005/092894	12/2004	WO	N/A
2006/066948	12/2005	WO	N/A
2006/095014	12/2005	WO	N/A
2006/097261	12/2005	WO	N/A
2006/135993	12/2005	WO	N/A
2007/047146	12/2006	WO	N/A
2007/095588	12/2006	WO	N/A
2007/113226	12/2006	WO	N/A
2007/144394	12/2006	WO	N/A
2008/009078	12/2007	WO	N/A
2008/028691	12/2007	WO	N/A
2008/112440	12/2007	WO	N/A
2010/027746	12/2009	WO	N/A
2010/087399	12/2009	WO	N/A
2010/121918	12/2009	WO	N/A
2011/002816	12/2010	WO	N/A
2011/016559	12/2010	WO	N/A
2011/031896	12/2010	WO	N/A
2011/082285	12/2010	WO	N/A
2011/093524	12/2010	WO	N/A
2011/148922	12/2010	WO	N/A
2011/153553	12/2010	WO	N/A
2012/016082	12/2011	WO	N/A
2012/041872	12/2011	WO	N/A
2012/054716	12/2011	WO	N/A
2012/061557	12/2011	WO	N/A
2012/174489	12/2011	WO	N/A

2013/064068	12/2012	WO	N/A
2013/096151	12/2012	WO	N/A
2013/096455	12/2012	WO	N/A
2013/106641	12/2012	WO	N/A
2013/140148	12/2012	WO	N/A
2013/155077	12/2012	WO	N/A
2013/155223	12/2012	WO	N/A
2013/184757	12/2012	WO	N/A
2014/011900	12/2013	WO	N/A
WO 2014030743	12/2013	WO	N/A
2014/071109	12/2013	WO	N/A
2014/143659	12/2013	WO	N/A
2014/152588	12/2013	WO	N/A
2014/159837	12/2013	WO	N/A
2014/201435	12/2013	WO	N/A
WO 2014210255	12/2013	WO	N/A
2015/003166	12/2014	WO	N/A
2015/017502	12/2014	WO	N/A
2015/054572	12/2014	WO	N/A
WO 2015027222	12/2014	WO	N/A
2015/108992	12/2014	WO	N/A
2015/132799	12/2014	WO	N/A
2015/143148	12/2014	WO	N/A
2015/144001	12/2014	WO	N/A
2015/184349	12/2014	WO	N/A
2016/044772	12/2015	WO	N/A
2016/049524	12/2015	WO	N/A
2016/049565	12/2015	WO	N/A
2016/049568	12/2015	WO	N/A
2016/118951	12/2015	WO	N/A
2016/164675	12/2015	WO	N/A
2016/168540	12/2015	WO	N/A
2017/015562	12/2016	WO	N/A
2017/058902	12/2016	WO	N/A
2017/058915	12/2016	WO	N/A
2017/068412	12/2016	WO	N/A
2017/070256	12/2016	WO	N/A
2017/087528	12/2016	WO	N/A
2017/100546	12/2016	WO	N/A
2017/172979	12/2016	WO	N/A
2018/140512	12/2017	WO	N/A
2020/028706	12/2019	WO	N/A
2020/086739	12/2019	WO	N/A
2020/113071	12/2019	WO	N/A

OTHER PUBLICATIONS

Appel et al., "Supramolecular Cross-Linked Networks via Host-Guest Complexation with Cucurbit[8]uril," *J. Am. Chem. Soc.* 132(40):14251-14260, 2010. cited by applicant

Arkin et al., "Binding of small molecules to an adaptive protein-protein interface," *PNAS* 100(4):1603-1608, 2003. cited by applicant

Bhatia et al., "The challenges posed by cancer heterogeneity," *Nature Biotechnology* 30(7):604-610, Jul. 2012. cited by applicant

Bos, "ras Oncogenes in Human Cancer: A Review," *Cancer Research* 49:4682-4689, 1989 (9 pages). cited by applicant

CAS Registry No. 5530-21-2, "1-Propanone, 1-[4-[2-(2-methoxy-4-propylphenoxy)acetyl]-piperazinyl]-," entered into STN Nov. 16, 1984, last updated Dec. 15, 2008, 6 pages. cited by applicant

CAS RN 1060281-01-7, "1,2,4-Triazolo[4,3-b]pyridazine, 3-(2-furanyl)-6-[4-[(2-(4-methylphenyl)ethenyl)sulfonyl]-1-piperazinyl]-," created on Oct. 12, 2008, 1 page. cited by applicant

CAS RN 1060345-23-4, "1,2,4-Triazolo[4,3-b]pyridazine, 3-(2-furanyl)-6-[4-[(2-phenylethenyl)sulfonyl]-1-piperazinyl]-," created on Oct. 13, 2008, 1 page. cited by applicant

CAS RN 1156156-36-3, "2-Propen-1-one, 1-(4-benzoyl-1-piperazinyl)-," created on Jun. 12, 2009, 3 pages. cited by applicant

CAS RN 1252566-45-2, "2-Pentenamide, N-[1-(2-fluorobenzoyl)-4-piperidinyl]-2-methyl-," created on Nov. 10, 2010, 3 pages. cited by applicant

CAS RN 1276218-20-2, "1,2,4-Triazolo[4,3-b]pyridazine, 3-(2-fluorophenyl)-6-[4-[(2-phenylethenyl)sulfonyl]-1-piperazinyl]-," created on Apr. 7, 2011, 1 page. cited by applicant

CAS RN 1312005-01-8, "2-Penten-1-one, 1-[4-(2,3-dimethoxybenzoyl)-1-piperazinyl]-2-methyl-," created on Jul. 7, 2011, 3 pages. cited by applicant

CAS RN 1346522-30-2, "Methanone, [4-(phenylsulfonyl)-1-piperazinyl][5-(2-thienyl)-1H-pyrazol-3-yl]-," created on Nov. 29, 2011, 3 pages. cited by applicant

CAS RN 1390243-63-6, "Benzenesulfonamide, N-[1-[(1-methyl-1H-pyrrol-2-yl)carbonyl]-4-piperidinyl]-," created on Aug. 12, 2012, 3 pages. cited by applicant

CAS RN 1394801-07-0, "2-Pentenamide, N-[1-(2-ethoxybenzoyl)-4-piperidinyl]-2-methyl-," created on Sep. 18, 2012, 3 pages. cited by applicant

CAS RN 921172-03-4, "2-Propen-1-one, 1-[4-[4-(4-ethoxyphenyl)-2-pyrimidinyl]-1-piperazinyl]-," created on Feb. 15, 2007, 1 page. cited by applicant

CAS RN 922358-69-8, "2-Propen-1-one, 1-[4-[4-[3-(trifluoromethyl)phenyl]-2-pyrimidinyl]-1-piperazinyl]-," created on Feb. 22, 2007, 1 page. cited by applicant

Chihara et al., "Kagakudaijiten," Tokyo Kagaku Dojin K.K., Tokyo, Japan, 1989, p. 69. (w/o English Translation). cited by applicant

Cho et al., "An Unnatural Biopolymer," *Science* 261:1303-1305, 1993. cited by applicant

DeWitt et al., "'Diversomers': An approach to nonpeptide, nonoligomeric chemical diversity," *Proc. Natl. Acad. Sci.* 90:6909-6913, 1993. cited by applicant

Duncan et al., "N-Dialkylaminoalkylbiphenylamines as Antimalarial and Atischistosomal Agents," *Journal of Medicinal Chemistry* 12:25-29, 1969. cited by applicant

Erlanson et al., "Site-directed ligand discovery," *Proc. Natl Acad. Sci. U.S.A.* 97(17):9367-9372, 2000. cited by applicant

Evans, "Isotopic Labeling with Carbon-14 and Tritium," *Principles of Radiopharmacology*, CRC Press Inc., Boca Raton, Florida, USA, 1979, pp. 11-13 and 24-25. (7 pages). cited by applicant

Gorfe et al., "Mapping the Nucleotide and Isoform-Dependent Structural and Dynamical Features of Ras Proteins," *Structure* 16:885-896, 2008. cited by applicant

Hall et al., "The structural basis for the transition from Ras-GTP to Ras-GDP," *PNAS* 99(19):12138-12142, 2002. cited by applicant

Hardy et al., "Discovery of an allosteric site in the caspases," *PNAS* 101(34):12461-12466, 2004. cited by applicant

Hattori et al., "Neutralizing monoclonal antibody against ras oncogene product p21 which impairs guanine nucleotide exchange," *Mol. Cell. Biol.* 7(5):1999-2002, 1987. cited by applicant

Ito et al., "Regional Polyesterism in the GTP-Bound Form of the Human c-Ha-Ras Protein,"

Biochemistry 36(30):9109-1119, 1997. cited by applicant

Jayanthi et al., "Biological Activities of Few Nitrogen Heterocycles using Pass and Lipophilicity using Adme Boxes," *Pharma Science Monitor: An International Journal of Pharmaceutical Sciences* 3(3, Suppl-1):1994-2006, 2012. cited by applicant

Johnson et al., "The Chemistry of β -Bromopropionyl Isocyanate. I. Synthesis of 1-Aryldihydrouracils," *The Journal of Organic Chemistry* 24(9):1391-1392, 1959. cited by applicant

Jordan, "Tamoxifen: A most unlikely pioneering medicine," *Nature Reviews* 2:205-213, 2003. cited by applicant

Kandoth et al., "Mutational landscape and significance across 12 major cancer types," *Nature* 502:333-339, Oct. 2013. (20 pages). cited by applicant

Kelly et al., "Synthesis of Isomeric 3-Piperidinyl and 3-Pyrrolidinyl Benzo[5,6]cyclohepta[1,2-b]pyridines: Sulfonamido Derivatives as Inhibitors of Ras Prenylation," *Bioorganic & Medicinal Chemistry* 6(6):673-686, 1998. cited by applicant

Lee et al., "The mutation spectrum revealed by paired genome sequences from a lung cancer patient," *Nature* 465:473-477, 2010. cited by applicant

Liang et al., "Parallel Synthesis and Screening of a Solid Phase Carbohydrate Library," *Science* 274:1520-1522, 1996. cited by applicant

Lone et al., "A substrate-free activity-based protein profiling screen for the discovery of selective PREPL inhibitors," *J. Am Chem Soc.* 133(30):11665-11674, 2011. (20 pages). cited by applicant

Margarit et al., "Structural Evidence for Feedback Activation by Ras·GTP of the Ras-Specific Nucleotide Exchange Factor SOS," *Cell* 112:685-695, 2003. cited by applicant

Maurer et al., "Small-molecule ligands bind to a distinct pocket in Ras and inhibit SOS-mediated nucleotide exchange activity," *PNAS* 109(14):5299-5304, 2012. cited by applicant

Milburn et al., "Molecular switch for signal transduction: structural differences between active and inactive forms of protooncogenic ras proteins," *Science* 247(4945):939-945, 1990. cited by applicant

Mirguet et al., "From ApoA1 upregulation to BET family bromodomain inhibition: Discovery of I-BET151," *Bioorganic & Medicinal Chemistry Letters* 22:2963-2967, 2012. cited by applicant

Pacold et al., "Crystal structure and functional analysis of Ras binding to its effector phosphoinositide 3-kinase gamma," *Cell* 103(6):931-943, 2000. cited by applicant

Palmioli et al., "Selective cytotoxicity of a bicyclic Ras inhibitor in cancer cells expressing K-RasG13D," *Biochemical and Biophysical Research Communications* 386(4):593-597, 2009. cited by applicant

Pautsch et al., "Crystal structure of the C3bot-RalA complex reveals a novel type of action of a bacterial exoenzyme," *The EMBO Journal* 24:3670-3680, 2005. cited by applicant

Peri et al., "Arabinose-derived bicyclic amino acids: synthesis, conformational analysis and construction of an $\alpha\beta$ 3-selective RGD peptide," *J. Am. Chem. Soc., Perkins Trans* 1(5):638-644, 2002. cited by applicant

Peri et al., "Sugar-Derived Ras Inhibitors: Group Epitope Mapping by NMR Spectroscopy and Biological Evaluation," *Eur. J. Org. Chem.* 2006(16):3707-3720, 2006. cited by applicant

Peri et al., "Synthesis of bicyclic sugar azido acids and their incorporation in cyclic peptides," *Chem. Commun.* 23:2303-2304, 2000. cited by applicant

Pinedo et al., "Aggressive combination therapy to cure patients with metastatic cancer," *The Lancet Oncology* 1:72-73, 2000. cited by applicant

Prior et al., "A Comprehensive Survey of Ras Mutations in Cancer," *Cancer Research* 72(10):2457-2467, May 2012. cited by applicant

PubChem Compound, "(Z)-1-(4-(5-(5-(Trifluoromethyl)-1,2,4-oxadiazol-3-yl)pyridin-2-yl)piperazin-1-yl)but-2-en-1-one," CID: 49702158, created on Nov. 27, 2010, 4 pages. cited by applicant

PubChem Compound, "1-[4-(5-Ethyl-1,2,4-oxadiazol-3-yl)phenyl]-4-[(2E)-3-(2-furyl)prop-2-

enoyl]piperazine,” CID: 53064614, created on Jun. 21, 2011, 4 pages. cited by applicant

PubChem Compound, “6-Benzothiazoleacetic acid, 2-phenyl-,” CID: 37535, created Aug. 8, 2005, 12 pages. cited by applicant

PubChem Compound, “ACILGBNJ,” CID: 768370, created Jul. 8, 2005, retrieved on Aug. 28, 2014 from <http://pubchem.ncbi.nlm.nih.gov/summary/summary.cgi?cid=768370>, 3 pages. cited by applicant

PubChem Compound, “Compound Summary for CID 21765509: Molecular Formula C₁₈H₂₁N₅O₈,” CID: 21765509, created on Dec. 5, 2007, retrieved on Aug. 5, 2014 from <http://pubchem.ncbi.nlm.nih.gov/summary/summary.cgi?cid=21765509#x304>, 4 pages. cited by applicant

PubChem Compound, “Compound Summary for CID 21765511: Molecular Formula C₁₈H₂₁N₅O₈,” CID: 21765511, created on Dec. 5, 2007, retrieved on Aug. 5, 2014 from <http://pubchem.ncbi.nlm.nih.gov/summary/summary.cgi?cid=21765511#x304>, 4 pages. cited by applicant

PubChem Compound, “Compound Summary for CID 60018735: Molecular Formula C₃₀H₃₀O₁₃,” CID: 60018735, created Aug. 20, 2012, 1 page. cited by applicant

PubChem Compound, “Compound Summary for CID 9897840: Molecular Formula C₅₀H₄₆O₂₀,” CID: 9897840, created Oct. 25, 2006, retrieved from <http://pubchem.ncbi.nlm.nih.gov/summary/summary.cgi?cid=9897840#x304>, 6 pages. cited by applicant

PubChem Compound, “MLS000416491,” CID: 2579941, deposited on Mar. 5, 2007, retrieved on May 15, 2017 from <https://pubchem.ncbi.nlm.nih.gov/substance/22405303#section=Top>, 7 pages. cited by applicant

PubChem Compound, “Substance Record for SID 44253980,” CID: 966800, deposited on Dec. 5, 2007, retrieved on May 11, 2017 from <https://pubchem.ncbi.nlm.nih.gov/substance/44253980#section=Top>, 5 pages. cited by applicant

Sasaki et al., “Selective Formation of Stable Triplexes Including a TA or a CG Interrupting Site with New Bicyclic Nucleoside Analogues (WNA),” *J. Am. Chem. Soc.* 126(2):516-528, 2004. cited by applicant

Schubbert et al., “Biochemical and Functional Characterization of Germ Line KRAS Mutations,” *Molecular and Cellular Biology* 27(22):7765-7770, 2007. cited by applicant

Silverman, “Prodrugs and Drug Delivery Systems,” *The Organic Chemistry of Drug Design and Drug Action*, p. 352-400, 1992. cited by applicant

Simone et al., “Oncology: Introduction,” *Cecil Textbook of Medicine*, Ed. Bennett et al., 20th Edition, vol. 1, pp. 1004-1010, 1996. cited by applicant

Singh et al., “A Gene Expression Signature Associated with “K-Ras Addiction” Reveals Regulators of EMT and Tumor Cell Survival,” *Cancer Cell* 15:489-500, 2009. cited by applicant

STN Registry No. 1066896-20-5, “Methanone, [3-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]-1-piperidinyl]-1-cyclopenten-1-yl-,” Oct. 27, 2008, 1 page. cited by applicant

STN Registry No. 1069510-87-7, “[1,4'-Bipiperidine]-4-carboxamide, l'-(1-cyclohexen-1-ylcarbonyl)-N-(2-pyridinylmethyl)-,” Nov. 2, 2008, 1 page. cited by applicant

STN Registry No. 1069530-02-4, “[1,4'-Bipiperidine]-4-carboxamide, l'-(1-cyclopenten-1-ylcarbonyl)-N-(2-pyridinylmethyl)-,” Nov. 2, 2008, 1 page. cited by applicant

STN Registry No. 1212098-43-5, “Methanone, 1-cyclohexen-1-yl [(3R,4R)-3-hydroxy-4-[4-(2-phenylethyl)-1-piperazinyl]-1-piperidinyl]-, rel-,,” Mar. 21, 2010, 1 page. cited by applicant

STN Registry Nos. 1179043-51-6; 1156698-93-9; 1156156-36-3; 1070748-55-8; 1070685-77-6; 1070667-53-6; 1070315-89-7; 1069961-90-5; 1069566-58-0; 1065560-16-8; 1065554-88-2; 1061025-62-4; 1060435-61-1; 958948-04-4; 958839-16-2, Jul. 4, 2012 to Dec. 19, 2001, retrieval date: Feb. 21, 2020, 6 pages. cited by applicant

Streuff et al., “First asymmetric aminohydroxylation of acrylamides,” *Tetrahedron: Asymmetry* 16(21):3492-3496, 2005. cited by applicant

Sunaga et al., "Knockdown of Oncogenic KRAS in Non-Small Cell Lung Cancers Suppresses Tumor Growth and Sensitizes Tumor Cells to Targeted Therapy," *Molecular Cancer Therapeutics* 10(2):336-346, 2011. cited by applicant

Tsubaki et al., "Reduction of metastasis, cell invasion, and adhesion in mouse osteosarcoma by YM529/ONO-5920-induced blockade of the Ras/MEK/ERK and Ras/PI3K/Akt pathway," *Toxicology and Applied Pharmacology* 259(3):402-410, 2012. cited by applicant

Tulshian et al., "Synthesis and Phosphodiesterase Activity of Carboxylic Acid Mimetics of Cyclic Guanosine 3',5'-Monophosphate," *J. Med. Chem.* 36(9):1210-1220, 1993. cited by applicant

Vetter et al., "The Guanine Nucleotide-Binding Switch in Three Dimensions," *Science* 294(5545):1299-1304, 2001. cited by applicant

Wistuba et al., "Methodological and practical challenges for personalized cancer therapies," *Nature Reviews* 8:135-141, Mar. 2011. cited by applicant

Wolff, (ed.), *Burger's Medicinal Chemistry and Drug Discovery, Fifth Edition, Volume 1: Principles and Practice*, San Diego, California, John Wiley & Sons, 1994, pp. 975-977. cited by applicant

Wu et al., "Stereoselective synthesis of dioxabicycles from 1-C-allyl-2-O-benzyl-glycosides—An intramolecular cyclization between 2-O-benzyl oxygen and the allyl double bond," *Can. J. Chem.* 84(1):597-602, 2006. cited by applicant

Young et al., "Oncogenic and Wild-type Ras Play Divergent Roles in the Regulation of Mitogen-Activated Protein Kinase Signaling," *Cancer Discovery* 3(1):112-123, 2013. cited by applicant

Adibekian et al., "Optimization and characterization of a triazole urea dual inhibitor for lysophospholipase 1 (LYPLA1) and lysophospholipase 2 (LYPLA2)," *Probe Reports from the NIH Molecular Libraries Program*, 2011, 42 pages. cited by applicant

Al-Muhammed et al., "In-vivo studies on dexamethasone sodium phosphate liposomes," *J. Microencapsulation* 13(3):293-306, 1996. cited by applicant

American Chemical Society, STN Database: Nov. 16, 1984, RN5530-21-2. cited by applicant

Appel et al., "Supramolecular Cross-Linked Networks via Host-Guest Complexation with Cucurbit[8]uril," *Am. Chem. Soc.* 132(40):14251-14260, Jul. 2010. cited by applicant

Arkin et al., "Binding of small molecules to an adaptive protein—protein interface," *PNAS* 100(4):1603-1608, Feb. 2003. cited by applicant

Bachovchin et al., "Identification of selective inhibitors of uncharacterized enzymes by high-throughput screening with fluorescent activity-based probes," *Nat. Biotechnol* 27(7):387-394, 2009. cited by applicant

Banker et al. (eds.), *Modern Pharmaceutics*, New York, Marcel Dekker, Inc., 1996, pp. 451 and 596. (3 pages). cited by applicant

Barbe et al., "Highly Chemoselective Metal-Free Reduction of Tertiary Amides," *J Am. Chem. Soc.* 130:18-19, 2008. cited by applicant

Bégué et al., "Ions α -Cetocarbenium. Influence De La Structure Sur L'Evolution Des Ions α -Cetocyclohexylcarbenium," *Tetrahedron* 31(20):2505-2511, 1975. (English Abstract Only). cited by applicant

Berge et al., "Pharmaceutical Salts," *Journal of Pharmaceutical Sciences* 66(1):1-19, 1977. cited by applicant

Berger et al., "N-Heteroaryl compounds with cyclic bridging unit as antiparasitic agents and their preparation and use for the treatment of parasitic diseases," *Chemical Abstracts* 156 (477498), 2012. (3 pages). cited by applicant

CAS RN 1156156-36-3, Jun. 12, 2009, "2-Propen-1-one, 1-(4-benzoyl-1-piperazinyl)-". cited by applicant

CAS RN 1252566-45-2, Nov. 10, 2010, "2-Pentenamide, N-[1-(2-fluorobenzoyl)-4-piperidinyl]-2-methyl-". cited by applicant

CAS RN 1312005-01-8, Jul. 7, 2011, "2-Penten-1-one, 1-[4-(2,3-dimethoxybenzoyl)-1-

piperazinyl]-2-methyl-". cited by applicant
CAS RN 1346522-30-2, Nov. 29, 2011, "Methanone, [4-(phenylsulfonyl)-1-piperazinyl][5-(2-thienyl)-1H-pyrazol-3-yl]-". cited by applicant
CAS RN 1390243-63-6, Aug. 12, 2012, "Benzenesulfonamide, N-[1-[(1-methyl-1H-pyrrol-2-yl)carbonyl]-4-piperidinyl]-". cited by applicant
CAS RN 1394801-07-0, Sep. 18, 2012, "2-Pentenamide, N-[1-(2-ethoxybenzoyl)-4-piperidinyl]-2-methyl-". cited by applicant
CAS RN 921172-03-4, Feb. 15, 2007, "2-Propen-1-one, 1-[4-[4-(4-ethoxyphenyl)-2-pyrimidinyl]-1-piperazinyl]-," 1 page. cited by applicant
CAS RN 922358-69-8, Feb. 22, 2007, "2-Propen-1-one, 1-[4-[3-(trifluoromethyl)phenyl]-2-pyrimidinyl]-1-piperazinyl]-," 1 page. cited by applicant
Chemcats Chemical Abstract, Accession No. 1301347730, Sep. 9, 2015, 2 pages. cited by applicant
Chemocare, "Taxol," retrieved from <http://www.chemocare.com/chemotherapy/drug-info/Taxol.aspx> on Feb. 22, 2017, 3 pages. cited by applicant
Cho et al., "An Unnatural Biopolymer," *Science* 261:1303-1305, Sep. 1993 (4 pages). cited by applicant
Chonn et al., "Recent advances in liposomal drug-delivery systems," *Current Opinion in Biotechnology* 6:698-708, 1995. cited by applicant
Choong et al., "Identification of Potent and Selective Small-Molecule Inhibitors of Caspase-3 through the Use of Extended Tethering and Structure-Based Drug Design," *J Med. Chem.* 45:5005-5022, 2002. cited by applicant
Cox et al., "Drugging the undruggable RAS: Mission Possible?," *Nat. Rev. Drug Discov.* 13(11):828-851, 2014. cited by applicant
DeWitt et al., "'Diversomers': An approach to nonpeptide, nonoligomeric chemical diversity," *Proc. Natl. Acad. Sci.* 90:6909-6913, Aug. 1993. cited by applicant
Duncan et al., "N-Dialkylaminoalkylbiphenylamines as Antimalarial and Atischiostosomal Agents," *Journal of Medicinal Chemistry* 12:25-29, Jan. 1969. cited by applicant
Erlanson et al., "Site-directed ligand discovery," *Proc. Natl Acad. Sci.* 97(17):9367-9372, Aug. 2000. cited by applicant
Forbes et al., "COSMIC 2005," *British Journal of Cancer* 94:318-322, 2006. cited by applicant
Furka et al., "General method for rapid synthesis of multicomponent peptide mixtures," *Int. J. Peptide Protein Res.* 37:487-493, 1991. cited by applicant
Gorfe et al., "Mapping the Nucleotide and Isoform-Dependent Structural and Dynamical Features of Ras Proteins," *Structure* 16:885-896, Jun. 2008. cited by applicant
Haggam et al., "Facile synthesis of some condensed 1,3-thiazines and thiazoles under conventional conditions: antitumor activity," *Research on Chemical Intermediates* 43 (11):6299-6315, 2017. cited by applicant
Hagihara et al., "Vinylogous Polypeptides: An Alternative Peptide Backbone," *J. Am. Chem. Soc.* 114(16):6568-6570, 1992. cited by applicant
Hall et al., "The Effect of Mg.SUP.2+ .on Guanine Nucleotide Exchange Rate of p21.SUP.N-ras.," *The Journal of Biological Chemistry* 261(24):10963-10965, 1986. cited by applicant
Hall et al., "The structural basis for the transition from Ras-GTP to Ras-GDP," *PNAS* 99(19):12138-12142, Sep. 2002. cited by applicant
Hansen et al., "The reactivity-driven biochemical mechanism of covalent KRAS.SUP.G12C .inhibitors," *Nature Structural& Molecular Biology* 25:454-462, 2018. (14 pages). cited by applicant
Hardy et al., "Discovery of an allosteric site in the caspases," *PNAS* 101(34):12461-12466, Aug. 2004. cited by applicant
Hattori et al., "Neutralizing Monoclonal Antibody against ras Oncogene Product p21 Which Impairs Guanine Nucleotide Exchange," *Mol. Cell. Biol.* 7(5):1999-2002, May 1987. cited by

applicant

Ito et al., "Regional Polysterism in the GTP-Bound Form of the Human c-Ha-Ras Protein," *Biochemistry* 36(30):9109-9119, 1997. cited by applicant

Johnson et al., "The Chemistry of β -Bromopropionyl Isocyanate. I. Synthesis of 1-Aryldihydrouracils," *The Journal of Organic Chemistry* 24(9):1391-1392, Sep. 1959. cited by applicant

Jones et al., "Increased frequency of the k-ras G12C mutation in MYH polyposis colorectal adenomas," *British Journal of Cancer* 90:1591-1593, 2004. cited by applicant

Jordan, "Tamoxifen: A most unlikely pioneering medicine," *Nature Reviews* 2:205-213, Mar. 2003. cited by applicant

Kelly et al., "Synthesis of Isomeric 3-Piperidinyl and 3-Pyrrolidinyl Benzo[5,6]cyclohepta[1,2-b]pyridines: Sulfonamido Derivatives as Inhibitors of Ras Prenylation," *Bioorganic & Medicinal Chemistry* 6(6):673-686, Jun. 1998. cited by applicant

Knochel et al., "Functionalization of heterocyclic compounds using polyfunctional magnesium and zinc reagents," *Beilstein Journal of Organic Chemistry* 7:1261-1277, 2011. cited by applicant

Kraulis et al., "Solution Structure and Dynamics of Ras p21-GDP Determined by Heteronuclear Three- and Four-Dimensional NMR Spectroscopy," *Biochemistry* 33:3515-3531, 1994. cited by applicant

Kumar et al., "Synthesis of 3-Sulfonylamino Quinolines from 1-(2-Aminophenyl) Propargyl Alcohols through a Ag(I)-Catalyzed Hydroamination, (2+3) Cycloaddition, and an Unusual Strain-Driven Ring Expansion," *Organic Letters* 17(9):2226-2229, Apr. 2015. cited by applicant

Kuroyanagi et al., "Structure-Activity Relationships of 1,3-Benzoxazole-4-carbonitriles as Novel Antifungal Agents with Potent in Vivo Efficacy," *Chem. Pharm. Bull.* 59(3):341-352, 2011. cited by applicant

Le Picard et al., "Design and Synthesis of Naphthalenic Derivatives as Potential Inhibitors of Hydroxyindole-O-methyltransferase," *Pharm. Pharmacol. Commun.* 5:183-188, 1999. cited by applicant

Lee et al., "Preparation of (S)-3-(pyrimidinyl- or pyridinylphenyl)-5-(acetylaminomethyl)-2-oxazolidinones as antibacterial agents," *Chemical Abstracts* 136(37590), 2001. (8 pages). cited by applicant

Lee et al., "The mutation spectrum revealed by paired genome sequences from a lung cancer patient," *Nature* 465:473-477, May 2010. cited by applicant

Lenzen et al., "[10]Analysis of Intrinsic and CDC25-Stimulated Guanine Nucleotide Exchange of p21.SUP.ras.—Nucleotide Complexes by Fluorescence Measurements," *Methods in Enzymology* 255:95-109, 1995. cited by applicant

Liang et al., "Parallel Synthesis and Screening of a Solid Phase Carbohydrate Library," *Science* 274:1520-1522, Nov. 1996. cited by applicant

Liu et al., "Polygonatum cyrtonelectin induces murine fibrosarcoma L929 cell apoptosis and autophagy via blocking Ras-Raf and PI3K-Akt signaling pathways," *Biochimie* 92 :1934-1938, 2010. cited by applicant

Loboda et al., "A gene expression signature of RAS pathway dependence predicts response to PI3K and RAS pathway inhibitors and expands the population of RAS pathway activated tumors," *BMC Medical Genomics* 3 (26):1-11, 2010. cited by applicant

Lone et al., "A substrate-free activity-based protein profiling screen for the discovery of selective PREPL inhibitors," *J. Am Chem Soc.* 133(30):11665-11674, Aug. 2011, 20 pages. cited by applicant

Long, "Taxol: An important compound with an impressive structure," Organic and General Chemistry at Flathead Valley Community College, Sep. 10, 2011, retrieved from <https://longscience.com/2011/09/10/taxol-an-organic-compound-you-should-know-about/> on Feb. 22, 2017, 4 pages. cited by applicant

Malani et al., "Synthesis, characterization and in vitro screening on bacterial, fungal and malarial strain of piprazinyl cyano biphenyl based compounds," *Bioorganic Chemistry* 51:16-23, 2013. cited by applicant

Margarit et al., "Structural Evidence for Feedback Activation by Ras•GTP of the Ras-Specific Nucleotide Exchange Factor SOS," *Cell* 112:685-695, Mar. 2003. cited by applicant

Mattos et al., "Multiple Binding Modes," in Kubinyi (ed.), *3D Qsar in Drug Design—Theory Methods and Applications*, vol. 2-3, Springer, Boston, Massachusetts, USA, 1998, pp. 243-244. (6 Pages). cited by applicant

Maurer et al., "Small-molecule ligands bind to a distinct pocket in Ras and inhibit SOS-mediated nucleotide exchange activity," *PNAS* 109(14):5299-5304, Apr. 2012. cited by applicant

McGregor et al., "Expanding the Scope of Electrophiles Capable of Targeting K-Ras Oncogenes," *Biochemistry* 56:3178-3183, 2017. cited by applicant

McMahon, "VEGF Receptor Signaling in Tumor Angiogenesis," *The Oncologist* 5(suppl. 1):3-10, 2000. cited by applicant

Milburn et al., "Molecular switch for signal transduction: structural differences between active and inactive forms of protooncogenic ras proteins," *Science* 247(4945):939-945, Feb. 1990. cited by applicant

Minto et al., "Pharmacokinetics and Pharmacodynamics of Nandrolone Esters in Oil Vehicle: Effects of Ester, Injection Site and Injection Volume," *The Journal of Pharmacology and Experimental Therapeutics* 281(1):93-102, 1997. cited by applicant

Noe et al., "Selective Inhibition of Aggrecanase in Osteoarthritis Treatment," U.S. Appl. No. 60/148,464, filed Aug. 12, 1999, 92 pages. cited by applicant

Ohnmacht, Jr. et al., "Antimalarials. 5. α -Dibutylaminomethyl- and α -(2-Piperidyl)-3-quinolinemethanols," *Journal of Medicinal Chemistry* 14(1):17-24, 1971. cited by applicant

Ostrem et al., "K-Ras(G12C) inhibitors allosterically control GTP affinity and effector interactions," *Nature* 503(7477):548-551, 2013. (14 pages). cited by applicant

Pacold et al., "Crystal structure and functional analysis of Ras binding to its effector phosphoinositide 3-kinase gamma," *Cell* 103(6):931-943, Dec. 2000. cited by applicant

Palmioli et al., "First experimental identification of Ras-inhibitor binding interface using a water-soluble Ras ligand," *Bioorganic and Medicinal Chemistry Letters* 19 :4217-4222, 2009. cited by applicant

Palmioli et al., "Selective cytotoxicity of a bicyclic Ras inhibitor in cancer cells expressing K-Ras.SUP.G13D.," *Biochemical and Biophysical Research Communications* 386(4):593-597, Sep. 2009. cited by applicant

Pardin et al., "Synthesis and evaluation of peptidic irreversible inhibitors of tissue transglutaminase," *Bioorg Med Chem* 14 (24):8379-8385, 2006. cited by applicant

Pathan et al., "Lead identification for the K-Ras protein: virtual screening and combinatorial fragment-based approaches," *OncoTargets and Therapy* 9:2575-2584, 2016. cited by applicant

Patricelli et al., "Selective Inhibition of Oncogenic KRAS Output with Small Molecules Targeting the Inactive State," *Cancer Discovery* 6(3)316-329, 2016. cited by applicant

Pautsch et al., "Crystal structure of the C3bot-RaIA complex reveals a novel type of action of a bacterial exoenzyme," *The EMBO Journal* 24:3670-3680, 2005. cited by applicant

Pédeboscq et al., "Synthesis and evaluation of apoptosis induction of thienopyrimidine compounds on KRAS and BRAF mutated colorectal cancer cell lines," *Bioorganic& Medicinal Chemistry* 20:6724-6731, 2012. cited by applicant

Peri et al., "Arabinose-derived bicyclic amino acids: synthesis, conformational analysis and construction of an α .SUB.v. β .SUB.3.-selective RGD peptide," *J. Am. Chem. Soc., Perkins Trans* 1(5):638-644, Feb. 2002. cited by applicant

Peri et al., "Sugar-Derived Ras Inhibitors: Group Epitope Mapping by NMR Spectroscopy and Biological Evaluation," *Eur. J. Org. Chem.* 2006 (16):3707-3720, Aug. 2006. cited by applicant

Peri et al., "Synthesis of bicyclic sugar azido acids and their incorporation in cyclic peptides," *Chem. Commun.* 23:2303-2304, Jan. 2000. cited by applicant

Pinedo et al., "Aggressive combination therapy to cure patients with metastatic cancer," *The Lancet Oncology* 1:72-73, Oct. 2000. cited by applicant

Pinedo et al., "Translational Research: The Role of VEGF in Tumor Angiogenesis," *The Oncologist* 5(suppl. 1):1-2, 2000. cited by applicant

Polinsky, "High-Speed Chemistry Libraries: Assessment of Drug-Likeness," in Wermuth (ed.), *The Practice of Medicinal Chemistry*, 2nd Ed., Academic Press, Cambridge, Massachusetts, USA, 2003, pp. 147-157. cited by applicant

PubChem CID 10375614, Create Date: Oct. 25, 2006, date accessed Sep. 18, 2017. cited by applicant

PubChem Compound "(Z)-1-(4-(5-(5-(Trifluoromethyl)-1,2,4-oxadiazol-3-yl)pyridin-2-yl)piperazin-1-yl)but-2-en-1-one" CID: 49702158, Create: Nov. 27, 2010, 4 pages. cited by applicant

PubChem Compound, "AKOS024742141," Nov. 27, 2010, retrieved from <http://pubchem.ncbi.nih.gov/compound/49702158#x304> on Nov. 20, 2015, 12 pages. cited by applicant

PubChem Compound, "(2S,6R)-hexahydrofuro[3,2-b]furan-2,6-diyl dicarbonochloridate:Compound Summary CID 53396983," Oct. 30, 2011, retrieved from <http://pubchem.ncbi.nlm.nih.gov/compound/53396983>, 6 pages. cited by applicant

PubChem Compound, "(4-hydroxypiperidin-1-yl)-pyridin-4-ylmethanone: AC1LGBNJ," retrieved on Feb. 17, 2014, from <http://pubchem.ncbi.nlm.nih.gov/summary/summary.cgi?cid=76837>, Jul. 8, 2005, 5 pages. cited by applicant

PubChem Compound, "Compound Summary for CID 21765509: Molecular Formula C18H21N5O8," Dec. 5, 2007, retrieved from <http://pubchem.ncbi.nlm.nih.gov/compound/21765509>, 4 pages. cited by applicant

PubChem Compound, "Compound Summary for CID 21765511: Molecular Formula C18H21N5O8," Dec. 5, 2007, retrieved from <http://pubchem.ncbi.nlm.nih.gov/compound/21765511>, 4 pages. cited by applicant

PubChem Compound, "Compound Summary for CID 60018735: Molecular Formula C30H30O13," Aug. 20, 2012, retrieved from <http://pubchem.ncbi.nlm.nih.gov/compound/60018735#section=Top>, 1 page. cited by applicant

PubChem Compound, "Compound Summary for CID 72623693: AGN-PC-0D83J7," Jan. 9, 2014, retrieved from <http://pubchem.ncbi.nlm.nih.gov/compound/72623693>, 6 pages. cited by applicant

PubChem Compound, "Compound Summary for CID 9897840: Molecular Formula C50H46O20," Oct. 25, 2006, retrieved from <http://pubchem.ncbi.nlm.nih.gov/compound/9897840>, 6 pages. cited by applicant

PubChem Compound, "NJUUFNUHACGHX-UHFFFAOYSA-N," Sep. 5, 2014, retrieved from <https://pubchem.ncbi.nlm.nih.gov/compound/77081212>, CID 77081212, 6 pages. cited by applicant

PubChem Compound, "SCHEMBL6674271," Dec. 1, 2012, retrieved from <http://pubchem.ncbi.nlm.nih.gov/compound/69861127#x304>, CID 69861127, 12 pages. cited by applicant

PubChem Compound, "SCHEMBL6797439," Dec. 1, 2012, retrieved from <http://pubchem.ncbi.nlm.nih.gov/compound/69898605#x304>, Cid 69898605, 12 pages. cited by applicant

PubChem Substance Record for SID 22405303, Mar. 5, 2007, CID 2579941 (MLS000416491), retrieved from <https://pubchem.ncbi.nlm.nih.gov/substance/22405303>, on May 15, 2017, 7 pages. cited by applicant

PubChem Substance Record for SID 44253980, Dec. 5, 2007, CID 966800 (1-BENZOYLPYRROLIDINE), retrieved from

<https://pubchem.ncbi.nlm.nih.gov/substance/44253980#section=Top> on May 11, 2017, 5 pages. cited by applicant

PubChem, Compound Summary for CID 53064614: Molecular Formula C₂₁H₂₂N₄O₃, Jun. 21, 2011, retrieved from <https://pubchem.ncbi.nlm.nih.gov/compound/53064614>, 11 pages. cited by applicant

PubChem, Compound Summary for CID 53064615: Molecular Formula C₂₁H₂₂N₄O₂S, Jun. 21, 2011, retrieved from <https://pubchem.ncbi.nlm.nih.gov/compound/53064615>, 11 pages. cited by applicant

PubChem, Compound Summary for CID 72080923: Molecular Formula C₂₁H₂₂N₄O₂S, Dec. 2, 2013, retrieved from <https://pubchem.ncbi.nlm.nih.gov/compound/72080923>, 10 pages. cited by applicant

PubChem, Compound Summary for CID 72080924: Molecular Formula C₂₁H₂₂N₄O₃, Dec. 2, 2013, retrieved from <https://pubchem.ncbi.nlm.nih.gov/compound/72080924>, 10 pages. cited by applicant

PubChem, Compound Summary for CID 72080937: Molecular Formula C₂₀H₂₀N₄O₃, Dec. 2, 2013, retrieved from <https://pubchem.ncbi.nlm.nih.gov/compound/72080937>, 10 pages. cited by applicant

Registry: 2-Propen-1-one, 1-[4-[4-(4-ethoxyphenyl)-2-pyrimidinyl]-1-piperaziny]-(CA Index Name), 1 page. cited by applicant

Rensland et al., "Substrate and Product Structural Requirements for Binding of Nucleotides to H-ras p21: The Mechanism of Discrimination between Guanosine and Adenosine Nucleotides," *Biochemistry* 34(2):593-599, 1995. cited by applicant

Sasaki et al., "Selective Formation of Stable Triplexes Including a TA or CG Interrupting Site with New Bicyclic Nucleoside Analogues (WNA)," *J. Am. Chem. Soc.* 126(2):516-528, Jan. 2004. cited by applicant

Schubbert et al., "Biochemical and Functional Characterization of Germ Line KRAS Mutations," *Molecular and Cellular Biology* 27(22):7765-7770, Nov. 2007. cited by applicant

Shima et al., "Discovery of small molecule Ras inhibitors that display antitumor activity by interfering with Ras, GTP-effector interaction," *Enzymes* 34(pt. b):1-23, 2013. cited by applicant

Singh et al., "A Gene Expression Signature Associated with "K-Ras Addiction" Reveals Regulators of EMT and Tumor Cell Survival," *Cancer Cell* 15:489-500, Jun. 2009. cited by applicant

Spiegel et al., "Small-molecule modulation of Ras signaling," *Nature Chemical Biology* 10:613-622, Aug. 2014. cited by applicant

Stefanachi et al., "1-, 3-, and 8-substituted-9-deazaxanthines as potent and selective antagonists at the human A.SUB.2B .adenosine receptor," *Bioorg Med Chem* 16(6):2852-2869, 2008. cited by applicant

STN Registry No. 1066881-73-9, "[1,4'-Bipiperidine]-4-carboxamide, 1'-(1-cyclohexen-1-ylcarbonyl)-N-(3-pyridinylmethyl)-," Oct. 27, 2008, 1 page. cited by applicant

STN Registry No. 1066896-20-5, "Methanone, [3-[4-(1,3-benzodioxo1-5-ylmethyl)-1-piperazinyl]-1-piperidinyl]-1-cyclopenten-1-yl-," Oct. 27, 2008, 1 page. cited by applicant

STN Registry No. 1066915-72-7, "[3-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]-1-piperidinyl]-1-cyclohexen-1-yl-," Oct. 27, 2008, 1 page. cited by applicant

STN Registry No. 1069510-87-7, "[1,4'-Bipiperidine]-4-carboxamide, 1'-(1-cyclohexen-1-ylcarbonyl)-N-(2-pyridinylmethyl)-," Nov. 2, 2008, 1 page. cited by applicant

STN Registry No. 1069530-02-4, "[1,4'-Bipiperidine]-4-carboxamide, 1'-(1-cyclopenten-1-ylcarbonyl)-N-(2-pyridinylmethyl)-," Nov. 2, 2008, 1 page. cited by applicant

STN Registry No. 1069736-48-6, "[1,4'-Bipiperidine]-4-carboxamide, 1'-(1-cyclopenten-1-ylcarbonyl)-N-(3-pyridinylmethyl)-," Nov. 2, 2008, 1 page. cited by applicant

STN Registry No. 1069909-93-8, "[1,4'-Bipiperidine]-4-carboxamide, 1'-(1-cyclopenten-1-ylcarbonyl)-N-(2-phenylethyl)-," Nov. 2, 2008, 1 page. cited by applicant

STN Registry No. 1212098-43-5, "Methanone, 1-cyclohexen-1-yl [(3R,4R)-3-hydroxy-4-[4-(2-phenylethyl)-1-piperazinyl]-1-piperidinyl]-, rel-," Mar. 21, 2010, 1 page. cited by applicant

STN Registry No. 1309145-12-7, "Methanone, 1-cyclopenten-1-yl [(3R,4R)-3-hydroxy-4-[4-(2-phenylethyl)-1-piperazinyl]-1-piperidinyl]-, rel-," Jun. 13, 2011, 1 page. cited by applicant

Streuff et al., "First asymmetric aminohydroxylation of acrylamides," *Tetrahedron: Asymmetry* 16 (21):3492-3496, Oct. 2005. cited by applicant

Sun et al., "Discovery of small molecules that bind to K-Ras and inhibit Sos-mediated activation," *Angew Chem Int Ed Engl.* 51(25):6140-6143, 2012. cited by applicant

Sunaga et al., "Knockdown of Oncogenic KRAS in Non-Small Cell Lung Cancers Suppresses Tumor Growth and Sensitizes Tumor Cells to Targeted Therapy," *Molecular Cancer Therapeutics* 10(2):336-346, Feb. 2011. cited by applicant

Sundberg et al., "The O-Styrylnitrene Route to 2-Substituted Indoles. Pyrolysis of O-Azidostyrenes," *Journal of Organic Chemistry* 37(5):719-724, 1972. cited by applicant

Sydor et al., "Transient Kinetic Studies on the Interaction of Ras and the Ras-Binding Domain of c-Raf-1 Reveal Rapid Equilibration of the Complex," *Biochemistry* 37:14292-14299, 1998. cited by applicant

Taveras et al., "Ras Oncoprotein Inhibitors: The Discovery of Potent, Ras Nucleotide Exchange Inhibitors and the Structural Determination of a Drug-Protein Complex," *Bioorganic and Medicinal Chemistry* 5(1):125-133, 1997. cited by applicant

Terfloth et al., "Electronic Screening: Lead Finding from Database Mining, " in Wermuth (ed.), *The Practice of Medicinal Chemistry*, 2nd Ed., Academic Press, Cambridge, Massachusetts, USA, 2003, pp. 131-145. cited by applicant

Tsubaki et al., "Reduction of metastasis, cell invasion, and adhesion in mouse osteosarcoma by YM529/ONO-5920-induced blockade of the Ras/MEK/ERK and Ras/PI3K/Akt pathway," *Toxicology and Applied Pharmacology* 259(3):402-410, Jan. 2012. cited by applicant

Tulshian et al., "Synthesis and Phosphodiesterase Activity of Carboxylic Acid Mimetics of Cyclic Guanosine 3',5'-Monophosphate," *J. Med. Chem.* 36(9):1210-1220, Jan. 1993. cited by applicant

Vetter et al., "The Guanine Nucleotide-Binding Switch in Three Dimensions," *Science* 294(5545):1299-1304, Nov. 2001. cited by applicant

Vippagunta et al., "Crystalline solids," *Advanced Drug Delivery Reviews* 48:3-26, 2001. cited by applicant

Wolff, (ed.), *Burger's Medicinal Chemistry and Drug Discovery, Fifth Edition, vol. 1: Principles and Practice*, San Diego, California, John Wiley & Sons, 1994, pp. 975-977. cited by applicant

Wu et al., "Stereoselective synthesis of dioxabicycles from 1-C-allyl-2-O-benzyl-glycosides—An intramolecular cyclization between 2-O-benzyl oxygen and the allyl double bond," *Can. J. Chem.* 84 (1):597-602, Jan. 2006. cited by applicant

Xu et al., "Design, Synthesis, and Biological Evaluation of 2-Oxo-3,4-dihydropyrimido[4,5-d]pyrimidinyl Derivatives as New Irreversible Epidermal Growth Factor Receptor Inhibitors with Improved Pharmacokinetic Properties," *Journal of Medicinal Chemistry* 56:8803-8813, 2013. cited by applicant

Yan et al., "Discovery and characterization of small molecules that target the GTPase Ral," *Nature* 515:443-447, Nov. 2014, 15 pages. cited by applicant

Yang et al., "Fragment-Based Discovery of Nonpeptidic BACE-1 Inhibitors Using Tethering," *Biochemistry* 48:4488-4496, 2009. cited by applicant

Young et al., "Oncogenic and Wild-type Ras Play Divergent Roles in the Regulation of Mitogen-Activated Protein Kinase Signaling," *Cancer Discovery* 3(1):112-123, Jan. 2013. cited by applicant

Zenkl et al., "Sugar-Responsive Fluorescent Nanospheres," *Macromol. Biosci.* 8:146-152, 2008. cited by applicant

U.S. Appl. No. 16/141,775, filed Sep. 25, 2018. cited by applicant

U.S. Appl. No. 16/281,905, filed Feb. 21, 2019. cited by applicant

U.S. Appl. No. 16/374,356, filed Apr. 3, 2019. cited by applicant
U.S. Appl. No. 16/421,186, filed May 23, 2019. cited by applicant
U.S. Appl. No. 16/445,683, filed Jun. 16, 2019. cited by applicant
U.S. Appl. No. 16/449,967, filed Jun. 24, 2019. cited by applicant
U.S. Appl. No. 16/480,195, filed Jul. 23, 2019. cited by applicant
U.S. Appl. No. 16/480,205, filed Jul. 23, 2019. cited by applicant
U.S. Appl. No. 16/480,211, filed Jul. 23, 2019. cited by applicant
U.S. Appl. No. 16/480,849, filed Jul. 25, 2019. cited by applicant
U.S. Appl. No. 16/480,861, filed Jul. 25, 2019. cited by applicant
U.S. Appl. No. 16/480,874, filed Jul. 25, 2019. cited by applicant
U.S. Appl. No. 16/514,675, filed Jul. 17, 2019. cited by applicant
U.S. Appl. No. 16/535,999, filed Aug. 8, 2019. cited by applicant
Boys et al., "Discovery of a series of 2-(1H-pyrazol-1-yl)pyridines as ALK5 inhibitors with potential utility in the prevention of dermal scarring," *Bioorganic & Medicinal Chemistry Letters* 22:3392-3397, 2012. cited by applicant
Canon et al., "The Clinical KRAS (G12C) Inhibitor AMG 510 Drives Anti-Tumor Immunity," *Nature* 575:217-223, 2019 (26 Pages). cited by applicant
Gura, "Cancer Models: Systems for Identifying New Drugs Are Often Faulty," *Science* 278(5340):1041-1042, 1997. cited by applicant
Horig et al., "From bench to clinic and back: perspective on the 1st IQPC translational research conference," *Journal of Translational Medicine* 2(44):1-8, 2004. cited by applicant
Jayanthi et al., "Biological Activities of Few Nitrogen Heterocycles using Pass and Lipophilicity using Adme Boxes," *Pharma Science Monitor: An International Journal of Pharmaceutical Sciences* 3(3, Suppl-1) 2012 (13 Pages). cited by applicant
Johnson et al., "Relationships between drug activity in NCI preclinical in vitro and in vivo models and early clinical trials," *Br. J. Cancer* 84(10):1424-1431, 2001. cited by applicant
Keith et al., "Heteroarylureas with spirocyclic diamine cores as inhibitors of fatty acid amide hydrolase," *Bioorganic & Medicinal Chemistry Letters* 24:737-741, 2014. cited by applicant
Liu et al., "Targeting the untargetable KRAS in cancer therapy," *Acta Pharmaceutica Sinica B* 9(5):871-879, 2019. cited by applicant
Pearce et al., "Chapter 18: Failure modes in anticancer drug discovery and development," *Cancer Drug Design and Discovery*, Ed. Stephen Neidle, pp. 424-435, 2008. cited by applicant
Reilly et al., "Examination of Diazaspiro Cores as Piperazine Bioisosteres in the Olaparib Framework Shows Reduced DNA Damage and Cytotoxicity," *J. Med. Chem.* 61:5367-5379, 2018. cited by applicant
Rytting, "Acute Leukemia," *Merck Manual*, Online Edition, retrieved Jul. 10, 2013, 6 pages. cited by applicant
Schafer et al., "Failure is an option: learning from unsuccessful proof-of-concept trials," *Drug Discovery Today* 13(21/22):913-916, 2008. cited by applicant
Silverman, "Prodrugs and Drug Delivery Systems," *The Organic Chemistry of Drug Design and Drug Action*, p. 352-399, 1992. cited by applicant
Simone, "Oncology: Introduction," *Cecil Textbook of Medicine*, Ed. J. Claude Bennett and Fred Plum, 20th Edition, vol. 1, pp. 1004-1010, 1996. cited by applicant
STN Registry No. 1027262-22-1, "Pentanoic acid, 5-fluoro-4-oxo-3-[[[(2S)-1-oxo-2-(4-oxo-6-phenyl-3(4H)-quinazolinyl)butyl]amino]-," Jun. 11, 2008, 1 page. cited by applicant
STN Registry Nos. 1179043-51-6; 1156698-93-9; 1070748-55-8; 1; 1070685-77-6; 1070667-53-6; 1070315-89-7; 1069961-90-5; 1069566-58-0; 1065560-16-8; 1065554-88-2; 1061025-62-4; 1060435-61-1; 958948-04-4; 958839-16-2, 6 pages. cited by applicant
Zheng et al., "The use of spirocyclic scaffolds in drug discovery," *Bioorganic & Medicinal*

Primary Examiner: Diamond; Alan D

Attorney, Agent or Firm: Seed IP Law Group LLP

Background/Summary

BACKGROUND

Technical Field

(1) Ras represents a group of closely related monomeric globular protein of 189 amino acids (21 kDa molecular mass) which is associated with the plasma membrane and which binds either GDP or GTP. Ras acts as a molecular switch. When Ras contains bound GDP it is in the resting or off position and is “inactive”. In response to exposure of the cell to certain growth promoting stimuli, Ras is induced to exchange its bound GDP for a GTP. With GTP bound, Ras is “switched on” and is able to interact with and activate other proteins (its “downstream targets”). The Ras protein itself has a very low intrinsic ability to hydrolyze GTP back to GDP, thus turning itself into the off state. Switching Ras off requires extrinsic proteins termed GTPase-activating proteins (GAPs) that interact with Ras and greatly accelerate the conversion of GTP to GDP. Any mutation in Ras which affects its ability to interact with GAP or to convert GTP back to GDP will result in a prolonged activation of the protein and consequently a prolonged signal to the cell telling it to continue to grow and divide. Because these signals result in cell growth and division, overactive Ras signaling may ultimately lead to cancer.

(2) Structurally, Ras proteins contain a G domain which is responsible for the enzymatic activity of Ras—the guanine nucleotide binding and the hydrolysis (GTPase reaction). It also contains a C-terminal extension, known as the CAAX box, which may be post-translationally modified and is responsible for targeting the protein to the membrane. The G domain is approximately 21-25 kDa in size and it contains a phosphate binding loop (P-loop). The P-loop represents the pocket where the nucleotides are bound in the protein, and this is the rigid part of the domain with conserved amino acid residues which are essential for nucleotide binding and hydrolysis (Glycine 12, Threonine 26 and Lysine 16). The G domain also contains the so called Switch I (residues 30-40) and Switch II (residues 60-76) regions, both of which are the dynamic parts of the protein which are often represented as the “spring-loaded” mechanism because of their ability to switch between the resting and loaded state. The key interaction is the hydrogen bonds formed by Threonine-35 and glycine-60 with the γ -phosphate of GTP which maintain Switch 1 and Switch 2 regions respectively in their active conformation. After hydrolysis of GTP and release of phosphate, these two relax into the inactive GDP conformation.

(3) The most notable members of the Ras subfamily are HRAS, KRAS and NRAS, mainly for being implicated in many types of cancer. However, there are many other members including DIRAS1; DIRAS2; DIRAS3; ERAS; GEM; MRAS; NKIRAS1; NKIRAS2; NRAS; RALA; RALB; RAP1A; RAP1B; RAP2A; RAP2B; RAP2C; RASD1; RASD2; RASL10A; RASL10B; RASL11A; RASL11B; RASL12; REM1; REM2; RERG; RERGL; RRAD; RRAS; RRAS2.

(4) Mutations in any one of the three main isoforms of RAS (H-Ras, N-Ras, or K-Ras) genes are among the most common events in human tumorigenesis. About 30% of all human tumors are found to carry some mutation in Ras genes. Remarkably, K-Ras mutations are detected in 25-30% of tumors. By comparison, the rates of oncogenic mutation occurring in the N-Ras and H-Ras family members are much lower (8% and 3% respectively). The most common K-Ras mutations

are found at residue G12 and G13 in the P-loop and at residue Q61.

(5) G12C is a frequent mutation of K-Ras gene (glycine-12 to cysteine). This mutation had been found in about 13% of cancer occurrences, about 43% of lung cancer occurrences, and in almost 100% of MYH-associated polyposis (familial colon cancer syndrome). However targeting this gene with small molecules is a challenge. Accordingly, there is a need in the art for small molecules for targeting Ras (e.g., K-Ras, H-Ras and/or N-Ras) and use of the same for treatment of various diseases, such as cancer. The present invention provides these and other related advantages.

BRIEF SUMMARY

(6) The present invention provides compounds which are capable of modulating G12C mutant K-Ras, H-Ras and/or N-Ras proteins. In some instances, the compound acts as an electrophile capable of forming a covalent bond with the cysteine residue at position 12 of a K-Ras, H-Ras or N-Ras G12C mutant protein.

(7) In some aspects of the invention, the compounds described herein are included in pharmaceutical compositions. In some embodiments, the pharmaceutical composition comprises a pharmaceutically acceptable carrier. In some aspects of the invention, the pharmaceutical composition is suitable for oral administration. In some embodiments, the pharmaceutical composition is suitable for injection.

(8) In one aspect, a method is provided. The method comprises a method of regulating activity of a K-Ras, H-Ras or N-Ras G12C mutant protein wherein the method comprises reacting the K-Ras, H-Ras or N-Ras G12C mutant protein with any of the compounds described herein. In some embodiments, the method inhibits proliferation of a cell population by contacting the cell population with any of the compounds described herein. In some embodiments, the method of inhibiting proliferation of a cell is measured as a decrease in cell viability of the cell population.

(9) In one aspect, a method of treating a disorder in a subject is provided. The method of treating a disorder in a subject comprises: (a) determining if the subject has a K-Ras, H-Ras or N-Ras G12C mutation; and (b) if the subject is determined to have the K-Ras, H-Ras or N-Ras G12C mutation then administering to the subject a therapeutically effective dose of a pharmaceutical composition comprising at least one compound described herein. In some embodiments, the disorder is a cancer. In some embodiments, the cancer is pancreatic cancer, colon cancer, MYH associated polyposis, colorectal cancer, lung cancer or NSCLC.

(10) In one aspect, a method of preparing a labeled K-Ras, H-Ras or N-Ras G12C mutant protein is provided. The method of preparing a labeled K-Ras, H-Ras or N-Ras G12C mutant comprises reacting the K-Ras, H-Ras or N-Ras G12C mutant with a compound described herein, resulting in a labeled K-Ras, H-Ras or N-Ras G12C protein.

(11) In one aspect, a compound of Formula I having is provided:

(12) ##STR00001##

(13) wherein X is O or NH, S, or CR.sub.23, R.sub.24; Y is CH.sub.2, CHR.sub.22, CO, SO, or SO.sub.2; n is an integer of value 1-6; R1 is aryl or heteroaryl, each of which is unsubstituted or substituted by one or more independent R2 substituents; R2 is halogen, —OH, oxo, alkoxy, alkyl, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene, each of which is unsubstituted or substituted by one or more independent R3 substituents; R3 is halogen, OH, cyano, alkyl, alkoxy, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene, each of which is unsubstituted or substituted by one or more independent R4 substituents; R4 is halogen, OH, cyano, alkyl, alkoxy, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene; C1 is alkyl, cycloalkyl, heterocycloalkyl, arylene, heteroarylene, or heterocycloalkylene; each of which may be substituted with a R5 substituent; C2 is a bond, cycloalkyl, heterocycloalkyl, arylene, heteroarylene, cycloalkylene, or heterocycloalkylene, wherein C, 1 and C2 may form a fused or spiro bicyclic ring; D is a bond, —NH—CH2—, —NH—, or —CH2—; R5 is OH, alkyl, or —CH2OH; and E is an electrophile capable of forming a covalent bond with the cysteine residue at position 12 of a K-Ras G12C mutant

protein.

(14) In some embodiments, the compound of Formula 1 has an O at position X. In other embodiments, the X represents a NH group.

(15) In some embodiments, the compound of Formula 1 has a bond at position D. In other embodiments, D is a NH group. In other embodiments, the compound of Formula 1 has —NH—CH₂- at position D connected to —NH—CH₂- connected to the carbonyl carbon.

(16) In some embodiments, the compound of Formula 1 has a group at position E that is selected from the group consisting of:

(17) ##STR00002##

(18) In some embodiments, the compound of Formula 1 has a phenyl moiety is at position R₁ and the phenyl moiety position R₁ is unsubstituted or substituted by one or more independent R₂ substituents. In other embodiments, the compound of Formula 1 has a benzothiadiazolyl moiety is at position R₁ and is unsubstituted or substituted by one or more independent R₂ substituents. In other embodiments, the compound of Formula 1 has a naphthalenyl moiety is at position R₁ and is unsubstituted or substituted by one or more independent R₂ substituents. In other embodiments, the compound of Formula 1 has an imidazopyridinyl moiety is at position R₁ and is unsubstituted or substituted by one or more independent R₂ substituents.

(19) In some embodiments, the compound of Formula 1 has a halogen at position R₂. In other embodiments, the compound of Formula 1 has an OH at position R₂. In other embodiments, the compound of Formula 1 has an OMe at position R₂. In other embodiments, the compound of Formula 1 has an aryl or heteroaryl moiety at position R₂ that is unsubstituted or substituted by one or more independent R₃ substituents. In other embodiments, the compound of Formula 1 has a phenyl, pyridinyl, or thiophenyl moiety at position R₂ that is unsubstituted or substituted by one or more independent R₃ substituents.

(20) In some embodiments, the compound of Formula 1 has a fused bicyclic ring structure at the —C₁-C₂- position. In some embodiments, the compound of Formula 1 has a

(21) ##STR00003##

at the —C₁-C₂ position. In other embodiments, the compound of Formula 1 has a

(22) ##STR00004##

at the —C₁-C₂ position. In other embodiments, the compound of Formula 1 has

(23) ##STR00005##

(24) at the —C₁-C₂ position.

(25) In some embodiments, the compound of Formula 1 has group at C.sub.1 that is selected from the group consisting of:

(26) ##STR00006##

(27) In some embodiments, the compound of Formula 1 has group at C.sub.2 that is selected from the group consisting of:

(28) ##STR00007##

(29) In some embodiments, the compound of Formula 1 is selected from the compounds shown in Table 1.

(30) In another aspect, a composition is provided; the composition comprises a compound of Formula II having the structure of

(31) ##STR00008##

wherein A is —CH₂-, —O—, or —NH—; R₆ is aryl or heteroaryl, each of which is unsubstituted or substituted by one or more independent R₇ substituents; R₇ is halogen, —OH, OR₁₀, NR₁₁R₁₂, alkyl, oxo, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene, each of which is unsubstituted or substituted by one or more independent R₈ substituents; R₈ is halogen, OH, cyano, alkyl, alkoxy, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene, each of which is unsubstituted or substituted by one or more independent R₉ substituents; R₁₀, R₁₁ and R₁₂ are independently hydrogen, alkyl, cycloalkyl, heteroalkyl,

heterocycloalkyl, aryl, or heteroaryl, each of which is unsubstituted or substituted by one or more independent R13 substituents; R9 and R13 are independently halogen, OH, cyano, alkyl, alkoxy, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene; E is an electrophile capable of forming a covalent bond with the cysteine residue at position 12 of a K-Ras G12C mutant protein.

(32) In some embodiments, the E of the Formula II compound represents

(33) ##STR00009##

(34) In some embodiments, the R1 of the Formula II compound represents an aryl, unsubstituted or substituted by one or more independent R2 substituents. In other embodiments, the R1 of the Formula II compound is a heteroaryl, unsubstituted or substituted by one or more independent R2 substituents. In some embodiments, the R1 of the Formula II compound represents a phenyl moiety unsubstituted or substituted by one or more independent R2 substituents. In some embodiments, the R1 of the Formula II compound represents a pyridinyl moiety unsubstituted or substituted by one or more independent R2 substituents.

(35) In some embodiments, the A of the Formula II compound is O. In other embodiments, the A of the Formula II compound is NH.

(36) In some embodiments, the Formula II compound is selected from the compounds shown in Table 2.

(37) In another aspect, a compound is provided; the compound of Formula III having the structure of:

(38) ##STR00010##

(39) wherein A is a bond, O, NH, or —C(O)— ; n is 0 or 2; R14 is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, each of which is unsubstituted or substituted by one or more independent R15 substituents; R15 is halogen, OR18, NR19R20, oxo, alkyl, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene, each of which is unsubstituted or substituted by one or more independent R16 substituents; R16 is halogen, OH, alkyl, alkoxy, cyano, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene, each of which is unsubstituted or substituted by one or more independent R17 substituents; R18, R19 and R20 are independently hydrogen, alkyl, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene, each of which is unsubstituted or substituted by one or more independent R21 substituents; R17 and R21 are independently halogen, OH, cyano, alkyl, alkoxy, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene; C1 is cycloalkyl or heterocycloalkyl; C2 is a bond, cycloalkyl, heterocycloalkyl, arylene, or heteroarylene; wherein C1 and C2 may form a fused or spiro bicyclic ring; and E is an electrophile capable of forming a covalent bond with the cysteine residue at position 12 of a K-Ras G12C mutant protein.

(40) In some embodiments, the A of the Formula III compound represents O. In other embodiments, the A of the Formula III compound represents —C(O)— . In other embodiments, the A of the Formula III compound represents a bond.

(41) In some embodiments, the n of the Formula III compound represents 0. In other embodiments, the n of the Formula III compound represents 1. In other embodiments, the n of the Formula III compound represents 2.

(42) In some embodiments, the C2 of the Formula III compound represents a bond. In some embodiments, the C2 of the Formula III compound represents a pyridinyl.

(43) In some embodiments the C1 of the Formula III compound is selected from the group consisting of:

(44) ##STR00011##

(45) In some embodiments, the R1 of the Formula III compound is an aryl or heteroaryl moiety unsubstituted or substituted by one or more independent R2 substituents. In other embodiments, the R1 of the Formula III compound is a phenyl moiety unsubstituted or substituted by one or more independent R2 substituents. In other embodiments, the R1 of the Formula III compound is

(46) ##STR00012##

In other embodiments, R1 of the Formula III compound represents

(47) ##STR00013##

In other embodiments, R1 represents a pyridinyl moiety unsubstituted or substituted by one or more independent R2 substituents. In other embodiments, the R1 of the Formula III compound represents

(48) ##STR00014##

In other embodiments, the R1 of the Formula III compound represents

(49) ##STR00015##

(50) In some embodiments, the R2 of the Formula III compound represents a halogen. In other embodiments, the R2 of the Formula III compound represents an alkyl.

(51) In other embodiments, the R2 of the Formula III compound represents a —CH₃.

(52) In some embodiments, the E of the Formula III compound represents:

(53) ##STR00016##

(54) In some embodiments, the Formula III compound is selected from the compounds shown in Table 3.

(55) In another aspect, a composition is provided; the composition comprises a compound shown in Table 4.

(56) In other embodiments, a compound of structure (V) is provided:

(57) ##STR00017##

or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein R^{sup.1}, L^{sup.1}, L^{sup.2}, E, G^{sup.1}, G^{sup.2}, G^{sup.3}, G^{sup.4}, R^{sup.30a}, R^{sup.30b}, R^{sup.31a}, R^{sup.31b}, R^{sup.32a}, R^{sup.32b}, R^{sup.33a}, R^{sup.33b}, n^{sup.1} n^{sup.2} n^{sup.3} and n^{sup.4} are as defined herein.

(58) Also provided in various different embodiments is a compound of structure (VI):

(59) ##STR00018##

or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein A, B, C, L^{sup.1a}, L^{sup.2}, E, G^{sup.3}, G^{sup.4}, R^{sup.32a}, R^{sup.32b}, R^{sup.33a}, R^{sup.33b}, R^{sup.37a}, R^{sup.37e}, n^{sup.3} n^{sup.4} and m are as defined herein,

(60) These and other aspects of the invention will be apparent upon reference to the following detailed description. To this end, various references are set forth herein which describe in more detail certain background information, procedures, compounds and/or compositions, and are each hereby incorporated by reference in their entirety.

Description

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

(1) The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

(2) FIG. 1 shows the enzymatic activity of Ras.

(3) FIG. 2 shows a signal transduction pathway for Ras.

(4) FIG. 3 shows some common oncogenes, their respective tumor type and cumulative mutation frequencies (all tumors).

(5) FIG. 4 shows the results of cell potency assay for the compound I-189.

(6) FIG. 5 shows the results of comparison of cell potency assay for the compound I-189, I-92 and I-94.

(7) FIG. 6 shows the results of comparison of cell potency assay for the compound I-92 and I-95.

(8) FIG. 7 shows the results of comparison of cell potency assay for the compound I-66, I-45 and I-91.

(9) FIG. 8 shows the results of comparison of cell potency assay for the compound I-47, I-42 and I-60.

DETAILED DESCRIPTION

(10) Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which this invention belongs. All patents and publications referred to herein are incorporated by reference.

(11) As used in the specification and claims, the singular form “a”, “an” and “the” include plural references unless the context clearly dictates otherwise.

(12) “Amino” refers to the —NH radical.

(13) “Carboxy” or “carboxyl” refers to the —CO radical.

(14) “Cyano” refers to the —CN radical.

(15) “Hydroxy” or “hydroxyl” refers to the —OH radical.

(16) “Imino” refers to the =NH substituent.

(17) “Nitro” refers to the —NO radical.

(18) “Oxo” refers to the =O substituent.

(19) “Thioxo” refers to the =S substituent.

(20) “Alkyl” refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, which is saturated or unsaturated (i.e., contains one or more double and/or triple bonds), having from one to twelve carbon atoms (C₁-C₁₂ alkyl), preferably one to eight carbon atoms (C₁-C₈ alkyl) or one to six carbon atoms (C₁-C₆ alkyl), and which is attached to the rest of the molecule by a single bond, e.g., methyl, ethyl, n-propyl, 1-methylethyl (iso-propyl), n-butyl, n-pentyl, 1,1-dimethylethyl (t-butyl), 3-methylhexyl, 2-methylhexyl, ethenyl, prop-1-enyl, but-1-enyl, pent-1-enyl, penta-1,4-dienyl, ethynyl, propynyl, butynyl, pentynyl, hexynyl, and the like. Alkyl includes alkenyls (one or more carbon-carbon double bonds) and alkynyls (one or more carbon-carbon triple bonds). Unless stated otherwise specifically in the specification, an alkyl group is optionally substituted.

(21) “Alkylene” or “alkylene chain” refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, which is saturated or unsaturated (i.e., contains one or more double and/or triple bonds), and having from one to twelve carbon atoms, e.g., methylene, ethylene, propylene, n-butylene, ethenylene, propenylene, n-butenylene, propynylene, n-butynylene, and the like. The alkylene chain is attached to the rest of the molecule through a single or double bond and to the radical group through a single or double bond. The points of attachment of the alkylene chain to the rest of the molecule and to the radical group can be through one carbon or any two carbons within the chain. Unless stated otherwise specifically in the specification, an alkylene chain is optionally substituted.

(22) “Alkenylene” is an alkylene, as defined above, which comprises one or more carbon-carbon double bonds. Unless stated otherwise specifically in the specification, an alkenylene is optionally substituted.

(23) “Alkylenecarbonyl” refers to a radical of the formula —C(=O)R , where R is an alkylene chain as defined above. Unless stated otherwise specifically in the specification, an alkylenecarbonyl is optionally substituted.

(24) “Alkenylenecarbonyl” refers to an alkylenecarbonyl, as defined above, which comprises one or more carbon-carbon double bonds. Unless stated otherwise specifically in the specification, an alkenylenecarbonyl is optionally substituted.

(25) “Alkoxy” refers to a radical of the formula —OR where R is an alkyl radical as defined above containing one to twelve carbon atoms. Unless stated otherwise specifically in the specification, an alkoxy group is optionally substituted.

(26) “Alkylamino” refers to a radical of the formula —NHR or —NR where

each R.sub.a is, independently, an alkyl radical as defined above containing one to twelve carbon atoms. Unless stated otherwise specifically in the specification, an alkylamino group is optionally substituted.

(27) “Aminoalkyl” refers to an alkyl group comprising at least one amino substituent. The amino substituent can be on a tertiary, secondary or primary carbon. Unless stated otherwise specifically in the specification, an aminoalkyl group is optionally substituted.

(28) “Alkylaminoalkyl” refers to an alkyl group comprising at least one alkylamino substituent. The alkylamino substituent can be on a tertiary, secondary or primary carbon. Unless stated otherwise specifically in the specification, an alkylaminoalkyl group is optionally substituted.

(29) “Aminocarbonyl” refers to a radical of the formula $\text{—C(=O)NR.sub.aR.sub.b}$, where R.sub.a and R.sub.b are each independently H or alkyl. Unless stated otherwise specifically in the specification, an alkoxy group is optionally substituted.

(30) “Aryl” refers to a hydrocarbon ring system radical comprising hydrogen, 6 to 18 carbon atoms and at least one aromatic ring. For purposes of this invention, the aryl radical is a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused or bridged ring systems. Aryl radicals include, but are not limited to, aryl radicals derived from aceanthrylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, fluoranthene, fluorene, as-indacene, s-indacene, indane, indene, naphthalene, phenalene, phenanthrene, pleiadene, pyrene, and triphenylene. Unless stated otherwise specifically in the specification, the term “aryl” or the prefix “ar-” (such as in “aralkyl”) is meant to include aryl radicals that are optionally substituted.

(31) “Arylene” refers to a divalent aryl group which links the rest of the molecule (e.g., compound of structure I-VI) to a radical group and/or to the rest of the molecule. Unless stated specifically otherwise, an arylene is optionally substituted.

(32) “Aralkyl” refers to a radical of the formula —R.sub.b—R.sub.e where R.sub.b is an alkylene chain as defined above and R, is one or more aryl radicals as defined above, for example, benzyl, diphenylmethyl and the like. Unless stated otherwise specifically in the specification, an aralkyl group is optionally substituted.

(33) “Carboxyalkyl” refers to a radical of the formula —R.sub.b—R.sub.e where R.sub.b is an alkylene chain as defined above and R.sub.c is a carboxy group as defined above. Unless stated otherwise specifically in the specification, carboxyalkyl group is optionally substituted.

(34) “Cyanoalkyl” refers to a radical of the formula —R.sub.b—R.sub.c where R.sub.b is an alkylene chain as defined above and R.sub.c is a cyano group as defined above. Unless stated otherwise specifically in the specification, a cyanoalkyl group is optionally substituted.

(35) “Cycloalkyl” or “carbocyclic ring” refers to a stable non-aromatic monocyclic or polycyclic hydrocarbon radical consisting solely of carbon and hydrogen atoms, which may include fused or bridged ring systems, having from three to fifteen carbon atoms, preferably having from three to ten carbon atoms, and which is saturated or unsaturated and attached to the rest of the molecule by a single bond. Monocyclic radicals include, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Polycyclic radicals include, for example, adamantyl, norbornyl, decalinyl, 7,7-dimethyl-bicyclo[2.2.1]heptanyl, and the like. A “cycloalkenyl” is a cycloalkyl comprising one or more carbon-carbon double bonds within the ring. Unless otherwise stated specifically in the specification, a cycloalkyl (or cycloalkenyl) group is optionally substituted.

(36) The term “bicycloalkyl” refers to a structure consisting of two cycloalkyl moieties, unsubstituted or substituted, that have two or more atoms in common. If the cycloalkyl moieties have exactly two atoms in common they are said to be “fused”. Examples include, but are not limited to, bicyclo[3.1.0]hexyl, perhydronaphthyl, and the like. If the cycloalkyl moieties have more than two atoms in common they are said to be “bridged”. Examples include, but are not limited to, bicyclo[3.2.1]heptyl (“norbornyl”), bicyclo[2.2.2]octyl, and the like.

(37) As used herein, the term “heteroatom” or “ring heteroatom” is meant to include oxygen (O),

nitrogen (N), sulfur (S), and phosphorus (P).

(38) The term “heteroalkyl,” by itself or in combination with another term, means, unless otherwise stated, a straight or branched chain, or cyclic hydrocarbon radical, or combinations thereof, comprising of at least one carbon atoms and at least one heteroatom selected from the group comprising of O, N, P, Si and S, and wherein the nitrogen, phosphorus, and sulfur atoms may optionally be oxidized and the nitrogen heteroatom may optionally be quaternized. The heteroatom(s) O, N, P and S and Si may be placed at any interior position of the heteroalkyl group or at the position at which alkyl group is attached to the remainder of the molecule. The alkyl portion of the moiety is unsubstituted or substituted.

(39) “Heteroalkylene” refers to an alkylene group comprising at least one heteroatom (e.g., N, O or S). In some embodiments, the heteroatom is within the alkylene chain (i.e., the heteroalkylene comprises at least one carbon-heteroatom-carbon bond. In other embodiments, the heteroatom is at a terminus of the alkylene and thus serves to join the alkylene to the remainder of the molecule (e.g., M1-H-A-M2, where M1 and M2 are portions of the a molecule, H is a heteroatom and A is an alkylene). Unless stated otherwise specifically in the specification, a heteroalkylene is optionally substituted.

(40) “Heteroalkylenecarbonyl” refers to a radical of the formula —C(=O)R.sub.a— , where R.sub.a is a heteroalkylene chain as defined above. Unless stated otherwise specifically in the specification, a heteroalkylenecarbonyl is optionally substituted.

(41) The term “heterobicycloalkyl” refers to a bicycloalkyl structure, which is unsubstituted or substituted, in which at least one carbon atom is replaced with a heteroatom independently selected from oxygen, nitrogen, and sulfur.

(42) The term “spiroalkyl” refers to a structure, which is unsubstituted or substituted, which comprises at least two cycloalkyl units joined at single carbon. In various embodiments the spiroalkyl rings can be 1-18 carbons.

(43) The term “heterospiroalkyl” refers to a spiroalkyl structure, which is unsubstituted or substituted, in which at least one carbon atom is replaced with a heteroatom independently selected from oxygen, nitrogen, and sulfur.

(44) “Cycloalkylalkyl” refers to a radical of the formula —R.sub.bR.sub.d— where R.sub.b is an alkylene chain as defined above and R.sub.d is a cycloalkyl radical as defined above. Unless stated otherwise specifically in the specification, a cycloalkylalkyl group is optionally substituted.

(45) “Fused” refers to any ring structure described herein which is fused to an existing ring structure in the compounds of the invention. When the fused ring is a heterocyclyl ring or a heteroaryl ring, any carbon atom on the existing ring structure which becomes part of the fused heterocyclyl ring or the fused heteroaryl ring is replaced with a nitrogen atom.

(46) “Halo” or “halogen” refers to bromo, chloro, fluoro or iodo.

(47) “Haloalkyl” refers to an alkyl radical, as defined above, that is substituted by one or more halo radicals, as defined above, e.g., trifluoromethyl, difluoromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 1,2-difluoroethyl, 3-bromo-2-fluoropropyl, 1,2-dibromoethyl, and the like. Unless stated otherwise specifically in the specification, a haloalkyl group is optionally substituted.

(48) “Heterocyclyl” or “heterocyclic ring” refers to a stable 3- to 18-membered non-aromatic ring radical which consists of two to twelve carbon atoms and from one to six heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. Unless stated otherwise specifically in the specification, the heterocyclyl radical is a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heterocyclyl radical is optionally oxidized; the nitrogen atom is optionally quaternized; and the heterocyclyl radical is partially or fully saturated. Examples of such heterocyclyl radicals include, but are not limited to, dioxolanyl, thienyl[1,3]dithianyl, decahydroisoquinolyl, imidazolynyl, imidazolidinyl, isothiazolidinyl, isoxazolidinyl, morpholynyl, octahydroindolyl, octahydroisoindolyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, oxazolidinyl,

piperidinyl, piperazinyl, 4-piperidonyl, pyrrolidinyl, pyrazolidinyl, quinuclidinyl, thiazolidinyl, tetrahydrofuryl, trithianyl, tetrahydropyranyl, thiomorpholinyl, thiamorpholinyl, 1-oxo-thiomorpholinyl, and 1,1-dioxo-thiomorpholinyl. Unless stated otherwise specifically in the specification. Unless stated otherwise specifically in the specification, a heterocyclyl group is optionally substituted.

(49) “Heterocycloalkylene” refers to a divalent saturated heterocyclyl group which links the rest of the molecule (e.g., compound of structure I-VI) to a radical group and/or to the rest of the molecule. Unless stated specifically otherwise, a heterocycloalkylene is optionally substituted.

(50) “Heterocycloalkylenecarbonyl” refers to a radical of the formula —R.sub.aC(=O)— , wherein R.sub.a is a heterocycloalkylene as defined above. Unless stated specifically otherwise, a heterocycloalkylenecarbonyl is optionally substituted.

(51) “N-heterocyclyl” refers to a heterocyclyl radical as defined above containing at least one nitrogen and where the point of attachment of the heterocyclyl radical to the rest of the molecule is through a nitrogen atom in the heterocyclyl radical. Unless stated otherwise specifically in the specification, a N-heterocyclyl group is optionally substituted.

(52) “Heterocyclylalkyl” or “heterocycloalkyl” refers to a radical of the formula —R.sub.bR.sub.c where R.sub.b is an alkylene chain as defined above and R.sub.c is a heterocyclyl radical as defined above, and if the heterocyclyl is a nitrogen-containing heterocyclyl, the heterocyclyl is optionally attached to the alkyl radical at the nitrogen atom. Unless stated otherwise specifically in the specification, a heterocyclylalkyl group is optionally substituted.

(53) “Heteroaryl” refers to a 5- to 14-membered ring system radical comprising hydrogen atoms, one to thirteen carbon atoms, one to six heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur, and at least one aromatic ring. For purposes of this invention, the heteroaryl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heteroaryl radical may be optionally oxidized; the nitrogen atom may be optionally quaternized. Examples include, but are not limited to, azepinyl, acridinyl, benzimidazolyl, benzothiazolyl, benzindolyl, benzodioxolyl, benzofuranyl, benzooxazolyl, benzothiazolyl, benzothiadiazolyl, benzo[b][1,4]dioxepinyl, 1,4-benzodioxanyl, benzonaphthofuranyl, benzoxazolyl, benzodioxolyl, benzodioxinyl, benzopyranyl, benzopyranonyl, benzofuranyl, benzofuranonyl, benzothienyl (benzothiophenyl), benzotriazolyl, benzo[4,6]imidazo[1,2-a]pyridinyl, carbazolyl, cinnolyl, dibenzofuranyl, dibenzothiophenyl, furanyl, furanonyl, isothiazolyl, imidazolyl, indazolyl, indolyl, indazolyl, isoindolyl, indolinyl, isoindolinyl, isoquinolyl, indoliziny, isoxazolyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, oxiranyl, 1-oxidopyridinyl, 1-oxidopyrimidinyl, 1-oxidopyrazinyl, 1-oxidopyridazinyl, 1-phenyl-1H-pyrrolyl, phenazinyl, phenothiazinyl, phenoxazinyl, phthalazinyl, pteridinyl, purinyl, pyrrolyl, pyrazolyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, quinazolinyl, quinoxalinyl, quinolinyl, quinuclidinyl, isoquinolinyl, tetrahydroquinolinyl, thiazolyl, thiadiazolyl, triazolyl, tetrazolyl, triazinyl, and thiophenyl (i.e. thienyl). Unless stated otherwise specifically in the specification, a heteroaryl group is optionally substituted.

(54) “Heteroarylene” refers to a divalent saturated heteroaryl group which links the rest of the molecule (e.g., compound of structure I-VI) to a radical group and/or to the rest of the molecule. Unless stated specifically otherwise, a heteroarylene is optionally substituted.

(55) “Heteroarylenecarbonyl” refers to a radical of the formula —R.sub.aC(=O)— , wherein R.sub.a is a heteroarylene as defined above. Unless stated specifically otherwise, a heteroarylenecarbonyl is optionally substituted.

(56) “N-heteroaryl” refers to a heteroaryl radical as defined above containing at least one nitrogen and where the point of attachment of the heteroaryl radical to the rest of the molecule is through a nitrogen atom in the heteroaryl radical. Unless stated otherwise specifically in the specification, an N-heteroaryl group is optionally substituted.

(57) “Heteroarylalkyl” refers to a radical of the formula —R.sub.bR.sub.f where R.sub.b is an

alkylene chain as defined above and R.sub.f is a heteroaryl radical as defined above. Unless stated otherwise specifically in the specification, a heteroarylalkyl group is optionally substituted.

(58) "Hydroxylalkyl" refers to an alkyl group comprising at least one hydroxyl substituent. The —OH substituent may be on a primary, secondary or tertiary carbon. Unless stated otherwise specifically in the specification, a hydroxylalkyl group is optionally substituted.

(59) "Thioalkyl" refers to a radical of the formula —SR.sub.a where R.sub.a is an alkyl radical as defined above containing one to twelve carbon atoms. Unless stated otherwise specifically in the specification, a thioalkyl group is optionally substituted.

(60) The term "substituted" used herein means any of the above groups (e.g., alkyl, alkylene, alkenylene, alkenylenecarbonyl, alkoxy, alkylamino, aminoalkyl, alkylaminoalkyl, thioalkyl, aryl, arylene, aralkyl, carboxyalkyl, cyanoalkyl, cycloalkyl, cycloalkylalkyl, heteroalkyl, heteroalkylene, heteroalkylenecarbonyl, heterobicycloalkyl, spiroalkyl, heterospiroalkyl, haloalkyl, heterocyclyl, heterocycloalkylene, heterocycloalkylenecarbonyl, N-heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylene, heteroarylenecarbonyl, N-heteroaryl, hydroxylalkyl, thioalkyl and/or heteroarylalkyl) wherein at least one hydrogen atom is replaced by a bond to a non-hydrogen atoms such as, but not limited to: a halogen atom such as F, Cl, Br, and I; an oxygen atom in groups such as hydroxyl groups, alkoxy groups, and ester groups; a sulfur atom in groups such as thiol groups, thioalkyl groups, sulfone groups, sulfonyl groups, and sulfoxide groups; a nitrogen atom in groups such as amines, amides, alkylamines, dialkylamines, arylamines, alkylarylamines, diarylamines, N-oxides, imides, and enamines; a silicon atom in groups such as trialkylsilyl groups, dialkylarylsilyl groups, alkylarylsilyl groups, and triarylsilyl groups; and other heteroatoms in various other groups. "Substituted" also means any of the above groups in which one or more hydrogen atoms are replaced by a higher-order bond (e.g., a double- or triple-bond) to a heteroatom such as oxygen in oxo, carbonyl, carboxyl, and ester groups; and nitrogen in groups such as imines, oximes, hydrazones, and nitriles. For example, "substituted" includes any of the above groups in which one or more hydrogen atoms are replaced

(61) with —NR.sub.gR.sub.h, —NR.sub.gC(=O)R.sub.h, —NR.sub.gC(=O)NR.sub.gR.sub.h, —NR.sub.gC(=O)OR.sub.h, —NR.sub.gSO.sub.2R.sub.h, —OC(=O)NR.sub.gR.sub.h, —OR.sub.g, —SR.sub.g, —SOR.sub.g, —SO.sub.2R.sub.g, —OSO.sub.2R.sub.g, —SO.sub.2OR.sub.g, —NSO.sub.2R.sub.g, and —SO.sub.2NR.sub.gR.sub.h. "Substituted also means any of the above groups in which one or more hydrogen atoms are replaced with —C(=O)R.sub.g, —C(=O)OR.sub.g, —C(=O)NR.sub.gR.sub.h, —CH.sub.2SO.sub.2R.sub.g, —CH.sub.2SO.sub.2NR.sub.gR.sub.h. In the foregoing, R.sub.g and R.sub.h are the same or different and independently hydrogen, alkyl, alkoxy, alkylamino, thioalkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, haloalkyl, heterocyclyl, N-heterocyclyl, heterocyclylalkyl, heteroaryl, N-heteroaryl and/or heteroarylalkyl. "Substituted" further means any of the above groups in which one or more hydrogen atoms are replaced by a bond to an amino, cyano, hydroxyl, imino, nitro, oxo, thioxo, halo, alkyl, alkoxy, alkylamino, thioalkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, haloalkyl, heterocyclyl, N-heterocyclyl, heterocyclylalkyl, heteroaryl, N-heteroaryl and/or heteroarylalkyl group. In addition, each of the foregoing substituents is optionally substituted with one or more of the above substituents.

(62) "Electrophile" or "electrophilic moiety" is any moiety capable of reacting with a nucleophile (e.g., a moiety having a lone pair of electrons, a negative charge, a partial negative charge and/or an excess of electrons, for example a —SH group). Electrophiles typically are electron poor or comprise atoms which are electron poor. In certain embodiments an electrophile contains a positive charge or partial positive charge, has a resonance structure which contains a positive charge or partial positive charge or is a moiety in which delocalization or polarization of electrons results in one or more atom which contains a positive charge or partial positive charge. In some embodiments, the electrophiles comprise conjugated double bonds, for example an α,β -unsaturated carbonyl or α,β -unsaturated thiocarbonyl compound.

(63) The term “effective amount” or “therapeutically effective amount” refers to that amount of a compound described herein that is sufficient to effect the intended application including but not limited to disease treatment, as defined below. The therapeutically effective amount may vary depending upon the intended treatment application (in vivo), or the subject and disease condition being treated, e.g., the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of ordinary skill in the art. The term also applies to a dose that will induce a particular response in target cells, e.g. reduction of platelet adhesion and/or cell migration. The specific dose will vary depending on the particular compounds chosen, the dosing regimen to be followed, whether it is administered in combination with other compounds, timing of administration, the tissue to which it is administered, and the physical delivery system in which it is carried.

(64) As used herein, “treatment” or “treating” refer to an approach for obtaining beneficial or desired results with respect to a disease, disorder or medical condition including but not limited to a therapeutic benefit and/or a prophylactic benefit. By therapeutic benefit is meant eradication or amelioration of the underlying disorder being treated. Also, a therapeutic benefit is achieved with the eradication or amelioration of one or more of the physiological symptoms associated with the underlying disorder such that an improvement is observed in the subject, notwithstanding that the subject may still be afflicted with the underlying disorder. In certain embodiments, for prophylactic benefit, the compositions are administered to a subject at risk of developing a particular disease, or to a subject reporting one or more of the physiological symptoms of a disease, even though a diagnosis of this disease may not have been made.

(65) A “therapeutic effect,” as that term is used herein, encompasses a therapeutic benefit and/or a prophylactic benefit as described above. A prophylactic effect includes delaying or eliminating the appearance of a disease or condition, delaying or eliminating the onset of symptoms of a disease or condition, slowing, halting, or reversing the progression of a disease or condition, or any combination thereof.

(66) The term “co-administration,” “administered in combination with,” and their grammatical equivalents, as used herein, encompass administration of two or more agents to an animal, including humans, so that both agents and/or their metabolites are present in the subject at the same time. Co-administration includes simultaneous administration in separate compositions, administration at different times in separate compositions, or administration in a composition in which both agents are present.

(67) “Pharmaceutically acceptable salt” includes both acid and base addition salts.

(68) “Pharmaceutically acceptable acid addition salt” refers to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which are formed with inorganic acids such as, but are not limited to, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as, but not limited to, acetic acid, 2,2-dichloroacetic acid, adipic acid, alginic acid, ascorbic acid, aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, camphoric acid, camphor-10-sulfonic acid, capric acid, caproic acid, caprylic acid, carbonic acid, cinnamic acid, citric acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glucoheptonic acid, gluconic acid, glucuronic acid, glutamic acid, glutaric acid, 2-oxo-glutaric acid, glycerophosphoric acid, glycolic acid, hippuric acid, isobutyric acid, lactic acid, lactobionic acid, lauric acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, mucic acid, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, propionic acid, pyroglutamic acid, pyruvic acid, salicylic acid, 4-aminosalicylic acid, sebacic acid, stearic acid, succinic acid, tartaric acid, thiocyanic acid, p-toluenesulfonic acid, trifluoroacetic acid, undecylenic acid, and the like.

(69) “Pharmaceutically acceptable base addition salt” refers to those salts which retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise undesirable. These salts are prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Preferred inorganic salts are the ammonium, sodium, potassium, calcium, and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as ammonia, isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, diethanolamine, ethanolamine, deanol, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, benethamine, benzathine, ethylenediamine, glucosamine, methylglucamine, theobromine, triethanolamine, tromethamine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like. Particularly preferred organic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline and caffeine.

(70) The terms “antagonist” and “inhibitor” are used interchangeably, and they refer to a compound having the ability to inhibit a biological function of a target protein, whether by inhibiting the activity or expression of the protein, such as K-Ras, H-Ras or N-Ras G12C. Accordingly, the terms “antagonist” and “inhibitors” are defined in the context of the biological role of the target protein. While preferred antagonists herein specifically interact with (e.g. bind to) the target, compounds that inhibit a biological activity of the target protein by interacting with other members of the signal transduction pathway of which the target protein is a member are also specifically included within this definition. A preferred biological activity inhibited by an antagonist is associated with the development, growth, or spread of a tumor.

(71) The term “agonist” as used herein refers to a compound having the ability to initiate or enhance a biological function of a target protein, whether by inhibiting the activity or expression of the target protein. Accordingly, the term “agonist” is defined in the context of the biological role of the target polypeptide. While preferred agonists herein specifically interact with (e.g. bind to) the target, compounds that initiate or enhance a biological activity of the target polypeptide by interacting with other members of the signal transduction pathway of which the target polypeptide is a member are also specifically included within this definition.

(72) As used herein, “agent” or “biologically active agent” refers to a biological, pharmaceutical, or chemical compound or other moiety. Non-limiting examples include a simple or complex organic or inorganic molecule, a peptide, a protein, an oligonucleotide, an antibody, an antibody derivative, antibody fragment, a vitamin derivative, a carbohydrate, a toxin, or a chemotherapeutic compound. Various compounds can be synthesized, for example, small molecules and oligomers (e.g., oligopeptides and oligonucleotides), and synthetic organic compounds based on various core structures. In addition, various natural sources can provide compounds for screening, such as plant or animal extracts, and the like.

(73) “Signal transduction” is a process during which stimulatory or inhibitory signals are transmitted into and within a cell to elicit an intracellular response. A modulator of a signal transduction pathway refers to a compound which modulates the activity of one or more cellular proteins mapped to the same specific signal transduction pathway. A modulator may augment (agonist) or suppress (antagonist) the activity of a signaling molecule.

(74) An “anti-cancer agent”, “anti-tumor agent” or “chemotherapeutic agent” refers to any agent useful in the treatment of a neoplastic condition. One class of anti-cancer agents comprises chemotherapeutic agents. “Chemotherapy” means the administration of one or more chemotherapeutic drugs and/or other agents to a cancer patient by various methods, including intravenous, oral, intramuscular, intraperitoneal, intravesical, subcutaneous, transdermal, buccal, or

inhalation or in the form of a suppository.

(75) The term “cell proliferation” refers to a phenomenon by which the cell number has changed as a result of division. This term also encompasses cell growth by which the cell morphology has changed (e.g., increased in size) consistent with a proliferative signal.

(76) The term “selective inhibition” or “selectively inhibit” refers to a biologically active agent refers to the agent's ability to preferentially reduce the target signaling activity as compared to off-target signaling activity, via direct or indirect interaction with the target.

(77) “Subject” refers to an animal, such as a mammal, for example a human. The methods described herein can be useful in both human therapeutics and veterinary applications. In some embodiments, the subject is a mammal, and in some embodiments, the subject is human.

(78) “Mammal” includes humans and both domestic animals such as laboratory animals and household pets (e.g., cats, dogs, swine, cattle, sheep, goats, horses, rabbits), and non-domestic animals such as wildlife and the like.

(79) “Radiation therapy” means exposing a subject, using routine methods and compositions known to the practitioner, to radiation emitters such as alpha-particle emitting radionuclides (e.g., actinium and thorium radionuclides), low linear energy transfer (LET) radiation emitters (i.e. beta emitters), conversion electron emitters (e.g. strontium-89 and samarium-153-EDTMP, or high-energy radiation, including without limitation x-rays, gamma rays, and neutrons.

(80) An “anti-cancer agent”, “anti-tumor agent” or “chemotherapeutic agent” refers to any agent useful in the treatment of a neoplastic condition. One class of anti-cancer agents comprises chemotherapeutic agents. “Chemotherapy” means the administration of one or more chemotherapeutic drugs and/or other agents to a cancer patient by various methods, including intravenous, oral, intramuscular, intraperitoneal, intravesical, subcutaneous, transdermal, buccal, or inhalation or in the form of a suppository.

(81) “Prodrug” is meant to indicate a compound that may be converted under physiological conditions or by solvolysis to a biologically active compound described herein (e.g., compound of structure (I)). Thus, the term “prodrug” refers to a precursor of a biologically active compound that is pharmaceutically acceptable. In some aspects, a prodrug is inactive when administered to a subject, but is converted in vivo to an active compound, for example, by hydrolysis. The prodrug compound often offers advantages of solubility, tissue compatibility or delayed release in a mammalian organism (see, e.g., Bundgard, H., *Design of Prodrugs* (1985), pp. 7-9, 21-24 (Elsevier, Amsterdam). A discussion of prodrugs is provided in Higuchi, T., et al., “Pro-drugs as Novel Delivery Systems,” A.C.S. Symposium Series, Vol. 14, and in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated in full by reference herein. The term “prodrug” is also meant to include any covalently bonded carriers, which release the active compound in vivo when such prodrug is administered to a mammalian subject. Prodrugs of an active compound, as described herein, are typically prepared by modifying functional groups present in the active compound in such a way that the modifications are cleaved, either in routine manipulation or in vivo, to the parent active compound. Prodrugs include compounds wherein a hydroxy, amino or mercapto group is bonded to any group that, when the prodrug of the active compound is administered to a mammalian subject, cleaves to form a free hydroxy, free amino or free mercapto group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of a hydroxy functional group, or acetamide, formamide and benzamide derivatives of an amine functional group in the active compound and the like.

(82) The term “in vivo” refers to an event that takes place in a subject's body.

(83) The invention disclosed herein is also meant to encompass all pharmaceutically acceptable compounds of structure (I) being isotopically-labeled by having one or more atoms replaced by an atom having a different atomic mass or mass number. Examples of isotopes that can be incorporated into the disclosed compounds include isotopes of hydrogen, carbon, nitrogen, oxygen,

phosphorous, fluorine, chlorine, and iodine, such as ²H, ³H, C, ¹³C, ¹⁴C, ¹³N, ¹⁵N, ¹⁵O, ¹⁷O, ¹⁸O, ³¹P, ³²P, ³⁵S, ¹⁸F, ³⁶Cl, ¹²³I, and ¹²⁵I, respectively. These radiolabelled compounds could be useful to help determine or measure the effectiveness of the compounds, by characterizing, for example, the site or mode of action, or binding affinity to pharmacologically important site of action. Certain isotopically-labeled compounds of structure (I), for example, those incorporating a radioactive isotope, are useful in drug and/or substrate tissue distribution studies. The radioactive isotopes tritium, i.e. ³H, and carbon-14, i.e. ¹⁴C, are particularly useful for this purpose in view of their ease of incorporation and ready means of detection.

(84) Substitution with heavier isotopes such as deuterium, i.e. ²H, may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased in vivo half-life or reduced dosage requirements, and hence are preferred in some circumstances.

(85) Substitution with positron emitting isotopes, such as ¹¹C, ¹⁸F, ¹⁵O and ¹³N, can be useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy. Isotopically-labeled compounds of structure (I) can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the Preparations and Examples as set out below using an appropriate isotopically-labeled reagent in place of the non-labeled reagent previously employed.

(86) The invention disclosed herein is also meant to encompass the in vivo metabolic products of the disclosed compounds. Such products may result from, for example, the oxidation, reduction, hydrolysis, amidation, esterification, and the like of the administered compound, primarily due to enzymatic processes. Accordingly, the invention includes compounds produced by a process comprising administering a compound of this invention to a mammal for a period of time sufficient to yield a metabolic product thereof. Such products are typically identified by administering a radiolabelled compound of the invention in a detectable dose to an animal, such as rat, mouse, guinea pig, monkey, or to human, allowing sufficient time for metabolism to occur, and isolating its conversion products from the urine, blood or other biological samples.

(87) “Stable compound” and “stable structure” are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

(88) Often crystallizations produce a solvate of the compound of the invention. As used herein, the term “solvate” refers to an aggregate that comprises one or more molecules of a compound of the invention with one or more molecules of solvent. In some embodiments, the solvent is water, in which case the solvate is a hydrate. Alternatively, in other embodiments, the solvent is an organic solvent. Thus, the compounds of the present invention may exist as a hydrate, including a monohydrate, dihydrate, hemihydrate, sesquihydrate, trihydrate, tetrahydrate and the like, as well as the corresponding solvated forms. In some aspects, the compound of the invention is a true solvate, while in other cases, the compound of the invention merely retains adventitious water or is a mixture of water plus some adventitious solvent.

(89) “Optional” or “optionally” means that the subsequently described event or circumstances may or may not occur, and that the description includes instances where said event or circumstance occurs and instances in which it does not. For example, “optionally substituted aryl” means that the aryl radical may or may not be substituted and that the description includes both substituted aryl radicals and aryl radicals having no substitution.

(90) A “pharmaceutical composition” refers to a formulation of a compound of the invention and a medium generally accepted in the art for the delivery of the biologically active compound to mammals, e.g., humans. Such a medium includes all pharmaceutically acceptable carriers, diluents or excipients therefor.

(91) “Pharmaceutically acceptable carrier, diluent or excipient” includes without limitation any adjuvant, carrier, excipient, glidant, sweetening agent, diluent, preservative, dye/colorant, flavor

enhancer, surfactant, wetting agent, dispersing agent, suspending agent, stabilizer, isotonic agent, solvent, or emulsifier which has been approved by the United States Food and Drug Administration as being acceptable for use in humans or domestic animals.

(92) The compounds of the invention, or their pharmaceutically acceptable salts may contain one or more asymmetric centers and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that are defined, in terms of absolute stereochemistry, as (R)- or (S)- or, as (D)- or (L)- for amino acids. The present invention is meant to include all such possible isomers, as well as their racemic and optically pure forms. Optically active (+) and (−), (R)- and (S)-, or (D)- and (L)-isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques, for example, chromatography and fractional crystallization. Conventional techniques for the preparation/isolation of individual enantiomers include chiral synthesis from a suitable optically pure precursor or resolution of the racemate (or the racemate of a salt or derivative) using, for example, chiral high pressure liquid chromatography (HPLC). When the compounds described herein contain olefinic double bonds or other centres of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both E and Z geometric isomers. Likewise, all tautomeric forms are also intended to be included.

(93) The present invention includes all manner of rotamers and conformationally restricted states of a compound of the invention.

(94) A “stereoisomer” refers to a compound made up of the same atoms bonded by the same bonds but having different three-dimensional structures, which are not interchangeable. The present invention contemplates various stereoisomers and mixtures thereof and includes “enantiomers”, which refers to two stereoisomers whose molecules are nonsuperimposable mirror images of one another.

(95) A “tautomer” refers to a proton shift from one atom of a molecule to another atom of the same molecule. The present invention includes tautomers of any said compounds.

(96) The chemical naming protocol and structure diagrams used herein are a modified form of the I.U.P.A.C. nomenclature system, using the ACD/Name Version 9.07 software program and/or ChemDraw Ultra Version 11.0.1 software naming program (CambridgeSoft). For complex chemical names employed herein, a substituent group is typically named before the group to which it attaches. For example, cyclopropylethyl comprises an ethyl backbone with a cyclopropyl substituent. Except as described below, all bonds are identified in the chemical structure diagrams herein, except for all bonds on some carbon atoms, which are assumed to be bonded to sufficient hydrogen atoms to complete the valency.

(97) Compounds

(98) In an aspect, the invention provides compounds which are capable of selectively binding to and/or modulating a G12C mutant K-Ras, H-Ras or N-Ras protein. In some embodiments, the compounds modulate the G12C mutant K-Ras, H-Ras or N-Ras protein by reaction with an amino acid. In some embodiment the compounds of the invention selectively react with the G12C mutant K-Ras, H-Ras or N-Ras proteins by forming an irreversible covalent bond with the cysteine at the 12 position. By binding to the Cystine 12 the compounds of the invention may lock the switch II of the G12C mutant K-Ras, H-Ras or N-Ras into an inactive stage. This inactive stage may be distinct from those observed for GTP and GDP bound K-Ras, H-Ras or N-Ras. Some compounds of the invention are also able to perturb the switch I conformation. Because effector binding to K-Ras, H-Ras or N-Ras is highly sensitive to the conformation of switch I and II, the irreversible binding of these compounds may disrupt K-Ras, H-Ras or N-Ras downstream signaling.

(99) In some embodiments, the invention provides a compound of Formula I

(100) ##STR00019##

(101) In various embodiments Y is —CH₂—, —CHR₂₂—, CO, SO or SO₂. In some embodiments Y is —CH₂—. In some embodiments Y is —CHR₂₂—. In some embodiments Y is CO. In some embodiments Y is SO. In some embodiments Y is SO₂.

(102) In various embodiments n is an integer with value 1-6. In some cases n is 1. In some cases n is 2. In some cases n is 3. In some cases n is 4. In some cases n is 5. In some cases n is 6.

(103) R.sub.1 is aryl or heteroaryl each of which is unsubstituted or substituted by one or more independent R.sub.2 substituents. In some embodiments R.sub.1 is substituted aryl. In some cases R.sub.1 is unsubstituted aryl. In some cases R.sub.1 is substituted heteroaryl. In some cases R.sub.1 is unsubstituted heteroaryl. In some embodiments R.sub.1 an aryl group substituted with one or more R.sub.3 groups. In some embodiments R.sub.1 a heteroaryl group substituted with one or more R.sub.3 groups. In some embodiments R.sub.1 is a substituted or unsubstituted phenyl group. In some embodiments R.sub.1 is an unsubstituted benzyl group. In some embodiments R.sub.1 is a phenyl group substituted with one or more independent R.sub.2 substituents. In some embodiments R.sub.1 is an unsubstituted benzothiadiazolyl group. In some embodiments R.sub.1 is a benzothiadiazolyl group substituted with one or more independent R.sub.2 substituents. In some embodiments R.sub.1 is an unsubstituted benzothiadiazolyl group. In some embodiments R.sub.1 is a benzothiadiazolyl group substituted with one or more independent R.sub.2 substituents. In some embodiments R.sub.1 is an unsubstituted naphthalenyl group. In some embodiments R.sub.1 is a naphthalenyl group substituted with one or more independent R.sub.2 substituents. In some embodiments R.sub.1 is an imidazopyridinyl naphthalenyl group. In some embodiments R.sub.1 is an imidazopyridinyl group substituted with one or more independent R.sub.2 substituents.

(104) In some embodiments, R.sub.1 is an unsubstituted phenyl group. In some embodiments, R.sub.1 is a substituted phenyl group (Formula Ia). In some embodiments R.sub.1 is a phenyl group with a halogen substituent at the 4-position of the phenyl ring. In some embodiments, R.sub.1 is a phenyl substituted with a substituted or unsubstituted aryl at the 5-position. In some embodiments, R.sub.1 is a phenyl substituted with a substituted or unsubstituted heteroaryl at the 5-position. In some embodiments, R.sub.1 is a phenyl substituted with a substituted or unsubstituted cycloalkyl at the 5-position. In some embodiments, R.sub.1 is a phenyl substituted with a substituted or unsubstituted heterocycloaryl (heteroaryl) at the 5-position. In some embodiments R.sub.1 is a phenyl with an alkoxy substituent at the 2 position of the phenyl ring. In some embodiments R.sub.1 is a phenyl with methoxy substituent at the 2 position of the phenyl ring. In some embodiments R.sub.1 is a phenyl group with a methoxy substituent at the 2 position, a halogen substituent at the 4 position and an aryl or heteroaryl substituent (both of which are substituted or unsubstituted) at the 5 position of the phenyl ring.

(105) ##STR00020##

(106) In some embodiments, R.sub.1 is capable of reversible interaction with K-Ras, H-Ras or N-Ras G12C mutant protein. In some embodiments R.sub.1 has high affinity towards K-Ras, H-Ras or N-Ras and is highly specific towards G12C K-Ras, H-Ras or N-Ras. In some embodiments R.sub.1 is capable of hydrophobic interaction with K-Ras, H-Ras or N-Ras G12C. In some embodiments R.sub.1 is able to form hydrogen bonds with various residues of G12C K-Ras, H-Ras or N-Ras protein. In some embodiments R.sub.1 interacts with one or more of G10, R68, Y71, Y96 or Q99 residues in K-Ras G12C (FIG. 1). In some embodiments R.sub.1 interacts with the G10 residue of K-Ras G12C. In some embodiments R.sub.1 interacts with the R68 residue of K-Ras G12C. In some embodiments R.sub.1 interacts with the Y71 residue of K-Ras G12C. In some embodiments R.sub.1 interacts with the Y96 residue of K-Ras G12C. In some embodiments R.sub.1 interacts with the Q99 residue of K-Ras G12C.

(107) R.sub.2 is, at each occurrence, independently halogen, oxo, hydroxy, substituted or unsubstituted alkoxy, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted heterocycloalkyl, aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted arylene, or substituted or unsubstituted arylene heteroarylene, each of which is unsubstituted or substituted by one or more independent R.sub.3 substituents. In some embodiments R.sub.2 is halogen. In some embodiments, R.sub.2 is hydroxy. In some embodiments R.sub.2 is an alkoxy group substituted with one or more

R.sub.3 substituents. In some embodiments R.sub.2 is an unsubstituted alkoxy group. In some embodiments R.sub.3 is an alkyl group substituted with one or more R.sub.3 groups. In some embodiments R.sub.2 is an unsubstituted alkyl group. In some embodiments R.sub.2 is a heteroalkyl group substituted with one or more R.sub.3 groups. In some embodiments R.sub.2 is an unsubstituted heteroalkyl group. In some embodiments R.sub.2 a cycloalkyl group substituted with one or more R.sub.3 groups. In some embodiments R.sub.2 is an unsubstituted cycloalkyl group. In some embodiments R.sub.2 is a heterocycloalkyl substituted with one or more R.sub.3 groups. In some embodiments R.sub.2 is an unsubstituted heterocycloalkyl group. In some embodiments R.sub.2 is an aryl substituted with one or more R.sub.3 groups. In some embodiments R.sub.2 is an unsubstituted aryl group. In some embodiments R.sub.2 is a heteroaryl group substituted with one or more R.sub.3 groups. In some embodiments R.sub.2 unsubstituted a heteroaryl group. In some embodiments R.sub.2 is an arylene substituted with one or more R.sub.3 groups. In some embodiments R.sub.2 is an unsubstituted arylene group. In some embodiments R.sub.2 is a heteroarylene group substituted with one or more R.sub.3 groups. In some embodiments R.sub.2 unsubstituted a heteroarylene group. In some embodiments R.sub.2 is halogen. In some embodiments R.sub.2 is hydroxy. In some embodiments R.sub.2 is alkoxy. In some embodiments R.sub.2 is methoxy. In some embodiments R2 is oxo.

(108) R.sub.3 is halogen, OH, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxy, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloheteroalkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted arylene, substituted or unsubstituted heteroarylene. In some embodiments R.sub.3 is halogen. In some embodiments R.sub.3 is hydroxy. In some embodiments R.sub.3 is cyano. In some embodiments R.sub.3 is unsubstituted alkyl. In some embodiments R.sub.3 is an alkyl substituted with one or more R.sub.4 groups. In some embodiments R.sub.3 is unsubstituted alkoxy. In some embodiments R.sub.3 is an alkoxy substituted with one or more R.sub.4 groups. In some embodiments R.sub.3 is unsubstituted cycloalkyl. In some embodiments R.sub.3 is a cycloalkyl substituted with one or more R.sub.4 groups. In some embodiments R.sub.3 is unsubstituted hetero alkyl. In some embodiments R.sub.3 is a hetero alkyl substituted with one or more R.sub.4 groups. In some embodiments R.sub.3 is unsubstituted heterocycloalkyl. In some embodiments R.sub.3 is a heterocycloalkyl substituted with one or more R.sub.4 groups. In some embodiments R.sub.3 is unsubstituted aryl. In some embodiments R.sub.3 is an aryl substituted with one or more R.sub.4 groups. In some embodiments R.sub.3 is unsubstituted heteroaryl. In some embodiments R.sub.3 is a heteroaryl substituted with one or more R.sub.4 groups. In some embodiments R.sub.3 is unsubstituted arylene. In some embodiments R.sub.3 is an arylene substituted with one or more R.sub.4 groups. In some embodiments R.sub.3 is unsubstituted heteroarylene. In some embodiments R.sub.3 is a heteroarylene substituted with one or more R.sub.4 groups. In some embodiments R.sub.3 is halogen. In some embodiments R.sub.3 is hydroxyl. In some embodiments R.sub.3 is cyano.

(109) In various embodiments R.sub.4 is halogen, OH, cyano, alkyl, alkoxy, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene moiety. In some embodiments R.sub.4 is halogen. In some embodiments R.sub.4 is hydroxyl. In some embodiments R.sub.4 is cyano. In some embodiments R.sub.4 is an alkyl. In some embodiments R.sub.4 is an alkoxy. In some embodiments R.sub.4 is a cycloalkyl. In some embodiments R.sub.4 is heteroalkyl. In some embodiments R.sub.4 is heterocycloalkyl. In some embodiments R.sub.4 is aryl. In some embodiments R.sub.4 is heteroaryl. In some embodiments R.sub.4 is arylene, in some embodiments R.sub.4 is heteroarylene.

(110) In various embodiments C.sub.1 is a substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkylene, substituted or unsubstituted heterocycloalkylene, substituted or unsubstituted arylene, or substituted or unsubstituted heteroarylene. In some embodiments C.sub.1 is a cycloalkylene substituted with one or more R.sub.5 groups. In some embodiments C.sub.1 is an

unsubstituted cycloalkylene. In some embodiments C.sub.1 is a heterocycloalkylene substituted with one or more R.sub.5 groups. In some embodiments C.sub.1 is an unsubstituted heterocycloalkylene. In some embodiments C.sub.1 is an arylene substituted with one or more R.sub.5 groups. In some embodiments C.sub.1 is an unsubstituted arylene. In some embodiments C.sub.1 is a heteroarylene substituted with one or more R.sub.5 groups. In some embodiments C.sub.1 is an unsubstituted heteroarylene. In some embodiments C.sub.1 is selected from:

(111) ##STR00021##

(112) In some embodiments, C.sub.1 is

(113) ##STR00022##

(114) In some embodiments R.sub.5 is —OH. In some embodiments R.sub.5 is —CH.sub.2OH. In some embodiment R.sub.5 is alkyl. In some embodiments R.sub.5 is methyl.

(115) In various embodiments C.sub.2 is a bond, a cycloalkylene, heterocycloalkylene, arylene, or heteroarylene. In some embodiments C.sub.2 is a bond. In some embodiments C.sub.2 is an unsubstituted cycloalkylene. In some embodiments C.sub.2 is an unsubstituted heterocycloalkylene. In some embodiments C.sub.2 is an unsubstituted arylene. In some embodiments C.sub.2 is an unsubstituted heteroarylene. In some embodiments C.sub.2 is selected from:

(116) ##STR00023##

(117) In some embodiments, C1 and C2 form a fused or spiro bicyclic ring. In some embodiments —C1-C2- form a fused bicyclic ring. In some embodiments —C1-C2- form a spiro bicyclic ring. In some embodiments —C1-C2- is

(118) ##STR00024##

In some embodiments —C1-C2- is

(119) ##STR00025##

In some embodiments —C1-C2- is

(120) ##STR00026##

(121) In various embodiments X is O, NH, S or CR.sub.23R.sub.24. In some embodiments X is O. In some embodiments X is NH. In various embodiments X is S. In various embodiments X is CR.sub.23R.sub.24.

(122) In various embodiments D is a bond, —NH—CH.sub.2—, —NH—, or —CH.sub.2—. In various embodiments D is a bond. In various embodiments D is —NH—CH.sub.2. In various embodiments D is —NH—. In various embodiments D is —CH.sub.2—.

(123) R.sub.22, R.sub.23 and R.sub.24 are each independently hydrogen, halogen, —OH, alkyl, cycloalkyl, aryl, heteroaryl, heteroalkyl, or heterocycloalkyl.

(124) In some embodiments E is an electrophile capable of bonding with a K-Ras, H-Ras or N-Ras protein comprising G12C mutation. In some embodiments, the electrophile E is capable of forming an irreversible covalent bond with a G12C mutant K-Ras, H-Ras or N-Ras protein. In some cases, the electrophile E binds with the cysteine residue at the position 12 of a G12C mutant K-Ras, H-Ras or N-Ras protein. In some cases E is selected from:

(125) ##STR00027##

(126) In some embodiments E is

(127) ##STR00028##

In some embodiments E is

(128) ##STR00029##

In some embodiments E is

(129) ##STR00030##

In some embodiments E is

(130) ##STR00031##

In some embodiments E is

(131) ##STR00032##

(132) In some embodiments the invention provides compounds of Formula I as shown in Table 1.

(133) In some embodiments, the invention provides compounds of Formula II

(134) ##STR00033##

(135) In various embodiments A is CH₂, O— or NH. In some embodiments A is CH₂. In some embodiments A is O. In some embodiments A is NH.

(136) In various embodiments R₆ is aryl or heteroaryl moiety, each of which is either unsubstituted or substituted with one or more R₇ groups. In some embodiments R₆ is an unsubstituted aryl. In some embodiments R₆ is an aryl substituted with one or more R₇ groups. In some embodiments R₆ is an unsubstituted heteroaryl. In some embodiments R₆ is a heteroaryl substituted with one or more R₇ groups. In some embodiments R₆ is a substituted or unsubstituted phenyl moiety. In some embodiments R₆ is an unsubstituted phenyl moiety. In some embodiments R₆ is a phenyl moiety substituted with one or more R₇ substituents. In some embodiments R₆ is a substituted or unsubstituted pyridinyl moiety. In some embodiments R₆ is an unsubstituted pyridinyl moiety. In some embodiments R₆ is a pyridinyl moiety substituted with one or more R₇ substituents.

(137) R₇ is halogen, —OH, OR₁₀, NR₁₁R₁₂, Oxo, unsubstituted or substituted alkyl, unsubstituted or substituted cycloalkyl, unsubstituted or substituted heteroalkyl, unsubstituted or substituted heterocycloalkyl, unsubstituted or substituted aryl, unsubstituted or substituted heteroaryl, unsubstituted or substituted arylene, or unsubstituted or substituted heteroarylene. In some embodiments R₇ is a halogen. In some embodiments R₇ is OH. In some embodiments R₇ is OR₁₀. In some embodiments R₇ is NR₁₁R₁₂. In some embodiments R₇ is unsubstituted alkyl. In some embodiments R₇ is a alkyl substituted with one or more R₈ substituents. In some embodiments R₇ is unsubstituted cycloalkyl. In some embodiments R₇ is a cycloalkyl substituted with one or more R₈ substituents. In some embodiments R₇ is unsubstituted heteroalkyl. In some embodiments R₇ is a heteroalkyl substituted with one or more R₈ substituents. In some embodiments R₇ is unsubstituted heterocycloalkyl. In some embodiments R₇ is a heterocycloalkyl substituted with one or more R₈ substituents. In some embodiments R₇ is unsubstituted aryl. In some embodiments R₇ is a aryl substituted with one or more R₈ substituents. In some embodiments R₇ is unsubstituted heteroaryl. In some embodiments R₇ is a heteroaryl substituted with one or more R₈ substituents. In some embodiments R₇ is unsubstituted arylene. In some embodiments R₇ is a arylene substituted with one or more R₈ substituents. In some embodiments R₇ is unsubstituted heteroarylene. In some embodiments R₇ is a heteroarylene substituted with one or more R₈ substituents.

(138) In various embodiments R₈ is halogen, OH, cyano, unsubstituted or substituted alkyl, unsubstituted or substituted alkoxy, unsubstituted or substituted cycloalkyl, unsubstituted or substituted heteroalkyl, unsubstituted or substituted heterocycloalkyl, unsubstituted or substituted aryl, unsubstituted or substituted heteroaryl, unsubstituted or substituted arylene, or unsubstituted or substituted heteroarylene. In some embodiments R₈ is halogen. In some embodiments R₈ is OH. In some embodiments R₈ is cyano. In some embodiments R₈ is unsubstituted alkyl. In some embodiments R₈ is a alkyl substituted with one or more R₉ groups. In some embodiments R₈ is unsubstituted alkoxy. In some embodiments R₈ is a alkoxy substituted with one or more R₉ groups. In some embodiments R₈ is unsubstituted cycloalkyl. In some embodiments R₈ is a cycloalkyl substituted with one or more R₉ groups. In some embodiments R₈ is unsubstituted heteroalkyl. In some embodiments R₈ is a heteroalkyl substituted with one or more R₉ groups. In some embodiments R₈ is unsubstituted heterocycloalkyl. In some embodiments R₈ is a heterocycloalkyl substituted with one or more R₉ groups. In some embodiments R₈ is unsubstituted aryl. In some embodiments R₈ is a aryl substituted with one or more R₉ groups. In some embodiments R₈ is unsubstituted heteroaryl. In some embodiments R₈ is a heteroaryl substituted with

(142) In various embodiments R.sub.9 is halogen, OH, cyano, alkyl, alkoxy, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene. In various embodiments R.sub.9 is halogen. In various embodiments R.sub.9 is OH. In various embodiments R.sub.9 is cyano. In various embodiments R.sub.9 is alkyl. In various embodiments R.sub.9 is cycloalkyl. In various embodiments R.sub.9 is heteroalkyl. In various embodiments R.sub.9 is heterocycloalkyl. In various embodiments R.sub.9 is aryl. In various embodiments R.sub.9 is heteroaryl. In various embodiments R.sub.9 is arylene. In various embodiments R.sub.9 is heteroarylene.

(143) In various embodiments R.sub.13 is halogen, OH, cyano, alkyl, alkoxy, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene. In various embodiments R.sub.13 is halogen. In various embodiments R.sub.13 is OH. In various embodiments R.sub.13 is cyano. In various embodiments R.sub.13 is alkyl. In various embodiments R.sub.13 is cycloalkyl. In various embodiments R.sub.13 is heteroalkyl. In various embodiments R.sub.13 is heterocycloalkyl. In various embodiments R.sub.13 is aryl. In various embodiments R.sub.13 is heteroaryl. In various embodiments R.sub.13 is arylene. In various embodiments R.sub.13 is heteroarylene.

(144) In some embodiments E is an electrophile capable of bonding with a K-Ras, H-Ras or N-Ras protein comprising G12C mutation. In some embodiments, the electrophile E is capable of forming an irreversible covalent bond with a G12C mutant K-Ras, H-Ras or N-Ras protein. In some cases, the electrophile E binds with the cysteine residue at the position 12 of a G12C mutant K-Ras, H-Ras or N-Ras protein.

(145) In some cases E has the general structure

(146) ##STR00034##

wherein:

(147) R.sub.25 is alkyl;

(148) R.sub.26 is cyano or alkyl or R.sub.26 joins with R.sub.27 to form a cycloalkene; and

(149) R.sub.27 is alkyl or R.sub.27 joins with R.sub.26 to form a cycloalkene.

(150) In some cases E is selected from:

(151) ##STR00035##

(152) In some embodiments E is

(153) ##STR00036##

In some embodiments E is

(154) ##STR00037##

In some embodiments E is

(155) ##STR00038##

In some embodiments E is

(156) ##STR00039##

(157) In some embodiments, R.sub.6 is capable of reversible interaction with K-Ras, H-Ras or N-Ras G12C mutant protein. In some embodiments R.sub.6 moiety has high affinity towards K-Ras, H-Ras or N-Ras and is highly specific towards G12C K-Ras, H-Ras or N-Ras. In some embodiments R.sub.6 is capable of hydrophobic interaction with K-Ras, H-Ras or N-Ras G12C. In some embodiments R.sub.6 is able to form hydrogen bonds with various residues of G12C K-Ras, H-Ras or N-Ras protein. In some embodiments R.sub.6 interacts with one or more of G10, R68, Y71, Y96 or Q99 residues in K-Ras G12C (FIG. 1). In some embodiments, R.sub.6 moiety interacts with the G10 residue of K-Ras G12C. In some embodiments R.sub.6 interacts with the R68 residue of K-Ras G12C. In some embodiments R.sub.6 interacts with the Y71 residue of K-Ras G12C. In some embodiments R.sub.6 interacts with the Y96 residue of K-Ras G12C. In some embodiments R.sub.6 interacts with the Q99 residue of K-Ras G12C.

(158) In some embodiments, the

(159) ##STR00040##

unit in Formula II provides proper length and geometry to the compound such that the electrophile E interacts with the cysteine residue at the 12 position in G12C K-Ras, H-Ras or N-Ras protein. In some embodiments the

(160) ##STR00041##

also interacts with other protein backbone residues.

(161) In some embodiments the invention provides compounds of Formula II as shown in Table 2.

(162) In some embodiments, the invention provides compounds of Formula III

(163) ##STR00042##

(164) In various embodiments A is a bond, O, NH or —(CO)—. In some embodiments A is a bond. In some embodiments A is O. In some embodiments A is NH. In some embodiments A is —C(O)—.

(165) In various embodiments n is 0 or 1. In some cases n is 0. In some cases n is 1.

(166) In various embodiments R.sub.14 is cycloalkyl, heterocycloalkyl, aryl or heteroaryl moiety. In some embodiments R.sub.14 is an unsubstituted cycloalkyl. In some embodiments R.sub.14 is cycloalkyl substituted with one or more R.sub.15 groups. In some embodiments R.sub.14 is an unsubstituted heterocycloalkyl. In some embodiments R.sub.14 is hetero cycloalkyl substituted with one or more R.sub.15 groups. In some embodiments R.sub.14 is an unsubstituted aryl. In some embodiments R.sub.14 is aryl substituted with one or more R.sub.15 groups. In some embodiments R.sub.14 is an unsubstituted heteroaryl. In some embodiments R.sub.14 is heteroaryl substituted with one or more R.sub.15 groups. In some embodiments R.sub.14 is an unsubstituted phenyl. In some embodiments R.sub.14 is phenyl substituted with one or more R.sub.15 groups.

(167) In various embodiments R.sub.15 is halogen, OR.sub.18, NR.sub.19R.sub.20, oxo, alkyl, alkoxy, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene, each of which is unsubstituted or substituted by one or more independent R.sub.3 substituents. In some embodiments R.sub.15 is halogen. In some embodiments R.sub.15 is cyano. In some embodiments R.sub.15 is NR.sub.19R.sub.20. In some embodiments R.sub.15 is oxo. In some embodiments R.sub.15 is OR.sub.18. In some embodiments R.sub.15 is unsubstituted alkyl. In some embodiments R.sub.15 is a alkyl substituted with one or more R.sub.16 groups. In some embodiments R.sub.15 is unsubstituted alkoxy. In some embodiments R.sub.16 is a alkoxy substituted with one or more R.sub.16 groups. In some embodiments R.sub.15 is unsubstituted cycloalkyl. In some embodiments R.sub.15 is a cycloalkyl substituted with one or more R.sub.16 groups. In some embodiments R.sub.15 is unsubstituted heteroalkyl. In some embodiments R.sub.15 is a heteroalkyl substituted with one or more R.sub.16 groups. In some embodiments R.sub.15 is unsubstituted heterocycloalkyl. In some embodiments R.sub.15 is a heterocycloalkyl substituted with one or more R.sub.16 groups. In some embodiments R.sub.15 is unsubstituted aryl. In some embodiments R.sub.15 is a aryl substituted with one or more R.sub.16 groups. In some embodiments R.sub.15 is unsubstituted heteroaryl. In some embodiments R.sub.15 is a heteroaryl substituted with one or more R.sub.16 groups. In some embodiments R.sub.15 is unsubstituted arylene. In some embodiments R.sub.15 is a arylene substituted with one or more R.sub.16 groups. In some embodiments R.sub.15 is unsubstituted heteroarylene. In some embodiments R.sub.15 is a heteroarylene substituted with one or more R.sub.16 groups

(168) In various embodiments R.sub.16 is halogen, OH, oxo, cyano alkyl, alkoxy, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene, each of which is unsubstituted or substituted by one or more independent R.sub.17 substituents. In some embodiments R.sub.16 is halogen. In some embodiments R.sub.16 is OH. In some embodiments R.sub.16 is cyano. In some embodiments R.sub.16 is unsubstituted alkyl. In some embodiments R.sub.16 is a alkyl substituted with one or more R.sub.17 groups. In some embodiments R.sub.16 is unsubstituted alkoxy. In some embodiments R.sub.16 is a alkoxy substituted with one or more R.sub.17 groups. In some embodiments R.sub.16 is unsubstituted cycloalkyl. In some embodiments R.sub.16 is a cycloalkyl substituted with one or more R.sub.17 groups. In some embodiments R.sub.16 is unsubstituted heteroalkyl. In some embodiments R.sub.16 is a heteroalkyl substituted with one or more R.sub.17 groups. In some embodiments R.sub.16 is unsubstituted heterocycloalkyl. In some embodiments R.sub.16 is a heterocycloalkyl substituted with one or more R.sub.17 groups. In some embodiments R.sub.16 is unsubstituted aryl. In some embodiments R.sub.16 is a aryl substituted with one or more R.sub.17 groups. In some embodiments R.sub.16 is unsubstituted heteroaryl. In some embodiments R.sub.16 is a heteroaryl substituted with one or more R.sub.17 groups. In some embodiments R.sub.16 is unsubstituted arylene. In some embodiments R.sub.16 is a arylene substituted with one or more R.sub.17 groups. In some

embodiments R.sub.16 is unsubstituted heteroarylene. In some embodiments R.sub.16 is a heteroarylene substituted with one or more R.sub.17 groups.

(169) In various embodiments, R.sub.18 is hydrogen, unsubstituted or substituted alkyl, unsubstituted or substituted cycloalkyl, unsubstituted or substituted heteroalkyl, unsubstituted or substituted heterocycloalkyl, unsubstituted or substituted aryl, or unsubstituted or substituted heteroaryl. In some embodiments R.sub.18 is hydrogen. In some embodiments R.sub.18 is unsubstituted alkyl. In some embodiments R.sub.18 is alkyl substituted with one or more R.sub.21 groups. In some embodiments R.sub.18 is unsubstituted cycloalkyl. In some embodiments R.sub.18 is cycloalkyl substituted with one or more R.sub.21 groups. In some embodiments R.sub.18 is unsubstituted heteroalkyl. In one embodiment R.sub.18 is heteroalkyl substituted with one or more R.sub.21 groups. In some embodiments R.sub.18 is unsubstituted heterocycloalkyl. In some embodiments R.sub.18 is heterocycloalkyl substituted with one or more R.sub.21 groups. In some embodiments R.sub.18 is unsubstituted aryl. In some embodiments R.sub.18 is aryl substituted with one or more R.sub.21 groups. In some embodiments R.sub.18 is unsubstituted heteroaryl. In some embodiments R.sub.18 is heteroaryl substituted with one or more R.sub.21 groups.

(170) In various embodiments, R.sub.19 is hydrogen, unsubstituted or substituted alkyl, unsubstituted or substituted cycloalkyl, unsubstituted or substituted heteroalkyl, unsubstituted or substituted heterocycloalkyl, unsubstituted or substituted aryl, or unsubstituted or substituted heteroaryl. In some embodiments R.sub.19 is hydrogen. In some embodiments R.sub.19 is unsubstituted alkyl. In some embodiments R.sub.19 is alkyl substituted with one or more R.sub.21 groups. In some embodiments R.sub.19 is unsubstituted cycloalkyl. In some embodiments R.sub.19 is cycloalkyl substituted with one or more R.sub.21 groups. In some embodiments R.sub.19 is unsubstituted heteroalkyl. In some embodiments R.sub.19 is heteroalkyl substituted with one or more R.sub.21 groups. In some embodiments R.sub.19 is unsubstituted heterocycloalkyl. In some embodiments R.sub.19 is heterocycloalkyl substituted with one or more R.sub.21 groups. In some embodiments R.sub.19 is unsubstituted aryl. In some embodiments R.sub.19 is aryl substituted with one or more R.sub.21 groups. In some embodiments R.sub.19 is unsubstituted heteroaryl. In some embodiments R.sub.19 is heteroaryl substituted with one or more R.sub.21 groups.

(171) In various embodiments R.sub.20 is hydrogen, unsubstituted or substituted alkyl, unsubstituted or substituted cycloalkyl, unsubstituted or substituted heteroalkyl, unsubstituted or substituted heterocycloalkyl, unsubstituted or substituted aryl, or unsubstituted or substituted heteroaryl. In some embodiments R.sub.20 is hydrogen. In some embodiments R.sub.20 is unsubstituted alkyl. In some embodiments R.sub.20 is alkyl substituted with one or more R.sub.21 groups. In some embodiments R.sub.20 is unsubstituted cycloalkyl. In some embodiments R.sub.20 is cycloalkyl substituted with one or more R.sub.13 groups. In some embodiments R.sub.20 is unsubstituted heteroalkyl. In some embodiments R.sub.20 is heteroalkyl substituted with one or more R.sub.21 groups. In some embodiments R.sub.20 is unsubstituted heterocycloalkyl. In some embodiments R.sub.20 is heterocycloalkyl substituted with one or more R.sub.21 groups. In some embodiments R.sub.20 is unsubstituted aryl. In some embodiments R.sub.20 is aryl substituted with one or more R.sub.21 groups. In some embodiments R.sub.20 is unsubstituted heteroaryl. In some embodiments R.sub.20 is heteroaryl substituted with one or more R.sub.21 groups.

(172) In various embodiments R.sub.17 is halogen, OH, cyano, alkyl, alkoxy, cycloalkyl, heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene. In various embodiments R.sub.17 is halogen. In various embodiments R.sub.17 is OH. In various embodiments R.sub.17 is cyano. In various embodiments R.sub.17 is alkyl. In various embodiments R.sub.17 is cycloalkyl. In various embodiments R.sub.17 is heteroalkyl. In various embodiments R.sub.17 is heterocycloalkyl. In various embodiments R.sub.17 is aryl. In various embodiments R.sub.17 is heteroaryl. In various embodiments R.sub.17 is arylene. In various embodiments R.sub.17 is heteroarylene.

(173) In various embodiments R.sub.21 is halogen, OH, cyano, alkyl, alkoxy, cycloalkyl,

heteroalkyl, heterocycloalkyl, aryl, heteroaryl, arylene, or heteroarylene. In various embodiments R.sub.21 is halogen. In various embodiments R.sub.21 is OH. In various embodiments R.sub.21 is cyano. In various embodiments R.sub.21 is alkyl. In various embodiments R.sub.21 is cycloalkyl. In various embodiments R.sub.21 is heteroalkyl. In various embodiments R.sub.21 is heterocycloalkyl. In various embodiments R.sub.21 is aryl. In various embodiments R.sub.21 is heteroaryl. In various embodiments R.sub.21 is arylene. In various embodiments R.sub.21 is heteroarylene.

(174) In various embodiments C.sub.1 is a substituted or unsubstituted alkyl, embodiments C.sub.1 is a substituted or unsubstituted cycloalkylene, substituted or unsubstituted heterocycloalkylene, substituted or unsubstituted arylene, or substituted or unsubstituted heteroarylene. In some embodiments C.sub.1 is a cycloalkylene substituted with one or more R.sub.5 groups. In some embodiments C.sub.1 is an unsubstituted cycloalkylene. In some embodiments C.sub.1 is a heterocycloalkylene substituted with one or more R.sub.5 groups. In some embodiments C.sub.1 is an unsubstituted heterocycloalkylene. In some embodiments C.sub.1 is an arylene substituted with one or more R.sub.5 groups. In some embodiments C.sub.1 is an unsubstituted arylene. In some embodiments C.sub.1 is a heteroarylene substituted with one or more R.sub.5 groups. In some embodiments C.sub.1 is an unsubstituted heteroarylene. In some embodiments C.sub.1 is selected from the group consisting of

(175) ##STR00043##

(176) In some embodiments, C.sub.1 is

(177) ##STR00044##

(178) In some embodiments R.sub.5 is —OH. In some embodiments R.sub.5 is —CH.sub.2OH, in some embodiments R5 is alkyl.

(179) In various embodiments C.sub.2 is a bond, a cycloalkylene, heterocycloalkylene, arylene, or heteroarylene. In some embodiments C.sub.2 is a bond. In some embodiments C.sub.2 is an unsubstituted cycloalkylene. In some embodiments C.sub.2 is an unsubstituted heterocycloalkylene. In some embodiments C.sub.2 is an unsubstituted arylene. In some embodiments C.sub.2 is an unsubstituted heteroarylene. In some embodiments C.sub.2 is selected from the group consisting of

(180) ##STR00045##

(181) In some embodiments, C1 and C2 form a fused or spiro bicyclic ring. In some embodiments —C1-C2- form a fused bicyclic ring. In some embodiments —C1-C2-form a spiro bicyclic ring. In some embodiments —C1-C2- is

(182) ##STR00046##

In some embodiments —C1-C2- is

(183) ##STR00047##

In some embodiments —C1-C2- is

(184) ##STR00048##

(185) In some embodiments E is an electrophile capable of bonding with a K-Ras, H-Ras or N-Ras protein comprising G12C mutation. In some embodiments, the electrophile E is capable of forming an irreversible covalent bond with a G12C mutant K-Ras, H-Ras or N-Ras protein. In some cases E is the general structure

(186) ##STR00049##

(187) R25 is alkyl. R.sub.26 is cyano or alkyl. R.sub.27 is alkyl. R.sub.26 and R.sub.27 can form cycloalkene.

(188) In some of the foregoing embodiments of compounds of Formula III, E is selected from:

(189) ##STR00050##

(190) In some embodiments E is

(191) ##STR00051##

In some embodiments E is

(192) ##STR00052##

In some embodiments E is

(193) ##STR00053##

In some embodiments E is

(194) ##STR00054##

(195) In some embodiments, R.sub.14 is capable of reversible interaction with K-Ras, H-Ras or N-Ras G12C mutant protein. In some embodiments R.sub.14 moiety has high affinity towards K-Ras, H-Ras or N-Ras and is highly specific towards G12C K-Ras, H-Ras or N-Ras. In some embodiments R.sub.14 is capable of hydrophobic interaction with K-Ras, H-Ras or N-Ras G12C. In some embodiments R.sub.14 is able to form hydrogen bonds with various residues of G12C K-Ras, H-Ras or N-Ras protein. In some embodiments R.sub.14 interacts with one or more of G10, R68, Y71, Y96 or Q99 residues in K-Ras G12C (FIG. 1). In some embodiments, R.sub.14 moiety interacts with the G10 residue of K-Ras G12C. In some embodiments R.sub.14 interacts with the R68 residue of K-Ras G12C. In some embodiments R.sub.14 interacts with the Y71 residue of K-Ras G12C. In some embodiments R.sub.14 interacts with the Y96 residue of K-Ras G12C. In some embodiments R.sub.14 interacts with the Q99 residue of K-Ras G12C.

(196) In some embodiments, the

(197) ##STR00055##

unit in Formula III provides proper length and geometry to the compound such that the electrophile E is able to interact with the cysteine residue at the 12 position in G12C K-Ras, H-Ras or N-Ras protein. In some embodiments the

(198) ##STR00056##

also interacts with other protein backbone residues.

(199) In some embodiments the invention provides compounds of Formula III as shown in Table 3.

(200) In some embodiments the invention provides compounds shown in Table 4.

(201) In still other embodiments, the invention provides a compound having the following structure (V):

(202) ##STR00057##

or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

(203) R.sup.1 is aryl or heteroaryl;

(204) R.sup.30a and R.sup.30b are, at each occurrence, independently H, —OH, —NH.sub.2, —CO.sub.2H, cyano, cyanoalkyl, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl; or R.sup.30a and R.sup.30b join to form a carbocyclic or heterocyclic ring; or R.sup.30a is H, —OH, —NH.sub.2, —CO.sub.2H, cyano, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R.sup.30b joins with R.sup.31b to form a carbocyclic or heterocyclic ring;

(205) R.sup.31a and R.sup.31b are, at each occurrence, independently H, —OH, —NH.sub.2, —CO.sub.2H, cyano, cyanoalkyl, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl; or R.sup.31a and R.sup.31b join to form a carbocyclic or heterocyclic ring; or R.sup.31a is H, —OH, —NH.sub.2, —CO.sub.2H, cyano, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R.sup.31b joins with R.sup.30b to form a carbocyclic or heterocyclic ring;

(206) R.sup.32a and R.sup.32b are, at each occurrence, independently H, —OH, —NH.sub.2, —CO.sub.2H, cyano, cyanoalkyl, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl; or R.sup.32a and R.sup.32b join to form a carbocyclic or heterocyclic ring; or R.sup.32a is H, —OH, —NH.sub.2, —CO.sub.2H, cyano, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R.sup.32b joins with R.sup.33b to form a carbocyclic or heterocyclic ring;

(207) R.sup.33a and R.sup.33b are, at each occurrence, independently H, —OH, —NH.sub.2, —CO.sub.2H, cyano, cyanoalkyl, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl,

aminoalkyl, carboxylalkyl or aminocarbonyl; or R.sup.33a and R.sup.33b join to form a carbocyclic or heterocyclic ring; or R.sup.33a is H, —OH, —NH.sub.2, —CO.sub.2H, cyano, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R.sup.33b joins with R.sup.32b to form a carbocyclic or heterocyclic ring;

(208) L.sup.1 is carbonyl, —NHC(=O)—, alkylene, alkenylene, heteroalkylene, heterocycloalkylene, heteroarylene, alkylenecarbonyl, alkenylenecarbonyl, heteroalkylenecarbonyl, heterocycloalkylenecarbonyl or heteroarylenecarbonyl;

(209) L.sup.2 is a bond or alkylene;

(210) G.sup.1, G.sup.2' G.sup.3 and G.sup.4 are each independently N or CR, where R is H, cyano, halo or C.sub.1-C.sub.6alkyl;

(211) n.sup.1, n.sup.2, n.sup.3 and n.sup.4 are each independently 1, 2 or 3; and

(212) E is an electrophilic moiety capable of forming a covalent bond with the cysteine residue at position 12 of a K-Ras, H-Ras or N-Ras G12C mutant protein.

(213) In some embodiments of the compounds of structure V, L.sup.1 is carbonyl, —NHC(=O)—, alkylene, heteroalkylene, alkylenecarbonyl or heteroalkylenecarbonyl;

(214) In some other embodiments, the compound has the following structure (Va):

(215) ##STR00058##

wherein:

(216) L.sup.1a is a bond, —NH—, alkylene, alkenylene, heteroalkylene, heterocycloalkylene or heteroarylene.

(217) In other embodiments of compound (Va), L.sup.1a is a bond, —NH—, alkylene or heteroalkylene

(218) In some more embodiments, the compound has the following structure (Vb):


(219) ##STR00059##

wherein:

(220) Q is —C(=O)—, —NR.sup.34C(=O)—, —S(=O).sub.2— or —NR.sup.34S(=O).sub.2—;

(221) R.sup.34 is H, C.sub.1-C.sub.6alkyl or hydroxylalkyl;

(222)  custom character is a carbon-carbon double bond or a carbon-carbon triple bond; and

(223) R.sup.35 and R.sup.36 are each independently H, cyano, C.sub.1-C.sub.6alkyl, aminoalkyl, alkylaminoalkyl, or hydroxylalkyl or R.sup.35 and R.sup.36 join to form a carbocyclic or heterocyclic ring when  custom character is a double bond; or R.sup.35 is absent and R.sup.36 is H, C.sub.1-C.sub.6alkyl, aminoalkyl, alkylaminoalkyl or hydroxylalkyl when  custom character is a triple bond.

(224) In some different embodiments, the compound has one of the following structures (Vc), (Vd), (Ve) or (Vf):

(225) ##STR00060##

(226) In still other embodiments, wherein the compound has one of the following structures (Vg), (Vh), (Vi) or (Vj):

(227) ##STR00061##

(228) In some other embodiments, the compound has one of the following structures (Vk), (Vl), (Vm), (Vn); (Vo) or (Vp):

(229) ##STR00062##

(230) In various other embodiments, R.sup.1 is aryl. For example, in some embodiments the aryl is bicyclic, such as a fused bicyclic aryl. In some more specific embodiments, the aryl is naphthyl.

(231) In various other embodiments, the aryl is monocyclic. For example, in some embodiments the aryl is phenyl.

(232) In some of the foregoing embodiments, the aryl is unsubstituted. In other of the foregoing embodiments, the aryl is substituted with one or more substituents. For example, in some embodiments the substituents are selected from halo, hydroxyl, cyano, aminocarbonyl, formyl, C.sub.1-C.sub.6alkyl, C.sub.1-C.sub.6alkylsulfonyl, C.sub.1-C.sub.6haloalkyl, C.sub.3-

C.sub.8cycloalkyl, C1-C.sub.6alkoxy, C.sub.1-C.sub.6hydroxyalkyl, C.sub.1-C.sub.6alkoxyalkyl, C.sub.1-C.sub.6aminoalkyl, aliphatic heterocyclyl, heteroaryl and aryl.

(233) In other embodiments, the aryl substituents are selected from fluoro, chloro, bromo, iodo, hydroxyl, cyano, methyl, ethyl, isopropyl, methylsulfonyl, methoxy, aminocarbonyl, trifluoromethyl, 2,2,2-trifluorethyl, cyclobutyl, cyclopropyl and phenyl, wherein the cyclopropyl and phenyl are optionally substituted with one or more substituents selected from C.sub.1-C.sub.6alkyl, halo, hydroxyl and cyano

(234) In some different embodiments, the substituents are selected from fluoro, chloro, bromo, iodo, hydroxyl, cyano, methyl, ethyl, methylsulfonyl, methoxy, aminocarbonyl, trifluoromethyl, cyclopropyl and phenyl, wherein the cyclopropyl and phenyl are optionally substituted with one or more substituents selected from halo, hydroxyl and cyano.

(235) In other exemplary embodiments, the aryl substituents are selected from fluoro, chloro, bromo, iodo, hydroxyl, methyl, ethyl, cyclobutyl and cyclopropyl, wherein the cyclopropyl is optionally substituted with one or more substituents selected from C.sub.1-C.sub.6alkyl, halo, hydroxyl and cyano

(236) In some more embodiments, the substituents are selected from fluoro, chloro, bromo, iodo, hydroxyl, methyl, ethyl and cyclopropyl, wherein the cyclopropyl is optionally substituted with one or more substituents selected from halo, hydroxyl and cyano.

(237) In still more embodiments, the substituents are selected from fluoro, chloro, bromo, hydroxyl and cyclopropyl, wherein the cyclopropyl is optionally substituted with one or more substituents selected from C.sub.1-C.sub.6alkyl, halo, hydroxyl and cyano.

(238) In some more specific embodiments, the substituents are selected from fluoro, chloro, bromo, hydroxyl and cyclopropyl, wherein the cyclopropyl is optionally substituted with one or more substituents selected from halo, hydroxyl and cyano. For example, in some embodiments the cyclopropyl comprises a geminal difluoro substitution.

(239) In still other embodiments, R^{sup.1} has one of the following structures:

(240) ##STR00063## ##STR00064## ##STR00065##

(241) In still other embodiments, R^{sup.1} has one of the following structures:

(242) ##STR00066## ##STR00067## ##STR00068## ##STR00069##

(243) In still other embodiments, R^{sup.1} is heteroaryl. For example, in some embodiments the heteroaryl is bicyclic, such as a fused bicyclic heteroaryl.

(244) In some more embodiments, the heteroaryl is monocyclic.

(245) In some of the foregoing embodiments, the heteroaryl comprises nitrogen, sulfur or a combination thereof. For example, in some embodiments the heteroaryl is dihydroquinoxaliny, indoleyl, benzoimidazolyl, pyridinyl or thiazolyl.

(246) In some embodiments, the heteroaryl is unsubstituted. In some other embodiments, the heteroaryl is substituted with one or more substituents. In some embodiments, the substituents are selected from C.sub.1-C.sub.6alkyl, halo and oxo. For example, in some embodiments the substituents are selected from halo and oxo. In other embodiments, the substituents are selected from ethyl and chloro. In some more specific embodiments, the substituents are chloro.

(247) In some embodiments of the foregoing compounds of structure (V), R^{sup.1} has one of the following structures:

(248) ##STR00070##

wherein R^{sup.1a} is, at each occurrence, independently H, C.sub.1-C.sub.6alkyl or halo.

(249) In various other embodiments, R^{sup.1} has one of the following structures:

(250) ##STR00071##

wherein R^{sup.1a} is, at each occurrence, independently H or halo.

(251) In still other embodiments of structure (V), R^{sup.1} has one of the following structures:

(252) ##STR00072##

(253) In some embodiments, Q is —C(=O)—. In some other embodiments, Q is —S(=O).sub.2—.

In still other embodiments, Q is $\text{—NR}^{\text{sup.34}}\text{C(=O)—}$. In still more other embodiments, Q is $\text{—NR}^{\text{sup.34}}\text{S(=O)}_2\text{—}$.

(254) In some more specific embodiments, $\text{R}^{\text{sup.34}}$ is H. For example, in some embodiments $\text{R}^{\text{sup.34}}$ is hydroxylalkyl, such as 2-hydroxylalkyl.

(255) In other of the foregoing embodiments, at least one of $\text{R}^{\text{sup.35}}$ or $\text{R}^{\text{sup.36}}$ is H. For example, in some embodiment search of $\text{R}^{\text{sup.35}}$ and $\text{R}^{\text{sup.36}}$ are H.

(256) In various other embodiments, $\text{R}^{\text{sup.36}}$ is alkylaminoalkyl. For example, in some embodiments $\text{R}^{\text{sup.36}}$ has the following structure:

(257) ##STR00073##

(258) In some different embodiments, $\text{R}^{\text{sup.36}}$ is hydroxylalkyl, for example 2-hydroxylalkyl

(259) In various other embodiments, $\text{R}^{\text{sup.35}}$ and $\text{R}^{\text{sup.36}}$ join to form a ring. In some of these embodiments, the ring is a cyclopentene, cyclohexene or phenyl ring.

(260) In other of the foregoing embodiments, E has one of the following structures:

(261) ##STR00074##

(262) In some embodiments, E is

(263) ##STR00075##

(264) In some more of the foregoing embodiments, $\text{L}^{\text{sup.1}}$ is heteroalkylene. In some more embodiments, the heteroalkylene is unsubstituted. In some different embodiments, the heteroalkylene is substituted.

(265) In various other embodiments, L is aminoalkylene. For example, in some embodiments $\text{L}^{\text{sup.1}}$ is $\text{—CH}_2\text{CH}_2\text{NH—}$.

(266) In other embodiments of the foregoing, $\text{L}^{\text{sup.1}}$ is heterocycloalkylene or heteroarylene. In some embodiments, the heterocycloalkylene or heteroarylene is unsubstituted. In other embodiments, the heterocycloalkylene or heteroarylene is substituted. In some further embodiments, $\text{L}^{\text{sup.1}}$ has one of the following structures:

(267) ##STR00076##

(268) In some different embodiments, $\text{L}^{\text{sup.1a}}$ is a bond.

(269) In some embodiments, $\text{L}^{\text{sup.1a}}$ is alkylene, alkenylene, heteroalkylene or heterocycloalkylene. In some other embodiments, $\text{L}^{\text{sup.1a}}$ is alkylene or heteroalkylene. In some of these embodiments, $\text{L}^{\text{sup.1a}}$ is substituted alkylene. In various other embodiments, $\text{L}^{\text{sup.1a}}$ is unsubstituted alkylene. For example, in some embodiments $\text{L}^{\text{sup.1a}}$ is

(270) ##STR00077##

(271) In some different embodiments, $\text{L}^{\text{sup.1a}}$ is substituted heteroalkylene. In some other embodiments, $\text{L}^{\text{sup.1a}}$ is unsubstituted heteroalkylene. In some of the foregoing embodiments, $\text{L}^{\text{sup.1a}}$ is aminoalkylene or thioalkylene, for example aminoalkylene. For example, in some embodiments $\text{L}^{\text{sup.1a}}$ has one of the following structures:

(272) ##STR00078##

(273) In other embodiments, $\text{L}^{\text{sup.1a}}$ is

(274) ##STR00079##

(275) In other embodiments, $\text{L}^{\text{sup.1a}}$ is substituted alkenylene. In different embodiments, $\text{L}^{\text{sup.1a}}$ is unsubstituted alkenylene. In some more specific embodiments, $\text{L}^{\text{sup.1a}}$ has the following structure:

(276) ##STR00080##

(277) In yet other embodiments, $\text{L}^{\text{sup.1a}}$ is substituted heterocycloalkylene. In some other embodiments, $\text{L}^{\text{sup.1a}}$ is unsubstituted heterocycloalkylene. For Example, in some embodiments, $\text{L}^{\text{sup.1a}}$ has the following structure:

(278) ##STR00081##

(279) In some of the foregoing embodiments, $\text{L}^{\text{sup.2}}$ is a bond.

(280) In various other embodiments, $\text{L}^{\text{sup.2}}$ is substituted alkylene. In still other embodiments, $\text{L}^{\text{sup.2}}$ is unsubstituted alkylene.

(281) In various embodiments of any of the foregoing compounds of structure (V):

(282) R.sup.30a and R.sup.30b are, at each occurrence, independently H, —OH, —NH.sub.2, —CO.sub.2H, cyano, cyanoalkyl, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl;

(283) R.sup.31a and R.sup.31b are, at each occurrence, independently H, —OH, —NH.sub.2, —CO.sub.2H, cyano, cyanoalkyl, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl;

(284) R.sup.32a and R.sup.32b are, at each occurrence, independently H, —OH, —NH.sub.2, —CO.sub.2H, cyano, cyanoalkyl, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl; and

(285) R.sup.33a and R.sup.33b are, at each occurrence, independently H, —OH, —NH.sub.2, —CO.sub.2H, cyano, cyanoalkyl, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl.

(286) In other embodiments, R.sup.30a, R.sup.30b, R.sup.31a, R.sup.31b, R.sup.32a, R.sup.32b, R.sup.33a and R.sup.33b are selected from H, C.sub.1-C.sub.6alkyl, hydroxylalkyl, cyano, cyanoalkyl and aminocarbonyl, for example H, C.sub.1-C.sub.6alkyl, hydroxylalkyl, cyano, and aminocarbonyl or in other embodiments H, C.sub.1-C.sub.6alkyl and hydroxylalkyl.

(287) In some of the foregoing embodiments, at least one of R.sup.30a, R.sup.30b, R.sup.31a, R.sup.31b, R.sup.32a, R.sup.32b, R.sup.33a or R.sup.33b is H. For example, in some embodiments each of R.sup.30a, R.sup.30b, R.sup.31a, R.sup.31b, R.sup.32a, R.sup.32b, R.sup.33a or R.sup.33b is H.

(288) In some other of the foregoing embodiments, at least one of R.sup.30a, R.sup.30b, R.sup.31a, R.sup.31b, R.sup.32a, R.sup.32b, R.sup.33a or R.sup.33b is hydroxylalkyl.

(289) In still other of the foregoing embodiments, at least one of R.sup.30a, R.sup.30b, R.sup.31a, R.sup.31b, R.sup.32a, R.sup.32b, R.sup.33a or R.sup.33b is cyano.

(290) In still more of the foregoing embodiments of compound (V), at least one of R.sup.30a, R.sup.30b, R.sup.31a, R.sup.31b, R.sup.32a, R.sup.32b, R.sup.33a or R.sup.33b is aminocarbonyl.

(291) In other embodiments, at least one of R.sup.30a, R.sup.30b, R.sup.31a, R.sup.31b, R.sup.32a, R.sup.32b, R.sup.33a or R.sup.33b is C.sub.1-C.sub.6alkyl.

(292) In some embodiments, R.sup.30a and R.sup.30b join to form a carbocyclic or heterocyclic ring. In different embodiments, R.sup.31a and R.sup.31b join to form a carbocyclic or heterocyclic ring. In more embodiments, R.sup.32a and R.sup.32b join to form a carbocyclic or heterocyclic ring. In yet other embodiments, R.sup.33a and R.sup.33b join to form a carbocyclic or heterocyclic ring.

(293) In even other embodiments, R.sup.30a is H, —OH, —NH.sub.2, —CO.sub.2H, cyano, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R.sup.30b joins with R.sup.31b to form a carbocyclic or heterocyclic ring.

(294) In more embodiments, R.sup.31a is H, —OH, —NH.sub.2, —CO.sub.2H, cyano, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R.sup.31b joins with R.sup.30b to form a carbocyclic or heterocyclic ring.

(295) In other embodiments, R.sup.32a is H, —OH, —NH.sub.2, —CO.sub.2H, cyano, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R.sup.32b joins with R.sup.33b to form a carbocyclic or heterocyclic ring.

(296) In still more embodiments, R.sup.33a is H, —OH, —NH.sub.2, —CO.sub.2H, cyano, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R.sup.33b joins with R.sup.32b to form a carbocyclic or heterocyclic ring.

(297) In some other embodiments, the compound is selected from a compound in Table 5.

(298) Compounds of structure V are prepared according to procedures well-known or derivable by one of ordinary skill in the art, for example by procedures analogous to those exemplified in Examples 8, 9, 18 and other examples provided below. Each of the compounds in Table 5 was

prepared in such a manner and analyzed by mass spectrometry and/or ¹H NMR. The mass spectrum ([M+H.^{sup.}+] or [M+Na.^{sup.}+]) and/or NMR spectrum was found to be consistent with the structures in Table V.

(299) General Reaction Scheme I illustrates an exemplary procedure for preparing compounds of structure (V).

(300) ##STR00082##

(301) Referring to General Reaction Scheme I, (V') and (VI') are available from commercial sources and/or are easily prepared according to procedures known in the art. All variables on (V') and (V''), with the exception of M.^{sup.}1, are as defined above. In some procedures, M.^{sup.}1 is NH. Briefly, an appropriately substituted acid (V') is activated and reacted with an appropriately substituted heterocycle (V'') under appropriate coupling conditions. The L.^{sup.}2-E moiety may be present in (V'') as illustrated or may be installed after coupling. For example L.^{sup.}2-E may be installed before or after coupling via acylation (or thioacylation) using a reagent such as an acid chloride or thionyl chloride.

(302) It should be noted that variations of the above procedure are possible, some of which are exemplified in the examples. For example, in some procedures (V'') is monocyclic and the second cyclic moiety is added after the coupling step. In other procedures, the acid moiety is present on the cyclic moiety (V'') and R.^{sup.}1 is appropriately substituted with a nucleophilic moiety to enable coupling to form (Va).

(303) Various other options are available to one of ordinary skill in the art to add various substituents and or modify or reorder the above described steps to arrive at different embodiments of compounds of structure V. It should also be noted that various substitutions on (V') and/or (V'') can be present during the coupling step (in protected or unprotected form) or the substituents can be added after (V') and (V'') are coupled. Methods for inclusion of these substituents are known in the art.

(304) It is understood that although an exemplary procedure for prepare (Va) is provided above, other compounds of structure (V) can be prepared by analogous methods. For example, the carbonyl of (Va) may be reduced to form compounds of structure (V) wherein L.^{sup.}1 does not comprise a carbonyl. Embodiments wherein L.^{sup.}1 is heterocycloalkylene or heteroarylene can be prepared from analogous methods, for example by use of Buchwald chemistry to include the heterocycloalkylene or heteroarylene portion. Other methods for preparation of different compounds of structure (V) are known in the art.

(305) Briefly, an appropriately substituted acid is reacted with an appropriately substituted heterocycle under amide coupling conditions. Acylation (or thioacylation) using a reagent such as an acid chloride or thionyl chloride results in compounds of structure V. Various options are available to one of ordinary skill in the art to add various substituents and/or modify or reorder the above described steps to arrive at different embodiments of compounds of structure V. The appropriate acid is purchased commercially or made according to well-known procedures.

(306) In still other embodiments, the invention provides a compound having the following structure (VI):

(307) ##STR00083##

or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

(308) A is CR.^{sup.}37b, N or NR.^{sup.}38a;

(309) B is CR.^{sup.}37c, N, NR.^{sup.}38b or S

(310) C is CR.^{sup.}37d, N, NR.^{sup.}38e or S

(311) G.^{sup.}3 and G.^{sup.}4 are each independently N or CR, wherein R is H, cyano, halo or C.^{sup.}1-C.^{sup.}6alkyl;

(312) L.^{sup.}1a is a bond, —NH—, alkylene or heteroalkylene

(313) L.^{sup.}2 is a bond or alkylene;

(314) R.^{sup.}32a and R.^{sup.}32b are, at each occurrence, independently H, —OH, —NH.^{sub.}2, —

CO.sub.2H, cyano, cyanoalkyl, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl; or R.sup.32a and R.sup.32b join to form a carbocyclic or heterocyclic ring; or R.sup.32a is H, —OH, —NH.sub.2, —CO.sub.2H, cyano, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R.sup.32b joins with R.sup.33b to form a carbocyclic or heterocyclic ring; (315) R.sup.33a and R.sup.33b are, at each occurrence, independently H, —OH, —NH.sub.2, —CO.sub.2H, cyano, cyanoalkyl, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl; or R.sup.33a and R.sup.33b join to form a carbocyclic or heterocyclic ring; or R.sup.33a is H, —OH, —NH.sub.2, —CO.sub.2H, cyano, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R.sup.33b joins with R.sup.32b to form a carbocyclic or heterocyclic ring; (316) R.sup.37a, R.sup.37, R.sup.37c, R.sup.37d and R.sup.37e are each independently H, halo, oxo, hydroxyl, cyano, aminocarbonyl, formyl, C.sub.1-C.sub.6alkyl, C.sub.1-C.sub.6alkylsulfonyl, C.sub.1-C.sub.6haloalkyl, C.sub.3-C.sub.8cycloalkyl, C.sub.1-C.sub.6alkoxy, C.sub.1-C.sub.6hydroxylalkyl, C.sub.1-C.sub.6alkoxyalkyl, C.sub.1-C.sub.6aminoalkyl, heterocyclyl or aryl;

(317) R.sup.38a, R.sup.38b and R.sup.38c are each independently H, C.sub.1-C.sub.6alkyl or aryl;

(318) n.sup.3 and n.sup.4 are each independently 1, 2 or 3

(319) m is 0 or 1;

(320)  is a single or double bond such that all valences are satisfied; and

(321) E is an electrophilic moiety capable of forming a covalent bond with the cysteine residue at position 12 of a K-Ras, H-Ras or N-Ras G12C mutant protein.

(322) In various other embodiments, the compound has one of the following structures (VIa), (VIb), (VIc), (VId), (VIf) or (VIg):

(323) ##STR00084## ##STR00085##


(324) In some different embodiments, the compound has one of the following structures (VIa'), (VIb'), (VIe'), (VId'), (VIf') or (VIg'):



(325) ##STR00086##

wherein:

(326) Q is —C(=O)—, —NR.sup.34C(=O)—, —S(=O).sub.2— or —NR.sup.34S(=O).sub.2—;

(327) R.sup.34 is H, C.sub.1-C.sub.6alkyl or hydroxylalkyl;

(328)  is a carbon-carbon double bond or a carbon-carbon triple bond; and

(329) R.sup.35 and R.sup.36 are each independently H, cyano, C.sub.1-C.sub.6alkyl, aminoalkyl, alkylaminoalkyl, or hydroxylalkyl or R.sup.35 and R.sup.36 join to form a carbocyclic or heterocyclic ring when  is a double bond; or R.sup.35 is absent and R.sup.36 is H, C.sub.1-C.sub.6alkyl, aminoalkyl, alkylaminoalkyl or hydroxylalkyl when  is a triple bond.

(330) In some specific embodiments of the foregoing compounds of structure (VI), and substructures thereof, R.sup.37a is halo, aryl or heteroaryl. In further such embodiments, R.sup.35 and R.sup.36 are each H.

(331) In various other embodiments, G.sup.3 is N and G.sup.4 is CR, for example CH.

(332) In some different embodiments, G.sup.3 is CR, for example, CH, and G.sup.4 is N.

(333) In still other embodiments, G.sup.3 is N and G.sup.4 is N.

(334) In various other embodiments, n.sup.3 is 2 and n.sup.4 is 2. In still other embodiments, n.sup.3 is 1 and n.sup.4 is 1. In some more embodiments, n.sup.3 is 2 and n.sup.4 is 1.

(335) In other of the foregoing embodiments, R.sup.37a, R.sup.37b, R.sup.37c, R.sup.37d and R.sup.37e are each independently H, —OH, halo, oxo, C.sub.1-C.sub.6alkyl, C.sub.1-C.sub.6alkoxy, heterocyclyl or aryl.

(336) In still other embodiments, R.sup.37a, R.sup.37b, R.sup.37c, R.sup.37d and R.sup.37e are each independently H, —OH, fluoro, chloro, bromo, iodo, oxo, methyl, methoxy, heteroaryl or

aryl.

(337) In some embodiments, R.sup.37a or R.sup.37e is aryl. In some more specific embodiments, R.sup.37a is aryl, such as phenyl.

(338) In some different embodiments, the aryl is unsubstituted. In some other embodiments, the aryl is substituted. For example, in some embodiments the aryl is substituted with one or more halo substituents. In some of these embodiments, the halo substituents are selected from fluoro and chloro.

(339) In still other embodiments, R.sup.37a is heteroaryl. In some of these embodiments, the heteroaryl is unsubstituted. In various other embodiments, the heteroaryl is substituted. In some more embodiments, the heteroaryl comprises nitrogen, sulfur or a combination thereof.

(340) In some more specific embodiments, the heteroaryl is thiophenyl.

(341) In other of the foregoing embodiments, R.sup.37a is halo. For example, in some embodiments halo is chloro, bromo or iodo.

(342) In some embodiments, R.sup.37a or R.sup.37e has one of the following structures:

(343) ##STR00087##

(344) In still other embodiments, R.sup.37a has one of the following structures:

(345) ##STR00088##

(346) In various different embodiments, R.sup.38a, R.sup.38b and R.sup.38c are each independently H or aryl. In still other embodiments, R.sup.38a, R.sup.38b and R.sup.38c are each independently H.

(347) In some other different embodiments, R.sup.38c is aryl. For example, in some embodiments the aryl is substituted with one or more halo substituents. In some of these embodiments, halo is chloro.

(348) In some other embodiments of the compounds of structure (VI), Q is —C(=O)—. In some other embodiments, Q is —S(=O).sub.2—. In still other embodiments, Q is —NR.sup.34C(=O)—. In still more other embodiments, Q is —NR.sup.34S(=O).sub.2—.

(349) In some more specific embodiments, R.sup.34 is H. For example, in some embodiments R.sup.34 is hydroxylalkyl, such as 2-hydroxylalkyl.

(350) In other of the foregoing embodiments, at least one of R.sup.35 or R.sup.36 is H. For example, in some embodiment search of R.sup.35 and R.sup.36 are H.

(351) In various other embodiments, R.sup.36 is alkylaminoalkyl. For example, in some embodiments R.sup.36 has the following structure:

(352) ##STR00089##

(353) In some different embodiments, R.sup.36 is hydroxylalkyl, for example 2-hydroxylalkyl

(354) In various other embodiments, R.sup.35 and R.sup.36 join to form a ring. In some of these embodiments, the ring is a cyclopentene, cyclohexene or phenyl ring.

(355) In other of the foregoing embodiments, E has one of the following structures:

(356) ##STR00090##

(357) In some embodiments, E is

(358) ##STR00091##

(359) In some more of the foregoing embodiments, L.sup.1 is heteroalkylene. In some more embodiments, the heteroalkylene is unsubstituted. In some different embodiments, the heteroalkylene is substituted.

(360) In various other embodiments, L is aminoalkylene. For example, in some embodiments L.sup.1 is —CH.sub.2CH.sub.2NH—.

(361) In some different embodiments, L.sup.1a is a bond.

(362) In some embodiments, L.sup.1a is alkylene, alkenylene, heteroalkylene or heterocycloalkylene. In some other embodiments, L.sup.1a is alkylene or heteroalkylene. In some of these embodiments, L.sup.1a is substituted alkylene. In various other embodiments, L.sup.1a is unsubstituted alkylene. For example, in some embodiments L.sup.1a is

(363) ##STR00092##

(364) In some different embodiments, L.sup.1a is substituted heteroalkylene. In some other embodiments, L.sup.1a is unsubstituted heteroalkylene. In some of the foregoing embodiments, L.sup.1a is aminoalkylene or thioalkylene, for example aminoalkylene. For example, in some embodiments L.sup.1a has one of the following structures:

(365) ##STR00093##

(366) In other embodiments, L.sup.1a is

(367) ##STR00094##

(368) In other embodiments, L.sup.1a is substituted alkenylene. In different embodiments, L.sup.1a is unsubstituted alkenylene. In some more specific embodiments, L.sup.1a has the following structure:

(369) ##STR00095##

(370) In yet other embodiments, L.sup.1a is substituted heterocycloalkylene. In some other embodiments, L.sup.1a is unsubstituted heterocycloalkylene. For Example, in some embodiments, L.sup.1a has the following structure:

(371) ##STR00096##

(372) In some of the foregoing embodiments, L.sup.2 is a bond.

(373) In various other embodiments, L.sup.2 is substituted alkylene. In still other embodiments, L.sup.2 is unsubstituted alkylene.

(374) In some embodiments of any of the foregoing compounds of structure (VI):

(375) R.sup.32a and R.sup.32b are, at each occurrence, independently H, —OH, —NH.sub.2, —CO.sub.2H, cyano, cyanoalkyl, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl; and

(376) R.sup.33a and R.sup.33b are, at each occurrence, independently H, —OH, —NH.sub.2, —CO.sub.2H, cyano, cyanoalkyl, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl.

(377) In other embodiments, R.sup.32a, R.sup.32b, R.sup.33a or R.sup.33b are selected from H, C.sub.1-C.sub.8alkyl, hydroxylalkyl, cyano, cyanoalkyl and aminocarbonyl, for example H, hydroxyl alkyl and cyano.

(378) In other of the foregoing embodiments, at least one of R.sup.32a, R.sup.32b, R.sup.33a or R.sup.33b is H. For example, in some embodiments each of R.sup.32a, R.sup.32b, R.sup.33a or R.sup.33b is H.

(379) In other of the foregoing embodiments, at least one of R.sup.32a, R.sup.32b, R.sup.33a or R.sup.33b is hydroxylalkyl.

(380) In still other embodiments, at least one of R.sup.32a, R.sup.32b, R.sup.33a or R.sup.33b is cyano.

(381) In some other different embodiments, least one of R.sup.32a, R.sup.32b, R.sup.33a or R.sup.33b is aminocarbonyl.

(382) In some embodiments, R.sup.32a and R.sup.32b join to form a carbocyclic or heterocyclic ring. In other embodiments, R.sup.33a and R.sup.33b join to form a carbocyclic or heterocyclic ring.

(383) In different embodiments, R.sup.32a is H, —OH, —NH.sub.2, —CO.sub.2H, cyano, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R.sup.32b joins with R.sup.33b to form a carbocyclic or heterocyclic ring.

(384) In still other embodiments, R.sup.33a is H, —OH, —NH.sub.2, —CO.sub.2H, cyano, C.sub.1-C.sub.6alkyl, C.sub.3-C.sub.8cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R.sup.33b joins with R.sup.32b to form a carbocyclic or heterocyclic ring.

(385) In some more specific embodiments, the compound is selected from a compound in Table 6.

(386) Compounds of structure VI are prepared according to procedures well-known or derivable by one of ordinary skill in the art, for example by procedures analogous to those exemplified in

Examples 13, 17, 19, 20, 22 and other examples provided below. Each of the compounds in Table 6 was prepared in such a manner and analyzed by mass spectrometry and/or ^1H NMR. The mass spectrum ($[\text{M}+\text{H}]^+$ or $[\text{M}+\text{Na}]^+$) and/or NMR spectrum was found to be consistent with the structures in Table VI.

(387) General Reaction Scheme II illustrates an exemplary procedure for preparing compounds of structure (VI).

(388) ##STR00097##

(389) Referring to General Reaction Scheme II, (VI') and (VI'') are available from commercial sources and/or are easily prepared according to procedures known in the art. All variables on (VI') and (VI''), with the exception of M_1 and M_2 , are as defined above. In some procedures, M_1 is NH and M_2 is absent. In other procedures M_1 is N or CH and M_2 is a precursor to L_1a which reacts with an activated acid. For example, in various procedures M_2 is $\text{NH}_{2.2}$, aminoalkyl or other heterosubstituted alkyl. Embodiments where M_2 comprises a carbanion (or M_1 is a carbanion) are also contemplated such that L_1 is alkylene. Briefly, an appropriately substituted acid (VI') is activated and reacted with an appropriately substituted heterocycle (VI'') under appropriate coupling conditions. The $\text{L}_2\text{-E}$ moiety may be present in (VI'') as illustrated or may be installed after coupling. For example $\text{L}_2\text{-E}$ may be installed before or after coupling via acylation (or thioacylation) using a reagent such as an acid chloride or thionyl chloride.

(390) It should be noted that variations of the above procedure are possible, some of which are exemplified in the examples. For example, in some procedures, the acid moiety is present on the cyclic moiety (VI') and (VI') is appropriately substituted with a nucleophilic moiety to enable coupling to form (VI). Other methods of bond formation, which do not require reaction of an activated acid are also available for preparation of the compounds. It should also be noted that various substitutions on (VI') and/or (VI'') can be present during the coupling step (in protected or unprotected form) or the substituents can be added after (VI') and (VI'') are coupled. Methods for inclusion of these substituents are known in the art.





















































































































































(391) Various options are available to one of ordinary skill in the art to add various substituents and/or modify or reorder the above described steps to arrive at different embodiments of compounds of structure VI. The appropriate acid is purchased commercially or made according to well-known procedures.

(392) It will also be appreciated by those skilled in the art that in the processes described herein (e.g., General Reaction Scheme I and II and the below examples) the functional groups of intermediate compounds may need to be protected by suitable protecting groups. Such functional groups include hydroxy, amino, mercapto and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl or diarylalkylsilyl (for example, t-butyldimethylsilyl, t-butyldiphenylsilyl or trimethylsilyl), tetrahydropyranyl, benzyl, and the like. Suitable protecting groups for amino, amidino and guanidino include t-butoxycarbonyl, benzyloxycarbonyl, and the like. Suitable protecting groups for mercapto include $-\text{C}(\text{O})-\text{R}''$ (where R'' is alkyl, aryl or arylalkyl), p-methoxybenzyl, trityl and the like. Suitable protecting groups for carboxylic acid include alkyl, aryl or arylalkyl esters. Protecting groups may be added or removed in accordance with standard techniques, which are known to one skilled in the art and as described herein. The use of protecting groups is described in detail in Green, T. W. and P. G. M. Wutz, Protective Groups in Organic Synthesis (1999), 3rd Ed., Wiley. As one of skill in the art would appreciate, the protecting group may also be a polymer resin such as a Wang resin, Rink resin or a 2-chlorotrityl-chloride resin.









(393) It will also be appreciated by those skilled in the art, although such protected derivatives of compounds of this invention may not possess pharmacological activity as such, they may be administered to a mammal and thereafter metabolized in the body to form compounds of the invention which are pharmacologically active. Such derivatives may therefore be described as























“prodrugs”. All prodrugs of compounds of this invention are included within the scope of the invention.












(394) It is also understood that one skilled in the art may be able to make these compounds by similar methods or by combining other methods known to one skilled in the art. It is further understood that one skilled in the art would be able to make, in a similar manner as described below, other compounds of the invention not specifically illustrated below by using the appropriate starting components and modifying the parameters of the synthesis as needed. In general, starting components may be obtained from sources such as Sigma Aldrich, Lancaster Synthesis, Inc., Maybridge, Matrix Scientific, TCI, and Fluorochem USA, etc. or synthesized according to sources known to those skilled in the art (see, for example, Advanced Organic Chemistry: Reactions, Mechanisms, and Structure, 5th edition (Wiley, December 2000)) or prepared as described in this invention.








(395) TABLE-US-00001 TABLE 1 Compound Number I- Structure 1  embedded image 2
 embedded image 3 00  embedded image 4 01  embedded image 5 02  embedded image 6 03
 embedded image 7 04  embedded image 8 05  embedded image 9 06  embedded image 10 07
 embedded image 11 08  embedded image 12 09  embedded image 13 0  embedded image 14
 embedded image 15  embedded image 16  embedded image 17  embedded image 18
 embedded image 19  embedded image 20  embedded image 21  embedded image 22
 embedded image 23 0  embedded image 24  embedded image 25  embedded image 26
 embedded image 27  embedded image 28  embedded image 29  embedded image 30
 embedded image 31  embedded image 32  embedded image 33 0  embedded image 34
 embedded image 35  embedded image 36  embedded image 37  embedded image 38
 embedded image 39  embedded image 40  embedded image 41  embedded image 42
 embedded image 43 0  embedded image 44  embedded image 45  embedded image 46
 embedded image 47  embedded image 48  embedded image 49  embedded image 50
 embedded image 51  embedded image 52  embedded image 53 0  embedded image 54
 embedded image 55  embedded image 56  embedded image 57  embedded image 58
 embedded image 59  embedded image 60  embedded image 61  embedded image 62
 embedded image 63 0  embedded image 64  embedded image 65  embedded image 66
 embedded image 67  embedded image 68  embedded image 69  embedded image 70
 embedded image 71  embedded image 72  embedded image 73 0  embedded image 74
 embedded image 75  embedded image 76  embedded image 77  embedded image 78
 embedded image 79  embedded image 80  embedded image 81  embedded image 82
 embedded image 83 0  embedded image 84  embedded image 85  embedded image 86
 embedded image 87  embedded image 88  embedded image 89  embedded image 90
 embedded image 91  embedded image 92  embedded image 93 0  embedded image 94
 embedded image 95  embedded image 96  embedded image 97  embedded image 98
 embedded image 99  embedded image 100  embedded image 101  embedded image 102
 embedded image 103 00  embedded image 104 01  embedded image 105 02  embedded image
106 03  embedded image 107 04  embedded image 108 05  embedded image 109 06
 embedded image 110 07  embedded image 111 08  embedded image 112 09  embedded image
113 0  embedded image 114  embedded image 115  embedded image 116  embedded image
117  embedded image 118  embedded image 119  embedded image 120  embedded image
121  embedded image 122  embedded image 123 0  embedded image 124  embedded image
125  embedded image 126  embedded image 127  embedded image 128  embedded image
129  embedded image 130  embedded image 131  embedded image 132  embedded image
133 0  embedded image 134  embedded image 135  embedded image 136  embedded image
137  embedded image 138  embedded image 139  embedded image 140  embedded image
141  embedded image 142  embedded image 143 0  embedded image 144  embedded image
145  embedded image 146  embedded image 147  embedded image 148  embedded image


































149  150  151  152  153 0  154  155  156  157  158  159  160  161  162  163 0  164  165  166  167  168  169  170  171  172  173 0  174  175  176  177  178  179  180  181  182  183 0  184  185  186  187  188  189  190  191  192  193 0  194  195  196  197  198  199  200  201  202  203 00  204 01  205 02  206 03  207 04  208 05  209 06  210 07  211 08  212 09  213 0  214  215  216  217  218  219  220  221  222  223 0  224  225  226  227  228  229  230  231  232  233 0  234  235  236  237  238  239  240  241  242  243 0  244  245  246  247  248  249  250 

(396) TABLE-US-00002 TABLE 2 Compound Number (II-) Structure 1  2  3  4  5 0  6  7  8 


(397) TABLE-US-00003 TABLE 3 Compound Number (III-) Structure 1  2  3  4  5  6  7 0  8  9  10  11  12  13  14  15  16  17 0  18  19  20  21  22 

















(398) TABLE-US-00004 TABLE 4 Compound Number (IV-) Structure 1  2  3  4  5 0  6  7  8  9  10  11 

(399) TABLE-US-00005 TABLE 5a Representative Compounds No. Structure Name V-1  1-(3-(4-(2-(4,5- dichloro-2- hydroxyphenyl- amino)acetyl) piperazin-1-yl)azetidin-1- yl)prop-2-en-1- one V-2  1-(3-(4-(2-(4,5- dichloro-2- hydroxyphenyl- amino)acetyl) piperazin-1- yl)pyrrolidin-1- yl)prop-2-en-1- one V-3  N-(1'-(2-(4,5- dichloro-2- hydroxyphenyl- amino)acetyl)- 1,3'-biazetidin-3-yl)acrylamide V-4 0  1-(4-(1-(2-(4,5- dichloro-2- hydroxyphenyl- amino)acetyl) pyrrolidin-3- yl)piperazin-1- yl)prop-2-en-1- one V-5  1-(3-(4-(2-(2,4- dichloro-5- methoxyphenyl- amino)acetyl) piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-6  1-(3-(4-(3-(4,5- dichloro-2- hydroxyphenyl) propanoyl) piperazin-1-yl) azetidin-1- yl)prop-2-en-1- one V-7  1-(3-(4-(2-(4- chloro-2- hydroxy-5-

methylphenyl- amino)acetyl) piperazin-1- yl)pyrrolidin-1- yl)prop-2-en-1- one V-8
 embedded image 1-(3-(4-(2-(5- chloro-2- hydroxy-4- methylphenyl- amino)acetyl) piperazin-1- yl)pyrrolidin- 1-yl)prop-2-en-1- one V-9  embedded image 5-(2-(4-(1- acryloylazetidin- 3- yl)piperazin-1- yl)-2- oxoethylamino)- 2,4- dichloro- benzonitrile V-10  embedded image 2-(2-(4- (1- acryloylazetidin- 3-yl)piperazin-1- yl)-2- oxoethylamino)- 4,5- dichlorobenzamide V-11
 embedded image 1-(3-(4-(2-(4- chloro-2- hydroxy-5- iodophenylamino) acetyl)piperazin-1- yl)pyrrolidin-1- yl)prop-2-en-1- one V-12  embedded image 1-(3-(4-(2-(4- chloro-5-ethyl-2- hydroxyphenyl- amino)acetyl) piperazin-1- yl)pyrrolidin-1- yl)prop-2-en-1- one V-13
 embedded image 1-(3-(4-(2',5',6- trichloro-4- methoxybiphenyl- carbonyl) piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-14 00  embedded image 1-(3-(4-(2-(5- chloro-4-fluoro-2- hydroxyphenyl- amino)acetyl) piperazin-1-yl)azetidin- 1-yl)prop-2-en-1- one V-15 01
 embedded image 1-(3-(4-(2-(4,5- dichloro-2- hydroxyphenyl- amino)ethyl) piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-16 02  embedded image 4-(2-(4-(1- acryloylazetidin- 3- yl)piperazin-1- yl)-2-oxoethyl)- 3,4- dihydroquinoxalin- 2(1H)-one V-17 03  embedded image 1- (1- acryloylazetidin- 3-yl)-N-(4,5- dichloro-2- hydroxybenzyl) piperidine-4- carboxamide V-18 04
 embedded image 1-(3-(4-(2-(5- bromo-4-chloro- 2- hydroxyphenyl- amino)acetyl) piperazin-1- yl)azetidin- 1-yl)prop-2-en-1- one V-19 05  embedded image 1-(3-(4-(2-(5- chloro-2- hydroxyphenyl- amino)acetyl) piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-20 06
 embedded image (E)-1-(3-(4-((4,5- dichloro-2- hydroxyphenyl) glycyloxy)piperazin-1- yl)azetidin- 1-yl)- 4- (dimethylamino) but-2-en-1-one V-21 07  embedded image 1-(3-(4-(1- acryloylazetidin- 3-yl)piperazin-1- yl)-3- oxopropyl)-5,6- dichloro-1H- benzo[d]imidazol- 2(3H)-one V-22 08
 embedded image 1-(3-(4-(2-(2,4,5- trichlorophenyl- amino)acetyl) piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-23 09  embedded image 1-(3-(4-(2-(2,4- dichloro-5- hydroxyphenyl- amino)acetyl) piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-24 0  embedded image 1-(3-(4-(2- (naphthalen-1- yl)acetyl)piperazin- 1-yl)azetidin-1- yl)prop-2-en-1- one V-25  embedded image 1- (3-(4-(2-(1H- indol-3- yl)acetyl)piperazin- 1-yl)azetidin- 1-yl)prop-2-en- 1-one V-26
 embedded image 1-(3-(4-(2-(4,5- dichloro-2- (trifluoromethyl) phenylamino) acetyl)piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-27  embedded image 1-(3-(4-(2-(3,4- dichloro-5- hydroxyphenyl- amino)acetyl) piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-28
 embedded image 1-(3-(4-(2-(4- bromo-5-chloro- 2- hydroxyphenyl- amino)acetyl) piperazin-1- yl)azetidin- 1-yl)prop-2-en-1- one V-29  embedded image 1-(3-(4-(2-(1H- indol-1- yl)acetyl) piperazin-1- yl)azetidin- 1-yl)prop-2-en- 1-one V-30  embedded image 1-(3-(4-(2-(5,6- dichloro- 1H- indol-1- yl)acetyl) piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-31  embedded image 1- (3-(4-(2-(4,5- dichloro-2- hydroxyphenyl- amino)propanoyl) piperazin-1- yl)azetidin-1- yl)prop-2- en-1- one V-32  embedded image 1-(3-(4-(2-(4- chloro-2- hydroxy-5- methylphenyl- amino)acetyl) piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-33  embedded image 1-(3-(4-(2- (3- chloro-5- hydroxyphenyl- amino)acetyl) piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-34 0
 embedded image 1-(3-(4-(2-(2- hydroxy-5- (methylsulfonyl) phenylamino) acetyl)piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-35  embedded image 1-(3-(4-(2-(4- chloro-5- cyclopropyl-2- hydroxyphenyl- amino)acetyl) piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-36
 embedded image N-(1'-(2-(5- bromo-4-chloro- 2- hydroxyphenyl- amino)acetyl)-1,3'- biazetidin- 3- yl)acrylamide V-37  embedded image 1-(3-(4-(2-(4- chloro-2- methoxy-5- (trifluoromethyl) phenylamino) acetyl)piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-38  embedded image 1-(3- (4-(2-(5- chlorothiazol-2- ylamino)acetyl) piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one V-39
 embedded image 1-(4-(1-(2-(4,5- dichloro-2- hydroxyphenyl- amino)acetyl) azetidin-3- yl)piperazin-1- yl)prop-2-en-1- one V-40  embedded image 2-(4,5-dichloro-2- hydroxyphenyl- amino)-1-(3-(4- (vinylsulfonyl) piperazin-1- yl)azetidin-1- yl)ethanone V-41  embedded image 2- (4,5-dichloro- 2- hydroxyphenyl- amino)-1-(4-(1- (vinylsulfonyl) pyrrolidin-3- yl)piperazin-1- yl)ethanone V-42  embedded image 4-(2-(4-(1- acryloylazetidin- 3-yl)piperazin-1- yl)-2- oxoethyl)- 3,4- dihydroquinoxalin- 2(1H)-one V-43  embedded image 1-(3-(4-(2-(3-

hydroxynaphthalen-2-ylamino)acetyl) piperazin-1-yl)prop-2-en-1-one V-44 0
 embedded image 5-(2-(4-(1-acryloylazetidin-3-yl)piperazin-1-yl)-2-oxoethylamino)-2,3-dichlorobenzamide V-45  embedded image N-(1'-(2-(4-chloro-5-cyclopropyl-2-hydroxyphenyl-amino)acetyl)-1,3'-biazetidin-3-yl)acrylamide V-46  embedded image 5-(2-(4-(1-acryloylazetidin-3-yl)piperazin-1-yl)-2-oxoethylamino)-2-chloro-4-methoxy-benzaldehyde V-47  embedded image 1-(3-(4-(4-chloro-5-cyclopropyl-2-methoxybenzoyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-48  embedded image 1-(3-(4-(2',5',6-trichloro-4-hydroxybiphenyl-carbonyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-49
 embedded image 1-(4-(1-acryloylazetidin-3-yl)piperazin-1-yl)-2-(4,5-dichloro-2-hydroxyphenyl-amino)butan-1-one V-50  embedded image 1-(3-(4-(2-(4-chloro-2-hydroxy-5-isopropylphenyl-amino)acetyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-51
 embedded image 1-(1-acryloylazetidin-3-yl)-4-(2-(4-chloro-5-cyclopropyl-2-hydroxyphenyl-amino)acetyl) piperazine-2-carboxamide V-52  embedded image 1-(3-(4-(2-(5-chloro-4-ethyl-2-hydroxyphenyl-amino)acetyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-53
 embedded image 1-(3-(4-(2-(4-chloro-5-cyclobutyl-2-hydroxyphenyl-amino)acetyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-54 0  embedded image 1-(3-(4-(2-(4-chloro-5-cyclopropyl-2-hydroxyphenyl-amino)acetyl)-2-(hydroxymethyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-55  embedded image 1-(3-(4-(2-(4-chloro-5-ethyl-2-hydroxyphenyl-amino)acetyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-56  embedded image 1-(3-(4-(2-(4-chloro-5-(2,2-difluoro-cyclopropyl)-2-hydroxyphenyl-amino)acetyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-57  embedded image 1-(3-(4-(2-(4-chloro-2-hydroxy-5-(2,2,2-trifluoroethyl) phenylamino) acetyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-58
 embedded image 1-(3-(4-(3-(4-chloro-5-cyclopropyl-2-hydroxyphenyl) propanoyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-59  embedded image 1-(3-(4-(2-(4-chloro-5-cyclobutyl-2-hydroxyphenyl-amino)propanoyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-60
 embedded image 1-(3-(4-(2-(4-chloro-5-cyclopropyl-2-hydroxyphenyl-amino)acetyl)-2-methylpiperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-61  embedded image 1-(3-(4-(2-(5,6-dichloro-1H-indol-3-yl)acetyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-62
 embedded image (E)-1-(4-(1-acryloylazetidin-3-yl)piperazin-1-yl)-3-(4-chloro-5-cyclopropyl-2-hydroxyphenyl) prop-2-en-1-one V-63  embedded image (S)-1-(3-(4-(2-(4-chloro-5-cyclopropyl-2-hydroxyphenyl-amino)acetyl)-2-(hydroxymethyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-64 0  embedded image 1-(3-(4-(2-(4-chloro-2-hydroxy-5-(1-methylcyclopropyl) phenylamino) acetyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-65
 embedded image 1-(3-(4-(2-(4-chloro-5-cyclopropyl-2-methoxyphenyl-thio)acetyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-66  embedded image 4-(1-acryloylazetidin-3-yl)-N-(5-bromo-4-chloro-2-hydroxybenzyl) piperazine-1-carboxamide V-67  embedded image (S)-1-(3-(4-(2-(4-chloro-5-cyclopropyl-2-hydroxyphenyl-amino)propanoyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-68
 embedded image (R)-1-(3-(4-(2-(4-chloro-5-cyclopropyl-2-hydroxyphenyl-amino)propanoyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-69
 embedded image (S)-1-(3-(4-(2-(4-chloro-5-ethyl-2-hydroxyphenyl-amino)propanoyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-70  embedded image (R)-1-(3-(4-(2-(4-chloro-5-ethyl-2-hydroxyphenyl-amino)propanoyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-71
 embedded image 2-(2-(4-(1-acryloylazetidin-3-yl)piperazin-1-yl)-2-oxoethylamino)-5-chloro-4-cyclopropyl-benzonitrile V-72  embedded image 1-(3-(4-(3-(4-chloro-5-ethyl-2-hydroxyphenyl)-1H-pyrazol-5-yl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-73
 embedded image 1-(3-(4-(1-(4-chloro-5-cyclopropyl-2-methoxyphenyl) pyrrolidine-2-carbonyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-74 0  embedded image 1-(3-(4-(2-(5-chloro-4-ethylpyridin-2-ylamino)acetyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-75
 embedded image 1-(3-(4-(2-(4,5-dichloro-7-methoxy-1H-indol-1-yl)acetyl) piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one V-76  embedded image 1-(3-(4-(1-(4-chloro-5-ethyl-2-

methoxyphenyl) piperidin-3- yl)piperazin-1- yl)azetidin-1- yl)prop-2-en-1- one
(400) TABLE-US-00006 TABLE 5b Experimental Mass Spectral Data for Compounds in Table 5
[M + [M + [M + [M + No. H.sup.+] No. H.sup.+] No. H.sup.+] No. H.sup.+] V-1 411.30 V-2
425.25 V-3 399.20 V-4 427.30 V-5 449.25* V-6 410.25 V-7 429.35* V-8 407.35 V-9 422.25 V-10
462.25* V-11 519.25 V-12 443.30* V-13 532.25* V-14 395.30 V-15 399.25 V-16 419.25 V-17
434.25* V-18 457.35 V-19 379.30 V-20 470.35 V-21 450.35 V-22 433.05 V-23 435.25* V-24
386.25* V-25 351.35 V-26 487.30* V-27 413.30 V-28 479.20* V-29 353.30 V-30 421.30 V-31
449.25* V-32 393.30 V-33 377.30.sup.+ V-34 423.35 V-35 441.30* V-36 445.20 V-37 461.30 V-38
368.30.sup.+ V-39 411.20 V-40 447.25.sup.+ V-41 463.20 V-42 382.40.sup.+ V-43 417.35* V-44
440.30 V-45 405.35 V-46 421.30 V-47 404.35 V-48 494.30 V-49 441.30 V-50 421.35 V-51 462.45
V-52 407.40 V-53 433.40 V-54 449.35 V-55 407.30 V-56 455.20 V-57 461.40 V-58 418.40 V-59
447.40 V-60 433.45 V-61 421.25 V-62 416.35 V-63 449.40 V-64 433.35 V-65 451.30 V-66 459.25
V-67 433.20 V-68 433.40 V-69 421.35 V-70 421.35 V-71 428.35 V-72 416.35 V-73 473.90 V-74
392.30 V-75 451.30 V-76 447.85 *[M + Na].sup.+ .sup.+[M - H].sup.-
(401) TABLE-US-00007 TABLE 6 Representative Compounds No. Structure Name [M + H].sup.+
VI-1  embedded image 1-(4-(2',6-dichloro-4- methoxybiphenyl- carbonyl)piperazin-1- yl)prop-
2-en-1-one 441.20* VI-2  embedded image 1-(4-(4-chloro-5-iodo- 2- methoxybenzoyl)
piperazin-1-yl)prop- 2-en-1-one 457.05* VI-3  embedded image 1-(4-(2',6-dichloro-4-
hydroxybiphenyl- carbonyl)piperazin-1- yl)prop-2-en-1-one 427.15 VI-4  embedded image 1-(4-
(2',6-dichloro-5'- fluoro-4- methoxybiphenyl- carbonyl)piperazin-1- yl)prop-2-en-1-one 459.15 VI-
5  embedded image 1-(4-(2',5',6-trichloro- 4- methoxybiphenyl- carbonyl)piperazin-1- yl)prop-
2-en-1-one 453.15 VI-6  embedded image 1-(2-(hydroxymethyl)- 4-(2',5',6-trichloro-4-
methoxybiphenyl- carbonyl)piperazin-1- yl)prop-2-en-1-one 483.20.sup.# VI-7
 embedded image 1-(3-(hydroxymethyl)- 4-(2',5',6-trichloro-4- methoxybiphenyl-
carbonyl)piperazin-1- yl)prop-2-en-1-one 485.20 VI-8  embedded image 1-(4-(4,5-dichloro-2-
hydroxybenzoyl) piperazin-1-yl)prop- 2-en-1-one 327.15 VI-9  embedded image 1-(4-(5-bromo-
4- chloro-2- hydroxybenzoyl) piperazin-1-yl)prop- 2-en-1-one 388* VI-10  embedded image
(E)-1-(4-(2',6- dichloro-4- methoxybiphenyl- carbonyl)piperazin- 1-yl)-4- (dimethylamino)but- 2-
en-1-one 476.23 VI-11  embedded image 1-(4-(3-(2- chlorophenyl)-1H- pyrazole-5-
carbonyl)piperazin-1- yl)prop-2-en-1-one 467.20* VI-12  embedded image 1-(4-(2',6-
dichlorobiphenyl- carbonyl)piperazin-1- yl)prop-2-en-1-one 389.20 VI-13  embedded image N-
(1-(2',6-dichloro-4- hydroxybiphenyl- carbonyl)azetidin-3- yl)acrylamide 389.30 VI-14
 embedded image N-(1-acryloylazetidin- 3-yl)-3-(2- chlorophenyl)-1H- pyrazole-5- carboxamide
331.15 VI-15  embedded image N-(1-(3-(2- chlorophenyl)-1H- pyrazole-5- carbonyl)azetidin-3-
yl)acrylamide 331.15 VI-16  embedded image 1-(4-(5-(thiophen-2- yl)-1H-pyrazole-3-
carbonyl)piperazin-1- yl)prop-2-en-1-one 339.15 VI-17  embedded image 1-(4-(2',5',6-trichloro-
4- hydroxybiphenyl- carbonyl)piperazin-1- yl)prop-2-en-1-one 438.30.sup.+ VI-18 0
 embedded image 1-(4-(2-(2- chlorophenyl)thiazole- 4-carbonyl)piperazin- 1-yl)prop-2-en-1-one
384.15.sup.+ VI-19  embedded image 1-(4-(4-(2- chlorophenyl)thiazole- 2-carbonyl)piperazin- 1-
yl)prop-2-en-1-one 384.15.sup.+ VI-20  embedded image 1-(4-(5-chloro-2-(2-
chlorophenyl)thiazole- 4-carbonyl)piperazin- 1-yl)prop-2-en-1-one 418.10* VI-21
 embedded image 1-(4-(4-(2- chlorophenyl)-1H- pyrrole-2- carbonyl)piperazin-1- yl)prop-2-en-1-
one 344.20 VI-22  embedded image 1-(4-(4-(2- chlorophenyl)-5- methylthiazole-2-
carbonyl)piperazin-1- yl)prop-2-en-1-one 376.25 VI-23  embedded image 1-(4-(4-(2-
chlorophenyl)-5- methyl-1H-pyrrole-2- carbonyl)piperazin-1- yl)prop-2-en-1-one 356.35.sup.+ VI-
24  embedded image (E)-1-(4-(2',6- dichloro-4- hydroxybiphenyl- carbonyl)piperazin- 1-yl)-4-
(dimethylamino)but- 2-en-1-one 462.35 VI-25  embedded image 1-(4-(2'-chloro-5-
hydroxybiphenyl- carbonyl)piperazin-1- yl)prop-2-en-1-one 371.25 VI-26  embedded image 1-(4-
(2'-chloro-4- hydroxybiphenyl- carbonyl)piperazin-1- yl)prop-2-en-1-one 371.25 VI-27
 embedded image 1-(4-(2',6-dichloro-5- hydroxybiphenyl- carbonyl)piperazin-1- yl)prop-2-en-1-

one 403.35.sup.+ VI-28 01  embedded image 5-(4- acryloylpiperazine-1- carbonyl)-1-(2,5- dichlorophenyl)-4- hydroxypyridin- 2(1H)-one 422.06 VI-29  embedded image 1-(4-(6-chloro-4- hydroxybiphenyl- carbonyl)piperazin-1- yl)prop-2-en-1-one 371.25 VI-30  embedded image 1-(4- (5-chloro-4-(2- chlorophenyl)-1H- pyrrole-2- carbonyl)piperazin-1- yl)prop-2-en-1-one 378.25.sup.+ VI-31  embedded image 1-(4-(4-(2,5- dichlorophenyl)-1H- pyrrole-2- carbonyl)piperazin-1- yl)prop-2-en-1-one 378.20 VI-32  embedded image 1-(4-(4-(2,4- dichlorophenyl)-1H- pyrrole-2- carbonyl)piperazin-1- yl)prop-2-en-1-one 377.90 VI-33  embedded image N-(1-(3,4- dichlorobenzoyl) piperidin-4- yl)ethenesulfonamide 363.02 VI-34  embedded image (3,4- dichlorophenyl)(4- (vinylsulfonyl) piperazin-1- yl)methanone 349.04 VI- 35  embedded image (S)-N-(1-(3,4- dichlorobenzoyl) piperidin-3- yl)ethenesulfonamide 363.06 VI-36  embedded image 1-(4-(3,4- dichlorobenzoyl) piperazin-1-yl)prop- 2-en-1-one 312.99 VI- 37  embedded image 1-acryloyl-4-(4',6- dichloro-4- hydroxybiphenyl- carbonyl) piperazine-2- carbonitrile 430.30 VI-38 00  embedded image 1-acryloyl-4-(2',5',6- trichloro-4- hydroxybiphenyl- carbonyl)piperazine-2- carbonitrile 464.30 VI-39 01  embedded image 1- acryloyl-4-(4,5- dichloro-2- hydroxybenzoyl) piperazine-2- carbonitrile 354.15 VI-40 02  embedded image 1-(4-(2-chloro-5- hydroxybiphenyl- carbonyl)piperazin-1- yl)prop-2-en-1-one 371.25 VI-41 03  embedded image 1-(4-(2,2'-dichloro-5- hydroxybiphenyl- carbonyl)piperazin-1- yl)prop-2-en-1-one 405.25 VI-42 04  embedded image 1-(4-(2,4'-dichloro-5- hydroxybiphenyl- carbonyl)piperazin-1- yl)prop-2-en-1-one 405.20 VI-42 05  embedded image 1-(4-(2,3'-dichloro- 5- hydroxybiphenyl- carbonyl)piperazin-1- yl)prop-2-en-1-one 405.25 *[M + Na].sup.+ .sup.+[M - H].sup.- .sup.#[M]

Pharmaceutical Compositions

(402) Other embodiments are directed to pharmaceutical compositions. The pharmaceutical composition comprises any one (or more) of the foregoing compounds and a pharmaceutically acceptable carrier. In some embodiments, the pharmaceutical composition is formulated for oral administration. In other embodiments, the pharmaceutical composition is formulated for injection. In still more embodiments, the pharmaceutical compositions comprise a compound as disclosed herein and an additional therapeutic agent (e.g., anticancer agent). Non-limiting examples of such therapeutic agents are described herein below.

(403) Suitable routes of administration include, but are not limited to, oral, intravenous, rectal, aerosol, parenteral, ophthalmic, pulmonary, transmucosal, transdermal, vaginal, otic, nasal, and topical administration. In addition, by way of example only, parenteral delivery includes intramuscular, subcutaneous, intravenous, intramedullary injections, as well as intrathecal, direct intraventricular, intraperitoneal, intralymphatic, and intranasal injections.

(404) In certain embodiments, a compound as described herein is administered in a local rather than systemic manner, for example, via injection of the compound directly into an organ, often in a depot preparation or sustained release formulation. In specific embodiments, long acting formulations are administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Furthermore, in other embodiments, the drug is delivered in a targeted drug delivery system, for example, in a liposome coated with organ-specific antibody. In such embodiments, the liposomes are targeted to and taken up selectively by the organ. In yet other embodiments, the compound as described herein is provided in the form of a rapid release formulation, in the form of an extended release formulation, or in the form of an intermediate release formulation. In yet other embodiments, the compound described herein is administered topically.

(405) The compounds according to the invention are effective over a wide dosage range. For example, in the treatment of adult humans, dosages from 0.01 to 1000 mg, from 0.5 to 100 mg, from 1 to 50 mg per day, and from 5 to 40 mg per day are examples of dosages that are used in some embodiments. An exemplary dosage is 10 to 30 mg per day. The exact dosage will depend upon the route of administration, the form in which the compound is administered, the subject to be

treated, the body weight of the subject to be treated, and the preference and experience of the attending physician.

(406) In some embodiments, a compound of the invention is administered in a single dose.

Typically, such administration will be by injection, e.g., intravenous injection, in order to introduce the agent quickly. However, other routes are used as appropriate. In some embodiments, a single dose of a compound of the invention is used for treatment of an acute condition.

(407) In some embodiments, a compound of the invention is administered in multiple doses. In some embodiments, dosing is about once, twice, three times, four times, five times, six times, or more than six times per day. In other embodiments, dosing is about once a month, once every two weeks, once a week, or once every other day. In another embodiment a compound of the invention and another agent are administered together about once per day to about 6 times per day. In another embodiment the administration of a compound of the invention and an agent continues for less than about 7 days. In yet another embodiment the administration continues for more than about 6, 10, 14, 28 days, two months, six months, or one year. In some cases, continuous dosing is achieved and maintained as long as necessary.

(408) Administration of the compounds of the invention may continue as long as necessary. In some embodiments, a compound of the invention is administered for more than 1, 2, 3, 4, 5, 6, 7, 14, or 28 days. In some embodiments, a compound of the invention is administered for less than 28, 14, 7, 6, 5, 4, 3, 2, or 1 day. In some embodiments, a compound of the invention is administered chronically on an ongoing basis, e.g., for the treatment of chronic effects.

(409) In some embodiments, the compounds of the invention are administered in dosages. It is known in the art that due to intersubject variability in compound pharmacokinetics, individualization of dosing regimen is necessary for optimal therapy. Dosing for a compound of the invention may be found by routine experimentation in light of the instant disclosure.

(410) In some embodiments, the compounds described herein are formulated into pharmaceutical compositions. In specific embodiments, pharmaceutical compositions are formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. Any pharmaceutically acceptable techniques, carriers, and excipients are used as suitable to formulate the pharmaceutical compositions described herein: Remington: The Science and Practice of Pharmacy, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pa. 1975; Liberman, H. A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and Pharmaceutical Dosage Forms and Drug Delivery Systems, Seventh Ed. (Lippincott Williams & Wilkins 1999).

(411) Provided herein are pharmaceutical compositions comprising a compound of structure (I) and a pharmaceutically acceptable diluent(s), excipient(s), or carrier(s). In certain embodiments, the compounds described are administered as pharmaceutical compositions in which compounds of structure (I) are mixed with other active ingredients, as in combination therapy. Encompassed herein are all combinations of actives set forth in the combination therapies section below and throughout this disclosure. In specific embodiments, the pharmaceutical compositions include one or more compounds of structure (I).

(412) A pharmaceutical composition, as used herein, refers to a mixture of a compound of structure (I) with other chemical components, such as carriers, stabilizers, diluents, dispersing agents, suspending agents, thickening agents, and/or excipients. In certain embodiments, the pharmaceutical composition facilitates administration of the compound to an organism. In some embodiments, practicing the methods of treatment or use provided herein, therapeutically effective amounts of compounds of structure (I) provided herein are administered in a pharmaceutical composition to a mammal having a disease, disorder or medical condition to be treated. In specific

embodiments, the mammal is a human. In certain embodiments, therapeutically effective amounts vary depending on the severity of the disease, the age and relative health of the subject, the potency of the compound used and other factors. The compounds described herein are used singly or in combination with one or more therapeutic agents as components of mixtures.

(413) In one embodiment, one or more compounds of structure (I) is formulated in an aqueous solutions. In specific embodiments, the aqueous solution is selected from, by way of example only, a physiologically compatible buffer, such as Hank's solution, Ringer's solution, or physiological saline buffer. In other embodiments, one or more compound of structure (I) is/are formulated for transmucosal administration. In specific embodiments, transmucosal formulations include penetrants that are appropriate to the barrier to be permeated. In still other embodiments wherein the compounds described herein are formulated for other parenteral injections, appropriate formulations include aqueous or nonaqueous solutions. In specific embodiments, such solutions include physiologically compatible buffers and/or excipients.

(414) In another embodiment, compounds described herein are formulated for oral administration. Compounds described herein are formulated by combining the active compounds with, e.g., pharmaceutically acceptable carriers or excipients. In various embodiments, the compounds described herein are formulated in oral dosage forms that include, by way of example only, tablets, powders, pills, dragees, capsules, liquids, gels, syrups, elixirs, slurries, suspensions and the like.

(415) In certain embodiments, pharmaceutical preparations for oral use are obtained by mixing one or more solid excipient with one or more of the compounds described herein, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as: for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methylcellulose, microcrystalline cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose; or others such as: polyvinylpyrrolidone (PVP or povidone) or calcium phosphate. In specific embodiments, disintegrating agents are optionally added. Disintegrating agents include, by way of example only, cross-linked croscarmellose sodium, polyvinylpyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

(416) In one embodiment, dosage forms, such as dragee cores and tablets, are provided with one or more suitable coating. In specific embodiments, concentrated sugar solutions are used for coating the dosage form. The sugar solutions, optionally contain additional components, such as by way of example only, gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs and/or pigments are also optionally added to the coatings for identification purposes. Additionally, the dyestuffs and/or pigments are optionally utilized to characterize different combinations of active compound doses.

(417) In certain embodiments, therapeutically effective amounts of at least one of the compounds described herein are formulated into other oral dosage forms. Oral dosage forms include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. In specific embodiments, push-fit capsules contain the active ingredients in admixture with one or more filler. Fillers include, by way of example only, lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In other embodiments, soft capsules, contain one or more active compound that is dissolved or suspended in a suitable liquid. Suitable liquids include, by way of example only, one or more fatty oil, liquid paraffin, or liquid polyethylene glycol. In addition, stabilizers are optionally added.

(418) In other embodiments, therapeutically effective amounts of at least one of the compounds described herein are formulated for buccal or sublingual administration. Formulations suitable for buccal or sublingual administration include, by way of example only, tablets, lozenges, or gels. In still other embodiments, the compounds described herein are formulated for parental injection,

including formulations suitable for bolus injection or continuous infusion. In specific embodiments, formulations for injection are presented in unit dosage form (e.g., in ampoules) or in multi-dose containers. Preservatives are, optionally, added to the injection formulations. In still other embodiments, the pharmaceutical compositions are formulated in a form suitable for parenteral injection as sterile suspensions, solutions or emulsions in oily or aqueous vehicles. Parenteral injection formulations optionally contain formulatory agents such as suspending, stabilizing and/or dispersing agents. In specific embodiments, pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. In additional embodiments, suspensions of the active compounds (e.g., compounds of structure (I)) are prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles for use in the pharmaceutical compositions described herein include, by way of example only, fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. In certain specific embodiments, aqueous injection suspensions contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension contains suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. Alternatively, in other embodiments, the active ingredient is in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

(419) In still other embodiments, the compounds of structure (I) are administered topically. The compounds described herein are formulated into a variety of topically administrable compositions, such as solutions, suspensions, lotions, gels, pastes, medicated sticks, balms, creams or ointments. Such pharmaceutical compositions optionally contain solubilizers, stabilizers, tonicity enhancing agents, buffers and preservatives.

(420) In yet other embodiments, the compounds of structure (I) are formulated for transdermal administration. In specific embodiments, transdermal formulations employ transdermal delivery devices and transdermal delivery patches and can be lipophilic emulsions or buffered, aqueous solutions, dissolved and/or dispersed in a polymer or an adhesive. In various embodiments, such patches are constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents. In additional embodiments, the transdermal delivery of the compounds of structure (I) is accomplished by means of iontophoretic patches and the like. In certain embodiments, transdermal patches provide controlled delivery of the compounds of structure (I). In specific embodiments, the rate of absorption is slowed by using rate-controlling membranes or by trapping the compound within a polymer matrix or gel. In alternative embodiments, absorption enhancers are used to increase absorption. Absorption enhancers or carriers include absorbable pharmaceutically acceptable solvents that assist passage through the skin. For example, in one embodiment, transdermal devices are in the form of a bandage comprising a backing member, a reservoir containing the compound optionally with carriers, optionally a rate controlling barrier to deliver the compound to the skin of the host at a controlled and predetermined rate over a prolonged period of time, and means to secure the device to the skin.

(421) In other embodiments, the compounds of structure (I) are formulated for administration by inhalation. Various forms suitable for administration by inhalation include, but are not limited to, aerosols, mists or powders. Pharmaceutical compositions of any of compound of structure (I) are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant (e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas). In specific embodiments, the dosage unit of a pressurized aerosol is determined by providing a valve to deliver a metered amount. In certain embodiments, capsules and cartridges of, such as, by way of example only, gelatin for use in an inhaler or insufflator are formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

(422) In still other embodiments, the compounds of structure (I) are formulated in rectal

compositions such as enemas, rectal gels, rectal foams, rectal aerosols, suppositories, jelly suppositories, or retention enemas, containing conventional suppository bases such as cocoa butter or other glycerides, as well as synthetic polymers such as polyvinylpyrrolidone, PEG, and the like. In suppository forms of the compositions, a low-melting wax such as, but not limited to, a mixture of fatty acid glycerides, optionally in combination with cocoa butter is first melted.

(423) In certain embodiments, pharmaceutical compositions are formulated in any conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. Any pharmaceutically acceptable techniques, carriers, and excipients are optionally used as suitable. Pharmaceutical compositions comprising a compound of structure (I) are manufactured in a conventional manner, such as, by way of example only, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or compression processes.

(424) Pharmaceutical compositions include at least one pharmaceutically acceptable carrier, diluent or excipient and at least one compound of structure (I), described herein as an active ingredient. The active ingredient is in free-acid or free-base form, or in a pharmaceutically acceptable salt form. In addition, the methods and pharmaceutical compositions described herein include the use of N-oxides, crystalline forms (also known as polymorphs), as well as active metabolites of these compounds having the same type of activity. All tautomers of the compounds described herein are included within the scope of the compounds presented herein. Additionally, the compounds described herein encompass unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. The solvated forms of the compounds presented herein are also considered to be disclosed herein. In addition, the pharmaceutical compositions optionally include other medicinal or pharmaceutical agents, carriers, adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure, buffers, and/or other therapeutically valuable substances.

(425) Methods for the preparation of compositions comprising the compounds described herein include formulating the compounds with one or more inert, pharmaceutically acceptable excipients or carriers to form a solid, semi-solid or liquid. Solid compositions include, but are not limited to, powders, tablets, dispersible granules, capsules, cachets, and suppositories. Liquid compositions include solutions in which a compound is dissolved, emulsions comprising a compound, or a solution containing liposomes, micelles, or nanoparticles comprising a compound as disclosed herein. Semi-solid compositions include, but are not limited to, gels, suspensions and creams. The form of the pharmaceutical compositions described herein include liquid solutions or suspensions, solid forms suitable for solution or suspension in a liquid prior to use, or as emulsions. These compositions also optionally contain minor amounts of nontoxic, auxiliary substances, such as wetting or emulsifying agents, pH buffering agents, and so forth.

(426) In some embodiments, pharmaceutical composition comprising at least one compound of structure (I) illustratively takes the form of a liquid where the agents are present in solution, in suspension or both. Typically when the composition is administered as a solution or suspension a first portion of the agent is present in solution and a second portion of the agent is present in particulate form, in suspension in a liquid matrix. In some embodiments, a liquid composition includes a gel formulation. In other embodiments, the liquid composition is aqueous.

(427) In certain embodiments, useful aqueous suspensions contain one or more polymers as suspending agents. Useful polymers include water-soluble polymers such as cellulosic polymers, e.g., hydroxypropyl methylcellulose, and water-insoluble polymers such as cross-linked carboxyl-containing polymers. Certain pharmaceutical compositions described herein comprise a mucoadhesive polymer, selected for example from carboxymethylcellulose, carbomer (acrylic acid polymer), poly(methylmethacrylate), polyacrylamide, polycarbophil, acrylic acid/butyl acrylate

copolymer, sodium alginate and dextran.

(428) Useful pharmaceutical compositions also, optionally, include solubilizing agents to aid in the solubility of a compound of structure (I). The term “solubilizing agent” generally includes agents that result in formation of a micellar solution or a true solution of the agent. Certain acceptable nonionic surfactants, for example polysorbate 80, are useful as solubilizing agents, as can ophthalmically acceptable glycols, polyglycols, e.g., polyethylene glycol 400, and glycol ethers.

(429) Furthermore, useful pharmaceutical compositions optionally include one or more pH adjusting agents or buffering agents, including acids such as acetic, boric, citric, lactic, phosphoric and hydrochloric acids; bases such as sodium hydroxide, sodium phosphate, sodium borate, sodium citrate, sodium acetate, sodium lactate and tris-hydroxymethylaminomethane; and buffers such as citrate/dextrose, sodium bicarbonate and ammonium chloride. Such acids, bases and buffers are included in an amount required to maintain pH of the composition in an acceptable range.

(430) Additionally, useful compositions also, optionally, include one or more salts in an amount required to bring osmolality of the composition into an acceptable range. Such salts include those having sodium, potassium or ammonium cations and chloride, citrate, ascorbate, borate, phosphate, bicarbonate, sulfate, thiosulfate or bisulfite anions; suitable salts include sodium chloride, potassium chloride, sodium thiosulfate, sodium bisulfite and ammonium sulfate.

(431) Other useful pharmaceutical compositions optionally include one or more preservatives to inhibit microbial activity. Suitable preservatives include mercury-containing substances such as merfen and thiomersal; stabilized chlorine dioxide; and quaternary ammonium compounds such as benzalkonium chloride, cetyltrimethylammonium bromide and cetylpyridinium chloride.

(432) Still other useful compositions include one or more surfactants to enhance physical stability or for other purposes. Suitable nonionic surfactants include polyoxyethylene fatty acid glycerides and vegetable oils, e.g., polyoxyethylene (60) hydrogenated castor oil; and polyoxyethylene alkylethers and alkylphenyl ethers, e.g., octoxynol 10, octoxynol 40.

(433) Still other useful compositions include one or more antioxidants to enhance chemical stability where required. Suitable antioxidants include, by way of example only, ascorbic acid and sodium metabisulfite.

(434) In certain embodiments, aqueous suspension compositions are packaged in single-dose non-reclosable containers. Alternatively, multiple-dose reclosable containers are used, in which case it is typical to include a preservative in the composition.

(435) In alternative embodiments, other delivery systems for hydrophobic pharmaceutical compounds are employed. Liposomes and emulsions are examples of delivery vehicles or carriers useful herein. In certain embodiments, organic solvents such as N-methylpyrrolidone are also employed. In additional embodiments, the compounds described herein are delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various sustained-release materials are useful herein. In some embodiments, sustained-release capsules release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization are employed.

(436) In certain embodiments, the formulations described herein comprise one or more antioxidants, metal chelating agents, thiol containing compounds and/or other general stabilizing agents. Examples of such stabilizing agents, include, but are not limited to: (a) about 0.5% to about 2% w/v glycerol, (b) about 0.1% to about 1% w/v methionine, (c) about 0.1% to about 2% w/v monothioglycerol, (d) about 1 mM to about 10 mM EDTA, (e) about 0.01% to about 2% w/v ascorbic acid, (f) 0.003% to about 0.02% w/v polysorbate 80, (g) 0.001% to about 0.05% w/v. polysorbate 20, (h) arginine, (i) heparin, (j) dextran sulfate, (k) cyclodextrins, (l) pentosan polysulfate and other heparinoids, (m) divalent cations such as magnesium and zinc; or (n) combinations thereof.

(437) In some embodiments, the concentration of one or more compounds provided in the

pharmaceutical compositions of the present invention is less than 100%, 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.4%, 0.3%, 0.2%, 0.1%, 0.09%, 0.08%, 0.07%, 0.06%, 0.05%, 0.04%, 0.03%, 0.02%, 0.01%, 0.009%, 0.008%, 0.007%, 0.006%, 0.005%, 0.004%, 0.003%, 0.002%, 0.001%, 0.0009%, 0.0008%, 0.0007%, 0.0006%, 0.0005%, 0.0004%, 0.0003%, 0.0002%, or 0.0001% w/w, w/v or v/v.

(438) In some embodiments, the concentration of one or more compounds of the invention is greater than 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, 19.75%, 19.50%, 19.25% 19%, 18.75%, 18.50%, 18.25% 18%, 17.75%, 17.50%, 17.25% 17%, 16.75%, 16.50%, 16.25% 16%, 15.75%, 15.50%, 15.25% 15%, 14.75%, 14.50%, 14.25% 14%, 13.75%, 13.50%, 13.25% 13%, 12.75%, 12.50%, 12.25% 12%, 11.75%, 11.50%, 11.25% 11%, 10.75%, 10.50%, 10.25% 10%, 9.75%, 9.50%, 9.25% 9%, 8.75%, 8.50%, 8.25% 8%, 7.75%, 7.50%, 7.25% 7%, 6.75%, 6.50%, 6.25% 6%, 5.75%, 5.50%, 5.25% 5%, 4.75%, 4.50%, 4.25%, 4%, 3.75%, 3.50%, 3.25%, 3%, 2.75%, 2.50%, 2.25%, 2%, 1.75%, 1.50%, 1.25%, 1%, 0.5%, 0.4%, 0.3%, 0.2%, 0.1%, 0.09%, 0.08%, 0.07%, 0.06%, 0.05%, 0.04%, 0.03%, 0.02%, 0.01%, 0.009%, 0.008%, 0.007%, 0.006%, 0.005%, 0.004%, 0.003%, 0.002%, 0.001%, 0.0009%, 0.0008%, 0.0007%, 0.0006%, 0.0005%, 0.0004%, 0.0003%, 0.0002%, or 0.0001% w/w, w/v, or v/v.

(439) In some embodiments, the concentration of one or more compounds of the invention is in the range from approximately 0.0001% to approximately 50%, approximately 0.001% to approximately 40%, approximately 0.01% to approximately 30%, approximately 0.02% to approximately 29%, approximately 0.03% to approximately 28%, approximately 0.04% to approximately 27%, approximately 0.05% to approximately 26%, approximately 0.06% to approximately 25%, approximately 0.07% to approximately 24%, approximately 0.08% to approximately 23%, approximately 0.09% to approximately 22%, approximately 0.1% to approximately 21%, approximately 0.2% to approximately 20%, approximately 0.3% to approximately 19%, approximately 0.4% to approximately 18%, approximately 0.5% to approximately 17%, approximately 0.6% to approximately 16%, approximately 0.7% to approximately 15%, approximately 0.8% to approximately 14%, approximately 0.9% to approximately 12%, approximately 1% to approximately 10% w/w, w/v or v/v.

(440) In some embodiments, the concentration of one or more compounds of the invention is in the range from approximately 0.001% to approximately 10%, approximately 0.01% to approximately 5%, approximately 0.02% to approximately 4.5%, approximately 0.03% to approximately 4%, approximately 0.04% to approximately 3.5%, approximately 0.05% to approximately 3%, approximately 0.06% to approximately 2.5%, approximately 0.07% to approximately 2%, approximately 0.08% to approximately 1.5%, approximately 0.09% to approximately 1%, approximately 0.1% to approximately 0.9% w/w, w/v or v/v.

(441) In some embodiments, the amount of one or more compounds of the invention is equal to or less than 10 g, 9.5 g, 9.0 g, 8.5 g, 8.0 g, 7.5 g, 7.0 g, 6.5 g, 6.0 g, 5.5 g, 5.0 g, 4.5 g, 4.0 g, 3.5 g, 3.0 g, 2.5 g, 2.0 g, 1.5 g, 1.0 g, 0.95 g, 0.9 g, 0.85 g, 0.8 g, 0.75 g, 0.7 g, 0.65 g, 0.6 g, 0.55 g, 0.5 g, 0.45 g, 0.4 g, 0.35 g, 0.3 g, 0.25 g, 0.2 g, 0.15 g, 0.1 g, 0.09 g, 0.08 g, 0.07 g, 0.06 g, 0.05 g, 0.04 g, 0.03 g, 0.02 g, 0.01 g, 0.009 g, 0.008 g, 0.007 g, 0.006 g, 0.005 g, 0.004 g, 0.003 g, 0.002 g, 0.001 g, 0.0009 g, 0.0008 g, 0.0007 g, 0.0006 g, 0.0005 g, 0.0004 g, 0.0003 g, 0.0002 g, or 0.0001 g.

(442) In some embodiments, the amount of one or more compounds of the invention is more than 0.0001 g, 0.0002 g, 0.0003 g, 0.0004 g, 0.0005 g, 0.0006 g, 0.0007 g, 0.0008 g, 0.0009 g, 0.001 g, 0.0015 g, 0.002 g, 0.0025 g, 0.003 g, 0.0035 g, 0.004 g, 0.0045 g, 0.005 g, 0.0055 g, 0.006 g, 0.0065 g, 0.007 g, 0.0075 g, 0.008 g, 0.0085 g, 0.009 g, 0.0095 g, 0.01 g, 0.015 g, 0.02 g, 0.025 g, 0.03 g, 0.035 g, 0.04 g, 0.045 g, 0.05 g, 0.055 g, 0.06 g, 0.065 g, 0.07 g, 0.075 g, 0.08 g, 0.085 g, 0.09 g, 0.095 g, 0.1 g, 0.15 g, 0.2 g, 0.25 g, 0.3 g, 0.35 g, 0.4 g, 0.45 g, 0.5 g, 0.55 g, 0.6 g, 0.65 g, 0.7 g, 0.75 g, 0.8 g, 0.85 g, 0.9 g, 0.95 g, 1 g, 1.5 g, 2 g, 2.5, 3 g, 3.5, 4 g, 4.5 g, 5 g, 5.5 g, 6 g, 6.5 g, 7 g, 7.5 g, 8 g, 8.5 g, 9 g, 9.5 g, or 10 g.

(443) In some embodiments, the amount of one or more compounds of the invention is in the range of 0.0001-10 g, 0.0005-9 g, 0.001-8 g, 0.005-7 g, 0.01-6 g, 0.05-5 g, 0.1-4 g, 0.5-4 g, or 1-3 g.

(444) Kits/Articles of Manufacture

(445) For use in the therapeutic applications described herein, kits and articles of manufacture are also provided. In some embodiments, such kits comprise a carrier, package, or container that is compartmentalized to receive one or more containers such as vials, tubes, and the like, each of the container(s) comprising one of the separate elements to be used in a method described herein. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers are formed from a variety of materials such as glass or plastic.

(446) The articles of manufacture provided herein contain packaging materials. Packaging materials for use in packaging pharmaceutical products include those found in, e.g., U.S. Pat. Nos. 5,323,907, 5,052,558 and 5,033,252. Examples of pharmaceutical packaging materials include, but are not limited to, blister packs, bottles, tubes, inhalers, pumps, bags, vials, containers, syringes, bottles, and any packaging material suitable for a selected formulation and intended mode of administration and treatment. For example, the container(s) includes one or more compounds described herein, optionally in a composition or in combination with another agent as disclosed herein. The container(s) optionally have a sterile access port (for example the container is an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). Such kits optionally comprising a compound with an identifying description or label or instructions relating to its use in the methods described herein.

(447) For example, a kit typically includes one or more additional containers, each with one or more of various materials (such as reagents, optionally in concentrated form, and/or devices) desirable from a commercial and user standpoint for use of a compound described herein. Non-limiting examples of such materials include, but not limited to, buffers, diluents, filters, needles, syringes; carrier, package, container, vial and/or tube labels listing contents and/or instructions for use, and package inserts with instructions for use. A set of instructions will also typically be included. A label is optionally on or associated with the container. For example, a label is on a container when letters, numbers or other characters forming the label are attached, molded or etched into the container itself, a label is associated with a container when it is present within a receptacle or carrier that also holds the container, e.g., as a package insert. In addition, a label is used to indicate that the contents are to be used for a specific therapeutic application. In addition, the label indicates directions for use of the contents, such as in the methods described herein. In certain embodiments, the pharmaceutical composition is presented in a pack or dispenser device which contains one or more unit dosage forms containing a compound provided herein. The pack for example contains metal or plastic foil, such as a blister pack. Or, the pack or dispenser device is accompanied by instructions for administration. Or, the pack or dispenser is accompanied with a notice associated with the container in form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the drug for human or veterinary administration. Such notice, for example, is the labeling approved by the U.S. Food and Drug Administration for prescription drugs, or the approved product insert. In some embodiments, compositions containing a compound provided herein formulated in a compatible pharmaceutical carrier are prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

(448) Methods

(449) The present invention provides a method of inhibiting Ras-mediated cell signaling comprising contacting a cell with an effective amount of one or more compounds disclosed herein. Inhibition of Ras-mediated signal transduction can be assessed and demonstrated by a wide variety of ways known in the art. Non-limiting examples include a showing of (a) a decrease in GTPase activity of Ras; (b) a decrease in GTP binding affinity or an increase in GDP binding affinity; (c) an increase in K_{off} of GTP or a decrease in K_{off} of GDP; (d) a decrease in the levels of signaling

transduction molecules downstream in the Ras pathway, such as a decrease in pMEK level; and/or (e) a decrease in binding of Ras complex to downstream signaling molecules including but not limited to Raf. Kits and commercially available assays can be utilized for determining one or more of the above.

(450) The invention also provides methods of using the compounds or pharmaceutical compositions of the present invention to treat disease conditions, including but not limited to conditions implicated by G12C K-Ras, H-Ras or N-Ras mutation, G12C H-Ras mutation and/or G12C N-Ras mutation (e.g., cancer).

(451) In some embodiments, a method for treatment of cancer is provided, the method comprising administering an effective amount of any of the foregoing pharmaceutical compositions comprising a compound of structure (I) to a subject in need thereof. In some embodiments, the cancer is mediated by a K-Ras, H-Ras or N-Ras G12C mutation. In other embodiments, the cancer is pancreatic cancer, colon cancer, MYH associated polyposis, colorectal cancer or lung cancer.

(452) In some embodiments the invention provides method of treating a disorder in a subject in need thereof, wherein the said method comprises determining if the subject has a K-Ras, H-Ras or N-Ras G12C mutation and if the subject is determined to have the K-Ras, H-Ras or N-Ras G12C mutation, then administering to the subject a therapeutically effective dose of at least one compound of structure (I) or a pharmaceutically acceptable salt, ester, prodrug, tautomer, solvate, hydrate or derivative thereof.

(453) The disclosed compounds strongly inhibit anchorage-independent cell growth and therefore have the potential to inhibit tumor metastasis. Accordingly, in another embodiment the disclosure provides a method for inhibiting tumor metastasis, the method comprising administering an effective amount a pharmaceutical composition of comprising any of the compounds disclosed herein and a pharmaceutically acceptable carrier to a subject in need thereof.

(454) K-Ras, H-Ras or N-Ras G12C mutations have also been identified in hematological malignancies (e.g., cancers that affect blood, bone marrow and/or lymph nodes). Accordingly, certain embodiments are directed to administration of a disclosed compounds (e.g., in the form of a pharmaceutical composition) to a patient in need of treatment of a hematological malignancy. Such malignancies include, but are not limited to leukemias and lymphomas. For example, the presently disclosed compounds can be used for treatment of diseases such as Acute lymphoblastic leukemia (ALL), Acute myelogenous leukemia (AML), Chronic lymphocytic leukemia (CLL), small lymphocytic lymphoma (SLL), Chronic myelogenous leukemia (CML), Acute monocytic leukemia (AMoL) and/or other leukemias. In other embodiments, the compounds are useful for treatment of lymphomas such as all subtypes of Hodgkins lymphoma or non-Hodgkins lymphoma.

(455) Determining whether a tumor or cancer comprises a G12C K-Ras, H-Ras or N-Ras mutation can be undertaken by assessing the nucleotide sequence encoding the K-Ras, H-Ras or N-Ras protein, by assessing the amino acid sequence of the K-Ras, H-Ras or N-Ras protein, or by assessing the characteristics of a putative K-Ras, H-Ras or N-Ras mutant protein. The sequence of wild-type human K-Ras, H-Ras or N-Ras is known in the art, (e.g. Accession No. NP203524).

(456) Methods for detecting a mutation in a K-Ras, H-Ras or N-Ras nucleotide sequence are known by those of skill in the art. These methods include, but are not limited to, polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) assays, polymerase chain reaction-single strand conformation polymorphism (PCR-SSCP) assays, real-time PCR assays, PCR sequencing, mutant allele-specific PCR amplification (MASA) assays, direct sequencing, primer extension reactions, electrophoresis, oligonucleotide ligation assays, hybridization assays, TaqMan assays, SNP genotyping assays, high resolution melting assays and microarray analyses. In some embodiments, samples are evaluated for G12C K-Ras, H-Ras or N-Ras mutations by real-time PCR. In real-time PCR, fluorescent probes specific for the K-Ras, H-Ras or N-Ras G12C mutation are used. When a mutation is present, the probe binds and fluorescence is detected. In some embodiments, the K-Ras, H-Ras or N-Ras G12C mutation is identified using a direct

sequencing method of specific regions (e.g., exon 2 and/or exon 3) in the K-Ras, H-Ras or N-Ras gene. This technique will identify all possible mutations in the region sequenced.

(457) Methods for detecting a mutation in a K-Ras, H-Ras or N-Ras protein are known by those of skill in the art. These methods include, but are not limited to, detection of a K-Ras, H-Ras or N-Ras mutant using a binding agent (e.g., an antibody) specific for the mutant protein, protein electrophoresis and Western blotting, and direct peptide sequencing.

(458) Methods for determining whether a tumor or cancer comprises a G12C K-Ras, H-Ras or N-Ras mutation can use a variety of samples. In some embodiments, the sample is taken from a subject having a tumor or cancer. In some embodiments, the sample is taken from a subject having a cancer or tumor. In some embodiments, the sample is a fresh tumor/cancer sample. In some embodiments, the sample is a frozen tumor/cancer sample. In some embodiments, the sample is a formalin-fixed paraffin-embedded sample. In some embodiments, the sample is processed to a cell lysate. In some embodiments, the sample is processed to DNA or RNA.

(459) The invention also relates to a method of treating a hyperproliferative disorder in a mammal that comprises administering to said mammal a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt, ester, prodrug, solvate, hydrate or derivative thereof. In some embodiments, said method relates to the treatment of cancer such as acute myeloid leukemia, cancer in adolescents, adrenocortical carcinoma childhood, AIDS-related cancers (e.g. Lymphoma and Kaposi's Sarcoma), anal cancer, appendix cancer, astrocytomas, atypical teratoid, basal cell carcinoma, bile duct cancer, bladder cancer, bone cancer, brain stem glioma, brain tumor, breast cancer, bronchial tumors, burkitt lymphoma, carcinoid tumor, atypical teratoid, embryonal tumors, germ cell tumor, primary lymphoma, cervical cancer, childhood cancers, chordoma, cardiac tumors, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myeloproliferative disorders, colon cancer, colorectal cancer, craniopharyngioma, cutaneous T-cell lymphoma, extrahepatic ductal carcinoma in situ (DCIS), embryonal tumors, CNS cancer, endometrial cancer, ependymoma, esophageal cancer, esthesioneuroblastoma, ewing sarcoma, extracranial germ cell tumor, extragonadal germ cell tumor, eye cancer, fibrous histiocytoma of bone, gall bladder cancer, gastric cancer, gastrointestinal carcinoid tumor, gastrointestinal stromal tumors (GIST), germ cell tumor, gestational trophoblastic tumor, hairy cell leukemia, head and neck cancer, heart cancer, liver cancer, hodgkin lymphoma, hypopharyngeal cancer, intraocular melanoma, islet cell tumors, pancreatic neuroendocrine tumors, kidney cancer, laryngeal cancer, lip and oral cavity cancer, liver cancer, lobular carcinoma in situ (LCIS), lung cancer, lymphoma, metastatic squamous neck cancer with occult primary, midline tract carcinoma, mouth cancer multiple endocrine neoplasia syndromes, multiple myeloma/plasma cell neoplasm, mycosis fungoides, myelodysplastic syndromes, myelodysplastic/myeloproliferative neoplasms, multiple myeloma, merkel cell carcinoma, malignant mesothelioma, malignant fibrous histiocytoma of bone and osteosarcoma, nasal cavity and paranasal sinus cancer, nasopharyngeal cancer, neuroblastoma, non-hodgkin lymphoma, non-small cell lung cancer (NSCLC), oral cancer, lip and oral cavity cancer, oropharyngeal cancer, ovarian cancer, pancreatic cancer, papillomatosis, paraganglioma, paranasal sinus and nasal cavity cancer, parathyroid cancer, penile cancer, pharyngeal cancer, pleuropulmonary blastoma, primary central nervous system (CNS) lymphoma, prostate cancer, rectal cancer, transitional cell cancer, retinoblastoma, rhabdomyosarcoma, salivary gland cancer, skin cancer, stomach (gastric) cancer, small cell lung cancer, small intestine cancer, soft tissue sarcoma, T-Cell lymphoma, testicular cancer, throat cancer, thymoma and thymic carcinoma, thyroid cancer, transitional cell cancer of the renal pelvis and ureter, trophoblastic tumor, unusual cancers of childhood, urethral cancer, uterine sarcoma, vaginal cancer, vulvar cancer, or Viral-Induced cancer. In some embodiments, said method relates to the treatment of a non-cancerous hyperproliferative disorder such as benign hyperplasia of the skin (e.g., psoriasis), restenosis, or prostate (e.g., benign prostatic hypertrophy (BPH)).

(460) In certain particular embodiments, the invention relates to methods for treatment of lung

cancers, the methods comprise administering an effective amount of any of the above described compound (or a pharmaceutical composition comprising the same) to a subject in need thereof. In certain embodiments the lung cancer is a non-small cell lung carcinoma (NSCLC), for example adenocarcinoma, squamous-cell lung carcinoma or large-cell lung carcinoma. In other embodiments, the lung cancer is a small cell lung carcinoma. Other lung cancers treatable with the disclosed compounds include, but are not limited to, glandular tumors, carcinoid tumors and undifferentiated carcinomas.

(461) Subjects that can be treated with compounds of the invention, or pharmaceutically acceptable salt, ester, prodrug, solvate, tautomer, hydrate or derivative of said compounds, according to the methods of this invention include, for example, subjects that have been diagnosed as having acute myeloid leukemia, acute myeloid leukemia, cancer in adolescents, adrenocortical carcinoma childhood, AIDS-related cancers (e.g. Lymphoma and Kaposi's Sarcoma), anal cancer, appendix cancer, astrocytomas, atypical teratoid, basal cell carcinoma, bile duct cancer, bladder cancer, bone cancer, brain stem glioma, brain tumor, breast cancer, bronchial tumors, burkitt lymphoma, carcinoid tumor, atypical teratoid, embryonal tumors, germ cell tumor, primary lymphoma, cervical cancer, childhood cancers, chordoma, cardiac tumors, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myeloproliferative disorders, colon cancer, colorectal cancer, craniopharyngioma, cutaneous T-cell lymphoma, extrahepatic ductal carcinoma in situ (DCIS), embryonal tumors, CNS cancer, endometrial cancer, ependymoma, esophageal cancer, esthesioneuroblastoma, ewing sarcoma, extracranial germ cell tumor, extragonadal germ cell tumor, eye cancer, fibrous histiocytoma of bone, gall bladder cancer, gastric cancer, gastrointestinal carcinoid tumor, gastrointestinal stromal tumors (GIST), germ cell tumor, gestational trophoblastic tumor, hairy cell leukemia, head and neck cancer, heart cancer, liver cancer, hodgkin lymphoma, hypopharyngeal cancer, intraocular melanoma, islet cell tumors, pancreatic neuroendocrine tumors, kidney cancer, laryngeal cancer, lip and oral cavity cancer, liver cancer, lobular carcinoma in situ (LCIS), lung cancer, lymphoma, metastatic squamous neck cancer with occult primary, midline tract carcinoma, mouth cancer multiple endocrine neoplasia syndromes, multiple myeloma/plasma cell neoplasm, mycosis fungoides, myelodysplastic syndromes, myelodysplastic/myeloproliferative neoplasms, multiple myeloma, merkel cell carcinoma, malignant mesothelioma, malignant fibrous histiocytoma of bone and osteosarcoma, nasal cavity and paranasal sinus cancer, nasopharyngeal cancer, neuroblastoma, non-hodgkin lymphoma, non-small cell lung cancer (NSCLC), oral cancer, lip and oral cavity cancer, oropharyngeal cancer, ovarian cancer, pancreatic cancer, papillomatosis, paraganglioma, paranasal sinus and nasal cavity cancer, parathyroid cancer, penile cancer, pharyngeal cancer, pleuropulmonary blastoma, primary central nervous system (CNS) lymphoma, prostate cancer, rectal cancer, transitional cell cancer, retinoblastoma, rhabdomyosarcoma, salivary gland cancer, skin cancer, stomach (gastric) cancer, small cell lung cancer, small intestine cancer, soft tissue sarcoma, T-Cell lymphoma, testicular cancer, throat cancer, thymoma and thymic carcinoma, thyroid cancer, transitional cell cancer of the renal pelvis and ureter, trophoblastic tumor, unusual cancers of childhood, urethral cancer, uterine sarcoma, vaginal cancer, vulvar cancer, or Viral-Induced cancer. In some embodiments subjects that are treated with the compounds of the invention include subjects that have been diagnosed as having a non-cancerous hyperproliferative disorder such as benign hyperplasia of the skin (e.g., psoriasis), restenosis, or prostate (e.g., benign prostatic hypertrophy (BPH)).

(462) The invention further provides methods of modulating a G12C Mutant K-Ras, H-Ras or N-Ras protein activity by contacting the protein with an effective amount of a compound of the invention. Modulation can be inhibiting or activating protein activity. In some embodiments, the invention provides methods of inhibiting protein activity by contacting the G12C Mutant K-Ras, H-Ras or N-Ras protein with an effective amount of a compound of the invention in solution. In some embodiments, the invention provides methods of inhibiting the G12C Mutant K-Ras, H-Ras

or N-Ras protein activity by contacting a cell, tissue, organ that express the protein of interest. In some embodiments, the invention provides methods of inhibiting protein activity in subject including but not limited to rodents and mammal (e.g., human) by administering into the subject an effective amount of a compound of the invention. In some embodiments, the percentage modulation exceeds 25%, 30%, 40%, 50%, 60%, 70%, 80%, or 90%. In some embodiments, the percentage of inhibiting exceeds 25%, 30%, 40%, 50%, 60%, 70%, 80%, or 90%.

(463) In some embodiments, the invention provides methods of inhibiting K-Ras, H-Ras or N-Ras G12C activity in a cell by contacting said cell with an amount of a compound of the invention sufficient to inhibit the activity of K-Ras, H-Ras or N-Ras G12C in said cell. In some embodiments, the invention provides methods of inhibiting K-Ras, H-Ras or N-Ras G12C activity in a tissue by contacting said tissue with an amount of a compound of the invention sufficient to inhibit the activity of K-Ras, H-Ras or N-Ras G12C in said tissue. In some embodiments, the invention provides methods of inhibiting K-Ras, H-Ras or N-Ras G12C activity in an organism by contacting said organism with an amount of a compound of the invention sufficient to inhibit the activity of K-Ras, H-Ras or N-Ras G12C in said organism. In some embodiments, the invention provides methods of inhibiting K-Ras, H-Ras or N-Ras G12C activity in an animal by contacting said animal with an amount of a compound of the invention sufficient to inhibit the activity of K-Ras, H-Ras or N-Ras G12C in said animal. In some embodiments, the invention provides methods of inhibiting K-Ras, H-Ras or N-Ras G12C activity in a mammal by contacting said mammal with an amount of a compound of the invention sufficient to inhibit the activity of K-Ras, H-Ras or N-Ras G12C in said mammal. In some embodiments, the invention provides methods of inhibiting K-Ras, H-Ras or N-Ras G12C activity in a human by contacting said human with an amount of a compound of the invention sufficient to inhibit the activity of K-Ras, H-Ras or N-Ras G12C in said human. The present invention provides methods of treating a disease mediated by K-Ras, H-Ras or N-Ras G12C activity in a subject in need of such treatment.

(464) The present invention also provides methods for combination therapies in which an agent known to modulate other pathways, or other components of the same pathway, or even overlapping sets of target enzymes are used in combination with a compound of the present invention, or a pharmaceutically acceptable salt, ester, prodrug, solvate, tautomer, hydrate or derivative thereof. In one aspect, such therapy includes but is not limited to the combination of one or more compounds of the invention with chemotherapeutic agents, therapeutic antibodies, and radiation treatment, to provide a synergistic or additive therapeutic effect.

(465) Many chemotherapeutics are presently known in the art and can be used in combination with the compounds of the invention. In some embodiments, the chemotherapeutic is selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites, intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, angiogenesis inhibitors, and anti-androgens.

(466) Non-limiting examples are chemotherapeutic agents, cytotoxic agents, and non-peptide small molecules such as Gleevec® (Imatinib Mesylate), Velcade® (bortezomib), Casodex (bicalutamide), Iressa® (gefitinib), and Adriamycin as well as a host of chemotherapeutic agents. Non-limiting examples of chemotherapeutic agents include alkylating agents such as thiopeta and cyclophosphamide (CYTOXAN™); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, trietylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabycin, carminomycin, carzinophilin,

Casodex™, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, androgens such as calusterone, dromostanolone propionate, epitostanol, mepitostane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfomithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK®; razoxane; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepe; taxanes, e.g. paclitaxel (TAXOL™, Bristol-Myers Squibb Oncology, Princeton, N.J.) and docetaxel (TAXOTERE™, Rhone-Poulenc Rorer, Antony, France); retinoic acid; esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included as suitable chemotherapeutic cell conditioners are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, (Nolvadex™), raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY 117018, onapristone, and toremifene (Fareston); and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; camptothecin-11 (CPT-11); topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO). Where desired, the compounds or pharmaceutical composition of the present invention can be used in combination with commonly prescribed anti-cancer drugs such as Herceptin®, Avastin®, Erbitux®, Rituxan®, Taxol®, Arimidex®, Taxotere®, ABVD, AVICINE, Abagovomab, Acridine carboxamide, Adecatumumab, 17-N-Allylamino-17-demethoxygeldanamycin, Alpharadin, Alvocidib, 3-Aminopyridine-2-carboxaldehyde thiosemicarbazone, Amonafide, Anthracenedione, Anti-CD22 immunotoxins, Antineoplastic, Antitumorigenic herbs, Apaziquone, Atiprimod, Azathioprine, Belotecan, Bendamustine, BIBW 2992, Biricodar, Brostallicin, Bryostatins, Buthionine sulfoximine, CBV (chemotherapy), Calyculin, cell-cycle nonspecific antineoplastic agents, Dichloroacetic acid, Discodermolide, Elsamitrucin, Enocitabine, Epothilone, Eribulin, Everolimus, Exatecan, Exisulind, Ferruginol, Forodesine, Fosfestrol, ICE chemotherapy regimen, IT-101, Imexon, Imiquimod, Indolocarbazole, Irofulven, Laniquidar, Larotaxel, Lenalidomide, Lucanthone, Lurtotecan, Mafosfamide, Mitozolomide, Nafoxidine, Nedaplatin, Olaparib, Ortataxel, PAC-1, Pawpaw, Pixantrone, Proteasome inhibitor, Rebeccamycin, Resiquimod, Rubitecan, SN-38, Salinosporamide A, Sapacitabine, Stanford V, Swainsonine, Talaporfin, Tariquidar, Tegafur-uracil, Temodar, Tesetaxel, Triplatin tetranitrate, Tris(2-chloroethyl)amine, Troxacitabine, Uramustine, Vadimezan, Vinflunine, ZD6126 or Zosuquidar.

(467) This invention further relates to a method for using the compounds or pharmaceutical compositions provided herein, in combination with radiation therapy for inhibiting abnormal cell growth or treating the hyperproliferative disorder in the mammal. Techniques for administering radiation therapy are known in the art, and these techniques can be used in the combination therapy described herein. The administration of the compound of the invention in this combination therapy

can be determined as described herein.

(468) Radiation therapy can be administered through one of several methods, or a combination of methods, including without limitation external-beam therapy, internal radiation therapy, implant radiation, stereotactic radiosurgery, systemic radiation therapy, radiotherapy and permanent or temporary interstitial brachytherapy. The term "brachytherapy," as used herein, refers to radiation therapy delivered by a spatially confined radioactive material inserted into the body at or near a tumor or other proliferative tissue disease site. The term is intended without limitation to include exposure to radioactive isotopes (e.g. At-211, I-131, I-125, Y-90, Re-186, Re-188, Sm-153, Bi-212, P-32, and radioactive isotopes of Lu). Suitable radiation sources for use as a cell conditioner of the present invention include both solids and liquids. By way of non-limiting example, the radiation source can be a radionuclide, such as I-125, I-131, Yb-169, Ir-192 as a solid source, I-125 as a solid source, or other radionuclides that emit photons, beta particles, gamma radiation, or other therapeutic rays. The radioactive material can also be a fluid made from any solution of radionuclide(s), e.g., a solution of I-125 or I-131, or a radioactive fluid can be produced using a slurry of a suitable fluid containing small particles of solid radionuclides, such as Au-198, Y-90. Moreover, the radionuclide(s) can be embodied in a gel or radioactive micro spheres.

(469) Without being limited by any theory, the compounds of the present invention can render abnormal cells more sensitive to treatment with radiation for purposes of killing and/or inhibiting the growth of such cells. Accordingly, this invention further relates to a method for sensitizing abnormal cells in a mammal to treatment with radiation which comprises administering to the mammal an amount of a compound of the present invention or pharmaceutically acceptable salt, ester, prodrug, solvate, hydrate or derivative thereof, which amount is effective in sensitizing abnormal cells to treatment with radiation. The amount of the compound, salt, or solvate in this method can be determined according to the means for ascertaining effective amounts of such compounds described herein.

(470) The compounds or pharmaceutical compositions of the invention can be used in combination with an amount of one or more substances selected from anti-angiogenesis agents, signal transduction inhibitors, antiproliferative agents, glycolysis inhibitors, or autophagy inhibitors.

(471) Anti-angiogenesis agents, such as MMP-2 (matrix-metalloproteinase 2) inhibitors, MMP-9 (matrix-metalloproteinase 9) inhibitors, and COX-11 (cyclooxygenase 11) inhibitors, can be used in conjunction with a compound of the invention and pharmaceutical compositions described herein. Anti-angiogenesis agents include, for example, rapamycin, temsirolimus (CCI-779), everolimus (RAD001), sorafenib, sunitinib, and bevacizumab. Examples of useful COX-II inhibitors include CELEBREX™ (alecoxib), valdecoxib, and rofecoxib. Examples of useful matrix metalloproteinase inhibitors are described in WO 96/33172 (published Oct. 24, 1996), WO 96/27583 (published Mar. 7, 1996), European Patent Application No. 97304971.1 (filed Jul. 8, 1997), European Patent Application No. 99308617.2 (filed Oct. 29, 1999), WO 98/07697 (published Feb. 26, 1998), WO 98/03516 (published Jan. 29, 1998), WO 98/34918 (published Aug. 13, 1998), WO 98/34915 (published Aug. 13, 1998), WO 98/33768 (published Aug. 6, 1998), WO 98/30566 (published Jul. 16, 1998), European Patent Publication 606,046 (published Jul. 13, 1994), European Patent Publication 931, 788 (published Jul. 28, 1999), WO 90/05719 (published May 31, 1990), WO 99/52910 (published Oct. 21, 1999), WO 99/52889 (published Oct. 21, 1999), WO 99/29667 (published Jun. 17, 1999), PCT International Application No. PCT/IB98/01113 (filed Jul. 21, 1998), European Patent Application No. 99302232.1 (filed Mar. 25, 1999), Great Britain Patent Application No. 9912961.1 (filed Jun. 3, 1999), U.S. Provisional Application No. 60/148,464 (filed Aug. 12, 1999), U.S. Pat. No. 5,863,949 (issued Jan. 26, 1999), U.S. Pat. No. 5,861,510 (issued Jan. 19, 1999), and European Patent Publication 780,386 (published Jun. 25, 1997), all of which are incorporated herein in their entireties by reference. Preferred MMP-2 and MMP-9 inhibitors are those that have little or no activity inhibiting MMP-1. More preferred, are those that selectively inhibit MMP-2 and/or MMP-9 relative to the other matrix-metalloproteinases (i.e., MMP-1, MMP-

3, MMP-4, MMP-5, MMP-6, MMP-7, MMP-8, MMP-10, MMP-11, MMP-12, and MMP-13). Some specific examples of MMP inhibitors useful in the invention are AG-3340, RO 32-3555, and RS 13-0830.

(472) Autophagy inhibitors include, but are not limited to chloroquine, 3-methyladenine, hydroxychloroquine (Plaquenil™), bafilomycin A1, 5-amino-4-imidazole carboxamide riboside (AICAR), okadaic acid, autophagy-suppressive algal toxins which inhibit protein phosphatases of type 2A or type 1, analogues of cAMP, and drugs which elevate cAMP levels such as adenosine, LY204002, N6-mercaptopurine riboside, and vinblastine. In addition, antisense or siRNA that inhibits expression of proteins including but not limited to ATG5 (which are implicated in autophagy), may also be used.

(473) The invention also relates to a method of and to a pharmaceutical composition for treating a cardiovascular disease in a mammal which comprises an amount of a compound of the invention, or a pharmaceutically acceptable salt, ester, prodrug, solvate, tautomer, hydrate or derivative thereof, or an isotopically-labeled derivative thereof, and an amount of one or more therapeutic agents use for the treatment of cardiovascular diseases.

(474) Exemplary agents for use in cardiovascular disease applications are anti-thrombotic agents, e.g., prostacyclin and salicylates, thrombolytic agents, e.g., streptokinase, urokinase, tissue plasminogen activator (TPA) and anisoylated plasminogen-streptokinase activator complex (APSAC), anti-platelets agents, e.g., acetyl-salicylic acid (ASA) and clopidogrel, vasodilating agents, e.g., nitrates, calcium channel blocking drugs, anti-proliferative agents, e.g., colchicine and alkylating agents, intercalating agents, growth modulating factors such as interleukins, transformation growth factor-beta and congeners of platelet derived growth factor, monoclonal antibodies directed against growth factors, anti-inflammatory agents, both steroidal and non-steroidal, and other agents that can modulate vessel tone, function, arteriosclerosis, and the healing response to vessel or organ injury post intervention. Antibiotics can also be included in combinations or coatings comprised by the invention. Moreover, a coating can be used to effect therapeutic delivery focally within the vessel wall. By incorporation of the active agent in a swellable polymer, the active agent will be released upon swelling of the polymer.

(475) In some embodiments, the compounds described herein are formulated or administered in conjunction with liquid or solid tissue barriers also known as lubricants. Examples of tissue barriers include, but are not limited to, polysaccharides, polyglycans, seprafilm, interceed and hyaluronic acid.

(476) In some embodiments, medicaments which are administered in conjunction with the compounds described herein include any suitable drugs usefully delivered by inhalation for example, analgesics, e.g. codeine, dihydromorphine, ergotamine, fentanyl or morphine; anginal preparations, e.g. diltiazem; antiallergics, e.g. cromoglycate, ketotifen or nedocromil; anti-infectives, e.g. cephalosporins, penicillins, streptomycin, sulphonamides, tetracyclines or pentamidine; antihistamines, e.g. methapyrilene; anti-inflammatories, e.g. beclomethasone, flunisolide, budesonide, tipredane, triamcinolone acetone or fluticasone; antitussives, e.g. noscapine; bronchodilators, e.g. ephedrine, adrenaline, fenoterol, formoterol, isoprenaline, metaproterenol, phenylephrine, phenylpropanolamine, pirbuterol, reproterol, rimiterol, salbutamol, salmeterol, terbutalin, isoetharine, tulobuterol, orciprenaline or (-)-4-amino-3,5-dichloro- α -[[[6-[2-(2-pyridinyl)ethoxy]hexyl]-amino]methyl]benzenemethanol; diuretics, e.g. amiloride; anticholinergics e.g. ipratropium, atropine or oxitropium; hormones, e.g. cortisone, hydrocortisone or prednisolone; xanthines e.g. aminophylline, choline theophyllinate, lysine theophyllinate or theophylline; and therapeutic proteins and peptides, e.g. insulin or glucagon. It will be clear to a person skilled in the art that, where appropriate, the medicaments are used in the form of salts (e.g. as alkali metal or amine salts or as acid addition salts) or as esters (e.g. lower alkyl esters) or as solvates (e.g. hydrates) to optimize the activity and/or stability of the medicament.

(477) Other exemplary therapeutic agents useful for a combination therapy include but are not

limited to agents as described above, radiation therapy, hormone antagonists, hormones and their releasing factors, thyroid and antithyroid drugs, estrogens and progestins, androgens, adrenocorticotrophic hormone; adrenocortical steroids and their synthetic analogs; inhibitors of the synthesis and actions of adrenocortical hormones, insulin, oral hypoglycemic agents, and the pharmacology of the endocrine pancreas, agents affecting calcification and bone turnover: calcium, phosphate, parathyroid hormone, vitamin D, calcitonin, vitamins such as water-soluble vitamins, vitamin B complex, ascorbic acid, fat-soluble vitamins, vitamins A, K, and E, growth factors, cytokines, chemokines, muscarinic receptor agonists and antagonists; anticholinesterase agents; agents acting at the neuromuscular junction and/or autonomic ganglia; catecholamines, sympathomimetic drugs, and adrenergic receptor agonists or antagonists; and 5-hydroxytryptamine (5-HT, serotonin) receptor agonists and antagonists.

(478) Therapeutic agents can also include agents for pain and inflammation such as histamine and histamine antagonists, bradykinin and bradykinin antagonists, 5-hydroxytryptamine (serotonin), lipid substances that are generated by biotransformation of the products of the selective hydrolysis of membrane phospholipids, eicosanoids, prostaglandins, thromboxanes, leukotrienes, aspirin, nonsteroidal anti-inflammatory agents, analgesic-antipyretic agents, agents that inhibit the synthesis of prostaglandins and thromboxanes, selective inhibitors of the inducible cyclooxygenase, selective inhibitors of the inducible cyclooxygenase-2, autacoids, paracrine hormones, somatostatin, gastrin, cytokines that mediate interactions involved in humoral and cellular immune responses, lipid-derived autacoids, eicosanoids, β -adrenergic agonists, ipratropium, glucocorticoids, methylxanthines, sodium channel blockers, opioid receptor agonists, calcium channel blockers, membrane stabilizers and leukotriene inhibitors.

(479) Additional therapeutic agents contemplated herein include diuretics, vasopressin, agents affecting the renal conservation of water, rennin, angiotensin, agents useful in the treatment of myocardial ischemia, anti-hypertensive agents, angiotensin converting enzyme inhibitors, β -adrenergic receptor antagonists, agents for the treatment of hypercholesterolemia, and agents for the treatment of dyslipidemia.

(480) Other therapeutic agents contemplated include drugs used for control of gastric acidity, agents for the treatment of peptic ulcers, agents for the treatment of gastroesophageal reflux disease, prokinetic agents, antiemetics, agents used in irritable bowel syndrome, agents used for diarrhea, agents used for constipation, agents used for inflammatory bowel disease, agents used for biliary disease, agents used for pancreatic disease. Therapeutic agents used to treat protozoan infections, drugs used to treat Malaria, Amebiasis, Giardiasis, Trichomoniasis, Trypanosomiasis, and/or Leishmaniasis, and/or drugs used in the chemotherapy of helminthiasis. Other therapeutic agents include antimicrobial agents, sulfonamides, trimethoprim-sulfamethoxazole quinolones, and agents for urinary tract infections, penicillins, cephalosporins, and other, β -lactam antibiotics, an agent comprising an aminoglycoside, protein synthesis inhibitors, drugs used in the chemotherapy of tuberculosis, Mycobacterium avium complex disease, and leprosy, antifungal agents, antiviral agents including nonretroviral agents and antiretroviral agents.

(481) Examples of therapeutic antibodies that can be combined with a compound of the invention include but are not limited to anti-receptor tyrosine kinase antibodies (cetuximab, panitumumab, trastuzumab), anti CD20 antibodies (rituximab, tositumomab), and other antibodies such as alemtuzumab, bevacizumab, and gemtuzumab.

(482) Moreover, therapeutic agents used for immunomodulation, such as immunomodulators, immunosuppressive agents, tolerogens, and immunostimulants are contemplated by the methods herein. In addition, therapeutic agents acting on the blood and the blood-forming organs, hematopoietic agents, growth factors, minerals, and vitamins, anticoagulant, thrombolytic, and antiplatelet drugs.

(483) For treating renal carcinoma, one may combine a compound of the present invention with sorafenib and/or avastin. For treating an endometrial disorder, one may combine a compound of the

present invention with doxorubicin, taxotere (taxol), and/or cisplatin (carboplatin). For treating ovarian cancer, one may combine a compound of the present invention with cisplatin (carboplatin), taxotere, doxorubicin, topotecan, and/or tamoxifen. For treating breast cancer, one may combine a compound of the present invention with taxotere (taxol), gemcitabine (capecitabine), tamoxifen, letrozole, tarceva, lapatinib, PD0325901, avastin, herceptin, OSI-906, and/or OSI-930. For treating lung cancer, one may combine a compound of the present invention with taxotere (taxol), gemcitabine, cisplatin, pemetrexed, Tarceva, PD0325901, and/or avastin.

(484) Further therapeutic agents that can be combined with a compound of the invention are found in Goodman and Gilman's "The Pharmacological Basis of Therapeutics" Tenth Edition edited by Hardman, Limbird and Gilman or the Physician's Desk Reference, both of which are incorporated herein by reference in their entirety.

(485) The compounds described herein can be used in combination with the agents disclosed herein or other suitable agents, depending on the condition being treated. Hence, in some embodiments the one or more compounds of the invention will be co-administered with other agents as described above. When used in combination therapy, the compounds described herein are administered with the second agent simultaneously or separately. This administration in combination can include simultaneous administration of the two agents in the same dosage form, simultaneous administration in separate dosage forms, and separate administration. That is, a compound described herein and any of the agents described above can be formulated together in the same dosage form and administered simultaneously. Alternatively, a compound of the invention and any of the agents described above can be simultaneously administered, wherein both the agents are present in separate formulations. In another alternative, a compound of the present invention can be administered just followed by and any of the agents described above, or vice versa. In some embodiments of the separate administration protocol, a compound of the invention and any of the agents described above are administered a few minutes apart, or a few hours apart, or a few days apart.

(486) The examples and preparations provided below further illustrate and exemplify the compounds of the present invention and methods of preparing such compounds. It is to be understood that the scope of the present invention is not limited in any way by the scope of the following examples and preparations. In the following examples, and throughout the specification and claims, molecules with a single chiral center, unless otherwise noted, exist as a racemic mixture. Those molecules with two or more chiral centers, unless otherwise noted, exist as a racemic mixture of diastereomers. Single enantiomers/diastereomers may be obtained by methods known to those skilled in the art.

EXAMPLES

Example 1

(487) ##STR00506##

5-Bromo-4-chloro-2-methoxyphenyl)methanol

(488) To a solution of methyl 5-bromo-4-chloro-2-methoxybenzoate (1.0 g, 3.6 mmol) in dry THF (10 mL) at 0° C., super-Hydride (1.0 M, 10 mL) was added and the resulting mixture was stirred at room temperature overnight. The mixture was partitioned between ethyl acetate and water. The organic layer was dried over Na.sub.2SO.sub.4, filtered and concentrated in vacuo to afford the desired product. The crude product was used directly in the next step without further purification.

5-Bromo-4-chloro-2-methoxybenzaldehyde

(489) To the solution of 5-bromo-4-chloro-2-methoxyphenyl)methanol in dry DCM (5 mL), Dess-Martin reagent (607 mg) was added and the resulting mixture was stirred at room temperature overnight. The mixture was partitioned between DCM and water. The organic layer was dried over Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified via Isolera One (silica cartridge, 0-60% ethyl acetate/hexanes) to afford the desired product (200 mg, 56% yield).
.sup.1H NMR (300 MHz, CDCl.sub.3) δ : 10.33 (s, 1H), 8.04 (s, 1H), 7.12 (s, 1H), 3.39 (s, 3H).

N-(5-bromo-4-chloro-2-methoxybenzyl)-1-(2,4-dimethoxyphenyl)methanamine

(490) The solution of 5-bromo-4-chloro-2-methoxybenzaldehyde (300 mg, 1.2 mmol) and 2,4-dimethoxybenzyl amine (167 mg, 1.2 mmol) in MeOH/DCM (4:1, 10 mL) was stirred at room temperature for 5 h. To this mixture, NaBH₄ (100 mg, 13 mmol) was added in two portions. The mixture stirred for 1 h and then partitioned between DCM and water. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo to afford the desired product. The crude product was used directly in the next step without further purification.

tert-Butyl 4-((5-bromo-4-chloro-2-methoxybenzyl)(2,4-dimethoxybenzyl)carbamoyl)piperidine-1-carboxylate

(491) The mixture of N-(5-bromo-4-chloro-2-methoxybenzyl)-1-(2,4-dimethoxyphenyl)methanamine (190 mg, 0.47 mmol), 1-(tert-butoxycarbonyl)piperidine-4-carboxylic acid (109 mg, 0.47 mmol), HATU (180 mg, 0.47 mmol) and Et₃N (101 mg, 1 mmol) in DMF (5 mL) was stirred at room temperature for 4 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo to afford the desired product. The crude product was used directly in the next step without further purification.

N-(5-Bromo-4-chloro-2-methoxybenzyl)piperidine-4-carboxamide

(492) The crude tert-Butyl 4-((5-bromo-4-chloro-2-methoxybenzyl)(2,4-dimethoxybenzyl)carbamoyl)piperidine-1-carboxylate was dissolved in 50% TFA in DCM (10 mL) and the resulting mixture was stirred at room temperature for 2 h. The mixture was concentrated in vacuo and the residue was partitioned between DCM and saturate NaHCO₃ aqueous solution. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo to afford the desired product.

1-Acryloyl-N-(5-bromo-4-chloro-2-methoxybenzyl)piperidine-4-carboxamide

(493) To a solution of N-(5-Bromo-4-chloro-2-methoxybenzyl)piperidine-4-carboxamide obtained from above step in DCM (10 mL) at 0° C., Et₃N (0.2 mL, 1.43 mmol) and acryloyl chloride (36.2 mg, 0.4 mmol) were added sequentially and the mixture was stirred at room temperature for 1 h. The mixture was partitioned between DCM and water. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo. The residue was purified via Isolera One (silica cartridge, 0-3% MeOH/DCM) to afford the desired product (13 mg). ¹H NMR (300 MHz, DMSO-d₆) δ: 8.27 (t, J=5.6 Hz, 1H), 7.37 (s, 1H), 7.26 (s, 1H), 6.81 (dd, J=10.4, 18.8 Hz, 1H), 6.08 (dd, J=2.4, 16.7 Hz, 1H), 5.66 (dd, J=2.4, 10.4 Hz, 1H), 4.40 (d, J=12.8 Hz, 1H), 4.17 (d, J=5.8 Hz, 2H), 4.07 (d, J=13.1 Hz, 1H), 3.83 (s, 3H), 3.07 (t, J=12.0 Hz, 1H), 2.68 (t, J=12.7 Hz, 1H), 2.48-2.50 (m, 1H), 1.75 (d, J=10.8 Hz, 2H), 1.47-1.51 (m, 2H). ESI-MS m/z: 417.05 [M+H]⁺.

Example 2

(494) ##STR00507##

Methyl 3-(5-bromo-4-methyl-2-oxopyridin-1(2H)-yl)propanoate

(495) To a solution of 5-bromo-4-methylpyridin-2(1H)-one (1.0 g, 5.3 mmol) in DMF (10 mL), K₂CO₃ (1.5 g, 10.6 mmol) was added, followed by addition of methyl acrylate (0.91 g, 10.6 mmol) at 80° C. The mixture was stirred at 80° C. for 1 h. The mixture was allowed to cool to room temperature and then partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo. The residue was used directly in the next step without further purification.

3-(5-Bromo-4-methyl-2-oxopyridin-1(2H)-yl)propanoic acid

(496) To a solution of the crude product from above step in MeOH (20 mL), aqueous KOH (2M, 5 mL) was added and the resulting mixture was stirred at room temperature overnight. The mixture was acidified with con. HCl and then extracted with ethyl acetate. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo. The crude product was used directly in the next step without further purification.

1-(3-(4-Acryloylpiperazin-1-yl)-3-oxopropyl)-5-bromo-4-methylpyridin-2(1H)-one
(497) A mixture of 3-(5-Bromo-4-methyl-2-oxopyridin-1(2H)-yl)propanoic acid (112 mg, 0.43 mmol), 1-(piperazin-1-yl)prop-2-en-1-one (80 mg, 0.43 mmol), HATU (163 mg, 0.43 mmol) and Et.sub.3N (0.2 mL, 1.43 mmol) in DMF (5 mL) was stirred at room temperature for 2 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified via Isolera One (silica cartridge, 0-6% MeOH/DCM) to afford the desired product (22.8 mg, 14 yield).
.sup.1H NMR (300 MHz, CDCl.sub.3) δ : 7.73 (s, 1H), 6.55 (dd, J=10.5, 16.7 Hz, 1H), 6.46 (s, 1H), 6.32 (dd, J=1.8, 16.8 Hz, 1H), 5.75 (dd, J=1.8, 10.5 Hz, 1H), 4.20 (t, J=6.0 Hz, 2H), 3.4-3.6 (m, 8H), 2.86 (t, J=6.0 Hz, 2H), 2.22 (s, 3H). ESI-MS m/z: 382.05 [M+H].sup.+.

Example 3

(498) ##STR00508## ##STR00509##

tert-Butyl 5-(vinylsulfonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

(499) To a stirred mixture of tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (1.0 g, 4.71 mmol) in DCM (20 mL) at 0° C., Et.sub.3N (1.43 g, 14.1 mmol) was added. The mixture was stirred at 0° C. for 5 min, and then a solution of 2-chloroethanesulfonyl chloride (0.77 g, 4.71 mmol) in DCM (5 mL) was added dropwise. The resulting mixture was stirred at room temperature for 0.5 h. The reaction mixture was poured into water (20 mL) and extracted with DCM (30 mL×3). The combined organic layer was washed with brine (15 mL×3), dried over anhydrous Na.sub.2SO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (10-50% ethyl acetate/petroleum ether) to afford the desired product (0.5 g, 35% yield) as a solid.

2-(Vinylsulfonyl)octahydropyrrolo[3,4-c]pyrrole

(500) A mixture of tert-butyl 5-(vinylsulfonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (150 mg, 0.50 mmol) in HCl/MeOH (20 mL, 2.86 M) was stirred at room temperature for 1 h. The mixture was concentrated in vacuo to afford the crude product (110 mg) as a solid which was used directly in next step without further purification.

4,5-Dichloro-2-nitrophenol

(501) To a solution of 3,4-dichlorophenol (30 g, 185 mmol) in DCM (300 mL) at -15° C., sulfuric acid (24 g, 278 mmol) was added. To this mixture, nitric acid (19 g, 194 mmol) was added dropwise (over 20 min) and the temperature was controlled between -15° C. to -5 OC. The resulting mixture was stirred at 0° C. for 1 h. The mixture was poured into ice and extracted with ethyl acetate. The combined organic layer was washed with water, saturate NaHCO.sub.3 aqueous solution and brine, dried over Na.sub.2SO.sub.4 and then concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=100:1) to afford the desired product (16 g, 42% yield) as a solid.

2-Amino-4,5-dichlorophenol

(502) To a stirred solution of 4,5-dichloro-2-nitrophenol (15 g, 72 mmol) in acetic acid (150 mL) and water (450 mL), iron powder (16 g, 285 mmol) was added in portions. The resulting mixture was stirred at 50° C. for 2 h. The mixture was allowed to cool to room temperature, filtered, and the cake was rinsed with ethyl acetate. The filtrate was extracted with ethyl acetate. The organic layer was washed with water, NaHCO.sub.3 (aq.) and brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo to afford the desired product (10 g, 78%).

Ethyl 2-((4,5-dichloro-2-hydroxyphenyl)amino)acetate

(503) To a solution of 2-amino-4,5-dichlorophenol (2.0 g, 11.3 mmol) and ethyl-2-oxoacetate (2.26 g, 12.42 mmol) in DCM (50 mL) at room temperature, AcOH (1 mL) was added and the resulting mixture was stirred for 1 h. To this mixture, NaBH(OAc).sub.3 (7.2 g, 33.9 mmol) was added and then stirred for 16 h. The mixture was concentrated in vacuo and the residue was suspended in DCM. The mixture was filtered through a pad of Celite and the filtrate was washed with brine. The organic layer was dried over anhydrous Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The

residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=10:1) to afford the desired product (2.4 g, 81% yield) as a solid.

2-((4,5-Dichloro-2-hydroxyphenyl)amino)acetic acid

(504) To a solution of ethyl 2-((4,5-dichloro-2-hydroxyphenyl)amino)acetate (2 g, 7.6 mmol) in of 4:1 mixture of tetrahydrofuran and water (30 mL) at room temperature, LiOH.H₂O (3.2 g, 76 mmol) were added and the resulting mixture was stirred for 30 min and then acidified with aqueous HCl (1 N) to adjust the pH to 3-5. The mixture was extracted with ethyl acetate (40 mL×3). The combined organic layer was washed with brine (40 mL×3), dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo to afford the crude product (2 g) which was used directly in the next step without further purification.

2-((4,5-dichloro-2-hydroxyphenyl)amino)-1-(5-(vinylsulfonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)ethanone

(505) To a stirred solution of 2-((4,5-dichloro-2-hydroxyphenyl)amino)acetic acid (90 mg, 0.385 mmol) in DMF (15 mL) at room temperature, 2-(vinylsulfonyl)octahydropyrrolo[3,4-c]pyrrole (110 mg, 0.46 mmol) was added followed by HOBt (78.05 mg, 0.58 mmol), EDCI.HCl (110.7 mg, 0.46 mmol) and Et₃N (116.7 mg, 1.2 mmol). The reaction mixture was stirred at room temperature for 1 h. The reaction mixture was partitioned between ethyl acetate and brine. The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (10-50% ethyl acetate/dichloroethane) to afford the desired product (20 mg, 10% yield) as a solid. ¹H NMR (400 MHz, DMSO-d₆) δ: 10.17 (s, 1H), 6.91 (dd, J=10.0, 16.4 Hz, 1H), 6.79 (s, 1H), 6.62 (s, 1H), 6.17-6.10 (m, 2H), 5.25 (t, J=4.0 Hz, 1H), 3.85-3.82 (m, 2H), 3.71-3.66 (m, 1H), 3.61-3.56 (m, 1H), 3.45-3.39 (m, 3H), 3.29-3.25 (m, 1H), 3.09-3.00 (m, 3H), 2.92-2.89 (m, 1H). ESI-MS m/z: 420.1 [M+H]⁺.

Example 4

(506) ##STR00510##

5-Chloro-2-nitrophenol

(507) To a solution of 2,4-dichloro-1-nitrobenzene (100 g, 0.52 mol) in DMSO (200 mL), aqueous solution of NaOH (41.6 g, 1.04 mol) in water (42 mL) was added and the resulting mixture was stirred 60° C. for 16 h. The mixture was allowed to cool to room temperature, poured to ice water, and then acidified with aqueous HCl (1 M) to adjusted the pH to 3-4. The mixture was extracted with ethyl acetate. The combined organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo. The residue was used directly in the next step (80 g, 88% yield).

4-Chloro-2-methoxy-1-nitrobenzene

(508) To a solution of 5-chloro-2-nitrophenol (40 g, 0.23 mol) in DMF (200 mL), K₂CO₃ (47.6 g, 0.345 mol) and iodomethane (49 g, 0.345 mol) were added and the resulting mixture was stirred at room temperature for 16 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over MgSO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether) to afford the desired product (30 g, 70% yield).

1-Chloro-2-iodo-5-methoxy-4-nitrobenzene

(509) To a solution of H₂SO₄ (600 mL, 90%), trifluoromethanesulfonic anhydride (11.3 g, 0.04 mol) and NIS (49.68 g, 0.22 mol) were added and resulting mixture was stirred at room temperature for 1 h. To this mixture, 4-chloro-2-methoxy-1-nitrobenzene (69 g, 0.368 mol) was added quickly. The mixture was stirred for 1 h, and then NIS (33.12 g, 0.148 mol) was slowly added to the mixture. The mixture was stirred at room temperature for 1 h and then was poured into ice-water. The precipitate collected by filtration, rinsed with water, aqueous Na₂SO₃ and NaHCO₃ solutions, and then dried in vacuo to afford the desired product (113 g, 98% yield).

4-Chloro-5-iodo-2-methoxybenzenamine

(510) To a solution of 1-chloro-2-iodo-5-methoxy-4-nitrobenzene (113 g, 0.361 mol) in acetic acid

(1 L) and water (50 mL) at 50° C., Fe (50.5 g, 0.903 mol) was added and the resulting mixture was stirred at 50° C. for 2 h. The mixture was allowed to cool to room temperature and then poured into ice-water. The precipitate was collected by filtration and rinsed with water. This crude product was dissolved with ethyl acetate (1 L) and filtered. The filtrate was washed with saturated NaHCO₃ solution and brine. The organic layer was dried over MgSO₄, filtered, and concentrated in vacuo to afford the desired product (87 g, 85% yield).

Ethyl 2-(4-chloro-5-iodo-2-methoxyphenylamino)acetate

(511) To a solution of 4-chloro-5-iodo-2-methoxybenzenamine (6 g, 21.2 mmol) in MeOH (50 mL) at room temperature, AcOH (3 drops) and ethyl glyoxalate (5.6 g, 27.5 mmol, 50% in toluene) were added. The mixture was stirred at room temperature for 2 h and then sodium cyanoborohydride (5.32 g, 84.8 mmol) was added to the mixture. The resulting mixture was stirred at 50° C. for 16 h. The mixture was allowed to cool to room temperature, and partitioned between ethyl acetate and water. The organic layer was dried over MgSO₄, filtered, and concentrated in vacuo to afford the crude product (8.6 g).

2-(4-Chloro-5-iodo-2-methoxyphenylamino)acetic acid

(512) To a solution of ethyl 2-(4-chloro-5-iodo-2-methoxyphenylamino)acetate (8.6 g, 22.9 mmol) in THF (50 mL) and water (50 mL), LiOH.H₂O (1.96 g, 45.9 mmol) was added and the resulting mixture was stirred at room temperature for 2 h. The mixture was washed with 20% ethyl acetate/petroleum ether. The aqueous layer was acidified with aqueous HCl (1 M) to adjust PH to 3-4 and extracted with ethyl acetate. The organic layer was dried over MgSO₄, filtered, and concentrated in vacuo to afford the desired product (5 g, 64% yield).

tert-Butyl 4-(2-(4-chloro-5-iodo-2-methoxyphenylamino)acetyl)piperazine-1-carboxylate

(513) To a solution of 2-(4-chloro-5-iodo-2-methoxyphenylamino)acetic acid (280 mg, 0.83 mmol) and tert-butyl piperazine-1-carboxylate (185 mg, 0.99 mmol) in DMF (10 mL) at room temperature, BOP (550 mg, 1.25 mmol) and DIEA (321 mg, 2.49 mmol) were added and the resulting mixture was stirred at room temperature for 1 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed brine, dried over MgSO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=3:1) to afford the desired product (350 mg, 68.7% yield). ESI-MS m/z: 510.3 [M+1]⁺.

tert-Butyl-4-(2-(5-(2-Chlorophenyl)-4-chloro-2-methoxyphenylamino)acetyl)piperazine-1-carboxylate

(514) To a solution of tert-butyl 4-(2-(4-chloro-5-iodo-2-methoxyphenylamino)acetyl)piperazine-1-carboxylate (200 mg, 0.4 mmol) and 2-chlorophenylboronic acid (69 mg, 0.44 mmol) in 1,4-dioxane (10 mL) and water (2 mL), Pd(PPh₃)₃ (40 mg, 0.035 mmol) and Na₂CO₃ (212 mg, 2 mmol) were added. The mixture was stirred at 80° C. for 16 h. The mixture was allowed to cool to room temperature and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=4:1) to afford the desired product (150 mg, 76% yield).

2-(5-(2-Chlorophenyl)-4-chloro-2-methoxyphenylamino)-1-(piperazin-1-yl)ethanone

(515) To a solution of tert-butyl-4-(2-(5-(2-chlorophenyl)-4-chloro-2-methoxyphenylamino)acetyl)piperazine-1-carboxylate (150 mg, 0.304 mmol) in DCM (2 mL), a solution of HCl in MeOH (10 mL, 29 mmol) was added. The mixture was stirred at room temperature for 1 h and then concentrated in vacuo to afford the crude product which was used in the next step without further purification.

1-(4-(2-(4-chloro-5-(2-Chlorophenyl)-2-methoxyphenylamino)acetyl)piperazin-1-yl)prop-2-en-1-one

(516) To a solution of the crude 2-(5-(2-Chlorophenyl)-4-chloro-2-methoxyphenylamino)-1-(piperazin-1-yl)ethanone (0.304 mmol) and Et₃N in DCM (5 mL) at 0° C., acryloyl chloride (27.5 mg, 0.304 mmol) was slowly added and the resulting mixture was stirred at room temperature

for 1 h. The mixture was quenched with saturated NaHCO₃ solution and extracted with DCM. The organic layer was washed with brine, dried over MgSO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel to afford the desired product (80 mg, 58.7% yield, 2 steps). ¹H NMR (400 MHz, DMSO-d₆) δ: 7.55-7.52 (m, 1H), 7.45-7.39 (m, 2H), 7.32-7.29 (m, 1H), 6.97 (s, 1H), 6.80 (dd, J=10.4, 16.4 Hz, 1H), 6.54 (s, 1H), 6.12 (dd, J=2.0, 16.8 Hz, 1H), 5.70 (dd, J=2.0, 10.4 Hz, 1H), 5.35 (bs., 1H), 3.93 (d, J=4.0 Hz, 2H), 3.89 (s, 1H), 3.55-3.49 (m, 8H). ESI-MS m/z: 448.2 [M+H]⁺.

Example 5

(517) ##STR00511##

1-(4-(2-(4-chloro-2-hydroxy-5-(2-Chlorophenyl)phenylamino)acetyl)piperazin-1-yl)prop-2-en-1-one

(518) To a solution of 1-(4-(2-(4-chloro-5-(2-Chlorophenyl)-2-methoxyphenylamino)acetyl)piperazin-1-yl)prop-2-en-1-one (70 mg, 0.147 mmol) in DCM (15 mL) -78° C. under argon, BBr₃ (187 mg, 0.754 mmol) was added. The mixture was stirred at room temperature for 1 h. The mixture was poured into ice water and extracted ethyl acetate. The organic layer was dried over MgSO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=40:1) to afford the desired product (15 mg, 24% yield). ¹H NMR (400 MHz, DMSO-d₆) δ: 10.08 (s, 1H), 7.53-7.51 (m, 1H), 7.41-7.37 (m, 2H), 7.30-7.28 (m, 1H), 6.84-6.77 (m, 2H), 6.48 (s, 1H), 6.12 (dd, J=2.4, 16.4 Hz, 1H), 5.70 (dd, J=2.0, 10.4 Hz, 1H), 5.24 (t, J=4.0 Hz, 1H), 3.90 (d, J=4.4 Hz, 2H), 3.56-3.49 (m, 8H). ESI-MS m/z: 434.2 [M+H]⁺.

Example 6

(519) ##STR00512##

4,5-Dichloro-2-vinylbenzenamine

(520) A mixture of 4,5-dichloro-2-iodobenzenamine (6 g, 20.8 mmol), tributyl(vinyl)stannane (6.6 g, 20.8 mmol), Pd(PPh₃)₄ (2.4 g, 2.1 mmol) in toluene (60 mL) was stirred at reflux under argon for 6 h. The mixture was allowed to cool to room temperature, quenched with aqueous KF solution and then extracted with ethyl acetate. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=4:1) to afford the desired product (3.0 g, 76.7% yield) as a colorless oil.

4,5-Dichloro-2-ethylbenzenamine

(521) A mixture of 4,5-dichloro-2-vinylbenzenamine (3.0 g, 15.95 mmol), Pd/C (3.3 g, 10%) in MeOH (20 mL) was stirred at room temperature under H₂ (1 atm) atmosphere for 4 h. The mixture was filtered and the filtrate was concentrated in vacuo to afford the desired product (2.4 g, 80% yield).

Ethyl 2-(4,5-dichloro-2-ethylphenylamino)acetate

(522) To a stirring solution of 4,5-dichloro-2-ethylbenzenamine (600 mg, 3.15 mmol) in DMF at 0° C., NaH (150 mg, 3.78 mmol) was added in portions. After stirring for 30 min, ethyl 2-bromoacetate (789 mg, 4.74 mmol) was added to the mixture. The resulting mixture was stirred at 120° C. for 16 h. The mixture was allowed to cool to room temperature, poured into ice-water and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=5:1) to afford the desired product (380 mg, 44.7% yield) as a yellow solid.

2-(4,5-Dichloro-2-ethylphenylamino)acetic acid

(523) A mixture of ethyl 2-(4,5-dichloro-2-ethylphenylamino)acetate (390 mg, 1.41 mmol) and LiOH·H₂O (592 mg, 14.1 mmol) in THF (8 mL) and water (2 mL) was stirred at room temperature for 2 h. The mixture was concentrated in vacuo and the residue was dissolved in H₂O and diluted with ethyl acetate. The mixture was acidified with aqueous HCl (10%) to

adjust the pH to 3-4 and then extracted with ethyl acetate. The organic layer was dried over Na.sub.2SO.sub.4, filtered and concentrated in vacuo to afford the desired product (300 mg, 85.75% yield) as a yellow solid. ESI-MS m/z: 246.2 [M-H].sup.-.

tert-Butyl 4-(2-(4,5-dichloro-2-ethylphenylamino)acetyl)piperazine-1-carboxylate

(524) To a stirred mixture of 2-(4,5-dichloro-2-ethylphenylamino)acetic acid (100 mg, 0.403 mmol), Et.sub.3N (122 mg, 1.21 mmol) in DCM (3 mL) at 0° C., EDCI.HCl (123 mg, 0.604 mmol) and HOBt (82 mg, 0.604 mmol) were added. The resulting mixture was stirred at 0° C. for 30 min, and then tert-butyl piperazine-1-carboxylate (90 mg, 0.483 mmol) was added. The mixture was stirred at room temperature for 16 h and partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=5:1) to afford the desired product (120 mg, 71.5% yield) as an off-white solid. ESI-MS m/z: 414.3 [M-H].sup.-.

2-(4,5-Dichloro-2-ethylphenylamino)-1-(piperazin-1-yl)ethanone hydrochloride

(525) A mixture of tert-butyl 4-(2-(4,5-dichloro-2-ethylphenylamino)acetyl)piperazine-1-carboxylate (120 mg, 0.288 mmol) in HCl/MeOH (2.86 M, 10 mL) was stirred at room temperature for 1 h. The mixture was concentrated in vacuo to afford the crude product (82 mg) as a yellow solid which was used directly in the next step without further purification.

1-(4-(2-(4,5-Dichloro-2-ethylphenylamino)acetyl)piperazin-1-yl)prop-2-en-1-one

(526) To a mixture of above crude 2-(4,5-dichloro-2-ethylphenylamino)-1-(piperazin-1-yl)ethanone hydrochloride (82 mg) and Et.sub.3N (69 mg, 0.686 mmol) in DCM (3 mL) 0° C., a solution of acryloyl chloride (23 mg, 0.251 mmol) in DCM (1 mL) was added. The resulting mixture was stirred at room temperature for 1 h and then was quenched with saturated NaHCO.sub.3 solution. The mixture was diluted with ethyl acetate, washed with saturated NaHCO.sub.3 solution and brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (DCM/MeOH=30:1) to afford the desired product (20 mg, 18.76% yield, 2 steps) as an off-white solid. .sup.1H NMR (400 MHz, CDCl.sub.3) δ: 7.11 (s, 1H), 6.58 (dd, J=10.4, 16.4 Hz, 1H), 6.54 (s, 1H), 6.36 (dd, J=1.6, 16.4 Hz, 1H), 5.79 (dd, J=1.6, 10.4 Hz, 1H), 5.08 (bs., 1H), 3.89-3.52 (m, 10H), 2.52 (q, J=7.2 Hz, 2H), 1.29-1.26 (t, J=7.6 Hz, 3H). ESI-MS m/z: 370.2 [M+H].sup.+.

Example 7

(527) ##STR00513##

2-(4-chloropyridin-2-yloxy)-4,5-dichlorobenzeneamine

(528) To a solution of 2-amino-4,5-dichlorophenol (1.5 g, 8.47 mmol) in THF (20 mL) at 0° C., NaH (60% dispersed in oil, 373 mg, 9.33 mmol) was added and the resulting mixture was stirred for 20 min. To this mixture, 4-chloro-2-fluoropyridine (1.67 g, 12.7 mmol) was added and the resulting mixture was stirred at reflux under argon for 15 h. The mixture was allowed to cool to room temperature and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=20:1 to 10:1) to afford the desired product (1.1 g, 45% yield). ESI-MS m/z: 287.0 [M-H].sup.-.

2-(4,5-Dichloro-2-iodophenoxy)-4-chloropyridine

(529) The mixture of 2-(4-chloropyridin-2-yloxy)-4,5-dichlorobenzeneamine (1.0 g, 3.47 mmol) in concentrated HCl (5 mL) and H.sub.2O (15 mL) was cooled to -10° C.-0 OC, NaNO.sub.2 (0.359 g, 5.21 mmol) was added and the resulting mixture was stirred at this temperature for 50 min. This mixture was added dropwise to the mixture of KI (10 g, 60.2 mmol), CuI (330 mg, 1.73 mmol) in H.sub.2O (40 mL) and then stirred at room temperature for 15 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=30:1) to afford the desired product (900 mg, 65% yield). ESI-MS m/z: 399.9 [M+H].sup.+.

2-(2-(4,5-Dichloro-2-iodophenoxy)pyridin-4-yloxy)ethanol

(530) A mixture of 2-(4,5-dichloro-2-iodophenoxy)-4-chloropyridine (0.45 g, 1.128 mmol), ethane-1,2-diol (4 mL, 71.89 mmol), K₂CO₃ (0.45 mg, 3.26 mmol) in acetone (6 mL) was stirred at 120° C. for 5 h. The mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=20:1 to 4:1) to afford the desired product (160 mg, 33% yield). ESI-MS m/z: 426.0 [M+H]⁺.

2-(2-(2-(4,5-Dichloro-2-iodophenoxy)pyridin-4-yloxy)ethyl)isoindoline-1,3-dione

(531) A mixture of 2-(2-(4,5-dichloro-2-iodophenoxy)pyridin-4-yloxy)ethanol (130 mg, 0.31 mmol), isoindoline-1,3-dione (54 mg, 0.37 mmol), PPh₃ (160 mg, 0.61 mmol) in THF (5 mL) at 0° C., DIAD (123 mg, 0.61 mmol) was added. The resulting mixture was allowed to warm to room temperature and stirred for 15 h. The mixture was concentrated in vacuo and the residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=20:1 to 5:1) to afford the desired product (120 mg, 71% yield). ESI-MS m/z: 555.0 [M+H]⁺.

2-(2-(4,5-Dichloro-2-iodophenoxy)pyridin-4-yloxy)ethanamine

(532) The mixture of 2-(2-(2-(4,5-dichloro-2-iodophenoxy)pyridin-4-yloxy)ethyl)isoindoline-1,3-dione (120 mg, 0.22 mmol), N₂H₄·H₂O (106 mg, 1.80 mmol) and EtOH (5 mL) was stirred at reflux for 1 h. The mixture was allowed to cool to room temperature and then concentrated in vacuo. The residue was slurried in DCM (5 mL) and MeOH (1 mL). The precipitate was removed by filtration, and the filtrate was concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (DCM/MeOH=10:1) to afford the desired product (35 mg, 38% yield).

N-(2-(2-(4,5-Dichloro-2-iodophenoxy)pyridin-4-yloxy)ethyl)ethanesulfonamide

(533) To a mixture of 2-(2-(4,5-dichloro-2-iodophenoxy)pyridin-4-yloxy)ethanamine (30 mg, 0.07 mmol), Et₃N (35.7 mg, 0.35 mmol) in DCM (5 mL), 2-chloroethanesulfonyl chloride (11.5 mg, 0.07 mmol) was added and the resulting mixture was stirred at room temperature for 1 h. Then solvent was removed under reduced pressure. The residue was taken in THF (2 mL) and H₂O (2 mL), then K₂CO₃ (100 mg) was added and the resulting mixture was stirred at room temperature for 1 h. The mixture was partitioned between DCM and water. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by flash column chromatography on silica gel (DCM/ethyl acetate=5:1) to afford the desired product (56 mg, 56% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.98 (d, J=5.6 Hz, 1H), 7.92 (s, 1H), 7.25 (s, 1H), 6.61-6.57 (m, 2H), 6.47 (d, J=2.0 Hz, 1H), 6.32 (d, J=16.4 Hz, 1H), 6.00 (d, J=10.0 Hz, 1H), 4.77 (t, J=5.2 Hz, 1H), 4.18 (t, J=5.2 Hz, 2H), 3.51-3.47 (m, 2H). ESI-MS m/z: 515.0 [M+H]⁺.

Example 8

(534) ##STR00514##

tert-Butyl 3-(4-(2-(4,5-dichloro-2-hydroxyphenylamino)acetyl)piperazin-1-yl)azetidine-1-carboxylate

(535) A mixture of 2-(4,5-dichloro-2-hydroxyphenylamino)acetic acid (500 mg, 2.12 mmol), tert-butyl 3-(piperazin-1-yl)azetidine-1-carboxylate (565 mg, 2.34 mmol), EDCI·HCl (488 mg, 2.54 mmol), HOBt (343 mg, 2.54 mmol), Et₃N (428 mg, 4.24 mmol) in DMF (20 mL) was stirred at room temperature for 15 h. The mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous NaHCO₃ solution and brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (DCM/MeOH=30:1) to afford the desired product (300 mg, 31% yield). ESI-MS m/z: 457.4 [M-H]⁻.

2-(4,5-Dichloro-2-hydroxyphenylamino)-1-(4-(azetidin-3-yl)piperazin-1-yl)ethanone hydrochloride

(536) A mixture of tert-butyl 3-(4-(2-(4,5-dichloro-2-hydroxyphenylamino)acetyl)piperazin-1-yl)azetidine-1-carboxylate (150 mg, 0.33 mmol) in HCl-MeOH (20 mL, 57 mmol) was stirred at

room temperature for 1 h. The mixture was concentrated in vacuo to afford the crude product (130 mg) which was used directly in the next step without further purification.

1-(3-(4-(2-(4,5-Dichloro-2-hydroxyphenylamino)acetyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one

(537) 2-(4,5-dichloro-2-hydroxyphenylamino)-1-(4-(azetidin-3-yl)piperazin-1-yl)ethanone hydrochloride (120 mg, 0.30 mmol) was added to the mixture of Et.sub.3N (0.2 mL, 1.44 mmol) in DCM (10 mL) followed by addition of DMF (1 drop). The mixture was stirred for 5 min and then acryloyl chloride (27 mg, 0.30 mmol) was added. The resulting mixture was stirred at room temperature for 1 h, poured into water and then extracted with MeOH/DCM. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated. The residue was purified by flash column chromatography on silica gel (DCM/MeOH/NH.sub.3.H.sub.2O=50:1:0.1 to 20:1:0.2) to afford the desired product (30 mg, 24% yield). .sup.1H NMR (400 MHz, DMSO-d.sub.6) δ : 10.17 (s, 1H), 6.78 (s, 1H), 6.71 (s, 1H), 6.30 (dd, J=10.4, 17.2 Hz, 1H), 6.09 (dd, J=2.0, 17.2 Hz, 1H), 5.67 (dd, J=2.4, 10.4 Hz, 1H), 5.32 (t, J=4.4 Hz, 1H), 4.26-4.22 (m, 1H), 4.11-4.04 (m, 1H), 3.93-3.91 (m, 3H), 3.79-3.75 (m, 3H), 3.52-3.51 (m, 4H), 3.19-3.16 (m, 1H), 2.36-2.30 (m, 4H). ESI-MS m/z: 411.2 [M-H].sup.-.

Example 9

(538) ##STR00515##

tert-Butyl 3-(3-(acrylamido)azetidin-1-yl)azetidine-1-carboxylate

(539) To a mixture of N-(azetidin-3-yl)acrylamide hydrochloride (500 mg, 3.40 mmol), tert-butyl 3-oxoazetidine-1-carboxylate (684 mg, 4.0 mmol), Et.sub.3N (343 mg, 3.40 mmol) and AcOH (100 mg, 0.167 mmol) in DCM (20 mL), NaBH(OAc).sub.3 (2.16 g, 10.2 mmol) was added, and the resulting mixture was stirred at room temperature for 16 h. The mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with saturated NaHCO.sub.3 solution and brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (DCM/MeOH=100:1 to 20:1) to afford the desired product (300 mg, 31% yield).

N-(1-(Azetidin-3-yl)azetidin-3-yl)acrylamide hydrochloride

(540) A mixture of tert-butyl 3-(3-(acrylamido)azetidin-1-yl)azetidine-1-carboxylate (300 mg, 1.07 mmol) in HCl-MeOH (30 mL, 86 mmol) was stirred at room temperature for 1 h. The mixture was concentrated in vacuo to afford the crude product (250 mg) which was used directly in the next step without further purification.

N-(1-(1-(2-(4,5-Dichloro-2-hydroxyphenylamino)acetyl)azetidin-3-yl)azetidin-3-yl)acrylamide

(541) A mixture of 2-(4,5-dichloro-2-hydroxyphenylamino)acetic acid (120 mg, 0.51 mmol), EDCI.HCl (147 mg, 0.77 mmol), HOBT (83 mg, 0.61 mmol), Et.sub.3N (154 mg, 1.53 mmol) in DMF (20 mL) was stirred at room temperature for 5 min and then N-(1-(azetidin-3-yl)azetidin-3-yl)acrylamide hydrochloride (150 mg, 0.69 mmol) was added. The resulting mixture was stirred at room temperature for 15 h. The mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with saturated NaHCO.sub.3 solution and brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (DCM/MeOH/NH.sub.3.H.sub.2O=100:10:1.5) to afford the desired product (6 mg, 3% yield). .sup.1H NMR (400 MHz, DMSO-d₆) δ : 10.19 (s, 1H), 8.59 (d, J=8.0 Hz, 1H), 6.79 (s, 1H), 6.55 (s, 1H), 6.20 (dd, J=10.0, 16.8 Hz, 1H), 6.09 (dd, J=2.0, 16.8 Hz, 1H), 5.62 (dd, J=2.0, 9.6 Hz, 1H), 5.20 (t, J=4.0 Hz, 1H), 4.40-4.35 (m, 1H), 4.19-4.15 (m, 1H), 3.96-3.88 (m, 2H), 3.73-3.69 (m, 3H), 3.53-3.45 (m, 3H), 3.00-2.96 (m, 2H). ESI-MS m/z: 399.2 [M+H].sup.+.

Example 10

(542) ##STR00516##

2,6-Diaza-spiro[3.4]octane-6-acryloyl-2-carboxylic acid tert-butyl ester

(543) To a mixture of 2,6-diaza-spiro[3.4]octane-2-carboxylic acid tert-butyl ester (80 mg, 0.38

mmol), Et.sub.3N (0.2 mL, 1.44 mmol) in DCM (20 mL), acryloyl chloride (34 mg, 0.38 mmol) was added and the resulting mixture was stirred at room temperature for 1 h. The mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with saturated NaHCO.sub.3 solution and brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (DCM/MeOH=40:1) to afford the desired product (50 mg, 50% yield). ESI-MS m/z: 289.2 [M+Na].sup.+.

1-(2,6-Diazaspiro[3.4]octan-6-yl)prop-2-en-1-one

(544) A mixture of 2,6-diaza-spiro[3.4]octane-6-acryloyl-2-carboxylic acid tert-butyl ester (50 mg, 0.19 mmol) in HCl/MeOH (10 mL, 29 mmol) was stirred at room temperature for 1 h. The mixture was concentrated in vacuo to afford the crude product (40 mg) which was used directly in the next step without further purification.

1-(2-((4,5-Dichloro-2-hydroxyphenyl)glycyl)-2,6-diazaspiro[3.4]octan-6-yl)prop-2-en-1-one

(545) The mixture of 2-(4,5-dichloro-2-hydroxyphenylamino)acetic acid (47 mg, 0.2 mmol), 1-(2,6-diazaspiro[3.4]octan-6-yl)prop-2-en-1-one (40 mg, 0.2 mmol), EDCI.HCl (46 mg, 0.24 mmol), HOBt (32 mg, 0.24 mmol) and Et.sub.3N (0.61 mg, 0.6 mmol) in DMF (10 mL) was stirred at room temperature for 2 h. The mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with saturated NaHCO.sub.3 solution and brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (DCM/MeOH=30:1) to afford the desired product (13 mg, 17% yield). .sup.1H NMR (400 MHz, DMSO-d₆) δ: 10.17 (s, 1H), 6.78 (s, 1H), 6.60-6.50 (m, 2H), 6.13 (dt, J=2.4, 16.4 Hz, 1H), 5.67 (dd, J=2.4, 10.4 Hz, 1H), 5.19 (dd, J=5.2, 10.0 Hz, 1H), 4.16-4.07 (m, 2H), 3.90-3.83 (m, 2H), 3.75-3.72 (m, 3H), 3.61-3.52 (m, 2H), 3.42-3.39 (m, 1H), 2.16-2.13 (m, 1H), 2.06-2.03 (m, 1H). ESI-MS m/z: 382.3 [M-H].sup.-.

Example 11

(546) ##STR00517##

3-(4,5-Dichloro-2-iodophenyl)phenol

(547) A mixture of 1,2-dichloro-4,5-diiodobenzene (1.5 g, 3.76 mmol), 3-hydroxyphenylboronic acid (0.52 g, 3.76 mmol), Na.sub.2CO.sub.3 (1.99 g, 18.8 mmol) and Pd(PPh.sub.3).sub.4 (0.35 g, 0.30 mmol), in 1,4-dioxane (10 mL) and water (2 mL) was stirred at reflux under argon for 16 h. The mixture was allowed to cool to room temperature and then concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=20:1) to afford the desired product (300 mg, 22% yield). ESI-MS m/z: 362.9 [M-H].sup.-.

2-[3-(4,5-Dichloro-2-iodo)phenoxy]ethanol

(548) A mixture of 3-(4,5-dichloro-2-iodophenyl)phenol (0.3 g, 0.82 mmol), 2-bromoethanol (0.20 g, 1.65 mmol), K.sub.2CO.sub.3 (0.57 g, 4.12 mmol) in DMF (5 mL) was stirred at 100° C. under nitrogen for 16 h. The mixture was allowed to cool to room temperature and partitioned between ethyl acetate and brine. The organic layer was dried over Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=4:1) to afford the desired product (150 mg, 45% yield).

N-(2-((4',5'-Dichloro-2'-iodo-[1,1'-biphenyl]-3-yl)oxy)ethyl)ethanesulfonamide

(549) The title compound was prepared from 2-[3-(4,5-dichloro-2-iodo)phenoxy]ethanol in three steps followed the procedure described in Example 28. .sup.1H NMR (400 MHz, CDCl.sub.3) δ: 8.01 (s, 1H), 7.37-7.33 (m, 2H), 6.94-6.91 (m, 2H), 6.81-6.80 (m, 1H), 6.57 (dd, J=10.0, 16.8 Hz, 1H), 6.30 (d, J=16.4 Hz, 1H), 5.97 (d, J=10.0 Hz, 1H), 4.77 (t, J=6.0 Hz, 1H), 4.14 (t, J=5.2 Hz, 2H), 3.49-3.45 (m, 2H). ESI-MS m/z: 496.0 [M-H].sup.-.

Example 12

(550) ##STR00518##

5-Bromo-4-chloro-2-iodobenzenamine

(551) To a solution of 3-bromo-4-chlorobenzenamine (10.0 g, 48.5 mmol) in CH.sub.3COOH (50 mL), NIS (10.9 g, 48.5 mmol) was added in portions and the resulting mixture was stirred at room

temperature for 16 h. The mixture was concentrated in vacuo. The residue was dissolved in ethyl acetate, washed with saturated NaHCO₃ solution and brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=50:1) to afford the desired product (2.5 g, 15.5% yield). ESI-MS m/z: 329.9 [M-H]⁺.

5-Bromo-4-chloro-2-vinylbenzenamine

(552) A mixture of 5-bromo-4-chloro-2-iodobenzenamine (2.5 g, 7.51 mmol), tributyl(vinyl)stannane (2.4 g, 7.51 mmol), Pd(PPh₃)₄ (867 mg, 0.75 mmol) in toluene (25 mL) was stirred at reflux under argon for 16 h. The reaction mixture was allowed to cool to room temperature and quenched with aqueous KF solution. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=20:1) to afford the desired product (1.0 g, 57.4% yield). ESI-MS m/z: 232.2 [M+H]⁺.

2',6-Dichloro-4-vinyl-[1,1'-biphenyl]-3-amine

(553) A mixture of 5-bromo-4-chloro-2-vinylbenzenamine (600 mg, 2.6 mmol), 2-chlorophenylboronic acid (1.25 g, 12.9 mmol), Pd(PPh₃)₄ (300 mg, 0.26 mmol), Na₂CO₃ (1.4 g, 13.0 mmol) in 1,4-dioxane (20 mL) and water (5 mL) was stirred at reflux under argon for 16 h. The mixture was allowed to cool to room temperature and then concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=30:1) to afford the desired product (500 mg, 73.1% yield). ESI-MS m/z: 263.1 [M+H]⁺.

2',6-Dichloro-4-ethyl-[1,1'-biphenyl]-3-amine

(554) A mixture of 2',6-dichloro-4-vinyl-[1,1'-biphenyl]-3-amine (500 mg, 1.9 mmol), CuCl (225 mg, 2.28 mmol) in MeOH (10 mL) at 0° C., NaBH₄ (722 mg, 19 mmol) was added in portions and the resulting mixture was stirred at room temperature for 20 min. The reaction was quenched with water and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=30:1) to afford the desired product (300 mg, 59.4% yield). ESI-MS m/z: 266.0 [M+H]⁺.

1-(4-(2-((2',6-Dichloro-4-ethyl-[1,1'-biphenyl]-3-yl)amino)acetyl)piperazin-1-yl)prop-2-en-1-one

(555) The title compound was prepared from 2',6-dichloro-4-ethyl-[1,1'-biphenyl]-3-amine in three steps followed the procedure described in Example 20 and the following procedure for the amide formation. 2-((2',6-dichloro-4-ethyl-[1,1'-biphenyl]-3-yl)amino)acetic acid (150 mg, 0.466 mmol) and Et₃N (235 mg, 2.33 mmol) in DMF (5 mL) at 0° C., EDCI.HCl (178 mg, 0.932 mmol) and HOBT (126 mg, 0.932 mmol) were added and the resulting mixture was stirred at 0° C. for 30 min. To this mixture, tert-butyl piperazine-1-carboxylate (123 mg, 0.698 mmol) was added and then stirred at room temperature for 16 h. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (DCM/MeOH=50:1) to afford the desired product (45 mg, 21.7% yield). ¹H NMR (400 MHz, DMSO-d₆) δ: 7.56-7.54 (m, 1H), 7.43-7.41 (m, 2H), 7.32-7.30 (m, 1H), 7.14 (s, 1H), 6.81 (dd, J=10.4, 16.4 Hz, 1H), 6.54 (s, 1H), 6.13 (dd, J=2.4, 16.8 Hz, 1H), 5.71 (dd, J=2.0, 10.4 Hz, 1H), 5.25 (bs., 1H), 3.96 (d, J=4.4 Hz, 2H), 3.56-3.52 (m, 8H), 2.54 (q, J=7.2 Hz, 2H), 1.23 (t, J=7.2 Hz, 3H). ESI-MS m/z: 444.3 [M-H]⁺.

Example 13

(556) ##STR00519##

2-Amino-5-bromo-3-methoxybenzoic acid

(557) To a solution of 2-amino-3-methoxybenzoic acid (5 g, 29.9 mmol) in MeOH (35 mL) at -5° C., NBS (5.59 g, 31.4 mmol) was added and the resulting mixture was stirred at 0° C. for 16 h. The

mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over MgSO_4 , filtered and concentrated in vacuo to afford the crude product (4 g, 54% yield). ESI-MS m/z : 244.2 $[\text{M}-\text{H}]^{\text{sup.}-}$.

3-Bromo-5-methoxybenzoic acid

(558) To a solution of 2-amino-5-bromo-3-methoxybenzoic acid (4 g, 16.3 mmol) in water (20 mL) at 0°C ., conc. HCl (7.5 mL, 90 mmol) and THF (20 mL) were added. The mixture was stirred for 30 min, and then NaNO_2 (3.16 g, 45.8 mmol) was added. The resulting mixture was stirred for 2 h and then hypophosphorous acid (5.1 g, 76 mmol, 50% in H_2O) was added to the reaction. The mixture was stirred at room temperature for 16 h. The precipitate was collected by filtration, washed with water and dried in vacuo to afford the desired product (3.2 g, 85% yield). ESI-MS m/z : 229.2 $[\text{M}-\text{H}]^{\text{sup.}-}$.

3-(2-Chlorophenyl)-5-methoxybenzoic acid

(559) To a solution of 3-bromo-5-methoxybenzoic acid (1 g, 4.06 mmol) and 2-chlorophenylboronic acid (1.27 g, 8.13 mmol) in 1,4-dioxane (10 mL) and water (2 mL), $\text{Pd}(\text{PPh}_3)_4$ (468 mg, 0.40 mmol) and Na_2CO_3 (2.15 g, 20.3 mmol) were added and the resulting mixture was stirred at 80°C . for 16 h. The mixture was allowed to cool to room temperature and acidified with aqueous HCl (1.0 M) to adjust the pH to 3-4. The mixture was extracted with ethyl acetate. The organic layer was dried over MgSO_4 , filtered and concentrated in vacuo to afford the desired product (800 mg, 75% yield) without further purification. ESI-MS m/z : 361.2 $[\text{M}-\text{H}]^{\text{sup.}-}$.

1-(4-(2'-Chloro-5-methoxy-[1,1'-biphenyl]-3-carbonyl)piperazin-1-yl)prop-2-en-1-one

(560) To a solution of tert-butyl 4-acryloylpiperazine-1-carboxylate (260 mg, 1.07 mmol) in DCM (2 mL), a solution of HCl in MeOH (10 mL, 28.6 mmol) was added and the resulting mixture was stirred at room temperature for 1 h. The mixture was concentrated in vacuo. The residue was added to the solution of 3-(2-chlorophenyl)-5-methoxybenzoic acid (280 mg, 1.07 mmol), HOBt (290 mg, 2.17 mmol), $\text{EDCI}\cdot\text{HCl}$ (410 mg, 2.17 mmol) and Et_3N (324 mg, 3.21 mmol) in DMF (10 mL). The resulting mixture was stirred at room temperature for 16 h and partitioned between DCM and saturated NaHCO_3 solution. The organic layer was dried over MgSO_4 , filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (200 mg, 52% yield). ESI-MS m/z : 385.2 $[\text{M}+\text{H}]^{\text{sup.}+}$.

1-(4-(2'-Chloro-5-hydroxy-[1,1'-biphenyl]-3-carbonyl)piperazin-1-yl)prop-2-en-1-one (VI-25)

(561) To a solution of 1-(4-(2'-chloro-5-methoxy-[1,1'-biphenyl]-3-carbonyl)piperazin-1-yl)prop-2-en-1-one (100 mg, 0.26 mmol) in DCM (15 mL) at -78°C ., BBr_3 (650 mg, 2.6 mmol) was added and the resulting mixture was stirred at room temperature for 1 h. The mixture was poured into ice-water, basified with sat NaHCO_3 aqueous solution to adjust the pH to 7-8 and extracted with ethyl acetate. The organic layer was dried over MgSO_4 , filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (25 mg, 26% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.96 (s, 1H), 7.58-7.56 (m, 1H), 7.42-7.40 (m, 3H), 6.89-6.75 (m, 4H), 6.14 (dd, $J=2.0, 16.8$ Hz, 1H), 5.71 (dd, $J=2.0, 10.0$ Hz, 1H), 3.68-3.44 (m, 8H). ESI-MS m/z : 371.2 $[\text{M}+\text{H}]^{\text{sup.}+}$.

Example 14

(562) ##STR00520##

tert-Butyl 4-(2-acrylamidoethyl)piperidine-1-carboxylate

(563) To a stirred mixture of tert-butyl 4-(2-aminoethyl)piperidine-1-carboxylate (1.0 g, 4.38 mmol) in DCM (20 mL) at 0°C ., Et_3N (1.33 g, 13.14 mmol) was added and the resulting mixture was stirred at 0°C . for 5 min. To this mixture, a solution of acryloyl chloride (0.39 g, 4.38 mmol) in DCM (5 mL) was added dropwise. The resulting mixture was stirred at room temperature for 30 min. The reaction mixture was poured into water (20 mL) and extracted with DCM (30

mL×3). The combined organic layer was washed with brine (15 mL×3), dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (10-50% ethyl acetate/dichloroethane) to afford the desired product (0.6 g, 50% yield) as a solid.

N-(2-(Piperidin-4-yl)ethyl)acrylamide

(564) A mixture of tert-butyl 4-(2-acrylamidoethyl)piperidine-1-carboxylate (600 mg, 2.13 mmol) in HCl/MeOH (60 mL, 2.86 M) was stirred at room temperature for 1 h. The mixture was concentrated in vacuo to yield the crude product (550 mg) as a solid which was used directly in the next step without further purification.

N-(2-(1-(2-((4,5-Dichloro-2-hydroxyphenyl)amino)acetyl)piperidin-4-yl)ethyl)acrylamide

(565) To a stirred solution of 2-((4,5-dichloro-2-hydroxyphenyl)amino)acetic acid (200 mg, 0.85 mmol) in DMF (30 mL) at room temperature, N-(2-(piperidin-4-yl)ethyl)acrylamide (222.8 mg, 1.02 mmol) was added followed by HOBt (172.9 mg, 1.28 mmol), EDCI.HCl (244.7 mg, 1.28 mmol) and Et₃N (257.9 mg, 2.55 mmol). The reaction mixture was stirred at room temperature for 1 h and then partitioned between ethyl acetate and brine. The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (1-3% methanol/dichloroethane) to afford the desired product (120 mg, 29% yield) as a solid. ¹H NMR (400 MHz, DMSO-d₆) δ: 10.15 (s, 1H), 8.07 (t, J=5.2 Hz, 1H), 6.78 (s, 1H), 6.72 (s, 1H), 6.18 (dd, J=10.0, 17.2 Hz, 1H), 6.06 (dd, J=2.4, 17.2 Hz, 1H), 5.56 (dd, J=2.4, 10.4 Hz, 1H), 5.34 (t, J=4.0 Hz, 1H), 4.36 (d, J=12.8 Hz, 1H), 3.94-3.87 (m, 3H), 3.20-3.15 (m, 2H), 2.98-2.92 (m, 1H), 2.63-2.57 (m, 1H), 1.74-1.69 (m, 2H), 1.57-1.52 (m, 1H), 1.41-1.35 (m, 2H), 1.12-0.95 (m, 2H). ESI-MS m/z: 400.4[M+H]⁺.

Example 15

(566) ##STR00521##

2',5',6-Trichloro-4-methoxy-[1,1'-biphenyl]-3-amine

(567) A mixture of 4-chloro-5-iodo-2-methoxybenzenamine (4.1 g, 14.5 mmol), 2,5-dichlorophenylboronic acid (3.3 g, 17.4 mmol), Pd(PPh₃)₄ (500 mg, 1.45 mmol) and Na₂CO₃ (4.7 g, 43.5 mmol) in 1,4-dioxane (150 mL) and water (15 mL) was stirred at 80° C. under argon for 16 h. The mixture was allowed to cool to room temperature, and then partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=20:1) to afford the desired product (3.3 g, 75.8% yield) as an off-white solid.

2,2',5'-Trichloro-5-iodo-4-methoxy-1,1'-biphenyl

(568) A mixture of 2',5',6-trichloro-4-methoxy-[1,1'-biphenyl]-3-amine (1.0 g, 3.3 mmol) in conc. HCl (10 mL) and water (10 mL) at 0° C., NaNO₂ (350 mg, 5 mmol) was added in portions and the resulting solution was stirred at -5-0° C. for 30 min. To this mixture, a solution of KI (2.2 g, 13.2 mmol) in H₂O (10 mL) and CuI (630 mg, 3.3 mmol) was added slowly. The resulting solution was stirred at room temperature for 45 min and then extracted with ethyl acetate. The organic layer was washed with saturated NaHCO₃ aqueous solution and brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=20:1) to afford the desired product (1.17 g, 87% yield) as a dark oil.

2,2',5'-Trichloro-4-methoxy-5-vinyl-1,1'-biphenyl

(569) A mixture of 2,2',5'-trichloro-5-iodo-4-methoxy-1,1'-biphenyl (1.17 g, 2.9 mmol), tributyl(vinyl)stannane (1.1 g, 3.5 mmol), Pd(PPh₃)₄ (670 mg, 0.6 mmol) in toluene (30 mL) was stirred at reflux under argon for 6 h. The reaction mixture was quenched with KF aqueous solution and then extracted with ethyl acetate. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/DCM=100:1) to afford the desired product (310 mg,

35% yield) as a white solid.

2-(2',5',6-Trichloro-4-methoxy-[1,1'-biphenyl]-3-yl)ethanol

(570) To a stirring solution of 2,2',5'-trichloro-4-methoxy-5-vinyl-1,1'-biphenyl (310 mg, 1 mmol) in THF (15 mL) under nitrogen at room temperature, BH.sub.3 (2 mL, 1 N) was added. After stirring for 8 h, a mixture of NaOH (160 mg, 4 mmol) in water (3 mL) and H.sub.2O.sub.2 (30% in H.sub.2O, 0.3 g, 4 mmol) was added and the resulting mixture was stirred for 6 h. The reaction was quenched with NaHCO.sub.3 aqueous solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=4:1) to afford the desired product (180 mg, 54% yield) as a colorless oil.

2-(2',5',6-Trichloro-4-methoxy-[1,1'-biphenyl]-3-yl)acetic acid

(571) To a solution of 2-(2',5',6-Trichloro-4-methoxy-[1,1'-biphenyl]-3-yl)ethanol (180 mg, 0.54 mmol) in CH.sub.3CN (20 mL) at 0° C., H.sub.5IO.sub.5\CrO.sub.3 (3.7 mL, 1.63 mmol, 0.44 M in water) was added and the resulting mixture was stirred at 0° C. for 1 h. The reaction was quenched with Na.sub.2HPO.sub.4 and diluted with ethyl acetate. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo to afford the product (110 mg, 59% yield) as a colorless oil. ESI-MS m/z: 343.1 [M-H].sup.-.

1-(4-(2-(2',5',6-Trichloro-4-methoxy-[1,1'-biphenyl]-3-yl)acetyl)piperazin-1-yl)prop-2-en-1-one

(572) To a solution of 2-(2',5',6-Trichloro-4-methoxy-[1,1'-biphenyl]-3-yl)acetic acid (110 mg, 0.318 mmol) Et.sub.3N (96 mg, 0.954 mmol) in DMF (3 mL) at 0° C., EDCI.HCl (92 mg, 0.477 mmol) and HOBt (65 mg, 0.477 mmol) were added and the resulting mixture was stirred at 0° C. for 30 min. To this mixture, 1-(piperazin-1-yl)prop-2-en-1-one (67 mg, 0.382 mmol) was added. The resulting mixture was stirred at room temperature for 16 h and then partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (DCM/MeOH=30:1) to afford the desired product (37 mg, 24.97% yield) as an off-white solid. .sup.1H NMR (400 MHz, DMSO-d6) δ: 7.61 (d, J=8.4 Hz, 1H), 7.52 (dd, J=2.4, 8.8 Hz, 1H), 7.40 (d, J=2.4 Hz, 1H), 7.20 (s, 1H), 7.10 (s, 1H), 6.81 (dd, J=10.8, 15.2 Hz, 1H), 6.13 (dd, J=2.4, 16.8 Hz, 1H), 5.71 (dd, J=2.0, 10.4 Hz, 1H), 3.84 (s, 3H), 3.67 (s, 2H), 3.54-3.41 (m, 8H). ESI-MS m/z: 467.1 [M+H].sup.+.

Example 16

(573) ##STR00522## ##STR00523##

Methyl 2-(4-chloro-5-iodo-2-methoxyphenylamino)acetate

(574) A mixture of tert-butyl 2-(4-chloro-5-iodo-2-methoxyphenylamino)acetic acid (3.0 g, 8.7 mmol), SOCl.sub.2 (3 mL) in MeOH (20 mL) was stirred at reflux for 2 h. The mixture was concentrated in vacuo to yield the crude product (3.1 g) as a yellow solid.

2-((2',5',6-Trichloro-4-methoxy-[1,1'-biphenyl]-3-yl)amino)acetic acid

(575) A mixture of methyl 2-(4-chloro-5-iodo-2-methoxyphenylamino)acetate (5.0 g, 14.08 mmol), 2,5-dichlorophenylboronic acid (4.03 g, 21.12 mmol), Pd(PPh.sub.3).sub.4 (1.626 g, 1.04 mmol), Na.sub.2CO.sub.3 (4.477 g, 42.24 mmol) in 1,4-dioxane (100 mL) and water (20 mL) was stirred at reflux under argon for 6 h. Then reaction mixture was allowed to cool to room temperature, quenched with water and acidified with HCl (10% in water) to adjust the pH to 3-4. The mixture was extracted with ethyl acetate, washed with brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (50% petroleum ether/ethyl acetate) to afford the desired product (3 g, 59% yield) as an off-white solid. ESI-MS m/z: 360.2 [M-H].sup.-.

1,4-di(tert-Butoxycarbonyl)piperazine-2-carboxylic acid

(576) To a solution of piperazine-2-carboxylic acid (21.2 g, 0.16 mol) in 1,4-dioxane at 0° C., NaOH (80 mL, 400 mmol) was added slowly (over 15 min) followed by (Boc).sub.2O (71 g, 33 mol) and the resulting mixture was stirred at room temperature for 16 h. The mixture was

concentrated in vacuo. The residue was dissolved in water (100 mL), acidified with conc. HCl at 0° C. to adjust the pH to 2-3 and then extracted with ethyl acetate. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo to afford the product (44.2 g, 83.7% yield) as an off-white solid.

di-tert-Butyl 2-carbamoylpiperazine-1,4-dicarboxylate

(577) To a solution of 1,4-di(tert-butoxycarbonyl)piperazine-2-carboxylic acid (8.21 g, 24.85 mmol) was dissolved in THF (50 mL) and Et.sub.3N (20 mL) at 0° C., ethyl chloroformate (2.8 g, 26.1 mmol) was added dropwise. After stirring at -5-0° C. for 1 h, NH.sub.4OH (20 mL) was added and the resulting solution was stirred at room temperature for 1 h. The mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with 1N NaOH and brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo to give the product (7.5 g, 91.7% yield) as an off-white solid.

Piperazine-2-carboxamide dihydrochloride

(578) A mixture of di-tert-butyl 2-carbamoylpiperazine-1,4-dicarboxylate (7.5 g, 22.79 mmol), in HCl/MeOH (75 mL, 2.86 N) was stirred at room temperature for 16 h. The mixture was concentrated in vacuo to yield the product (4.58 g, 100% yield) as a yellow solid.

tert-Butyl 3-carbamoylpiperazine-1-carboxylate

(579) To a solution of piperazine-2-carboxamide dihydrochloride (2.02 g, 10 mmol) and Et.sub.3N (3.03 g, 30 mmol) in DCM (40 mL) at 0° C., (Boc).sub.2O (2.18 g, 10 mmol) was added dropwise (over 1 h). The resulting solution was stirred at room temperature for 16 h and then partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (DCM/MeOH=30:1) to afford the desired product (1.1 g, 48.0% yield).

tert-Butyl 4-acryloyl-3-carbamoylpiperazine-1-carboxylate

(580) To a solution of tert-butyl 3-carbamoylpiperazine-1-carboxylate (300 mg, 1.31 mmol) and Et.sub.3N (396 mg, 3.93 mmol) in DCM (5 mL) at 0° C., acryloyl chloride (130 mg, 1.44 mmol) in DCM (1 mL) was added and the resulting mixture was stirred at room temperature for 1.5 h. The mixture was partitioned between DCM and saturated NaHCO.sub.3 aqueous solution. The organic layer was washed with saturated NaHCO.sub.3 and brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (DCM/MeOH=30:1) to afford the desired product (200 mg, 53.9% yield) as an off-white solid.

1-Acryloylpiperazine-2-carboxamide hydrochloride

(581) A mixture of tert-butyl 4-acryloyl-3-carbamoylpiperazine-1-carboxylate (200 mg, 0.706 mmol) in HCl/MeOH (20 mL, 2.86 N) was stirred at room temperature for 1 h. The mixture was concentrated in vacuo to yield the crude product (160 mg) as a yellow solid which was used directly in next step without further purification.

1-Acryloyl-4-(2-((2',5',6-trichloro-4-methoxy-[1,1'-biphenyl]-3-yl)amino)acetyl)piperazine-2-carboxamide

(582) 2-((2',5',6-Trichloro-4-methoxy-[1,1'-biphenyl]-3-yl)amino)acetic acid (231 mg, 0.641 mmol) and Et.sub.3N (592 mg, 2.564 mmol) in DMF (3 mL) at 0° C., EDCI.HCl (184 mg, 0.961 mmol) and HOBt (134 mg, 0.961 mmol) were added and the resulting mixture was stirred at 0° C. for 30 min. To this mixture, the above 1-acryloylpiperazine-2-carboxamide hydrochloride (160 mg) was added and stirred at room temperature for 16 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (DCM/MeOH=30:1) to afford the desired product (10 mg, 2.97% yield) as an off-white solid.

.sup.1H NMR (400 MHz, DMSO-d6) δ: 7.60-7.38 (m, 3H), 6.70-6.99 (m, 1H), 6.86-6.78 (m, 1H), 6.54-6.49 (m, 1H), 6.16-6.10 (m, 1H), 5.76-5.69 (m, 1H), 5.32-5.31 (m, 1H), 4.93-4.84 (m, 1H), 4.93-4.84 (m, 1H), 4.70-4.62 (m, 1H), 4.35-4.32 (m, 1H), 4.04-3.96 (m, 2H), 3.89-3.63 (m, 1H),

2.90-2.86 (m, 1H). ESI-MS m/z: 525.2 [M+H].sup.+.

Example 17

(583) ##STR00524##

1-(4-(2',6-dichloro-4-methoxy-[1,1'-biphenyl]-3-carbonyl)piperazin-1-yl)prop-2-en-1-one

(584) tert-Butyl 4-(2',6-dichloro-4-methoxy-[1,1'-biphenyl]-3-carbonyl)piperazine-1-carboxylate (200 mg, 0.43 mmol) was stirred in HCl in MeOH (2.86 M, 10 mL) for 1 h. The mixture was concentrated in vacuo to yield the crude product. The residue was dissolved in DCM (15 mL), triethylamine (0.5 mL), acryloyl chloride (40 mg, 0.43 mmol) was added to the mixture. The reaction mixture was stirred at room temperature for 30 min, poured into water, and extracted with DCM. The organic layer was washed with water and brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (16 mg, 10% yield) as white solid. ESI-MS m/z: 419.2 [M+H].sup.+.

1-(4-(2',6-Dichloro-4-hydroxy-[1,1'-biphenyl]-3-carbonyl)piperazin-1-yl)prop-2-en-1-one (VI-3)

(585) To a solution of 1-(4-(2',6-Dichloro-4-methoxy-[1,1'-biphenyl]-3-carbonyl)piperazin-1-yl)prop-2-en-1-one (200 mg, 0.48 mmol) in DCM (15 mL) at -60° C., BBr.sub.3 (0.6 g, 2.4 mmol) was added dropwise and the resulting mixture was stirred at room temperature for 1 h. The mixture was poured into ice-water, basified with saturated NaHCO.sub.3 solution to adjust the pH to 8-9, and extracted with DCM. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=20:1) to afford the desired product (10 mg, 5% yield). .sup.1H NMR (400 MHz, DMSO-d6) δ: 10.6 (s, 1H), 7.57-7.33 (m, 5H), 7.12 (s, 1H), 7.05 (s, 1H), 6.80 (m, 1H), 6.15-6.11 (dd, J=2, 16.8 Hz, 1H), 5.72-5.70 (m, 1H), 3.6 (m, 8H). ESI-MS m/z: 405.3 [M+H].sup.+.

Example 18

(586) ##STR00525## ##STR00526##

4,5-Dichloro-2-methoxybenzoic acid

(587) A mixture of 4-chloro-2-methoxybenzoic acid (10 g, 53.6 mmol) and NCS (35 g, 19.2 mmol) in acetonitrile (200 mL) was stirred at room temperature for 48 h. The mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with water and brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo to get the crude product (23.3 g).

Methyl 4,5-dichloro-2-methoxybenzoate

(588) A mixture of 4,5-dichloro-2-methoxybenzoic acid (8.2 g, 37 mmol) and K.sub.2CO.sub.3 (11.8 g, 111 mmol) in DMF (100 mL), CH.sub.3I (6.3 g, 44 mmol) was added dropwise and the resulting mixture was stirred at room temperature for 16 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with water and brine, dried over Na.sub.2SO.sub.4, and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=10:1) to afford the desired product. (4,5-Dichloro-2-methoxyphenyl)methanol

(589) To a mixture of LiAlH.sub.4 (2.42 g, 64 mmol) in THF (40 mL) at -40° C. under argon, a solution of methyl 4,5-dichloro-2-methoxybenzoate (6 g, 26 mmol) in THF (50 mL) was added dropwise. The reaction mixture was stirred at -5° C. to 5° C. for 1 h. The mixture was cooled to -20° C. and then water (2 mL) and NaOH (15%) aqueous were added. The resulting mixture was stirred for 15 min. The solid was filtered, and the cake rinsed with ethyl acetate. The combined filtrate was dried over Na.sub.2SO.sub.4 and concentrated in vacuo to afford the crude product (4.6 g).

2-(4,5-Dichloro-2-methoxybenzyl)isoindoline-1,3-dione

(590) To a mixture of 4,5-dichloro-2-methoxyphenyl)methanol (4.5 g, 22 mmol), isoindoline-1,3-dione (9.6 g, 65 mmol) and PPh.sub.3 (17 g, 65 mmol) in THF (100 mL) at room temperature, DIAD (13 g, 65 mmol) was added and the resulting mixture was stirred at room temperature for 16

h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na_2SO_4 , concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=10:1) to afford the desired product.

(4,5-Dichloro-2-methoxyphenyl)methanamine

(591) To a solution of 2-(4,5-dichloro-2-methoxybenzyl)isoindoline-1,3-dione (1.8 g, 5 mmol) in EtOH (5 mL), hydrazine hydrate (1.34 g, 27 mmol) was added and the resulting mixture was stirred at reflux for 1 h. The mixture was concentrated in vacuo and the residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=20:1) to afford the desired product (0.8 g, 78% yield).

tert-Butyl 4-((4,5-dichloro-2-methoxybenzyl)carbamoyl)piperidine-1-carboxylate

(592) The mixture of (4,5-dichloro-2-methoxyphenyl)methanamine (0.8 g, 3.90 mmol), 1-(tert-butoxycarbonyl)piperidine-4-carboxylic acid (0.88 g, 3.84 mmol), BOP (2 g, 1.16 mmol) and DIEA (1.6 g, 2.91 mmol) in DMF (20 mL) was stirred at room temperature for 1 h. The mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with water and brine, dried over Na_2SO_4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=100:1) to afford the desired product (0.987 g, 62% yield). ESI-MS m/z : 415.4 $[\text{M}-\text{H}]^+$.

N-(4,5-Dichloro-2-methoxybenzyl)piperidine-4-carboxamide

(593) The mixture of 4-((4,5-dichloro-2-methoxybenzyl)carbamoyl)piperidine-1-carboxylate (987 mg, 2.37 mmol) in HCl/MeOH (20 mL, 57.2 mmol) was stirred at room temperature for 1 h. Then the solvent was evaporated in vacuo and the residue was dissolved with dichloromethane (5 mL). To this mixture, NaH (85 mg, 3.55 mmol) was added. Then the resulting mixture was stirred at room temperature for 30 min. The solvent was removed under reduced pressure to yield the crude product (800 mg).

tert-butyl 3-(4-(4,5-Dichloro-2-methoxybenzylcarbamoyl)piperidin-1-yl)azetidine-1-carboxylate

(594) A mixture of N-(4,5-dichloro-2-methoxybenzyl)piperidine-4-carboxamide (750 mg, 2.37 mmol), tert-butyl 3-oxoazetidine-1-carboxylate (607 mg, 3.55 mmol), AcOH (1 mL) and MeOH (5 mL) was stirred at reflux for 2 h. To this mixture, NaBH_3CN (0.74 g, 11.85 mmol) was added and the resulting mixture was stirred at 60° C. for 16 h. The mixture was allowed to cool to room temperature and partitioned between NH_4Cl aqueous solution and ethyl acetate. The organic layer was washed with water and brine, dried over Na_2SO_4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (220 mg, 18% yield). ESI-MS m/z : 472.3 $[\text{M}+\text{H}]^+$.

1-(1-Acryloylazetidin-3-yl)-N-(4,5-dichloro-2-methoxybenzyl)piperidine-4-carboxamide

(595) A mixture of tert-butyl 3-(4-(4,5-dichloro-2-methoxybenzylcarbamoyl)piperidin-1-yl)azetidine-1-carboxylate (210 mg, 0.44 mmol) in HCl/MeOH (10 mL, 2.86 M) was stirred at room temperature for 1 h. The mixture was concentrated in vacuo to yield the crude residue. The residue was dissolved in DCM (5 mL), triethylamine (0.5 mL) and acryloyl chloride (40 mg, 0.43 mmol) were added. The reaction mixture was stirred at room temperature for 30 min and then partitioned between DCM and water. The organic layer was washed with water and brine, dried over Na_2SO_4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (150 mg, 82% yield).

N-(4,5-Dichloro-2-hydroxybenzyl)-1-(1-acryloylazetidin-3-yl)piperidine-4-carboxamide

(596) To a solution of 1-(1-acryloylazetidin-3-yl)-N-(4,5-dichloro-2-methoxybenzyl)piperidine-4-carboxamide (150 mg, 0.35 mmol) in DCM (15 mL) at -60° C., BBr_3 (0.6 g, 2.4 mmol) was added dropwise. The resulting mixture was allowed to warm to room temperature and stirred for 1 h. The mixture was poured into ice-water, basified with saturated NaHCO_3 solution to adjust

the pH to 8-9, and then extracted with DCM. The organic layer was dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=20:1) to afford the desired product (34 mg, 24% yield).
.sup.1H NMR (400 MHz, DMSO-d₆) δ: 10.34 (s, 1H), 8.2-8.25 (m, 1H), 7.2 (s, 1H), 7.0 (s, 1H), 6.91 (s, 1H), 6.33-6.27 (m, 1H), 6.12-6.07 (dd, J=2.4, 12.4 Hz, 1H), 5.68-5.65 (dd, J=2.4, 10.4 Hz, 1H), 4.24-4.20 (m, 1H), 4.17-4.14 (m, 2H), 4.14-3.99 (m, 1H), 3.94-3.90 (m, 1H), 3.73-3.70 (m, 1H), 3.10 (s, 1H), 2.84-2.80 (m, 2H), 2.22 (m, 1H), 1.80 (s, 2H), 1.73-1.71 (m, 2H), 1.63-1.57 (m, 2H). ESI-MS m/z: 412.2 [M+H].sup.+.

Example 19

(597) ##STR00527##

Methyl 4-chloro-5-iodo-2-methoxybenzoate

(598) A mixture of 4-chloro-5-iodo-2-methoxybenzoic acid (2 g, 6.41 mmol) concentrated sulfuric acid (1.5 mL) in MeOH (50 mL) was stirred at reflux for 16 h. The mixture was allowed to cool to room temperature and partitioned between water and ethyl acetate. The organic layer was washed with water and brine, dried over Na₂SO₄ and concentrated in vacuo to afford the desired product (1.85 g, 85% yield) as a yellow oil.

Methyl 2',5',6-trichloro-4-methoxy-[1,1'-biphenyl]-3-carboxylate

(599) A mixture of Methyl 4-chloro-5-iodo-2-methoxybenzoate (1.8 g, 5.51 mmol), (2,5-dichlorophenyl)boronic acid (2.1 g, 11.03 mmol), Pd(PPh₃)₄ (403 mg, 0.55 mmol), Na₂CO₃ (1.75 g, 16.54 mmol) in 1,4-dioxane (50 mL) and water (5 mL) was stirred at reflux under argon for 16 h. The mixture was filtered and filtrate was concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=20:1) to afford the desired product (1.6 g, 85% yield).

1-(3-(Hydroxymethyl)-4-(2',5',6-trichloro-4-methoxy-[1,1'-biphenyl]-3-carbonyl)piperazin-1-yl)prop-2-en-1-one (VI-7)

(600) The title compound was prepared from methyl 2',5',6-trichloro-4-methoxy-[1,1'-biphenyl]-3-carboxylate in three steps followed the procedure described in Example 20.

(601) tert-butyl 3-(hydroxymethyl)-4-(2',5',6-trichloro-4-methoxy-[1,1'-biphenyl]-3-carbonyl)piperazine-1-carboxylate (420 mg, 0.79 mmol) was stirred in HCl in MeOH (2.85 N). The solvent was removed under reduced pressure to yield the crude residue which was dissolved in DMF (20 mL), acrylic acid (57 mg, 0.79 mmol), BOP (421 mg, 0.95 mmol) and DIEA (409 mg, 3.17 mmol) were added. The reaction was stirred at room temperature for 1 h. The resulting mixture was poured into water, extracted with ethyl acetate and washed with water and brine. The combined organic layer was dried over Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography on silica gel (dichloromethane/methanol=60:1) to afford the desired product (92 mg, 24% yield).
.sup.1H NMR (400 MHz, DMSO-d₆) δ: 7.64-7.20 (m, 5H), 6.83-6.70 (m, 1H), 6.16-6.11 (d, 1H), 5.74-5.71 (d, 1H), 6.91 (s, 1H), 5.08-4.01 (m, 3H), 3.90-3.86 (d, 3H), 3.49-3.22 (m, 2H), 2.93-2.74 (m, 2H), 2.89-2.67 (m, 2H). ESI-MS m/z: 451.2 [M-H].sup.-.

Example 20

(602) ##STR00528##

tert-Butyl 4-(4-bromo-1H-pyrrole-2-carbonyl)piperazine-1-carboxylate

(603) To a mixture of 4-bromo-1H-pyrrole-2-carboxylic acid (800 mg, 4.21 mmol), tert-butylpiperazine-1-carboxylate (822 mg, 4.42 mmol), BOP (2.2 g, 5.05 mmol) in DMF (5 mL), DIEA (1.63 g, 12.63 mmol) was added and the resulting mixture was stirred at room temperature for 1 h. The mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with water and brine, dried over Na₂SO₄ and concentrated in vacuo to afford the crude product (920 mg, 61% yield) which was used directly in the next step without purification.

tert-Butyl 4-(4-(2,4-dichlorophenyl)-1H-pyrrole-2-carbonyl)piperazine-1-carboxylate

(604) A mixture of tert-butyl 4-(4-bromo-1H-pyrrole-2-carbonyl)piperazine-1-carboxylate (350 mg, 0.98 mmol), (2,4-dichlorophenyl) boronic acid (280 mg, 1.47 mmol), Pd(PPh₃)₄ (116 mg,

0.1 mmol), Na.sub.2CO.sub.3 (312 mg, 2.94 mmol) in 1,4-dioxane (10 mL) and water (2 mL) was stirred at reflux under argon for 16 h. The mixture was allowed to cool to room temperature and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=30:1) to afford the desired product (273 mg, 59% yield).

1-(4-(4-(2,4-Dichlorophenyl)-1H-pyrrole-2-carbonyl)piperazin-1-yl)prop-2-en-1-one (VI-32) (605) The mixture of tert-butyl 4-(4-(2,4-dichlorophenyl)-1H-pyrrole-2-carbonyl)piperazine-1-carboxylate (270 mg, 0.64 mmol) in HCl/MeOH (20 mL, 57.2 mmol) was stirred for 1 h. The mixture was concentrated in vacuo and the residue was dissolved in DMF (5 mL). To this mixture, acrylic acid (50 mg, 0.7 mmol), BOP (437 mg, 0.72 mmol) and DIEA (248 mg, 1.92 mmol) were added. The reaction mixture was stirred at room temperature for 1 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=60:1) to afford the desired product (40 mg, 27% yield) as a white solid. ¹H NMR (400 MHz, DMSO-d₆) δ: 11.88 (s, 1H), 7.66-7.63 (m, 2H), 7.43 (m, 1H), 7.38 (s, 1H), 6.91 (s, 1H), 6.85-6.78 (m, 1H), 6.18-6.13 (dd, J=2.4, 12.4 Hz, 1H), 5.74-5.71 (dd, J=2.4, 10.4 Hz, 1H), 3.76 (s, 4H), 3.68-3.63 (m, 4H). ESI-MS m/z: 377.3 [M-H].sup.-.

Example 21

(606) ##STR00529##

2,2',5'-Trichloro-4-methoxy-5-vinylbiphenyl

(607) To a stirred solution of 2,2',5'-trichloro-5-iodo-4-methoxybiphenyl (3.4 g, 8.3 mmol) in toluene (100 mL) at room temperature, tributyl(vinyl)stannane (3.1 g, 9.9 mmol) was added followed by Pd(PPh.sub.3).sub.4 (2 g, 1.7 mmol). The reaction mixture was degassed and back-filled with nitrogen (several cycles) and then stirred at reflux for 16 h. The mixture was allowed to cool to room temperature, a solution of KF (1.72 g, 29.7 mmol) in H.sub.2O (10 mL) was added and then stirred for 1 h. The mixture was partitioned between ethyl acetate and water. The organic layer was dried over anhydrous Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (1-2% ethyl acetate/petroleum ether) to afford the desired product (2.4 g, 92% yield) as a white solid

2-(2',5',6-Trichloro-4-methoxybiphenyl-3-yl)acetaldehyde

(608) To a mixture of 2,2',5'-trichloro-4-methoxy-5-vinylbiphenyl (1.0 g, 3.2 mmol) in anhydrous DMF (10 mL), PdCl.sub.2 (1.14 g, 6.4 mmol) was added and the resulting mixture was stirred at room temperature under nitrogen for 16 h. To this mixture, water (0.5 mL) was added and the resulting mixture was stirred at room temperature for another 6 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with water, dried over anhydrous Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (1-3% ethyl acetate/petroleum ether) to afford the desired product (300 mg, 29% yield) as a white solid.

tert-Butyl 4-(2-(2',5',6-trichloro-4-methoxybiphenyl-3-yl)ethyl)piperazine-1-carboxylate

(609) To a stirred solution of 2-(2',5',6-trichloro-4-methoxybiphenyl-3-yl)acetaldehyde (300 mg, 0.92 mmol) in DCM (20 mL) at room temperature, tert-butyl piperazine-1-carboxylate (205 mg, 1.1 mmol) was added followed by drops of AcOH. The reaction mixture was stirred at room temperature for 1 h, and then NaBH(OAc).sub.3 (1.95 g, 9.2 mmol) was added. The resulting mixture was stirred at reflux for 16 h. The mixture was allowed to cool to room temperature, diluted with water and extracted with DCM. The organic layer was dried over anhydrous Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (1-33% ethyl acetate/petroleum ether) to afford the desired product (350 mg, 77% yield) as a white solid.

1-(4-(2-(2',5',6-Trichloro-4-hydroxybiphenyl-3-yl)ethyl)piperazin-1-yl)prop-2-en-1-one

(610) The title compound was prepared from tert-butyl 4-(2-(2',5',6-trichloro-4-methoxybiphenyl-

3-yl)ethyl)piperazine-1-carboxylate in three steps following the procedure described in Example 18. ¹H NMR (400 MHz, DMSO-d₆) δ: 10.96 (bs., 1H), 7.54-7.59 (d, J=8.6 Hz, 1H), 7.47-7.50 (m, 1H), 7.39-7.40 (d, J=2.6 Hz, 1H), 7.08 (s, 1H), 6.94 (s, 1H), 6.77-6.84 (m, 1H), 6.09 (dd, J=2.1, 17.0 Hz, 1H), 5.65 (dd, J=2.6, 10.7 Hz, 1H), 3.50 (m, 4H), 2.72-2.76 (m, 2H), 2.54-2.57 (m, 2H), 2.47-2.50 (m, 4H). ESI-MS m/z: 439.1 [M+H].⁺.

Example 22

(611) ##STR00530##

tert-Butyl 4-(2',6-dichloro-4-methoxy-[1,1'-biphenyl]-3-carbonyl)piperazine-1-carboxylate (612) To a stirred solution of 2',6-dichloro-4-methoxy-[1,1'-biphenyl]-3-carboxylic acid (500 mg, 1.68 mmol) in DMF (10 mL) at room temperature, tert-butyl piperazine-1-carboxylate (345 mg, 1.85 mmol), BOP (892 mg, 2.02 mmol) and DIEA (542 mg, 4.2 mmol) were added and the resulting mixture was stirred at room temperature for 2 h. The reaction mixture was diluted with ethyl acetate and washed with brine. The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (550 mg, 70% yield). ESI-MS m/z: 465.4 [M+H].⁺.

(E)-1-(4-(2',6-dichloro-4-methoxy-[1,1'-biphenyl]-3-carbonyl)piperazin-1-yl)-4-(dimethylamino)but-2-en-1-one

(613) A mixture of tert-Butyl 4-(2',6-dichloro-4-methoxy-[1,1'-biphenyl]-3-carbonyl)piperazine-1-carboxylate (550 mg, 1.18 mmol) in HCl/MeOH (20 mL, 57.2 mmol) was stirred at room temperature for 1 h. The solvent was removed under reduced pressure to yield the crude product. The crude residue was dissolved with DMF (10 mL), 4-(dimethylamino)but-2-enoic acid (215 mg, 0.47 mmol), BOP (627 mg, 1.42 mmol) and DIEA (610 mg, 4.73 mmol) was added. The resulting mixture was stirred at room temperature for 2 h. The reaction mixture was diluted with ethyl acetate and washed with brine. The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (450 mg, 80% yield, 2 steps). ESI-MS m/z: 476.4 [M+H].⁺.

(E)-1-(4-(2',6-dichloro-4-hydroxy-[1,1'-biphenyl]-3-carbonyl)piperazin-1-yl)-4-(dimethylamino)but-2-en-1-one (VI-24)

(614) A solution of (E)-1-(4-(2',6-dichloro-4-methoxy-[1,1'-biphenyl]-3-carbonyl)piperazin-1-yl)-4-(dimethylamino)but-2-en-1-one (270 mg, 0.57 mmol) in DCM (5 mL) at -78° C., BBr₃ (1.43 g, 5.7 mmol) was added dropwise. The resulting mixture was stirred at room temperature for 2 h. The mixture was poured into ice-water, basified with the aqueous NaHCO₃ to adjust the pH to 7 and then extracted with DCM (3×20 mL). The organic layer was dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (150 mg, 57% yield). ¹H NMR (400 MHz, DMSO-d₆) δ: 13.12 (s, 1H), 8.02 (s, 1H), 7.81 (s, 1H), 6.81 (m, 1H), 6.13 (dd, J=2.8, 16.8 Hz, 1H), 5.71 (dd, J=2.0, 10.4 Hz, 1H), 4.10 (s, 2H), 3.50-3.60 (m, 8H). ESI-MS m/z: 462.4 [M+H].⁺.

Example 23

(615) ##STR00531##

4,5-Dichloro-2-nitrobenzaldehyde

(616) A mixture of 1,2-dichloro-4-methyl-5-nitrobenzene (2 g, 9.71 mmol) and DMF-DMA (3 g, 25.2 mmol) in DMF (50 mL) was stirred at 140° C. for 14 h. The dark solution was cooled to 0° C., and then added to a solution of NaIO₄ (10.8 g, 50.5 mmol) in DMF/water (1:4, 25 mL) at 0° C. After stirring at room temperature for 8 h, the mixture was filtered and the cake was rinsed with ethyl acetate. The filtrate was diluted with ethyl acetate and washed with brine. The organic layer was dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by column chromatography on silica gel (ethyl acetate/petroleum ether=1:20) to afford the desired product (480

mg, 23% yield, 2 steps)

2-(5,6-Dichloro-1H-indazol-3-yl)acetic acid

(617) To a stirred mixture of 4,5-dichloro-2-nitrobenzaldehyde (1.1 g, 5.00 mmol) and malonic acid (676 mg, 6.5 mmol) in formic acid (3 mL) at 40° C., ammonium formate (768 mg, 12.5 mmol) was added. The resulting mixture was stirred at 65° C. for 1 h and then at 95° C. for 4 h. To this mixture, conc. HCl (2.5 mL) was added and then stirred at 95° C. for 1 h. The mixture was allowed to cool to room temperature, quenched with water (5 mL) and extracted with isobutyl ketone. The aqueous layer was adjusted the pH to 4 with KOH (50%) and filtered to afford the crude solid (500 mg). The solid was dissolved in a mixture of NaOH aqueous solution (8 mL, 5%) and N.sub.2H.sub.4.H.sub.2O (90 mg, 1.79 mmol) and heated to 85° C. To this mixture, Raney-Ni was added and the resulting mixture was stirred at 85° C. for 30 min. The mixture was allowed to cool to room temperature and filtered. The aqueous layer was acidified with aqueous HCl (6 M) to adjust the pH to 2. The precipitate was collected by filtration and dried in vacuo to afford the desired product (250 mg, yield 22%, 2 steps).

1-(4-(2-(5,6-Dichloro-1H-indazol-3-yl)acetyl)piperazin-1-yl)prop-2-en-1-one

(618) The title compound was prepared from 2-(5,6-dichloro-1H-indazol-3-yl)acetic acid in three steps followed the procedure described in Example 40. ¹H NMR (400 MHz, DMSO-d₆) δ: 7.56 (m, 1H), 7.33-7.45 (m, 3H), 7.12 (s, 1H), 7.07 (s, 1H), 6.62 (s, 2H), 3.60 (m, 8H), 3.10 (s, 2H), 2.20 (s, 6H). ESI-MS m/z: 367.2 [M+H].sup.+.

Example 24

(619) ##STR00532##

tert-Butyl 4-(2-bromoethyl)piperazine-1-carboxylate

(620) A mixture of tert-butyl piperazine-1-carboxylate (5.0 g, 26.9 mmol), 1,2-dibromoethane (25 mL), DIPEA (3.5 g, 26.9 mmol) was stirred at 30° C. under argon for 72 h. The solvent was removed under reduced pressure and the residue was purified by column chromatography on silica gel (1-2% methanol/dichloroethane) to afford the desired product (2.8 g, 36% yield) as a solid. ESI-MS m/z: 293.1 [M+1].sup.+.

tert-Butyl 4-(2-((4-chloro-5-iodo-2-methoxyphenyl)amino)ethyl)piperazine-1-carboxylate

(621) To a stirred solution of 4-chloro-5-iodo-2-methoxyaniline (968 mg, 3.42 mmol) in anhydrous THF (20 mL) at 0° C., NaH (60% in mineral oil, 205.2 mg, 5.13 mmol,) was added and the resulting mixture was stirred at reflux under nitrogen for 1 h. To this mixture, tert-butyl 4-(2-bromoethyl)piperazine-1-carboxylate (500 mg, 1.71 mmol) was added and the resulting mixture was stirred at room temperature for 15 h. The mixture was concentrated in vacuo and the residue was partitioned between ethyl acetate and brine. The organic layer was dried over anhydrous Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (10-30% ethyl acetate/petroleum ether) to afford the desired product (180 mg, 21% yield) as a solid.

tert-Butyl 4-(2-((2',5',6-trichloro-4-methoxy-[1,1'-biphenyl]-3-yl)amino)ethyl)piperazine-1-carboxylate

(622) A mixture of tert-butyl 4-(2-((4-chloro-5-iodo-2-methoxyphenyl)amino)ethyl)piperazine-1-carboxylate (180 mg, 0.36 mmol), (2,5-dichlorophenyl)boronic acid (138.9 mg, 0.72 mmol), Na.sub.2CO.sub.3 (114.5 mg, 1.08 mmol), Pd(PPh.sub.3).sub.4 (42 mg, 0.036 mmol) in 1,4-dioxane (20 mL) and water (5 mL) was stirred at 90° C. under argon for 16 h. The mixture was allowed to cool to room temperature and then concentrated in vacuo. The residue was purified by column chromatography on silica gel (petroleum ether/ethyl acetate=5:1) to afford the desired product (135 mg, 73% yield) as a solid. ESI-MS m/z: 514.3 [M+H].sup.+.

1-(4-(2-((2',5',6-Trichloro-4-hydroxy-[1,1'-biphenyl]-3-yl)amino)ethyl)piperazin-1-yl)prop-2-en-1-one

(623) The title compound was prepared from tert-butyl 4-(2-((2',5',6-trichloro-4-methoxy-[1,1'-biphenyl]-3-yl)amino)ethyl)piperazine-1-carboxylate in three steps followed the procedure described in Example 18. ¹H NMR (400 MHz, DMSO-d₆) δ: 10.04 (s, 1H), 7.56 (d, J=8.4 Hz,

1H), 7.48-7.45 (m, 1H), 7.37 (d, J=2.8 Hz, 1H), 6.79 (dd, J=10.4, 16.8 Hz, 1H), 6.77 (s, 1H), 6.40 (s, 1H), 6.09 (dd, J=2.4, 16.4 Hz, 1H), 5.67 (dd, J=2.4, 10.4 Hz, 1H), 4.88 (bs., 1H), 3.54-3.51 (m, 4H), 3.11 (d, J=4.8 Hz, 2H), 2.55 (t, J=6.0 Hz, 2H), 2.39-2.38 (m, 4H). ESI-MS m/z: 454.1[M+H].sup.+.

Example 25

(624) ##STR00533##

tert-butyl 5-(3,4-dichlorobenzoyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

(625) To a solution of acid 3,4-dichlorobenzoic acid (0.45 g, 2.36 mmol), tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (0.5 g, 2.36 mmol) and HOBt (0.4 g, 2.83 mmol) in DMF (15 mL), EDCI.HCl (0.54 g, 2.83 mmol) and DIEA (1.5 g, 11.5 mmol) were added. The mixture was stirred at room temperature overnight and then partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4, filtered and concentrated in vacuo to afford the crude product which was used directly in the next step.

(3,4-dichlorophenyl)(hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone

(626) tert-Butyl 5-(3,4-dichlorobenzoyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (crude product from above step) was dissolved in 50% TFA in DCM (10 mL) and the resulting mixture was stirred at room temperature for 3 h. The mixture was concentrated in vacuo. The residue was dissolved in ethyl acetate and washed with saturated NaHCO.sub.3 aqueous solution. The organic layer was dried over Na.sub.2SO.sub.4, filtered and concentrated in vacuo to afford the crude product. The crude product was used directly in the next step without further purification.

(3,4-Dichlorophenyl)(5-(vinylsulfonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone

(627) To a solution of (3,4-dichlorophenyl)(hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone (100 mg, 0.35 mmol) in DCM (10 mL) at 0° C., Et.sub.3N (106 mg, 1 mmol) was added followed by sulfonyl chloride (60 mg, 0.37 mmol). The resulting mixture was stirred at room temperature overnight. The mixture was partitioned between DCM and water. The organic layer was dried over Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified by Isolera One (silica gel cartridge, 10-100% ethyl acetate/hexanes) to afford the desired product (14 mg, 11%). .sup.1H NMR (300 MHz, CDCl.sub.3) δ: 7.60 (d, J=2.0 Hz, 1H), 7.51 (d, J=8.2 Hz, 1H), 7.35 (dd, J=2.0, 8.2 Hz, 1H), 6.50 (dd, J=9.9, 16.5 Hz, 1H), 6.30 (d, J=16.6 Hz, 1H), 6.09 (d, J=9.9 Hz, 1H), 2.90-4.10 (m, 10H). ESI-MS m/z: 375.0 [M+H].sup.+.

Example 26

(628) ##STR00534##

Example 27

(629) ##STR00535##

Example 28

(630) ##STR00536##

Example 29

(631) ##STR00537##

Example 30

(632) ##STR00538##

Example 31

(633) ##STR00539##

Example 32

(634) ##STR00540##

Example 33

(635) ##STR00541##

Example 34

(636) ##STR00542##

Example 35

(637) ##STR00543##

Example 36

(638) ##STR00544##

tert-Butyl 3-(4-(2-(4-Chloro-5-iodo-2-methoxyphenylamino)acetyl)piperazin-1-yl)azetidine-1-carboxylate

(639) To a solution of 2-(4-chloro-5-iodo-2-methoxyphenylamino)acetic acid (2.0 g, 5.88 mmol), tert-butyl 3-(piperazin-1-yl)azetidine-1-carboxylate (1.84 g, 7.64 mmol), EDCI.HCl (2.26 g, 11.76 mmol), and HOBt (1.59 g, 11.76 mmol) in DMF (3 mL) at 0° C., Et.sub.3N (3.28 mL, 23.52 mmol) was added. The resulting mixture was stirred at RT for 16 h and then partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was washed by a mixture of ethyl acetate/petroleum ether=1:5 to afford the desired product (2.24 g, 67% yield) as a white solid. ESI-MS m/z: 565.4 [M+H].sup.+.

tert-Butyl 3-(4-(2-(4-chloro-5-cyclobutyl-2-methoxyphenylamino)acetyl)piperazin-1-yl)azetidine-1-carboxylate

(640) A mixture of tert-butyl 3-(4-(2-(4-chloro-5-iodo-2-methoxyphenylamino)acetyl)piperazin-1-yl)azetidine-1-carboxylate (697 mg, 1.24 mmol), cyclobutylzinc bromide (4.46 mL, 2.23 mmol, 0.5 M in THF), Pd(OAc).sub.2 (56 mg, 0.248 mmol), and S-Phos (102 mg, 0.248 mmol) in THF (15 mL) was stirred at 65° C. under argon for 16 h. The mixture was allowed to cool to RT, quenched with aqueous NH.sub.4Cl solution and then extracted with ethyl acetate. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (methanol/dichloromethane=1:30) to afford the desired product (596 mg, 98% yield) as a brown oil. ESI-MS m/z: 493.5 [M+H].sup.+.

1-(3-(4-(2-(4-Chloro-5-cyclobutyl-2-hydroxyphenylamino)acetyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (V-53)

(641) The title compound was prepared from tert-butyl 3-(4-(2-(4-chloro-5-cyclobutyl-2-methoxyphenylamino)acetyl)piperazin-1-yl)azetidine-1-carboxylate in three steps according to the procedure described in Example 17. .sup.1H NMR (400 MHz, DMSO-d6) δ: 9.66 (s, 1H), 6.63 (s, 1H), 6.52 (s, 1H), 6.31 (dd, J=10.2, 16.9 Hz, 1H), 6.10 (dd, J=2.1, 16.8 Hz, 1H), 5.68 (dd, J=2.1, 10.2 Hz, 1H), 5.16 (t, J=4.4 Hz, 1H), 4.27-4.23 (m, 1H), 4.08-4.04 (m, 1H), 3.97-3.93 (m, 3H), 3.80-3.76 (m, 1H), 3.65-3.59 (m, 1H), 3.56-3.54 (m, 4H), 3.20-3.14 (m, 1H), 2.40-2.25 (m, 4H), 2.20-2.15 (m, 2H), 2.09-2.05 (m, 2H), 1.97-1.90 (m, 1H), 1.80-1.74 (m, 1H). ESI-MS m/z: 433.4 [M+H].sup.+.

Example 37

(642) ##STR00545##

Methyl 2-(4-chloro-5-iodo-2-methoxyphenylamino)propanoate

(643) A mixture of tert-butyl 4-chloro-5-iodo-2-methoxybenzenamine (2 g, 7.07 mmol), methyl 2-bromopropanoate (1.17 g, 7.07 mmol), K.sub.2CO.sub.3 (1.94 g, 14.14 mmol) and KI (0.235 g, 1.414 mmol) in DMF (25 mL) was stirred at 100° C. for 16 h. The mixture was allowed to cool to RT, quenched with aqueous NaHCO.sub.3 solution and then extracted with ethyl acetate. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (ethyl acetate/petroleum ether=1:20) to afford the desired product (1.12 g, 43% yield) as a yellow solid. ESI-MS m/z: 370.1 [M+H].sup.+.

2-(4-Chloro-5-iodo-2-methoxyphenylamino)propanoic acid

(644) To a solution of methyl 2-(4-chloro-5-iodo-2-methoxyphenylamino)propanoate (1.12 g, 3.04 mmol) in mixture of tetrahydrofuran (20 mL) and water (10 mL) at RT, LiOH.H.sub.2O (0.51 g, 12.16 mmol) was added and the resulting mixture was stirred for 1 h. The aqueous phase was washed with TBME and then acidified with aqueous HCl (1 N) to adjust the pH to 5. The mixture was extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous Na.sub.2SO.sub.4, filtered and concentrated in vacuo to afford the crude product (760 mg) which was used directly in the next step without further purification. ESI-MS m/z: 356.1 [M+H].sup.+.

Tert-Butyl 3-(4-(2-((4-Chloro-5-iodo-2-methoxyphenyl)amino)propanoyl)piperazin-1-yl)azetidine-1-carboxylate

(645) To a solution of 2-(4-chloro-5-iodo-2-methoxyphenylamino)propanoic acid (760 mg, 2.13 mmol), tert-butyl 3-(piperazin-1-yl)azetidine-1-carboxylate (669 mg, 2.78 mmol), EDCI.HCl (818 mg, 4.26 mmol), HOBT (575 mg, 4.26 mmol) in DMF (8 mL) at 0° C., Et.sub.3N (861 mg, 8.52 mmol) was added. The resulting mixture was stirred at RT for 16 h and then partitioned between ethyl acetate and water. The organic layer was washed with saturated NaHCO.sub.3 solution and brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (ethyl acetate/petroleum ether=1:1) to afford the desired product (673 mg, 55% yield) as a white solid. ESI-MS m/z: 579.4 [M+H].sup.+.

Tert-Butyl 3-(4-(2-((4-Chloro-5-cyclobutyl-2-methoxyphenyl)amino)propanoyl)piperazin-1-yl)azetidine-1-carboxylate

(646) A mixture of tert-butyl 3-(4-(2-((4-chloro-5-iodo-2-methoxyphenyl)amino)propanoyl)piperazin-1-yl)azetidine-1-carboxylate (673 mg, 1.162 mmol), cyclobutylzinc bromide (5.11 mL, 2.556 mmol, 0.5 M in THF), Pd(OAc).sub.2 (52 mg, 0.23 mmol), S-Phos (95 mg, 0.23 mmol) in THF (10 mL) was stirred at 65° C. under argon for 16 h. The mixture was allowed to cool to RT, quenched with aqueous NH.sub.4Cl solution and then extracted with ethyl acetate. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (ethyl acetate/petroleum ether=1:1) to afford the desired product (565 mg, 96% yield) as a light yellow solid. ESI-MS m/z: 507.6 [M+H].sup.+.

1-(3-(4-(2-((4-Chloro-5-cyclobutyl-2-hydroxyphenyl)amino)propanoyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (V-59)

(647) The title compound was prepared from tert-butyl 3-(4-(2-((4-chloro-5-cyclobutyl-2-methoxyphenyl)amino)propanoyl)piperazin-1-yl)azetidine-1-carboxylate in three steps according to the procedure described in Example 17. .sup.1H NMR (400 MHz, DMSO-d6) δ: 9.63 (s, 1H), 6.62 (s, 1H), 6.49 (s, 1H), 6.30 (dd, J=10.1, 16.8 Hz, 1H), 6.10 (d, J=18.7 Hz, 1H), 5.68 (d, J=10.4 Hz, 1H), 4.86 (d, J=9.2 Hz, 1H), 4.69-4.63 (m, 1H), 4.27-4.23 (m, 1H), 4.07-4.03 (m, 1H), 3.97-3.62 (m, 1H), 3.82-3.76 (m, 2H), 3.64-3.55 (m, 3H), 3.77-3.11 (m, 1H), 2.44-2.15 (m, 6H), 2.08-1.90 (m, 4H), 1.80-1.72 (m, 2H), 1.97-1.90 (m, 1H), 1.24 (d, J=6.4 Hz, 3H). ESI-MS m/z: 447.4 [M+H].sup.+.

Example 38

(648) ##STR00546##

1-Benzhydrylazetidin-3-yl methanesulfonate

(649) A mixture of 1-benzhydrylazetidin-3-ol (20.0 g, 83.68 mmol) and Et.sub.3N (12.68 g, 125.52 mmol) in DCM (200 mL) at 0° C., MsCl (11.447 mg, 100.41 mmol) was added in portions and the resulting solution was stirred at RT for 1 h. The reaction mixture was diluted with ethyl acetate and washed with brine. The organic layer was dried over anhydrous Na.sub.2SO.sub.4, filtered and concentrated in vacuo to afford the desired product (26.526 g, 100% yield).

tert-Butyl 4-(1-benzhydrylazetidin-3-yl)piperazine-1-carboxylate

(650) A mixture of 1-benzhydrylazetidin-3-yl methanesulfonate (26.53 g, 83.68 mmol), tert-butyl piperazine-1-carboxylate (18.68 g, 100.41 mmol) and K.sub.2CO.sub.3 (23.09 g, 163.36 mmol) in CH.sub.3CN (200 mL) was stirred at 80° C. for 16 h. The reaction mixture was cooled to RT and diluted with ethyl acetate and washed with brine. The organic layer was dried over anhydrous Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (25.5 g, 80% yield).

tert-Butyl 4-(azetidi-3-yl)piperazine-1-carboxylate

(651) A mixture of tert-butyl 4-(1-benzhydrylazetidin-3-yl)piperazine-1-carboxylate (10.0 g, 24.57 mmol) and 10% Pd/C (2.5 g) in MeOH (100 mL) was stirred under H.sub.2 atmosphere at 50° C.

for 48 h. The reaction mixture was cooled to RT and filtered. The filtrate was diluted concentrated in vacuo to afford a crude desire product (6.7 g) as a colorless oil.

tert-Butyl 4-(1-acryloylazetid-3-yl)piperazine-1-carboxylate

(652) A mixture of tert-butyl 4-(azetidi-3-yl)piperazine-1-carboxylate (6.7 g, 27.80 mmol) and Et.sub.3N (8.43 g, 83.40 mmol) in DCM (100 mL) at 0° C., acryloyl chloride (3.77 g, 41.7 mmol) was added in portions and the resulting solution was stirred at RT for 1 h. The reaction mixture was diluted with DCM and washed with brine. The organic layer was dried over anhydrous Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified by column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (3.6 g, 49.66% yield, 2 steps). ¹H NMR (400 MHz, DMSO-d₆) δ: 6.30 (dd, J=10.4, 16.8 Hz, 1H), 6.09 (dd, J=2.4, 17.2 Hz, 1H), 5.67 (dd, J=2.4, 10.4 Hz, 1H), 4.22 (t, J=8, 1H), 4.03-4.00 (m, 1H), 3.94-3.90 (m, 1H), 3.75-3.70 (m, 1H), 3.32 (t, J=8.8 Hz, 4H), 3.10-3.18 (m, 1H), 2.22-2.30 (m, 1H), 1.40 (s, 9H).

Example 39

(653) ##STR00547##

Piperazine-2-carbonitrile

(654) To a mixture of tert-butyl 3-cyanopiperazine-1-carboxylate (200 mg, 0.95 mmol) in dichloromethane (10 mL), CF₃COOH (2 mL) was added and the resulting was stirred at RT for 1 h. The mixture was concentrated in vacuo to afford the crude product.

4-(4',6-Dichloro-4-methoxy-[1,1'-biphenyl]-3-carbonyl)piperazine-2-carbonitrile

(655) To a mixture of 4',6-dichloro-4-methoxy-[1,1'-biphenyl]-3-carboxylic acid (309 mg, 1.04 mmol), EDCI (272 mg, 1.43 mmol), HOBT (195 mg, 1.43 mmol), Et.sub.3N (288 mg, 2.85 mmol) in dichloromethane (10 mL) at 0° C., piperazine-2-carbonitrile was added at 0° C. and the resulting mixture was stirred at RT for 8 h. The mixture was partitioned between dichloromethane and water. The organic layer was washed brine, dried over MgSO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (225 mg, 61% yield). ESI-MS m/z: 444.3 [M+H].sup.+.

1-Acryloyl-4-(4',6-dichloro-4-hydroxy-[1,1'-biphenyl]-3-carbonyl)piperazine-2-carbonitrile (VI-37)

(656) The title compound was prepared from 4-(4',6-dichloro-4-methoxy-[1,1'-biphenyl]-3-carbonyl)piperazine-2-carbonitrile in two steps according to the procedure described in Example 17. ¹H NMR (400 MHz, CDCl₃) δ: 9.24 (s, 1H), 7.44-7.33 (m, 5H), 7.20 (s, 1H), 6.57-6.45 (m, 2H), 6.79 (s, 1H), 5.94-5.91 (m, 1H), 5.75 (s, 1H), 4.62-4.61 (m, 1H), 4.50-4.46 (m, 1H), 4.06 (s, 1H), 3.61 (s, 1H), 3.36-3.33 (m, 1H), 3.16-3.10 (m, 1H). ESI-MS m/z: 428.4 [M+H].sup.+.

Example 40

(657) ##STR00548## ##STR00549##

5-Bromo-4-chloro-2-methoxybenzenamine

(658) The title compound was prepared from 4-chloro-2-methoxy-1-nitrobenzene in two steps according to the procedure described in Example 4.

1-Bromo-2-chloro-5-iodo-4-methoxybenzene

(659) To a mixture of 5-bromo-4-chloro-2-methoxyaniline (3 g, 12.7 mmol) in 6N HCl (60 mL, 360 mmol) at 0° C., a solution of NaNO₂ (963 mg, 13.9 mmol) in water (20 mL) was added dropwise while keeping the internal temperature around 0° C. KI (10.5 g, 63.4 mmol) and CuI (4.8 g, 25.4 mmol) were dissolved in water (20 mL) and added to the stirred reaction mixture. The reaction was kept at 5° C. for 2 h. The reaction mixture was extracted with ethyl acetate. The combined organic layer was washed with water, Na.sub.2SO.sub.3 (aq, 10%) and brine, dried over anhydrous Na.sub.2SO.sub.4, an concentrated. The residue was purified by flash column chromatography on silica gel (ethyl acetate/petroleum ether=1:100) to afford the desired product (3.2 g, 73% yield).

3-(5-Bromo-4-chloro-2-methoxyphenyl)propanal

(660) A mixture of 1-bromo-2-chloro-5-iodo-4-methoxybenzene (3.2 g, 9.2 mmol), prop-2-en-1-ol (1.3 g, 23.0 mmol), Pd(OAc)₂ (206 mg, 0.9 mmol), TBAC (2.56 g, 9.2 mmol), NaHCO₃ (2.3 g, 27.6 mmol) in DMF (50 mL) was stirred under Argon at 60° C. for 16 h. The mixture was allowed to cool to RT, and then partitioned between ethyl acetate and water. The combined organic layer was washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by flash column chromatography on silica gel (ethyl acetate/petroleum ether=1:20) to afford the desired product (860 mg, 34% yield).

3-(5-Bromo-4-chloro-2-methoxyphenyl)propanoic acid

(661) To a stirred solution of Jones reagent (3 mL, 5.4 mmol, 2.8 M) in acetone (20 mL), 3-(5-bromo-4-chloro-2-methoxyphenyl)propanal (860 mg, 3.1 mmol) was added. The reaction was stirred at RT for 12 h, quenched with iso-propylalcohol and then stirred for 10 min. The resulting mixture was diluted with water, extracted with ethyl acetate. The combined organic layer was washed with water and brine, dried over Na₂SO₄ and concentrated. The residue was purified by flash column chromatography on silica gel (ethyl acetate/petroleum ether=1:1) to afford the desired product (358 mg, 38% yield). ESI-MS m/z: 291.1 [M+H]⁺.

tert-Butyl 3-(4-(3-(5-bromo-4-chloro-2-methoxyphenyl)propanoyl)piperazin-1-yl)azetidine-1-carboxylate

(662) To a stirred solution of 3-(5-bromo-4-chloro-2-methoxyphenyl)propanoic acid (350 mg, 1.2 mmol) in DMF (30 mL) at RT, tert-butyl 3-(piperazin-1-yl)azetidine-1-carboxylate (317 mg, 1.3 mmol), BOP (731 mg, 1.4 mmol) and DIEA (461 mg, 3.6 mmol) were added and the mixture was stirred at RT for 1 h. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (285 mg, 46% yield).

tert-Butyl 3-(4-(3-(4-chloro-5-cyclopropyl-2-hydroxyphenyl)propanoyl)piperazin-1-yl)azetidine-1-carboxylate

(663) A mixture of tert-butyl 3-(4-(3-(5-bromo-4-chloro-2-methoxyphenyl)propanoyl)piperazin-1-yl)azetidine-1-carboxylate (280 mg, 0.54 mmol), cyclopropylboronic acid (185 mg, 2.2 mmol), K₃PO₄ (444 mg, 1.9 mmol), tricyclohexylphosphine (30 mg, 0.1 mmol), Pd(OAc)₂ (24 mg, 0.11 mmol) in toluene (10 mL) and water (1 mL) was stirred at reflux under argon for 16 h. The mixture was allowed to cool to RT and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=60:1) to afford the desired product (194 mg, 75% yield). ESI-MS m/z: 477.3 [M+H]⁺.

1-(3-(4-(3-(4-Chloro-5-cyclopropyl-2-hydroxyphenyl)propanoyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (V-58)

(664) The title compound was prepared from tert-butyl 3-(4-(3-(4-chloro-5-cyclopropyl-2-hydroxyphenyl)propanoyl)piperazin-1-yl)azetidine-1-carboxylate in three steps according to the procedure described in Example 17. ¹H NMR (400 MHz, DMSO-d₆) δ: 9.68 (s, 1H), 6.79 (s, 1H), 6.74 (s, 1H), 6.33-6.26 (m, 1H), 6.12-6.07 (dd, J=1.9, 17.2 Hz, 1H), 5.68-5.65 (dd, J=2.0, 10.2 Hz, 1H), 4.24-4.20 (m, 1H), 4.05-4.01 (m, 1H), 3.94-3.90 (m, 1H), 3.76-3.72 (m, 1H), 3.45-3.42 (m, 4H), 3.13-3.11 (m, 1H), 2.69-2.65 (m, 2H), 2.53-2.51 (m, 2H), 2.25-2.23 (bs, 4H), 1.97-1.93 (m, 1H), 0.90-0.86 (m, 1H), 0.60-0.55 (m, 1H). ESI-MS m/z: 418.4 [M+H]⁺.

Example 41

(665) ##STR00550## ##STR00551##

Methyl 5-acetamido-2-chloro-4-methoxybenzoate

(666) To a mixture of methyl 5-amino-2-chloro-4-methoxybenzoate (3.6 g, 16.7 mmol), Et₃N (6.7 g, 66.8 mmol) and DCM (100 mL) at RT, acetyl chloride (1.57 g, 20.1 mmol) was added dropwise and the resulting mixture was stirred for 12 h. The reaction mixture was partitioned

between dichloromethane and water. The organic layer was washed with saturated NaHCO₃ solution and brine, dried over anhydrous Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (ethyl acetate/petroleum ether=1:1) to afford the desired product (2.7 g, 63% yield).

N-(4-Chloro-5-(2-hydroxypropan-2-yl)-2-methoxyphenyl)acetamide

(667) To a solution of methyl 5-acetamido-2-chloro-4-methoxybenzoate (2.7 g, 11.1 mmol), in THF (40 mL) at -40° C. under Argon, methylmagnesium bromide (21 mL, 21 mmol, 1M in ether) was added dropwise while keeping the internal temperature at -40° C. Then the mixture was allowed to warm to RT, and stirred for 2 h. The reaction mixture was poured into ice-cooled NH₄Cl (10%) solution, and extracted with ethyl acetate. The combined organic layer was washed with water and brine, dried over anhydrous Na₂SO₄ and concentrated in vacuo to afford the desired product (2.3 g, 80% yield).

N-(4-Chloro-2-methoxy-5-(prop-1-en-2-yl)phenyl)acetamide

(668) To a solution of N-(4-chloro-5-(2-hydroxypropan-2-yl)-2-methoxyphenyl)acetamide (3.2 g, 12.4 mmol) in DCM (20 mL) at -5° C., SOCl₂ (3.7 g, 37.25 mmol) was added dropwise. The mixture was warmed to RT, and then stirred at reflux for 2 h. The reaction mixture was concentrated and the residue was purified by flash column chromatography on silica gel (ethyl acetate/petroleum ether=3:1) to afford the desired product (1.9 g, 64% yield).

N-(4-Chloro-2-methoxy-5-(1-methylcyclopropyl)phenyl)acetamide

(669) To a solution of N-(4-chloro-2-methoxy-5-(prop-1-en-2-yl)phenyl)acetamide (1.0 g, 4.17 mmol) in toluene (20 mL) at 0° C., CH₂I₂ (5.6 g, 20.86 mmol) and Et₂Zn (41.7 mL, 41.7 mmol, 1.0 M in hexane) was added. The mixture was kept at 0° C. for 30 min, and then stirred at RT for 16 h. The reaction mixture was quenched with saturated NH₄Cl solution and stirred for 15 min. The mixture was concentrated in vacuo to remove toluene and the resulting mixture was extracted with dichloromethane. The combined organic layer was washed with water and brine, dried over anhydrous Na₂SO₄, concentrated to afford the desired product (820 mg, 77% yield).

4-Chloro-2-methoxy-5-(1-methylcyclopropyl)aniline

(670) A mixture of N-(4-chloro-2-methoxy-5-(1-methylcyclopropyl)phenyl)acetamide (820 mg, 3.23 mmol), KOH (1.8 g, 32.3 mmol), ethanol (40 mL) and water (20 mL) was stirred at reflux for 12 h. The reaction mixture was extracted with ethyl acetate. The combined organic layer was washed with water and brine, dried over anhydrous Na₂SO₄ and filtered. The filtrate was concentrated in vacuo and the residue was purified by flash column chromatography on silica gel (ethyl acetate/petroleum ether=20:1) to afford the desired product (460 mg, 67% yield). ESI-MS m/z: 212.4 [M+H]⁺.

Ethyl 2-((4-chloro-2-methoxy-5-(1-methylcyclopropyl)phenyl)amino)acetate

(671) To a solution of 4-chloro-2-methoxy-5-(1-methylcyclopropyl)aniline (450 mg, 2.13 mmol) in MeOH (20 mL) at RT, AcOH (3 drops) and ethyl glyoxalate (326 mg, 3.19 mmol, 50% in toluene) were added. The mixture was stirred at RT for 2 h and then sodium cyanoborohydride (403 mg, 6.39 mmol) was added to the mixture. The resulting mixture was stirred at 50° C. for 16 h. The mixture was allowed to cool to RT, and partitioned between ethyl acetate and water. The organic layer was dried over MgSO₄, filtered, and concentrated in vacuo to afford the crude product (636 mg). ESI-MS m/z: 298.2 [M+H]⁺.

2-((4-Chloro-2-methoxy-5-(1-methylcyclopropyl)phenyl)amino)acetic acid

(672) To a solution of ethyl 2-((4-chloro-2-methoxy-5-(1-methylcyclopropyl)phenyl)amino)acetate (630 mg, 2.12 mmol) in THF (15 mL) and water (5 mL), LiOH.H₂O (889 mg, 21.2 mmol) was added and the resulting mixture was stirred at RT for 2 h. The mixture was washed with 20% ethyl acetate/petroleum ether. The aqueous layer was acidified with aqueous HCl (1 N) to adjust pH to 3-4 and extracted with ethyl acetate. The organic layer was dried over MgSO₄, filtered and concentrated in vacuo to afford the desired product (200 mg, 33% yield).

tert-Butyl 3-(4-(2-((4-chloro-2-methoxy-5-(1-methylcyclopropyl)phenyl)amino)acetyl)piperazin-1-yl)azetidine-1-carboxylate

(673) To a solution of 2-((4-chloro-2-methoxy-5-(1-methylcyclopropyl)phenyl)amino)acetic acid (110 mg, 0.41 mmol) and tert-butyl 3-(piperazin-1-yl)azetidine-1-carboxylate (118 mg, 0.49 mmol) in DMF (15 mL) at RT, BOP (217 mg, 0.49 mmol) and DIEA (159 mg, 1.23 mmol) were added and the resulting mixture was stirred at RT for 1 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over MgSO_4 , filtered and concentrated in vacuo to afford the desired product (192 mg, 95% yield).

1-(3-(4-(2-((4-Chloro-2-hydroxy-5-(1-methylcyclopropyl)phenyl)amino)acetyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (V-64)

(674) The title compound was prepared from tert-butyl 3-(4-(2-((4-chloro-2-methoxy-5-(1-methylcyclopropyl)phenyl)amino)acetyl)piperazin-1-yl)azetidine-1-carboxylate in three steps according to the procedure described in Example 17. ^1H NMR (400 MHz, DMSO- d_6) δ : 9.70 (s, 1H), 6.64 (s, 1H), 6.51 (s, 1H), 6.35-6.28 (m, 1H), 6.13-6.08 (dd, $J=1.9, 17.9$ Hz, 1H), 5.69-5.66 (dd, $J=2.1, 10.1$ Hz, 1H), 5.13-5.11 (m, 1H), 4.25-4.23 (m, 1H), 4.08-4.05 (m, 1H), 3.95-3.91 (m, 3H), 3.80-3.76 (m, 1H), 3.53 (bs, 4H), 3.18-3.16 (m, 1H), 2.38-2.31 (m, 4H), 1.26 (s, 3H), 0.72-0.64 (m, 4H). ESI-MS m/z : 434.4 $[\text{M}+\text{H}]^+$.

Example 42

(675) ##STR00552##

5-Chloro-4-iodo-2-methoxybenzoic acid

(676) To a stirred solution of 4-amino-5-chloro-2-methoxybenzoic acid (5 g, 24.8 mmol) in water (10 mL) at 0°C ., concentrated sulfuric acid (50 mL) was added. Then a solution of NaNO_2 (1.9 g, 27.3 mmol) in water (10 mL) was added dropwise while keeping the internal temperature around 0°C . KI (4.5 g, 27.3 mmol) and I_2 (3.5 g, 13.64 mmol) were dissolved in water and added dropwise to the stirred reaction mixture. The reaction was stirred at 5°C . for 2 h and then extracted with ethyl acetate. The organic layer was washed with water, Na_2SO_3 (aq, 10%) and brine, dried over anhydrous Na_2SO_4 and concentrated in vacuo to afford desired product (1.55 g, 19% yield). ESI-MS m/z : 311.1 $[\text{M}+\text{H}]^+$.

tert-Butyl 4-(5-chloro-4-iodo-2-methoxybenzoyl)piperazine-1-carboxylate

(677) To a stirred solution of 5-chloro-4-iodo-2-methoxybenzoic acid (1.55 g, 4.9 mmol) in DMF (30 mL) at RT, tert-butyl piperazine-1-carboxylate (1.02 g, 5.5 mmol), BOP (2.63 g, 25.9 mmol) and DIEA (1.92 g, 14.9 mmol) were added and the resulting mixture was stirred at RT for 1 h. The reaction mixture was extracted with ethyl acetate and washed with brine. The organic layer was dried over anhydrous Na_2SO_4 , filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (1.96 g, 76% yield).

tert-Butyl-4-(2-chloro-5-methoxy-[1,1'-biphenyl]-4-carbonyl)piperazine-1-carboxylate

(678) A mixture of tert-butyl 4-(5-chloro-4-iodo-2-methoxybenzoyl)piperazine-1-carboxylate (300 mg, 0.56 mmol), phenylboronic acid (82 mg, 0.67 mmol), $\text{Pd}(\text{PPh}_3)_4$ (129 mg, 0.1 mmol), Na_2CO_3 (180 mg, 1.68 mmol) in 1,4-dioxane (10 mL) and water (2 mL) was stirred at reflux under argon for 16 h. The mixture was allowed to cool to room temperature and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=30:1) to afford the desired product (219 mg, 80% yield).

1-(4-(2-Chloro-5-hydroxy-[1,1'-biphenyl]-4-carbonyl)piperazin-1-yl)prop-2-en-1-one (VI-42)

(679) The title compound was prepared from tert-butyl-4-(2-chloro-5-methoxy-[1,1'-biphenyl]-4-carbonyl)piperazine-1-carboxylate in three steps according to the procedure described in Example 17. ^1H NMR (400 MHz, DMSO- d_6) δ : 10.40 (s, 1H), 7.50-7.41 (m, 5H), 7.35 (s, 1H), 6.89 (s, 1H), 6.84 (m, 1H), 6.17-6.13 (d, 1H), 5.73-5.71 (m, 1H), 3.63 (s, 6H), 3.30 (s, 2H). ESI-MS m/z : 371.2 $[\text{M}+\text{H}]^+$.

Example 43

(680) ##STR00553## ##STR00554##

4-Chloro-2-methoxy-5-(prop-1-en-2-yl)benzenamine

(681) A mixture of 4-chloro-5-iodo-2-methoxybenzenamine (1.0 g, 3.53 mmol), 4,4,5,5-tetramethyl-2-(prop-1-en-2-yl)-1,3,2-dioxaborolane (889 mg, 5.29 mmol), Pd(PPh.sub.3).sub.4 (363 mg, 0.353 mmol), Na.sub.2CO.sub.3 (1.12 g, 10.6 mmol) in DME (10 mL) and water (3 mL) was stirred at reflux under argon for 6 h. The reaction mixture was allowed to cool to RT and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (5% petroleum ether/ethyl acetate) to afford the desired product (173 mg, 25% yield) as an off-white solid. ESI-MS m/z: 198.5[M+H].sup.+.

4-Chloro-5-isopropyl-2-methoxybenzenamine

(682) A mixture of 4-chloro-2-methoxy-5-(prop-1-en-2-yl)benzenamine (160 mg, 0.81 mmol), Raney-Ni (20 mg) in MeOH (5 mL) was stirred at RT under H.sub.2 (1 atm) atmosphere for 8 h. The mixture was filtered and the filtrate was concentrated in vacuo to afford the desired product (150 mg, 93% yield).

tert-Butyl 3-(4-(2-(4-chloro-5-isopropyl-2-methoxyphenylamino)acetyl)piperazin-1-yl)azetidine-1-carboxylate

(683) The title compound was prepared from 4-chloro-5-isopropyl-2-methoxybenzenamine in three steps according to the procedure described in Example 42.

1-(3-(4-(2-(4-Chloro-5-isopropyl-2-methoxyphenylamino)acetyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one

(684) A mixture of tert-butyl 3-(4-(2-(4-chloro-5-isopropyl-2-methoxyphenylamino)acetyl)piperazin-1-yl)azetidine-1-carboxylate (102 mg, 0.212 mmol) in HCl/MeOH (2.86 M, 5 mL) was stirred at RT for 1 h. The mixture was concentrated in vacuo to afford the crude product, the crude was dissolved in DMF (5 mL) at RT, acrylic acid (17 mg, 0.233 mmol), BOP (113 mg, 0.254 mmol) and DIEA (82 mg, 0.636 mmol) were added and the resulting mixture was stirred at RT for 1 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed brine, dried over MgSO.sub.4, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (77 mg, 85% yield, 2 steps). ESI-MS m/z: 435.4 [M+H].sup.+.

1-(3-(4-(2-(4-Chloro-2-hydroxy-5-isopropylphenylamino)acetyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (V-50)

(685) To a solution of 1-(3-(4-(2-(4-chloro-5-isopropyl-2-methoxyphenylamino)acetyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (77 mg, 0.18 mmol) in DCM (15 mL) at -60° C., BBr.sub.3 (443 mg, 1.8 mmol) was added dropwise and the resulting mixture was stirred at RT for 1 h. The mixture was cooled to -60° C., MeOH was added dropwise and then basified with Et.sub.3N to adjust the pH to 8-9. The mixture was poured into water and extracted with dichloromethane. The organic layer was dried over anhydrous Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (25 mg, 33% yield). .sup.1H NMR (400 MHz, DMSO-d6) δ: 9.50 (bs, 1H), 6.62 (s, 1H), 6.470 (s, 1H), 6.30 (m, 1H), 6.10 (dd, J=2.4, 17.2 Hz, 1H), 5.68 (dd, J=2.0, 10.4 Hz, 1H), 5.14 (m, 1H), 4.25 (m, 1H), 4.06 (m, 1H), 3.96 (m, 1H), 3.91 (m, 2H), 3.78 (m, 1H), 3.54 (m, 4H), 3.17 (m, 2H), 2.35 (m, 4H), 1.21 (m, 6H). ESI-MS m/z: 421.4 [M+H].sup.+.

Example 44

(686) ##STR00555##

2-Amino-5-chloro-4-cyclopropylphenol

(687) To a mixture of 2-amino-5-chloro-4-iodophenol (500 mg, 1.9 mmol), PdCl.sub.2(dppf) (136 mg, 0.19 mmol) in THF (10 mL) under argon at RT, cyclopropylmagnesium bromide (16 mL, 11.4 mmol, 0.7 M in THF) was added and the mixture was stirred at reflux for 15 h. The mixture was

allowed to cool to RT, and partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated. The residue was purified by flash column chromatography on silica gel (10-20% ethyl acetate/hexanes) to afford the desired product (220 mg, 63% yield) as a brown solid. .sup.1H NMR (400 MHz, DMSO-d6) δ : 9.27 (s, 1H), 6.62 (s, 1H), 6.22 (s, 1H), 4.53 (s, 2H), 1.89-1.93 (m, 1H), 0.83-0.87 (m, 2H), 0.46-0.49 (m, 2H).

Ethyl 2-(4-chloro-5-cyclopropyl-2-hydroxyphenylamino)acetate

(688) To a solution of 2-amino-5-chloro-4-cyclopropylphenol (200 mg, 1.01 mmol) in MeOH (20 mL) at RT, AcOH (3 drops) and ethyl glyoxalate (416 mg, 2.02 mmol, 50% in toluene) were added. The mixture was stirred at RT for 2 h and then sodium cyanoborohydride (190 mg, 3.03 mmol) was added to the mixture. The resulting mixture was stirred at 40° C. for 15 h. The mixture was allowed to cool to RT and partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated. The residue was purified by flash column chromatography on silica gel (10-20% methanol/dichloromethane) to afford the desired product (290 mg, 100% yield) as a solid. .sup.1H NMR (400 MHz, DMSO-d6) δ : 9.63 (s, 1H), 6.66 (s, 1H), 5.93 (s, 1H), 5.07 (t, J=6.4 Hz, 1H), 4.12 (q, J=7.2 Hz, 2H), 3.91 (d, J=6.4 Hz, 2H), 1.92-1.97 (m, 1H), 1.20 (t, J=6.8 Hz, 2H), 0.84-0.87 (m, 2H), 0.51-0.55 (m, 2H).

2-(4-Chloro-5-cyclopropyl-2-hydroxyphenylamino)acetic acid

(689) To a solution of ethyl 2-(4-chloro-5-cyclopropyl-2-hydroxyphenylamino)acetate (290 mg, 0.89 mmol) in of 4:1 mixture of tetrahydrofuran and water (30 mL) at RT, LiOH.H.sub.2O (226 mg, 5.34 mmol) was added and the resulting mixture was stirred for 2 h at 60° C. The mixture was acidified with aqueous HCl (1 N) to adjust the pH to 3-5 and then extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous Na.sub.2SO.sub.4, filtered and concentrated in vacuo to afford the product (100 mg, 47% yield). .sup.1H NMR (400 MHz, DMSO-d6) δ : 9.64 (s, 1H), 6.66 (s, 1H), 5.96 (s, 1H), 3.81 (s, 2H), 1.89-1.96 (m, 1H), 0.84-0.87 (m, 2H), 0.54-0.56 (m, 2H).

tert-Butyl 4-(1-benzhydrylazetidin-3-yl)-3-carbamoylpiperazine-1-carboxylate

(690) A mixture of 1-benzhydrylazetidin-3-yl methanesulfonate (2.69 g, 8.5 mmol), K.sub.2CO.sub.3 (1.76 g, 12.8 mmol), tert-butyl 3-carbamoylpiperazine-1-carboxylate (1.95 g, 8.5 mmol) in CH.sub.3CN (40 mL) was stirred at reflux for 16 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with water and brine, dried over Na.sub.2SO.sub.4, and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product. (2.08 g, 54% yield).

tert-Butyl 4-(azetidin-3-yl)-3-carbamoylpiperazine-1-carboxylate

(691) A mixture of 4-chloro-2-methoxy-5-(prop-1-en-2-yl)benzenamine (1 g, 2.22 mmol), Pd/C (300 mg) in MeOH (25 mL) was stirred at 50° C. under H.sub.2 (1 atm) atmosphere for 12 h. The mixture was cooled and filtered. The filtrate was concentrated in vacuo to afford the desired product (640 mg, 100% yield).

tert-Butyl 4-(1-acryloylazetidin-3-yl)-3-carbamoylpiperazine-1-carboxylate

(692) To a solution of tert-butyl 4-(azetidin-3-yl)-3-carbamoylpiperazine-1-carboxylate (640 mg, 2.22 mmol) and Et.sub.3N (463 mg, 4.58 mmol) in DCM (10 mL) at 0° C., acryloyl chloride (248 mg, 2.74 mmol) was added dropwise and the resulting mixture was stirred at RT for 1.5 h. The mixture was partitioned between dichloromethane and saturated NaHCO.sub.3 solution. The organic layer was washed with saturated brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (350 mg, 47% yield).

1-(1-acryloylazetidin-3-yl)-4-(2-(4-chloro-5-cyclopropyl-2-hydroxyphenylamino)acetyl)piperazine-2-carboxamide (V-51)

(693) A mixture of tert-butyl 4-(1-acryloylazetidin-3-yl)-3-carbamoylpiperazine-1-carboxylate (120

mg, 0.35 mmol) in HCl/MeOH (2.86 M, 10 mL) was stirred at RT for 1 h. The mixture was concentrated in vacuo to afford the crude residue. It was dissolved in DMF (5 mL) at 0° C., 2-(4-chloro-5-cyclopropyl-2-hydroxyphenylamino)acetic acid (31 mg, 0.427 mmol), BOP (206 mg, 0.466 mmol) and K₂CO₃ (150 mg, 1.164 mmol) were added and the resulting mixture was stirred at RT for 1 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed brine, dried over MgSO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=20:1) to afford the desired product (126 mg, 75% yield, 2 steps). ¹H NMR (400 MHz, DMSO-d₆) δ: 9.63 (bs, 1H), 7.53 (d, 1H), 7.26-6.94 (m, 2H), 6.67 (s, 1H), 6.34-6.27 (m, 1H), 6.10 (m, 2H), 5.68 (d, J=10.4, 1H), 5.08 (m, 1H), 4.30 (m, 2H), 3.93 (m, 6H), 3.52 (m, 2H), 3.29 (m, 1H), 3.15 (m, 1H), 3.06 (m, 1H), 2.47 (m, 1H), 1.98 (m, 1H), 0.87 (m, 2H), 0.64 (m, 2H). ESI-MS m/z: 462.5 [M+H]⁺.

Example 45

(694) ##STR00556##

4-(2-(4,5-Dichloro-2-methoxyphenylamino)acetyl)piperazine-2-carbonitrile

(695) To a mixture of 2-(4,5-dichloro-2-methoxyphenylamino)acetic acid (260 mg, 1.04 mmol), EDCI (273 mg, 1.43 mmol), HOBt (194 mg, 1.43 mmol) and Et₃N (288 mg 2.85 mmol) in DCM (10 mL) at 0° C., piperazine-2-carbonitrile was added. The resulting mixture was stirred at RT for 8 h. The mixture was partitioned between dichloromethane and water. The organic layer was washed with brine, dried over MgSO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the desired product (108 mg, 30% yield).

1-acryloyl-4-(2-(4,5-dichloro-2-hydroxyphenylamino)acetyl)piperazine-2-carbonitrile (VI-39)

(696) The title compound was prepared from 4-(2-(4,5-dichloro-2-methoxyphenylamino)acetyl)piperazine-2-carbonitrile in two steps according to the procedure described in Example 17. ¹H NMR (400 MHz, DMSO-d₆) δ: 10.3 (s, 1H), 6.90 (m, 1H), 6.87 (s, 1H), 6.73 (s, 1H), 6.28 (d, J=16.4 Hz, 1H), 5.88 (d, J=10.4 Hz, 1H), 5.68 (m, 1H), 5.33 (m, 1H), 4.66 (m, 1H), 4.40 (m, 1H), 4.02 (m, 2H), 3.22 (m, 1H), 3.04 (m, 1H), 2.91 (m, 1H). ESI-MS m/z: 383.2 [M+H]⁺.

Example 46

(697) ##STR00557##

1-tert-Butyl 3-methyl 4-(1-benzhydrylazetidin-3-yl)piperazine-1,3-dicarboxylate

(698) A mixture of 1-benzhydrylazetidin-3-yl methanesulfonate (2.4 g, 7.56 mmol), tert-butyl methyl piperazine-1,3-dicarboxylate (1.85 g, 7.56 mmol), K₂CO₃ (1.6 g, 11.34 mmol) in CH₃CN (40 mL) was stirred at reflux for 16 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with water and brine, dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (10% petroleum ether/ethyl acetate) to afford the desired product (1.85 g, 51% yield).

tert-Butyl 4-(1-benzhydrylazetidin-3-yl)-3-(hydroxymethyl)piperazine-1-carboxylate

(699) To a mixture of LiAlH₄ (500 mg, 13.5 mmol) in THF (40 mL) at -40° C. under argon, a solution of 1-tert-butyl 3-methyl 4-(1-benzhydrylazetidin-3-yl) piperazine-1,3-dicarboxylate (1.8 g, 3.87 mmol) in THF (10 mL) was added dropwise. The reaction mixture was stirred at -5° C. to 5° C. for 1 h and cooled to -20° C. Then water (2 mL) and NaOH (15%) aqueous were added. The resulting mixture was stirred for 15 min. The solid was filtered, and the cake rinsed with ethyl acetate. The combined filtrate was dried over Na₂SO₄ and concentrated in vacuo to afford the product (1.6 g, 94% yield).

1-(3-(4-(2-(4-Chloro-5-cyclopropyl-2-hydroxyphenylamino)acetyl)-2-(hydroxymethyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (V-54)

(700) The title compound was prepared from 4-(2-(4,5-dichloro-2-

methoxyphenylamino)acetyl)piperazine-2-carbonitrile in four steps according to the procedure described in Example 44. ¹H NMR (400 MHz, DMSO-d₆) δ: 9.68 (bs, 1H), 6.67 (s, 1H), 6.35-6.27 (m, 1H), 6.12-6.05 (m, 2H), 5.67 (dd, J=1.6, 10.4 Hz, 1H), 5.11 (m, 1H), 4.82-4.63 (m, 1H), 4.24 (m, 1H), 4.13 (m, 1H), 3.95 (m, 1H), 3.88 (m, 2H), 3.85 (m, 1H), 3.77-3.67 (m, 2H), 3.20 (m, 1H), 3.15 (m, 1H), 2.76-2.60 (m, 2H), 2.40 (m, 1H), 1.95 (m, 1H), 0.87 (m, 2H), 0.62 (m, 2H). ESI-MS m/z: 449.4 [M+H]⁺.

Example 47

(701) ##STR00558##

Ethyl 2-(5,6-dichloro-1H-indol-3-yl)acetate

(702) To a mixture of 5,6-dichloro-1H-indole (1.0 g, 5.37 mmol), Cu(OTf)₂ (194 mg, 0.537 mmol) in DCM (15 mL) at RT, ethyl 2-diazoacetate (918 mg, 8.05 mmol) was added dropwise. The resulting mixture was stirred at RT for 16 h, quenched with water, and then extracted dichloromethane. The organic layer was dried over Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by Prep-HPLC to afford the desired product (120 mg, 8.2% yield) as light yellow solid. ESI-MS m/z: 272.1 [M+H]⁺.

2-(5,6-Dichloro-1H-indol-3-yl)acetic acid

(703) A mixture of ethyl 2-(5,6-dichloro-1H-indol-3-yl)acetate (120 mg, 0.44 mmol), LiOH (90 mg, 2.20 mmol) in THF (3 mL) and H₂O (1 mL) was stirred at RT for 16 h. The solution was poured into water, adjusted pH to 3-4 with 1N HCl and extracted with ethyl acetate. The organic layer was dried over Na₂SO₄ and concentrated in vacuo to afford the desired product (90 mg, 84.5% yield) as a yellow solid.

1-(3-(4-(2-(5,6-Dichloro-1H-indol-3-yl)acetyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (V-61)

(704) A mixture of 2-(5,6-dichloro-1H-indol-3-yl)acetic acid (90 mg, 0.372 mmol), 1-(3-(piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (87 mg, 0.446 mmol), EDCI.HCl (107 mg, 0.558 mmol), HOBT (75 mg, 0.558 mmol) in DMF (3 mL) at 0° C., Et₃N (112 mg, 1.11 mmol) was added. The resulting mixture was stirred at RT for 16 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=30:1) to afford the desired product (12 mg, 7.66% yield) as an off-white solid. ¹H NMR (400 MHz, DMSO-d₆) δ: 11.19 (bs, 1H), 7.79 (s, 1H), 7.59 (s, 1H), 7.35 (d, 1H), 6.28 (dd, J=9.6, 16.8 Hz, 1H), 6.09 (dd, J=2.4, 17.2 Hz, 1H), 5.66 (dd, J=2.4, 10.4 Hz, 1H), 4.21-4.18 (m, 1H), 4.02-3.98 (m, 1H), 3.93-3.88 (m, 1H), 3.78 (s, 2H), 3.74-3.70 (m, 1H), 3.53-3.47 (m, 4H), 3.10-3.07 (m, 1H), 2.25-2.19 (m, 4H). ESI-MS m/z: 423.3 [M+1]⁺.

Example 48

(705) ##STR00559##

tert-Butyl 3-(4-(2-(4-bromo-5-chloro-2-methoxyphenylamino)acetyl)piperazin-1-yl)azetidine-1-carboxylate

(706) The title compound was prepared from 4-bromo-5-chloro-2-methoxybenzenamine in three steps according to the procedure described in Example 4.

tert-Butyl 3-(4-(2-(5-chloro-4-ethyl-2-methoxyphenylamino)acetyl)piperazin-1-yl)azetidine-1-carboxylate

(707) To a mixture of tert-butyl 3-(4-(2-(4-bromo-5-chloro-2-methoxyphenylamino)acetyl)piperazin-1-yl)azetidine-1-carboxylate (100 mg, 0.193 mmol), Pd(dppf)₂Cl₂ (29 mg, 0.04 mmol) and K₂CO₃ (55 mg, 0.386 mmol) in DMF (10 mL) at RT, Et₂Zn (0.8 mL, 0.8 mmol, 1.0 M in hexane) was added. The resulting mixture was stirred at 80° C. for 16 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (dichloromethane/methanol=50:1) to afford the crude product (100 mg). ESI-MS m/z: 467.5 [M+1]⁺.

1-(3-(4-(2-(5-Chloro-4-ethyl-2-hydroxyphenylamino)acetyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (V-52)

(708) The title compound was prepared from tert-butyl 3-(4-(2-(5-chloro-4-ethyl-2-methoxyphenylamino)acetyl)piperazin-1-yl)azetidine-1-carboxylate in 3 steps according to the procedure described in Example 17. ¹H NMR (400 MHz, DMSO-d₆) δ: 9.6 (s, 1H), 6.6 (s, 1H), 6.5 (s, 1H), 6.3 (dd, J=10.4, 17.2 Hz, 1H), 6.1 (dd, J=2.4, 17.2 Hz, 1H), 5.7 (dd, J=2.4, 10.4 Hz, 1H), 5.1 (t, J=4.4 Hz, 1H), 4.2 (t, J=8 Hz, 1H), 4.1 (dd, J=4.8, 8.8 Hz, 1H), 3.95 (dd, J=7.2, 10.0 Hz, 1H), 3.9 (d, J=4.4 Hz, 2H), 3.8 (dd, J=4.8, 10.4 Hz, 1H), 3.6-3.5 (m, 4H), 3.2-3.1 (m, 1H), 3.1-3.0 (m, 1H), 2.5-2.3 (m, 4H), 1.1 (t, J=7.2 Hz, 3H); ESI-MS m/z: 407.4 [M+H].⁺

Example 49

(709) ##STR00560##

tert-Butyl3-(4-(2-((4-chloro-5-ethyl-2-methoxyphenyl)amino)acetyl)piperazin-1-yl) azetidine-1-carboxylate

(710) The title compound was prepared from tert-butyl 3-(4-(2-(4-chloro-5-iodo-2-methoxyphenylamino)acetyl)piperazin-1-yl)azetidine-1-carboxylate in one step according to the procedure described in Example 50.

1-(3-(4-(2-(4-chloro-5-ethyl-2-hydroxyphenylamino)acetyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (V-55)

(711) The title compound was prepared from tert-butyl3-(4-(2-((4-chloro-5-ethyl-2-methoxyphenyl)amino)acetyl)piperazin-1-yl) azetidine-1-carboxylate in three steps according to the procedure described in Example 17. ¹H NMR (400 MHz, DMSO-d₆) δ: 9.67 (s, 1H), 6.66 (s, 1H), 6.47 (s, 1H), 6.30 (dd, J=10.5, 16.9 Hz, 1H), 6.12 (dd, J=1.7, 16.7 Hz, 1H), 5.69 (dd, J=1.7, 16.7 Hz, 1H), 5.1 (m, 1H), 4.26 (m, 1H), 4.07 (m, 1H), 3.96 (m, 1H), 3.88 (d, J=4.4, 2H), 3.78 (m, 1H), 3.53 (m, 4H), 3.17 (m, 1H), 2.54 (m, 2H), 2.37 (m, 4H), 1.14 (m, 3H). ESI-MS m/z: 407.3[M+H].⁺

Example 50

(712) ##STR00561##

(2-Chloro-4-methoxy-5-nitrophenyl)methanol

(713) To a solution of 2-chloro-4-methoxy-5-nitrobenzaldehyde (6.0 g, 29 mmol) in MeOH at 0° C. (50 mL), sodium borohydride (4.45 g, 117 mmol) was added in portions and the resulting mixture was stirred at RT for 30 min. The mixture was concentrated in vacuo. The residue was dissolved in ethyl acetate, washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=10:1) to afford the desired product (5.0 g, 78.4% yield).

1-(Bromomethyl)-2-chloro-4-methoxy-5-nitrobenzene

(714) To a solution of (2-chloro-4-methoxy-5-nitrophenyl)methanol (5.0 g, 23 mmol) in dichloromethane (50 mL) at 0° C., tribromophosphine (3.08 g, 11.5 mmol) was added in portions and the resulting mixture was stirred at RT for 2 h. The mixture was poured into ice-water and extracted with dichloromethane. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=10:1) to afford the desired product (3.5 g, 54.2% yield).

1-Chloro-5-methoxy-4-nitro-2-(2,2,2-trifluoroethyl)benzene

(715) A mixture of (2-chloro-4-methoxy-5-nitrophenyl)methanol (3.5 g, 12.5 mmol), methyl 2,2-difluoro-2-(fluorosulfonyl)acetate (4.8 g, 25 mmol), copper iodide (617 mg, 3.25 mmol) in NMP (20 mL) was stirred at 80° C. for 24 h under Argon. After cooled to RT, the reaction mixture was dissolved in ethyl acetate, washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=100:1) to afford the desired product (1.2 g, 36.4% yield). ¹H NMR (400 MHz, DMSO-d₆) δ: 8.15 (s, 1H), 7.60 (s, 1H), 3.98 (s, 3H), 3.89 (dd, J=1.7, 11.2 Hz, 2H).

4-Chloro-2-methoxy-5-(2,2,2-trifluoroethyl)aniline

(716) A mixture of 1-chloro-5-methoxy-4-nitro-2-(2,2,2-trifluoroethyl)benzene (1.2 g, 4.51 mmol), tin(II) chloride dehydrate (5.0 g, 22.5 mmol) in EtOH (20 mL) was stirred at reflux for 2 h. After cooled to RT, the reaction mixture was added saturated NaHCO₃ solution to adjusted pH to 7-8 and then extracted with ethyl acetate. The organic layer washed with brine, dried over Na₂SO₄ and concentrated in vacuo to afford the desired product (900 mg, 85% yield). ¹H NMR (400 MHz, DMSO-d₆) δ: 7.29 (s, 1H), 7.11 (s, 1H), 5.41 (s, 2H), 4.21 (s, 3H), 3.99 (dd, J=1.7, 11.2 Hz, 2H).

1-(3-(4-(2-((4-Chloro-2-hydroxy-5-(2,2,2-trifluoroethyl)phenyl)amino)acetyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (V-57)

(717) The title compound was prepared from 4-chloro-2-methoxy-5-(2,2,2-trifluoroethyl)aniline in six steps according to the procedure described in Example 41. ¹H NMR (400 MHz, DMSO-d₆) δ: 10.08 (s, 1H), 6.74 (s, 1H), 6.59 (s, 1H), 6.34 (dd, J=10.5, 16.9 Hz, 1H), 6.12 (dd, J=1.7, 16.7 Hz, 1H), 5.69 (dd, J=1.7, 16.7 Hz, 1H), 5.22 (m, 1H), 4.24 (m, 1H), 4.04 (m, 1H), 3.94 (m, 1H), 3.88 (d, J=4.4 Hz, 2H), 3.78 (m, 1H), 3.57 (m, 2H), 3.54 (m, 4H), 3.18 (m, 1H), 2.37 (m, 4H). ESI-MS m/z: 461.2[M+H]⁺.

Example 51

(718) ##STR00562##

4-Chloro-5-cyclopropyl-2-methoxybenzenamine

(719) A mixture of 4-chloro-5-iodo-2-methoxyaniline (5.0 g, 17.6 mmol), cyclopropylboronic acid (1.8 g, 21.1 mmol), Pd(OAc)₂ (314 mg, 1.4 mmol), tricyclohexylphosphine (500 mg, 17.6 mmol), K₃PO₄·3H₂O (16.4 g, 61.6 mmol) in toluene (62.5 mL) and H₂O (3 mL) was stirred at 80° C. under argon for 16 h. The mixture was allowed to cool to RT, and then partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=10:1) to afford the desired product (3.1 g, 88.5% yield). ESI-MS m/z: 198.2[M+H]⁺.

1-Chloro-2-cyclopropyl-4-iodo-5-methoxybenzene

(720) To a mixture of 4-chloro-5-cyclopropyl-2-methoxyaniline (2.2 g, 11.05 mmol), conc. HCl (12 mL) and water (12 mL) at 0° C., the solution of sodium nitrate (762.8 mg, 11.05 mmol) in water (2.5 mL) was added dropwise. After stirring at 0° C. for 15 min, a solution of KI (1.83 g, 11.05 mmol) in water (5 mL) was added dropwise. The resulting mixture was stirred at RT for 4 h, poured into water (20 mL) and then extracted with dichloromethane. The organic layer was washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (0-10% ethyl acetate/petroleum ether) to afford the desired product (680 mg, 20% yield) as a solid. ¹H NMR (400 MHz, DMSO-d₆) δ: 7.37 (s, 1H), 7.08 (s, 1H), 3.84 (s, 3H), 2.00 (m, 1H), 0.89 (m, 2H), 0.65 (m, 1H).

(E)-1-(4-(1-Acryloylazetidin-3-yl)piperazin-1-yl)-3-(4-chloro-5-cyclopropyl-2-methoxyphenyl)prop-2-en-1-one

(721) A mixture of 1-chloro-2-cyclopropyl-4-iodo-5-methoxybenzene (300 mg, 0.974 mmol), tert-butyl 3-(4-acryloylpiperazin-1-yl)azetidine-1-carboxylate (431 mg, 1.46 mmol), Pd(OAc)₂ (54.6 mg, 0.243 mmol), sodium acetate (239 mg, 292 mmol), tetrabutylammonium chloride (539 mg, 1.95 mmol) in DMF (7 mL) was stirred at 100° C. for 24 h. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo to purified by silica gel (dichloromethane/methanol=40:1) to afford the desired product (350 mg, 84% yield). ESI-MS m/z: 476.2 [M+H]⁺.

(E)-1-(4-(1-Acryloylazetidin-3-yl)piperazin-1-yl)-3-(4-chloro-5-cyclopropyl-2-hydroxyphenyl)prop-2-en-1-one (V-62)

(722) The title compound was prepared from (E)-1-(4-(1-acryloylazetidin-3-yl)piperazin-1-yl)-3-

(4-chloro-5-cyclopropyl-2-methoxyphenyl)prop-2-en-1-one in three steps according to the procedure described in Example 17. ¹H NMR (400 MHz, DMSO-d₆) δ: 10.3 (s, 1H), 7.71 (m, 1H), 7.30 (s, 1H), 7.22 (m, 1H), 6.93 (s, 1H), 6.34 (dd, J=10.5, 16.9 Hz, 1H), 6.12 (dd, J=1.7, 16.7 Hz, 1H), 5.66 (dd, J=1.7, 16.7 Hz, 1H), 4.26 (m, 1H), 4.08 (m, 1H), 3.94 (m, 1H), 3.79 (m, 1H), 3.69 (m, 2H), 3.58 (m, 2H), 3.18 (m, 1H), 2.33 (m, 4H), 1.99 (m, 1H), 0.92 (m, 2H), 0.73 (m, 2H). ESI-MS m/z: 416 [M+H].sup.+.

Example 52

(723) ##STR00563##

Methyl 2-((4-chloro-5-cyclopropyl-2-methoxyphenyl)thio)acetate

(724) A mixture of 1-chloro-2-cyclopropyl-4-iodo-5-methoxybenzene (380 mg, 1.23 mmol), Pd.sub.2(dba).sub.3 (56 mg, 0.061 mmol), methyl 2-mercaptoacetate (196 mg, 1.85 mmol), 1,1'-Bis(diphenylphosphino)ferrocene (136 mg, 0.246 mmol), Et.sub.3N (372 mg, 3.69 mmol) in NMP (8 mL) was stirred under argon at 80° C. for 24 h. After cooled to RT, the reaction mixture was extracted with ethyl acetate. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate=20:1) to afford the desired product (340 mg, 92% yield). .sup.1H NMR (400 MHz, DMSO-d₆) δ: 7.05 (s, 1H), 6.81 (s, 1H), 3.84 (s, 2H), 3.82 (s, 3H), 3.62 (s, 3H), 2.00 (m, 1H), 0.94 (m, 2H), 0.64 (m, 2H).

1-(3-(4-(2-(4-Chloro-5-cyclopropyl-2-methoxyphenylthio)acetyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (V-65)

(725) The title compound was prepared from methyl 2-((4-chloro-5-cyclopropyl-2-methoxyphenyl)thio)acetate in four steps according to the procedure described in Example 40.

.sup.1H NMR (400 MHz, DMSO-d₆) δ: 7.03 (s, 1H), 6.93 (s, 1H), 6.31 (dd, J=10.5, 16.9 Hz, 1H), 6.11 (dd, J=1.7, 16.7 Hz, 1H), 5.68 (dd, J=1.7, 16.7 Hz, 1H), 4.25 (m, 1H), 4.04 (m, 1H), 3.88 (s, 2H), 3.81 (s, 3H), 3.75 (m, 1H), 3.52 (m, 4H), 3.16 (m, 1H), 2.36-2.25 (m, 4H), 2.02 (m, 1H), 0.93 (m, 2H), 0.66 (m, 2H). ESI-MS m/z: 450 [M+H].sup.+.

Example 53

(726) ##STR00564##

(S)-2-(5-Bromo-4-chloro-2-methoxyphenylamino)propanoic acid

(727) A mixture of 1-bromo-2-chloro-5-iodo-4-methoxybenzene (3 g, 8.64 mmol), (S)-2-aminopropanoic acid (769 mg, 8.64 mmol), CuI (164 mg, 0.864 mmol), 2-hydroxybenzaldehyde phenylhydrazone (366 mg, 1.73 mmol), K.sub.3PO.sub.4.3H.sub.2O (4.6 g, 17.28 mmol) in DMF (10 mL) was stirred under argon at 80° C. for 16 h. The mixture was allowed to cool to RT, H.sub.2O and Et.sub.2O were added to the solution. The resulting solution was partitioned into two phases, the aqueous phase was separated, and the organic layer was extracted with 5% NaOH. The combined aqueous phase was acidified to pH 4 with 20% HCl, and then extracted with Et.sub.2O. The resulting organic layer was dried over MgSO.sub.4 and concentrated in vacuo to afford the desired product (1.7 g, 64% yield). ESI-MS m/z: 306.1 [M+H].sup.-

(S)-tert-Butyl 3-(4-(2-((5-bromo-4-chloro-2-methoxyphenyl)amino)propanoyl)piperazin-1-yl)azetidine-1-carboxylate

(728) To a solution of (S)-2-(5-bromo-4-chloro-2-methoxyphenylamino)propanoic acid (1.6 g, 5.21 mmol), tert-butyl 3-(piperazin-1-yl)azetidine-1-carboxylate (1.88 g, 7.82 mmol), EDCI.HCl (2.0 g, 10.42 mmol), HOBt (1.41 g, 10.42 mmol) in DMF (20 mL) at 0° C., Et.sub.3N (1.58 g, 15.63 mmol) was added. The resulting mixture was stirred at RT for 16 h and then partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (methanol/dichloroethane=1:50) to afford the desired product (2.1 g, 76% yield). ESI-MS m/z: 531.3 [M+H].sup.+.

((S)-tert-Butyl 3-(4-(2-((4-chloro-5-cyclopropyl-2-methoxyphenyl)amino)propanoyl)piperazin-1-yl)azetidine-1-carboxylate

(729) A mixture of (S)-tert-butyl 3-(4-(2-((5-bromo-4-chloro-2-methoxyphenyl)amino)propanoyl)piperazin-1-yl)azetidine-1-carboxylate (700 mg, 1.32 mmol), cyclopropylboronic acid (114 mg, 1.32 mmol), Pd(OAc)₂ (15 mg, 0.066 mmol), tricyclohexylphosphine (37 mg, 0.132 mmol), K₃PO₄·H₂O (974 mg, 4.62 mmol) in DMF (10 mL) and H₂O (0.5 mL) was stirred under argon at 80° C. for 16 h. The mixture was allowed to cool to RT, and then partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (methanol/dichloroethane=1:100) to afford the desired product (400 mg, 62%). ESI-MS m/z: 493.2[M+H]⁺.

(S)-1-(3-(4-(2-((4-Chloro-5-cyclopropyl-2-hydroxyphenyl)amino)propanoyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (V-67)

(730) The title compound was prepared from (S)-tert-butyl 3-(4-(2-((4-chloro-5-cyclopropyl-2-methoxyphenyl)amino)propanoyl)piperazin-1-yl)azetidine-1-carboxylate in three steps according to the procedure described in Example 17. ¹H NMR (400 MHz, DMSO-d₆) δ: 9.69 (s, 1H), 6.66 (s, 1H), 6.29 (dd, J=10.5, 16.9 Hz, 1H), 6.12 (dd, J=1.7, 16.7 Hz, 1H), 6.05 (s, 1H), 5.68 (dd, J=1.7, 16.7 Hz, 1H), 4.84 (m, 1H), 4.61 (m, 1H), 4.24 (m, 1H), 4.06 (m, 1H), 3.94 (m, 1H), 3.78 (m, 4H), 3.55 (m, 1H), 2.43-2.17 (m, 4H), 1.97 (m, 1H), 0.88 (m, 2H), 0.63 (m, 2H). ESI-MS m/z: 433.3 [M+H]⁺.

Example 54

(731) ##STR00565##

(S)-tert-Butyl 3-(4-(2-((4-chloro-5-ethyl-2-methoxyphenyl)amino)propanoyl)piperazin-1-yl)azetidine-1-carboxylate

(732) To a solution of (S)-tert-butyl 3-(4-(2-((5-bromo-4-chloro-2-methoxyphenyl)amino)propanoyl)piperazin-1-yl)azetidine-1-carboxylate (400 mg, 0.75 mmol), PdCl₂(dppf) (95 mg, 0.13 mmol) in THF (20 mL) at RT, Et₂Zn (2.86 mL, 2.86 mmol, 1.0 M in hexane) was added. The resulting mixture was stirred under argon at 80° C. for 4 h and then partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (methanol/dichloroethane=1:80) to afford the desired product (250 mg, 69% yield). ESI-MS m/z: 481.2 [M+H]⁺.

(S)-1-(3-(4-(2-((4-Chloro-5-ethyl-2-hydroxyphenyl)amino)propanoyl)piperazin-1-yl)azetidin-1-yl)prop-2-en-1-one (V-69)

(733) The title compound was prepared from (S)-tert-butyl 3-(4-(2-((4-chloro-5-ethyl-2-methoxyphenyl)amino)propanoyl)piperazin-1-yl)azetidine-1-carboxylate in three steps according to the procedure described in Example 17. ¹H NMR (400 MHz, DMSO-d₆) δ: 9.61 (s, 1H), 6.64 (s, 1H), 6.48 (s, 1H), 6.29 (dd, J=10.5, 16.9 Hz, 1H), 6.12 (dd, J=1.7, 16.7 Hz, 1H), 5.68 (dd, J=1.7, 16.7 Hz, 1H), 4.89 (m, 1H), 4.61 (m, 1H), 4.25 (m, 1H), 4.06 (m, 1H), 3.94 (m, 1H), 3.72-3.53 (m, 4H), 3.16 (m, 1H), 2.5 (m, 2H), 2.43-2.17 (m, 4H), 1.21 (dd, 3H), 1.15 (m, 3H). ESI-MS m/z: 406.2 [M+H]⁺.

Example 55

(734) ##STR00566##

tert-Butyl 4-(1-acryloylazetidin-3-yl)-3-methylpiperazine-1-carboxylate

(735) The title compound was prepared from tert-butyl 3-methylpiperazine-1-carboxylate in three steps according to the procedure described in Example 38.

1-(3-(2-Methylpiperazin-1-yl)azetidin-1-yl)prop-2-en-1-one hydrochloride

(736) The mixture of tert-butyl 4-(1-acryloylazetidin-3-yl)-3-methylpiperazine-1-carboxylate (62 mg, 0.199 mmol) in MeOH/HCl (20 mL, 2.9 M) was stirred at RT for 1 h. The mixture was concentrated in vacuo to afford the crude product (59 mg). The crude product was used directly in the next step without further purification.

1-(3-(4-(2-((4-Chloro-5-cyclopropyl-2-hydroxyphenyl)amino)acetyl)-2-methylpiperazin-1-

yl)azetidin-1-yl)prop-2-en-1-one (V-60)

(737) To the mixture of 2-((4-chloro-5-cyclopropyl-2-hydroxyphenyl)amino)acetic acid (30 mg, 0.124 mmol) and NMM (50 mg, 0.496 mmol) in dry THF (30 mL) at -10°C ., ethyl chloroformate (15 mg, 0.136 mmol) was added and the resulting mixture was stirred at -10°C . for 45 min. Then it was added a mixture of 1-(3-(2-methylpiperazin-1-yl)azetidin-1-yl)prop-2-en-1-one hydrochloride (37 mg, 0.149 mmol), Et.sub.3N (50 mg, 0.496 mmol) and dichloromethane (3 mL). The resulting mixture was stirred at RT for 30 min. The mixture was partitioned between ethyl acetate and water. The organic layer was washed with saturated NaHCO.sub.3 solution and brine, dried over Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified with column chromatography on silica gel (dichloromethane/methanol=40:1) to afford the desired product (10 mg, 186% yield) as a white solid. ¹H NMR (400 MHz, DMSO-d₆) δ : 9.65 (s, 1H), 6.66 (s, 1H), 6.34-6.27 (m, 1H), 6.10-6.07 (m, 2H), 5.68-5.65 (d, J=10.4 Hz, 1H), 5.12 (m, 1H), 4.29-4.19 (m, 1H), 4.12-4.10 (m, 1H), 4.08-3.81 (m, 4H), 3.78 (s, 4H), 2.63 (m, 2H), 2.25 (m, 1H), 1.96 (m, 1H), 1.24 (s, 1H), 0.96-0.87 (m, 5H), 0.63 (m, 2H). ESI-MS m/z: 433.5 [M+H].sup.+.

Example 56

(738) ##STR00567##

4-Bromo-5-chloro-2-nitrobenzamide

(739) A mixture of 4-bromo-5-chloro-2-nitrobenzoic acid (1.3 g, 4.63 mmol), Et.sub.3N (1.4 g, 13.9 mmol) in THF (20 mL) at 0°C ., ethyl chloroformate (1.5 g, 13.9 mmol) was added. The resulting mixture was stirred at 0°C . for 1 h. Then NH.sub.3.H.sub.2O (4 mL) was added and stirred for 0.5 h. The mixture was extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous Na.sub.2SO.sub.4, filtered and concentrated in vacuo to afford the crude product (900 mg).

2-Amino-4-bromo-5-chlorobenzamide

(740) To a solution of 4-bromo-5-chloro-2-nitrobenzamide (900 mg, 3.2 mmol) in AcOH (20 mL) and water (5 mL) at 70°C ., Fe powder (900 mg, 16.1 mmol) was added and the resulting mixture was stirred at 70°C . for 1 h. The mixture was allowed to cool to RT and poured into ice-water. The precipitate was collected by filtration and rinsed with water. This crude product was dissolved with ethyl acetate and filtered. The filtrate was washed with saturated NaHCO.sub.3 solution and brine. The organic layer was dried over MgSO.sub.4, filtered, and concentrated in vacuo to afford the desired product (770 mg, 97% yield). ESI-MS m/z: 250.1 [M+H].sup.+.

tert-Butyl 3-(4-(2-((5-bromo-2-carbamoyl-4-chlorophenyl)amino)acetyl)piperazin-1-yl)azetidine-1-carboxylate

(741) The title compound was prepared from 2-amino-4-bromo-5-chlorobenzamide in three steps according to the procedure described in Example 41. ESI-MS m/z: 532.5 [M+H].sup.+.

tert-Butyl 3-(4-(2-((2-carbamoyl-4-chloro-5-cyclopropylphenyl)amino)acetyl)piperazin-1-yl)azetidine-1-carboxylate

(742) To a solution of tert-butyl 3-(4-(2-((5-bromo-2-carbamoyl-4-chlorophenyl)amino)acetyl)piperazin-1-yl)azetidine-1-carboxylate (350 mg, 0.66 mmol) and cyclopropylboronic acid (226 mg, 2.64 mmol) in toluene (10 mL) and water (2 mL), Pd(OAc).sub.2 (15 mg, 0.07 mmol), PCy.sub.3 (37 mg, 0.132 mmol) and K.sub.3PO.sub.4 (487 mg, 2.31 mmol) were added. The mixture was stirred at 80°C . for 16 h. The mixture was allowed to cool to RT and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (1-5% methanol/dichloroethane) to afford the desired product (150 mg, 46% yield) as a solid.

2-((2-(4-(1-Acryloylazetidin-3-yl)piperazin-1-yl)-2-oxoethyl)amino)-5-chloro-4-cyclopropylbenzamide

(743) The title compound was prepared from tert-butyl 3-(4-(2-((2-carbamoyl-4-chloro-5-cyclopropylphenyl)amino)acetyl)piperazin-1-yl)azetidine-1-carboxylate in two steps according to the procedure described in Example 17. ESI-MS m/z: 446.4 [M+H].sup.+.

2-((2-(4-(1-Acryloylazetid-3-yl)piperazin-1-yl)-2-oxoethyl)amino)-5-chloro-4-cyclopropylbenzonitrile (V-71)

(744) A mixture of 2-((2-(4-(1-acryloylazetid-3-yl)piperazin-1-yl)-2-oxoethyl)amino)-5-chloro-4-cyclopropylbenzamide (30 mg, 0.067 mmol) and Et.sub.3N (41 mg, 0.404 mmol) in DCM (10 mL) at RT, trifluoroacetic anhydride (56 mg, 0.268 mmol) was added. The resulting mixture was stirred at RT for 0.5 h, poured into water and then extracted with dichloromethane. The organic layer was washed with brine, dried over Na.sub.2SO.sub.4 and concentrated. The residue was purified by flash column chromatography on silica gel (1-4% methanol/dichloroethane) to afford the desired product (20 mg, 72% yield). ¹H NMR (400 MHz, DMSO-d₆) δ: 7.60 (s, 1H), 6.34-6.30 (m, 1H), 6.27 (s, 1H), 6.12-6.07 (m, 1H), 6.01-5.99 (t, J=4 Hz, 1H), 5.69-5.65 (m, 1H), 4.26-4.22 (m, 1H), 4.07-4.04 (m, 3H), 3.96-3.92 (m, 1H), 3.80-3.76 (m, 1H), 3.53-3.51 (m, 4H), 3.19-3.13 (m, 1H), 2.45-2.30 (m, 4H), 2.16-2.09 (m, 1H), 1.08-1.03 (m, 2H), 0.87-0.80 (m, 2H). ESI-MS m/z: 428.4 [M+H].sup.+.

Example 57

(745) ##STR00568##

2-Chloro-4-methoxy-1-vinylbenzene

(746) To a suspension of phosphonium salt (2.05 g, 5 mmol) in THF (50 mL), was added t-BuOK (0.84 g, 7.5 mmol). The mixture turned to yellow and was kept stirring at RT for 1 h. 2-Chloro-4-methoxybenzaldehyde (0.85 g, 5 mmol) was added to the mixture. The mixture was stirred for 24 h, diluted with sat. NaHCO.sub.3 and then extracted with hexane. Organic layer was dried over Na.sub.2SO.sub.4, filtered and concentrated in vacuo. The residue was purified by Isolera One (100% hexanes to afford the desired product (0.45 g, 53% yield). ¹H NMR (CDCl.sub.3, δ): 7.49 (d, J=6.8 Hz, 1H), 7.03 (dd, J=8.8, 14.0 Hz, 1H), 6.90 (d, J=2.0, 1H), 6.79 (dd, J=2.0, 6.8 Hz, 1H), 5.62 (d, J=14.0 Hz, 1H), 5.26 (d, J=8.8 Hz, 1H), 3.80 (s, 3H).

2-Chloro-1-(2,2-difluorocyclopropyl)-4-methoxybenzene

(747) The solution of 2-chloro-4-methoxy-1-vinylbenzene (290 mg, 1.72 mmol) in dry THF (4 mL) was degassed, and then TMS-CF.sub.3 and NaI were added. The mixture was stirred at 80° C. overnight. TLC (100% Hexane) showed the reaction as complete. The mixture was diluted with hexane (20 mL). The inorganic salt was removed by filtration. The filtrate was concentrate in vacuo. The residue was purified via Isolera One (Hexane=100%).

1-Chloro-2-(2,2-difluorocyclopropyl)-5-methoxy-4-nitrobenzene

(748) To a solution of 2-chloro-1-(2,2-difluorocyclopropyl)-4-methoxybenzene (328 mg, 1.5 mmol) in Ac.sub.2O (2 mL), was added HNO.sub.3 (10 drops) at 0° C. The mixture was stirred from 0° C. to rt. Ac.sub.2O was removed in vacuo. The residue was diluted with DCM and washed with water. The organic layer was dried over Na.sub.2SO.sub.4. The solvent was removed in vacuo. The residue was purified via Isolera One (EtOAc/Hexane=0-15%) to afford the desired product. ¹H NMR (CDCl.sub.3, δ): 7.77 (s, 1H), 7.16 (s, 1H), 3.98 (s, 3H), 2.78-2.90 (m, 1H), 1.90-1.98 (m, 1H), 1.60-1.68 (m, 1H). ESI-MS m/z: 264.1 [M+H].sup.+.

4-Chloro-5-(2,2-difluorocyclopropyl)-2-methoxyaniline

(749) The above obtained 1-chloro-2-(2,2-difluorocyclopropyl)-5-methoxy-4-nitrobenzene was dissolved in 10 mL of co-solvent of AcOH/i-PrOH (1:5). Zn dust was added to the mixture. The mixture was stirred at 60° C. for 30 min. The solvent was removed in vacuo. The residue was diluted with DCM and the inorganic salt was removed by filtration. The filtrate was concentrated to give crude product which was used in next step without further purification.

1-(3-(4-((4-chloro-5-(2,2-difluorocyclopropyl)-2-hydroxyphenyl)glycyl)piperazin-1-yl)azetid-1-yl)prop-2-en-1-one (V-56)

(750) The title compound was prepared from 4-chloro-5-(2,2-difluorocyclopropyl)-2-methoxyaniline in 6 steps according to the procedure described in Example 41. ¹H NMR (CDCl.sub.3, δ): 9.90 (s, 1H), 6.73 (s, 1H), 6.40 (s, 1H), 6.30 (dd, J=8.4, 13.6 Hz, 1H), 6.10 (dd, J=1.6, 12.0 Hz, 1H), 5.66 (dd, J=1.6, 8.4 Hz, 1H), 5.18 (t, J=3.2, 3.6 Hz, 1H), 4.24 (t, J=6.0, 6.8

H_z, 1H), 4.03-4.08 (m, 1H), 3.86-3.97 (m, 3H), 3.74-3.80 (m, 1H), 3.52 (br. s, 4H), 3.13-3.20 (m, 1H), 2.77-2.87 (m, 1H), 2.25-2.43 (m, 4H), 1.87-1.97 (m, 2H). ESI-MS m/z: 455.2 [M+H].sup.-.

Example 58

Biochemical Assay of the Compounds

(751) Test compounds were prepared as 10 mM stock solutions in DMSO (Fisher cat #BP-231-100). KRAS G12C 1-169, his-tagged protein, GDP-loaded was diluted to 2 μM in buffer (20 mM Hepes, 150 mM NaCl, 1 mM MgCl₂). Compounds were tested for activity as follows:

(752) Compounds were diluted to 50× final test concentration in DMSO in 96-well storage plates. Compound stock solutions were vortexed before use and observed carefully for any sign of precipitation. Dilutions were as follow: For 100 μM final compound concentration, compounds were diluted to 5000 μM (5 μl 10 mM compound stock+5 μl DMSO and mixed well by pipetting. For 30 μM final compound concentration, compounds were diluted to 1500 μM (3 μl 10 mM compound stock+17 μl DMSO) and mixed well by pipetting. For 10 μM final compound concentration, compounds were diluted to 500 μM (2 μl 10 mM compound stock+38 μl DMSO) and mixed well by pipetting.

49 μl of the stock protein solution was added to each well of a 96-well PCR plate (Fisher cat #1423027). 1 μl of the diluted 50× compounds were added to appropriate wells in the PCR plate using 12-channel pipettor. Reactions were mixed carefully and thoroughly by pipetting up/down with a 200 μl multi-channel pipettor. The plate was sealed well with aluminum plate seal, and stored in a drawer at room temperature for 2 hrs or 24 hrs. 5 μl of 2% formic acid (Fisher cat #A117) in DI H₂O was then added to each well followed by mixing with a pipette. The plate was then resealed with aluminum seal and stored on dry ice until analyzed as described below.

(753) The above described assays were analyzed by mass spectrometry according to the following procedure:

(754) The MS instrument is set to positive polarity, 2 GHz resolution, and low mass (1700) mode and allowed to equilibrate for 30 minutes. The instrument is then calibrated, switched to acquisition mode and the appropriate method loaded.

(755) After another 30 minute equilibration time, a blank batch (i.e., buffer) is run to ensure equipment is operating properly. The samples are thawed at 37° C. for 10 minutes, briefly centrifuged, and transfer to the bench top. Wells A1 and H12 are spiked with 1 uL 500 uM internal standard peptide, and the plates centrifuged at 2000×g for 5 minutes. The method is then run and masses of each individual well recorded.

(756) The masses (for which integration data is desired) for each well are pasted into the platemap and exported from the analysis. Masses for the internal standards are exported as well. The data at 50 ppm is extracted for the +19 charge state, and identity of well A1 is assigned using the internal standard spike and integrated. Peak data is exported as a TOF list and the above steps are repeated individually, for the +20, 21, 22, 23, 24, and 25 charge states.

(757) Other in vitro analyses are as follows:

(758) Inhibition of Cell Growth:

(759) The ability of the subject compounds to inhibit Ras-mediated cell growth is assessed and demonstrated as follows. Cells expressing a wildtype or a mutant Ras are plated in white, clear bottom 96 well plates at a density of 5,000 cells per well. Cells are allowed to attach for about 2 hours after plating before a compound disclosed herein is added. After certain hours (e.g., 24 hours, 48 hours, or 72 hours of cell growth), cell proliferation is determined by measuring total ATP content using the Cell Titer Glo reagent (Promega) according to manufacturer's instructions. Proliferation EC₅₀s is determined by analyzing 8 point compound dose responses at half-log intervals decreasing from 100 M.

(760) Inhibition of Ras-Mediated Signaling Transduction:

(761) The ability of the compounds disclosed herein in inhibiting Ras-mediated signaling is assessed and demonstrated as follows. Cells expressing wild type or a mutant Ras (such as G12C,

G12V, or G12A) are treated with or without (control cells) a subject compound. Inhibition of Ras signaling by one or more subject compounds is demonstrated by a decrease in the steady-state level of phosphorylated MEK, and/or Raf binding in cells treated with the one or more of the subject compounds as compared to the control cells.

(762) Inhibition of Ras-Mediated Signaling Transduction:

(763) The ability of the compounds disclosed herein in inhibiting Ras-mediated signaling is assessed and demonstrated as follows. Cells expressing wild type or a mutant Ras (such as G12C, G12V, or G12A) are treated with or without (control cells) a subject compound. Inhibition of Ras signaling by one or more subject compounds is demonstrated by percentage binding of compound to the G12C mutated Ras protein in cells treated with the one or more of the subject compounds as compared to the control cells.

(764) Inhibition of Ras-Mediated Signaling Transduction:

(765) The ability of the compounds disclosed herein in inhibiting Ras-mediated signaling is assessed and demonstrated as follows. Cells expressing wild type or a mutant Ras (such as G12C, G12V, or G12A) are treated with or without (control cells) a subject compound. Inhibition of Ras signaling by one or more subject compounds is demonstrated by a decrease in binding of Ras complex to downstream signaling molecules (for example Raf) in cells treated with the one or more of the subject compounds as compared to the control cells.

(766) The compounds in Tables 1-6 were tested according to the above procedures. Results for the compounds of Tables 5 and 6 are presented below. Each of the compounds in Tables 5 and 6 were found to covalently bind to KRAS G12C to the extent of at least about 5% (i.e., at least about 5% of the protein present in the well was found to be covalently bound to test compound).

(767) TABLE-US-00008 TABLE 7a Activity of Representative Compounds* Binding Binding Binding No. % No. % No. % No. % V-1 ++++ V-2 +++ V-3 ++++ V-4 +++ V-5 + V-6 + V-7 ++ V-8 ++ V-9 + V-10 + V-11 ++++ V-12 +++ V-13 ++ V-14 +++ V-15 + V-16 + V-17 ++ V-18 ++++ V-19 +++ V-20 ++ V-21 + V-22 + V-23 + V-24 + V-25 + V-26 + V-27 ++ V-28 +++ V-29 + V-30 + V-31 ++++ V-32 ++++ V-33 + V-34 + V-35 ++++ V-36 ++++ V-37 ++ V-38 + V-39 +++ V-40 ++++ V-41 ++++ V-42 + V-43 ++++ V-44 + V-45 ++++ V-46 + V-47 + V-48 ++ V-49 ++++ N/A N/A N/A N/A N/A *Binding activity determined at 24 hrs. + indicates binding activity from 5% to 15% ++ indicates binding activity greater than 15% and up to 25% +++ indicates binding activity greater than 25% and up to 50% ++++ indicates binding activity greater than 50%

(768) TABLE-US-00009 TABLE 7b Activity of Representative Compounds* Binding Binding Binding No. % No. % No. % No. % V-50 ++ V-51 ++ V-52 + V-53 ++ V-54 ++ V-55 ++ V-56 ++ V-57 ++ V-58 + V-59 ++ V-60 ++ V-61 + V-62 + V-63 ++ V-64 ++ V-65 + V-66 + V-67 ++ V-68 ++ V-69 ++ V-70 ++ V-71 + V-72 + V-73 + V-74 + V-75 + V-76 N/A N/A N/A *Binding activity determined at 2 hrs. + indicates binding activity from 5% to 20% ++ indicates binding activity greater than 20%

(769) TABLE-US-00010 TABLE 8a Activity of Representative Compounds* Binding Binding Binding No. % No. % No. % No. % VI-1 + VI-2 ++ VI-3 ++++ VI-4 ++ VI-5 ++++ VI-6 ++ VI-7 +++ VI-8 + VI-9 + VI-10 + VI-11 + VI-12 +++ VI-13 + VI-14 + VI-15 + VI-16 + VI-17 ++++ VI-18 + VI-19 + VI-20 + VI-21 ++ VI-22 + VI-23 +++ VI-24 ++++ VI-25 +++ VI-26 ++ VI-27 +++ VI-28 ++ VI-29 +++ VI-30 ++ VI-31 + VI-32 ++++ VI-33 + VI-34 +++ VI-35 ++ VI-36 + *Binding activity determined at 24 hrs. + indicates binding activity from 5% to 10% ++ indicates binding activity greater than 10% and up to 20% +++ indicates binding activity greater than 20% and up to 30% ++++ indicates binding activity greater than 30%

(770) TABLE-US-00011 TABLE 8b Activity of Representative Compounds* Binding Binding Binding No. % No. % No. % No. % VI-37 ++ VI-38 ++ VI-39 ++ VI-40 + VI-41 + VI-42 + VI-43 + N/A N/A *Binding activity determined at 2 hrs. + indicates binding activity from 5% to 20% ++ indicates binding activity greater than 20%

Example 59

Assessing Inhibition of Cell Proliferation by Compound I-189

(771) Two cancer cell lines, NCI H441 (human lung adenocarcinoma cells comprising a G12V mutation) and MIA paca-2 (human pancreatic carcinoma comprising a G12C mutation) were used in this experiment. Both the cell lines were treated with compound I-189 at a concentration of 100 μ M, 30 μ M, 10 μ M and 3 μ M and cell potency was measured as described in Example 1. The results of this experiment are shown in FIG. 4.

Example 60

Comparison of Cell Proliferation Inhibition by Compound I-189, I-92 and I-94

(772) The results of these SAR studies are shown in FIG. 5. Two cancer cell lines, NCI H441 (human lung adenocarcinoma cells comprising a G12V mutation) and MIA paca-2 (human pancreatic carcinoma comprising a G12C mutation) were used in this experiment. Both the cell lines were treated with compound I-189 at a concentration of 100 M, 30 M, 10 μ M and 3 μ M and cell potency was measured. Similar experiments were performed with compounds I-192 and I-94.

Example 61

Comparison of Cell Proliferation Inhibition by Compound I-92 and I-95

(773) The results of these studies are shown in FIG. 6. Two cancer cell lines, NCI H441 (human lung adenocarcinoma cells comprising a G12V mutation) and MIA paca-2 (human pancreatic carcinoma comprising a G12C mutation) were used in this experiment. Both the cell lines were treated with compound I-192 at a concentration of 100 M, 30 M, 10 μ M and 3 μ M and cell potency was measured. Similar experiments were performed with compound I-95.

Example 62

Comparison of Cell Proliferation Inhibition by Compound I-66, I-45 and I-91

(774) The results of these studies are shown in FIG. 7. Two cancer cell lines, NCI H441 (human lung adenocarcinoma cells comprising a G12V mutation) and MIA paca-2 (human pancreatic carcinoma comprising a G12C mutation) were used in this experiment. Both the cell lines were treated with compound I-66 at a concentration of 100 M, 30 M, 10 μ M and 3 μ M and cell potency was measured. Similar experiments were performed with compounds I-45 and I-91.

Example 63

Comparison of Cell Proliferation Inhibition by Compound I-47, I-42 and I-60


(775) The results of these studies are shown in FIG. 8. Three cell lines, NCI H441 (human lung adenocarcinoma cells), NCI 1568 (lung adenocarcinoma cells) and MIA paca-2 (human pancreatic carcinoma) were used in this experiment. Both the cell lines were treated with compound I-66 at a concentration of 100 M, 30 M, 10 μ M and 3 M and cell potency was measured. Similar experiments were performed with compounds I-47, I-42 and I-60.

(776) The various embodiments described above can be combined to provide further embodiments. All of the U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet, including but not limited to U.S. Patent Application No. 61/852,123, filed Mar. 15, 2013 and U.S. Patent Application No. 61/889,480, filed Oct. 10, 2013, are incorporated herein by reference, in their entirety. Aspects of the embodiments can be modified, if necessary to employ concepts of the various patents, applications and publications to provide yet further embodiments. These and other changes can be made to the embodiments in light of the above-detailed description. In general, in the following claims, the terms used should not be construed to limit the claims to the specific embodiments disclosed in the specification and the claims, but should be construed to include all possible embodiments along with the full scope of equivalents to which such claims are entitled. Accordingly, the claims are not limited by the disclosure.

(777) From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be

made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

Claims

1. A method for treating a cancer mediated by a K-Ras G12C, H-Ras G12C or N-Ras G12C mutation, the method comprising administering an effective amount of a compound having the following structure (VI): ##STR00569## or a pharmaceutically acceptable salt, tautomer, stereoisomer, adduct, solvate, isotopically-labeled derivative or prodrug thereof, to a subject in need thereof, wherein: A is CR^{37b}, N or NR^{38a}; B is CR^{37c}, N, NR^{38b} or S; C is CR^{37d}, N, NR^{38c} or S; G³ and G⁴ are each independently N or CR, wherein R is H, cyano, halo or $[-C^{1-6}alkyl]$; $C^{1-6}alkyl$; L^{1a} is a bond, —NH—, alkylene or heteroalkylene; L² is a bond or alkylene; R^{32a} and R^{32b} are, at each occurrence, independently H, —OH, —NH₂, —CO₂H, cyano, cyanoalkyl, C¹⁻⁶alkyl, C³⁻⁸cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl; or R^{32a} and R^{32b} join to form a carbocyclic or heterocyclic ring; or R^{32a} is H, —OH, —NH₂, —CO₂H, cyano, C¹⁻⁶alkyl, C³⁻⁸cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R^{32b} joins with R^{33b} to form a carbocyclic or heterocyclic ring; R^{33a} and R^{33b} are, at each occurrence, independently H, —OH, —NH₂, —CO₂H, cyano, cyanoalkyl, C¹⁻⁶alkyl, C³⁻⁸cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl; or R^{33a} and R^{33b} join to form a carbocyclic or heterocyclic ring; or R^{33a} is H, —OH, —NH₂, —CO₂H, cyano, C¹⁻⁶alkyl, C³⁻⁸cycloalkyl, hydroxylalkyl, aminoalkyl, carboxylalkyl or aminocarbonyl and R^{33b} joins with R^{32b} to form a carbocyclic or heterocyclic ring; R^{37a}, R^{37b}, R^{37c}, R^{37d} and R^{37e} are each independently H, $[-halo, oxo, hydroxyl, cyano, aminocarbonyl, formyl, C^{1-6}alkyl, C^{1-6}alkylsulfonyl, C^{1-6}haloalkyl, C^{3-8}cycloalkyl, C^{1-6}alkoxy, C^{1-6}hydroxylalkyl, C^{1-6}alkoxyalkyl, C^{1-6}aminoalkyl, heterocyclyl or aryl]$; $-OH, fluoro, chloro, bromo, iodo, oxo, methyl, methoxy, heteroaryl or aryl$; R^{38a}, R^{38b} and R^{38c} are each independently H, C¹⁻⁶alkyl or aryl; n³ and n⁴ are each independently 1, 2 or 3; m is 0 or 1;  custom character is a single or double bond such that all valences are satisfied; and E is an electrophilic moiety capable of forming a covalent bond with the cysteine residue at position 12 of a K-Ras, H-Ras or N-Ras G12C mutant protein.
2. The method of claim 1, wherein the compound has one of the following structures (VIa), (VIb), (VIc), (VId), (VId'), (VId'), (VId'), (VId'), (VId'), (VId') or (VId'): ##STR00570## ##STR00571##
3. The method of claim 1, wherein the compound has one of the following structures (VIa'), (VIb'), (VIc'), (VId'), (VId'), (VId'), (VId') or (VId'): ##STR00572## ##STR00573## wherein: Q is C(=O)—, NR^{34C}(=O)—, —S(=O)₂— or —NR^{34S}(=O)₂—; R³⁴ is H, C¹⁻⁶alkyl or hydroxylalkyl;  custom character is a carbon-carbon double bond or a carbon-carbon triple bond; and R³⁵ and R³⁶ are each independently H, cyano, C¹⁻⁶alkyl, aminoalkyl, alkylaminoalkyl, or hydroxylalkyl or R³⁵ and R³⁶ join to form a carbocyclic or heterocyclic ring when  custom character is a double bond; or R³⁵ is absent and R³⁶ is H, C¹⁻⁶alkyl, aminoalkyl, alkylaminoalkyl or hydroxylalkyl when  custom character is a triple bond.
4. The method of claim 1, wherein G³ is N and G⁴ is CR.
5. The method of claim 1, wherein G³ is CR and G⁴ is N.
6. The method of claim 1, wherein G³ is N and G⁴ is N.
7. The method of claim 1, wherein n³ is 2 and n⁴ is 2.

8. The method of claim 1, wherein n.sup.3 is 1 and n.sup.4 is 1.
9. The method of claim 1, wherein n.sup.3 is 2 and n is 1.
10. The method of claim 1, wherein R.sup.37a, R.sup.37b, R.sup.37c, R.sup.37d and R.sup.37e are each independently H, —OH, halo, oxo, C.sub.1-C.sub.6alkyl, C.sub.1-C.sub.6alkoxy, heterocyclyl or aryl..].
11. The method of claim 10, wherein R.sup.37a, R.sup.37b, R.sup.37c, R.sup.37d and R.sup.37e are each independently H, —OH, fluoro, chloro, bromo, iodo, oxo, methyl, methoxy, heteroaryl or aryl..].
12. The method of claim 1, wherein R.sup.37a or R.sup.37e is aryl.
13. The method of claim 12, wherein aryl is phenyl.
14. The method of claim 12, where the aryl is substituted with one or more halo substituents.
15. The method of claim 14, wherein the halo substituents are selected from fluoro and chloro.
16. The method of claim 1, wherein R.sup.37a is heteroaryl.
17. The method of claim 16, wherein the heteroaryl is thiophenyl.
18. The method of claim 1, wherein R.sup.37a or R.sup.37e, or both, is fluoro, chloro, bromo or iodo.
19. The method of claim 18, wherein halo is chloro, bromo or iodo.
20. The method of claim 1, wherein R.sup.37a or R.sup.37e has one of the following structures:
##STR00574##
21. The method of claim 1, wherein R.sup.38a, R.sup.38b and R.sup.38c are each independently H or aryl.
22. The method of claim 21, wherein R.sup.38a, R.sup.38b and R.sup.38c are each independently H.
23. The method of claim 1, wherein R.sup.38c is aryl.
24. The method of claim 23, wherein the aryl is substituted with one or more halo substituents.
25. The method of claim 24, wherein halo is chloro.
26. The method of claim 3, wherein Q is —C(=O)—.
27. The method of claim 3, wherein each of R.sup.35 and R.sup.36 are H.
28. The method of claim 1, wherein E has one of the following structures: ##STR00575##
29. The method of claim 1, wherein L.sup.1a is a bond.
30. The method of claim 1, wherein L.sup.2 is a bond.
31. The method of claim 1, wherein at least one of R.sup.32a, R.sup.32b, R.sup.33a, or R.sup.33b is H.
32. The method of claim 1, wherein each of R.sup.32a, R.sup.32b, R.sup.33a or R.sup.33b is H.
33. The method of claim 1, wherein the compound has one of the following structures:
##STR00576## ##STR00577## ##STR00578## ##STR00579## ##STR00580## ##STR00581##
34. The method of claim 1, wherein the compound is formulated in a pharmaceutical composition comprising the compound and a pharmaceutically acceptable carrier.
35. The method of claim 34, wherein the pharmaceutical composition is formulated for oral administration.
36. The method of claim 34, wherein the pharmaceutical composition is formulated for injection.
37. The method of claim 1, further comprising administering an additional anticancer agent to the subject.
38. The method of claim 1, wherein the cancer is a hematological cancer, pancreatic cancer, MYH associated polyposis, colorectal cancer or lung cancer.
39. The method of claim 1, wherein the compound has the following structure:
##STR00582## .Iaddend.
40. The method of claim 39, wherein R.sup.32a, R.sup.32b, R.sup.33a and R.sup.33b are, at each occurrence, independently H, cyano, hydroxyalkyl or C.sub.1-C.sub.6alkyl..Iaddend.

.Iadd.41. The method of claim 39, wherein R.sup.32a, R.sup.32b, R.sup.33a and R.sup.33b are, at each occurrence, independently H, cyano, -CH.sub.2OH or -CH.sub.3..Iaddend.
