

(12) **Patent Application Publication**
LEE et al.

(43) **Pub. Date:** **Aug. 14, 2025**

Oct. 31, 2022 (KR) 10-2022-0142612

The present invention relates to a *Corynebacterium* sp. mutant microorganism producing L-glutamic acid and a method of producing L-glutamic acid using the same, and more specifically, to a novel nicotinamide mononucleotide transporter variant involved in the L-glutamic acid biosynthetic pathway, a polynucleotide, and a transformant, as well as a method of producing L-glutamic acid using the same. The nicotinamide mononucleotide transporter variant according to the present invention is obtained by substituting one or more amino acids in the amino acid sequence constituting nicotinamide mononucleotide transporter to change the enzymatic activity of the nicotinamide mononucleotide transporter, and a recombinant microorganism comprising the nicotinamide mononucleotide transporter variant is capable of efficiently producing L-glutamic acid.

Specification includes a Sequence Listing.





Fig. 1



**MODIFIED MICROORGANISM OF GENUS
CORYNEBACTERIUM PRODUCING
L-GLUTAMIC ACID AND METHOD FOR
PRODUCING L-GLUTAMIC ACID USING
SAME**

TECHNICAL FIELD

[0001] The present invention relates to a *Corynebacterium* sp. mutant microorganism producing L-glutamic acid and a method of producing L-glutamic acid using the same, and more specifically, to a novel nicotinamide mononucleotide transporter variant involved in the L-glutamic acid biosynthetic pathway, a polynucleotide, and a transformant, as well as a method of producing L-glutamic acid using the same.

BACKGROUND ART

[0002] L-glutamic acid is a typical amino acid that is produced by microbial fermentation. Monosodium L-glutamate (MSG) may increase the preference of foods such as meat, fish, chicken, vegetables, sauces, soups and seasonings by balancing and harmonizing the overall taste of the food, may enhance the taste of low-salt foods having a salt content reduced up to 30%, and thus is widely used as a household seasoning and a seasoning for the production of processed food.

[0003] In brief, regarding the pathway of L-glutamic acid fermentation, glucose mainly undergoes the glycolytic pathway (EMP), but a portion thereof is metabolized into two pyruvic acid molecules through the pentose phosphate pathway. Among these molecules, one molecule combines with CO₂ to form oxaloacetic acid, and the other molecule combines with acetyl CoA from pyruvic acid to form citric acid. Then, oxaloacetic acid and citric acid enter the citric acid cycle (TCA cycle) to form α -ketoglutaric acid. Here, since the TCA cycle lacks the metabolic pathway for the oxidation of α -ketoglutaric acid to succinic acid and isocitrate dehydrogenase and glutamate dehydrogenase are closely involved therein, reductive amination of α -ketoglutaric acid efficiently occurs, thus producing L-glutamic acid.

[0004] For the production of L-glutamic acid, either naturally occurring wild-type strains or mutant strains modified from the wild-type strains so as to have an increased ability to produce glutamic acid may be used. In recent years, in order to improve the efficiency of production of L-glutamic acid, there has been development of a variety of recombinant strains or mutant strains having excellent L-glutamic acid productivity by applying genetic recombination technology to microorganisms such as *Escherichia coli* and *Corynebacterium*, which are widely used in the production of useful substances such as amino acids and nucleic acids, and methods of producing L-glutamic acid using the same. In particular, there have been attempts to increase the production of L-glutamic acid by targeting genes such as enzymes, transcription factors and transport proteins, which are involved in the biosynthetic pathways of L-glutamic acid, or by inducing mutations in promoters that regulate the expression of these genes. However, there are dozens of types of proteins such as enzymes, transcription factors and transport proteins, which are involved directly or indirectly in the production of L-glutamic acid, and thus much research is still needed on the increase in L-glutamic acid productivity by changes in the activity of these proteins.

PRIOR ART DOCUMENTS

Patent Documents

[0005] U.S. Pat. No. 6,852,516

[0006] U.S. Pat. No. 6,962,805

DISCLOSURE

Technical Problem

[0007] An object of the present invention is to provide a novel nicotinamide mononucleotide transporter variant.

[0008] Another object of the present invention is to provide a polynucleotide encoding the variant.

[0009] Still another object of the present invention is to provide a transformant comprising the variant or polynucleotide.

[0010] Yet another object of the present invention is to provide a method of producing L-glutamic acid using the transformant.

Technical Solution

[0011] One aspect of the present invention provides a nicotinamide mononucleotide transporter variant consisting of the amino acid sequence of SEQ ID NO: 2 in which alanine at position 163 in the amino acid sequence of SEQ ID NO: 4 is substituted with threonine.

[0012] As used in the present invention, the term “nicotinamide mononucleotide transporter” is a transmembrane protein having homology to nicotinamide ribonucleoside (NR) uptake permease (PnuC), which is responsible for the phosphorylation, transport, and control of transcription of nicotinamide ribonucleoside (NR), and it may be involved in NAD⁺ (re)generation, thereby increasing the energy flux in central metabolic pathways such as the glycolytic pathway and the TCA cycle. The nicotinamide mononucleotide transporter may be a gene encoding nicotinamide mononucleotide transporter or sequence having substantial identity thereto. As used herein, the term “substantial identity” means that, when each gene sequence, i.e., a base sequence or nucleotide sequence, and any other nucleotide sequence are aligned to correspond to each other as much as possible and analyzed, the other nucleotide sequence has a sequence homology of at least 70%, at least 80%, at least 90%, or at least 98% with each nucleotide sequence.

[0013] The nicotinamide mononucleotide transporter in the present invention is encoded by the ncg10063 gene and comprises the amino acid sequence of SEQ ID NO: 4.

[0014] According to one embodiment of the present invention, the amino acid sequence of SEQ ID NO: 4 may be derived from a wild-type *Corynebacterium* sp. microorganism.

[0015] More specifically, the *Corynebacterium* sp. microorganism may be *Corynebacterium glutamicum*.

[0016] As used in the present invention, the term “variant” refers to a polypeptide which is obtained by conservative substitution, deletion, modification or addition of one or more amino acids at the N-terminus, C-terminus of and/or within the amino acid sequence of a specific protein and has an amino acid sequence different from that of the protein before mutation, but retains functions or properties of the protein before mutation. As used herein, the term “conservative substitution” means substituting one amino acid with another amino acid having similar structural and/or chemical

properties. The conservative substitution may have little or no impact on the activity of the protein or polypeptide. The amino acid is selected from among alanine (Ala), isoleucine (Ile), valine (Val), leucine (Leu), methionine (Met), asparagine (Asn), cysteine (Cys), glutamine (Gln), serine (Ser), threonine (Thr), phenylalanine (Phe), tryptophan (Trp), tyrosine (Tyr), aspartic acid (Asp), glutamic acid (Glu), arginine (Arg), histidine (His), lysine (Lys), glycine (Gly), and proline (Pro).

[0017] In addition, some variants include those in which one or more portions, such as an N-terminal leader sequence or transmembrane domain, have been removed, or those in which a portion has been removed from the N- and/or C-terminus of a mature protein.

[0018] The variant may have increased (enhanced), unchanged, or decreased (weakened) ability compared to that of the protein before mutation. Here, the term “increased or enhanced” includes: a case in which the activity of the protein itself has increased compared to the activity of the protein before mutation; a case in which the overall activity of the protein in the cell is higher than that in the wild-type strain or the strain expressing the protein before mutation due to increased expression or translation of the gene encoding the protein; and a combination thereof. In addition, the term “decreased or weakened” includes: a case in which the activity of the protein itself has decreased compared to the activity of the protein before mutation; a case in which the overall activity of the protein in the cell is lower than that in the wild-type strain or the strain expressing the protein before mutation due to reduced expression or translation of the gene encoding the protein; and a combination thereof. In the present invention, the term “variant” may be used interchangeably with terms such as variant type, modification, variant polypeptide, mutated protein, mutation, and the like.

[0019] The variant in invention may be a nicotinamide mononucleotide transporter variant consisting of the amino acid sequence of SEQ ID NO: 2 in which the amino acid alanine at position 163 in the amino acid sequence of SEQ ID NO: 4 is substituted with threonine.

[0020] Another aspect of the present invention provides a polynucleotide encoding the nicotinamide mononucleotide transporter variant.

[0021] As used in the present invention, the term “polynucleotide” refers to a DNA or RNA strand having a certain length or more, which is a long-chain polymer of nucleotides formed by linking nucleotide monomers via covalent bonds. More specifically, the term “polynucleotide” refers to a polynucleotide fragment encoding the variant.

[0022] The polynucleotide may comprise a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 2.

[0023] According to one embodiment of the present invention, the polynucleotide may comprise the nucleotide sequence represented by SEQ ID NO: 1.

[0024] Still another aspect of the present invention provides a vector comprising a polynucleotide encoding the nicotinamide mononucleotide transporter variant.

[0025] Yet another aspect of the present invention provides a transformant comprising the nicotinamide mononucleotide transporter variant or the polynucleotide.

[0026] As used in the present invention, the term “vector” refers to any type of nucleic acid sequence transfer structure that is used as a means for transferring and expressing a gene

of interest in a host cell. Unless otherwise specified, the term “vector” may mean one allowing the nucleic acid sequence contained therein to be expressed after insertion into the host cell genome and/or one allowing the nucleic acid sequence to be expressed independently. This vector comprises essential regulatory elements operably linked so that an inserted gene can be expressed. As used herein, the term “operably linked” means that a gene of interest and regulatory sequences thereof are functionally linked together in a manner enabling gene expression, and the “regulatory elements” include a promoter for initiating transcription, any operator sequence for regulating transcription, a sequence encoding suitable mRNA ribosome-binding sites, and a sequence for regulating termination of transcription and translation.

[0027] The vector in the present invention is not particularly limited as long as it may replicate in a host cell, and any vector known in the art may be used. Examples of the vector include a natural or recombinant plasmid, cosmid, virus and bacteriophage. Examples of a phage vector or cosmid vector include, but are not limited to, pWE15, M13, λ MBL3, λ MBL4, λ IXII, λ ASHII, λ APII, λ t10, λ t11, Charon4A, and Charon21A, and examples of a plasmid vector include, but are not limited to, pBR series, pUC series, pBluescriptII series, pGEM series, pTZ series, pCL series, and pET series.

[0028] The vector may typically be constructed as a vector for cloning or as a vector for expression. The vector for expression may be a conventional vector that is used in the art to express a foreign gene or protein in a plant, animal, or microorganism, and may be constructed through various methods known in the art.

[0029] As used in the present invention, the term “recombinant vector” may be transformed into a suitable host cell, and then may replicate regardless of the genome of the host cell or may be integrated into the genome itself. In this case, the “suitable host cell” may contain a replication origin, which is a particular nucleotide sequence which enables the vector to replicate in the suitable host cell and from which replication starts. For example, when the vector used is an expression vector and uses a prokaryotic cell as a host, the vector generally comprises a strong promoter capable of promoting transcription (e.g., pL λ promoter, CMV promoter, trp promoter, lac promoter, tac promoter, T7 promoter, etc.), a ribosome binding site for initiation of translation, and a transcription/translation termination sequence. When a eukaryotic cell is used as a host, the vector comprises a replication origin operating in the eukaryotic cell, and examples of the replication origin include, but are not limited to, an f1 replication origin, an SV40 replication origin, a pMB1 replication origin, an adeno replication origin, an AAV replication origin, and a BBV replication origin. In addition, the recombinant vector may comprise a promoter derived from the genome of a mammalian cell (e.g., metallothionein promoter) or a promoter derived from a mammalian virus (e.g., adenovirus late promoter, vaccinia virus 7.5K promoter, SV40 promoter, cytomegalovirus promoter, HSV-tk promoter, etc.), and generally has a polyadenylation sequence as a transcription termination sequence.

[0030] The recombinant vector may comprise a selection marker. The selection marker serves to select a transformant (host cell) transformed with the vector, and since only cells expressing the selection marker can survive in the medium treated with the selection marker, it is possible to select

transformed cells. Representative examples of the selection marker include, but are not limited to, kanamycin, streptomycin, and chloramphenicol.

[0031] The transformant may be produced by inserting the recombinant vector into a host cell, and the transformant may be obtained by introducing the recombinant vector into an appropriate host cell. The host cell is a cell capable of stably and continuously cloning or expressing the expression vector, and any host cell known in the art may be used.

[0032] Where the vector is transformed into prokaryotic cells to generate recombinant microorganisms, examples of host cells that may be used include, but are not limited to, *E. coli* sp. strains such as *E. coli* JM109, *E. coli* BL21, *E. coli* RR1, *E. coli* LE392, *E. coli* B, *E. coli* X 1776, *E. coli* W3110, and *E. coli* XL1-Blue, *Bacillus* sp. strains such as *Bacillus subtilis* and *Bacillus thuringiensis*, *Corynebacterium* sp. strains, and various *Enterobacteriaceae* strains such as *Salmonella typhimurium*, *Serratia marcescens*, and *Pseudomonas* species.

[0033] Where the vector is transformed into eukaryotic cells to generate recombinant microorganisms, examples of host cells that may be used include, but are not limited to, yeast (e.g., *Saccharomyces cerevisiae*), insect cells, plant cells and animal cells, such as Sp2/0, CHO K1, CHO DG44, PER.C6, W138, BHK, COS7, 293, HepG2, Huh7, 3T3, RIN, and MDCK cell lines.

[0034] As used in the present invention, the term “transformation” refers to a phenomenon in which external DNA is introduced into a host cell, thereby artificially causing genetic changes, and the term “transformant” refers to a host cell into which external DNA has been introduced and in which the expression of the gene of interest is stably maintained.

[0035] The transformation may be performed using a suitable vector introduction technique selected depending on the host cell, so that the gene of interest or a recombinant vector comprising the same may be expressed in the host cell. For example, introduction of the vector may be performed by electroporation, heat-shock, calcium phosphate (CaPO_4) precipitation, calcium chloride (CaCl_2) precipitation, microinjection, polyethylene glycol (PEG) method, DEAE-dextran method, cationic liposome method, lithium acetate-DMSO method, or any combination thereof, without being limited thereto. As long as the transformed gene may be expressed in the host cell, it may be inserted into the chromosome of the host cell, or may exist extrachromosomally, without being limited thereto.

[0036] The transformant may include a cell transfected, transformed, or infected with the recombinant vector of the present invention in vivo or in vitro, and may be used in the same sense as a recombinant host cell, a recombinant cell, or a recombinant microorganism.

[0037] According to one embodiment of the present invention, the transformant may be a *Corynebacterium* sp. microorganism.

[0038] More specifically, examples of the *Corynebacterium* sp. microorganism include, but are not limited to, *Corynebacterium glutamicum*, *Corynebacterium crudilactis*, *Corynebacterium deserti*, *Corynebacterium callunae*, *Corynebacterium suranareeae*, *Corynebacterium lubricantis*, *Corynebacterium doosanense*, *Corynebacterium efficiens*, *Corynebacterium uterequi*, *Corynebacterium stationis*, *Corynebacterium pacaense*, *Corynebacterium singulare*, *Corynebacterium humireducens*, *Corynebacte-*

rium marinum, *Corynebacterium halotolerans*, *Corynebacterium spheniscorum*, *Corynebacterium freiburgense*, *Corynebacterium striatum*, *Corynebacterium canis*, *Corynebacterium ammoniagenes*, *Corynebacterium renale*, *Corynebacterium pollutisoli*, *Corynebacterium imitans*, *Corynebacterium caspium*, *Corynebacterium testudinoris*, *Corynebacterium pseudopelargi*, and *Corynebacterium flavescentis*.

[0039] The transformant in the present invention may be a strain either comprising the above-described nicotinamide mononucleotide transporter variant or a polynucleotide encoding the same or comprising the vector comprising the same, a strain expressing the nicotinamide mononucleotide transporter variant or the polynucleotide, or a strain having activity the for nicotinamide mononucleotide transporter variant, without being limited thereto.

[0040] According to one embodiment of the present invention, the transformant may have the ability to produce L-glutamic acid.

[0041] The transformant may naturally have the ability to produce L-glutamic acid or may be one artificially endowed with the ability to produce L-glutamic acid.

[0042] According to one embodiment of the present invention, the transformant may have an increased ability to produce L-glutamic acid, due to a change in nicotinamide mononucleotide transporter activity.

[0043] As used in the present invention, the term “increased ability to produce” means that L-glutamic acid productivity has increased compared to that of the parent strain. As used herein, the term “parent strain” refers to a wild-type strain or mutant strain to be mutated, and includes a strain that is to be mutated directly or to be transformed with a recombinant vector or the like. In the present invention, the parent strain may be a wild-type *Corynebacterium* sp. microorganism or a *Corynebacterium* sp. microorganism mutated from the wild-type microorganism.

[0044] The transformant according to the present invention exhibits an increased ability to produce L-glutamic acid compared to the parent strain, due to the change in nicotinamide mononucleotide transporter activity caused by introduction of the nicotinamide mononucleotide transporter variant thereinto. More specifically, the amount of L-glutamic acid produced by the transformant may be at least 1%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% higher than that produced by the parent strain, or may be 1.1-fold, 1.5-fold, 2-fold, 2.5-fold, 3-fold, 3.5-fold, 4-fold, 4.5-fold, 5-fold, 5.5-fold, 6-fold, 6.5-fold, 7-fold, 7.5-fold, 8-fold, 8.5-fold, 9-fold, 9.5-fold, or 10-fold higher than that produced by the parent strain, without being limited thereto. For example, the amount of L-glutamic acid produced by the transformant comprising the nicotinamide mononucleotide transporter variant may be at least 5%, specifically 5 to 50% (preferably 7 to 30%) higher than that produced by the parent strain.

[0045] Still yet another aspect of the present invention provides a method for producing L-glutamic acid, comprising steps of: culturing the transformant in a medium; and recovering L-glutamic acid from the transformant or the medium in which the transformant has been cultured.

[0046] The culturing may be performed using a suitable medium and culture conditions known in the art, and any person skilled in the art may easily adjust and use the medium and the culture conditions. Specifically, the medium

may be a liquid medium, without being limited thereto. Examples of the culturing method include, but are not limited to, batch culture, continuous culture, fed-batch culture, or a combination thereof.

[0047] According to one embodiment of the present invention, the medium should meet the requirements of a specific strain in a proper manner, and may be appropriately modified by a person skilled in the art. For culture media for *Corynebacterium* sp. microorganisms, reference may be made to, but not limited to, a known document (Manual of Methods for General Bacteriology, American Society for Bacteriology, Washington D.C., USA, 1981).

[0048] According to one embodiment of the present invention, the medium may contain various carbon sources, nitrogen sources, and trace element components. Examples of carbon sources that may be used include: sugars and carbohydrates such as glucose, sucrose, lactose, fructose, maltose, starch, and cellulose; oils and fats such as soybean oil, sunflower oil, castor oil, and coconut oil; fatty acids such as palmitic acid, stearic acid, and linoleic acid; alcohols such as glycerol and ethanol; and organic acids such as acetic acid. These substances may be used individually or as a mixture, without being limited thereto. Examples of nitrogen sources that may be used include peptone, yeast extract, meat extract, malt extract, corn steep liquor, soybean meal, urea, or inorganic compounds such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate, and ammonium nitrate. The nitrogen sources may also be used individually or as a mixture, without being limited thereto. Examples of phosphorus sources that may be used include, but are not limited to, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts. In addition, the culture medium may contain, but is not limited to, metal salts such as magnesium sulfate or iron sulfate, which are required for growth. In addition, the culture medium may contain essential growth substances such as amino acids and vitamins. Moreover, suitable precursors may be used in the culture medium. The medium or individual components may be added to the culture medium batchwise or in a continuous manner by a suitable method during culturing, without being limited thereto.

[0049] According to one embodiment of the present invention, the pH of the culture medium may be adjusted by adding compounds such as ammonium hydroxide, potassium hydroxide, ammonia, phosphoric acid and sulfuric acid to the microorganism culture medium in an appropriate manner during the culturing. In addition, during the culturing, foaming may be suppressed using an anti-foaming agent such as a fatty acid polyglycol ester. Additionally, to keep the culture medium in an aerobic condition, oxygen or an oxygen-containing gas (for example, air) may be injected into the culture medium. The temperature of the culture medium may be generally 20° C. to 45° C., for example, 25° C. to 40° C. The culturing may be continued until a desired amount of a useful substance is produced. For example, the culturing time may be 10 hours to 160 hours.

[0050] According to one embodiment of the present invention, in the step of recovering L-glutamic acid from the cultured transformant or the medium in which the transformant has been cultured, the produced L-glutamic acid may be collected or recovered from the medium using a suitable method known in the art depending on the culture method. Examples of a method that may be used to recover the

produced L-glutamic acid include, but are not limited to, centrifugation, filtration, extraction, spraying, drying, evaporation, precipitation, crystallization, electrophoresis, fractional dissolution (e.g., ammonium sulfate precipitation), chromatography (e.g., ion exchange, affinity, hydrophobicity and size exclusion), and the like.

[0051] According to one embodiment of the present invention, the step of recovering L-glutamic acid may be performed by centrifuging the culture medium at a low speed to remove biomass and separating the obtained supernatant through ion-exchange chromatography.

[0052] According to one embodiment of the present invention, the step of recovering L-glutamic acid may include a process of purifying the L-glutamic acid.

Advantageous Effects

[0053] The nicotinamide mononucleotide transporter variant according to the present invention is obtained by substituting one or more amino acids in the amino acid sequence constituting nicotinamide mononucleotide transporter to change the enzymatic activity of the nicotinamide mononucleotide transporter, and a recombinant microorganism comprising the nicotinamide mononucleotide transporter variant is capable of efficiently producing L-glutamic acid.

BRIEF DESCRIPTION OF DRAWINGS

[0054] FIG. 1 shows the structure of plasmid pk19msb according to one embodiment of the present invention.

MODE FOR INVENTION

[0055] Hereinafter, the present invention will be described in more detail. However, this description is merely presented by way of example to facilitate the understanding of the present invention, and the scope of the present invention is not limited by this exemplary description.

Example 1. Construction of Vector for Expression of Nicotinamide Mononucleotide Transporter Variant

[0056] A vector for expression of a variant having a substitution of threonine (T) for alanine (A) at position 163 in the amino acid sequence of nicotinamide mononucleotide transporter (SEQ ID NO: 4) was constructed.

[0057] Using the gDNA of wild-type *Corynebacterium glutamicum* ATCC13869 as a template, PCR reactions were performed using a primer pair of primer 1 and primer 2 and a primer pair of primer 3 and primer 4, respectively. Thereafter, using a mixture of the two PCR products as a template, overlapping PCR was performed using primer 1 and primer 4 to obtain a fragment. Here, Takara PrimeSTAR Max DNA polymerase was used as polymerase, and PCR amplification was performed under the following conditions: denaturation at 95° C. for 5 min, and then 30 cycles, each consisting of 95° C. for 30 sec, 58° C. for 30 sec, and 72° C. for 1 min and 30 sec, followed by reaction at 72° C. for 5 min. A pk19msb vector was treated with *sma*I and ligated with the PCR product (fragment) obtained above, and the resulting plasmid was named pK_pnuC (A163T).

[0058] The primer sequences used for vector construction are shown in Table 1 below.

TABLE 1

Primer name	SEQ ID NO.	Primer sequence (5'→3')
Primer 1	6	CAGTTTCAACCAGGGCGGTTTCAT
Primer 2	7	CATCCGCGAGTCATTCCGTAGG
Primer 3	8	CCTACGGAATGACTCGCGGATG
Primer 4	9	GCCTCGCCCAAGGGTGTGA

Example 2. Construction of Mutant Strain into Which Nicotinamide Mononucleotide Transporter Variant Has Been Introduced

[0059] *Corynebacterium glutamicum* U3 (KCCM13218P) was used as a parent strain into which the nicotinamide mononucleotide transporter variant was to be introduced, and an electrocompetent cell preparation method, a modification of the method of van der Rest et al., was used as a method for transformation of the U3 strain.

[0060] First, the U3 strain was primarily cultured in 100 mL of 2YT medium (containing 16 g/l of tryptone, 10 g/l of yeast extract, and 5 g/l of sodium chloride) supplemented with 2% glucose, thus preparing a seed culture. Thereafter, isonicotinic acid hydrazine at a concentration of 1 mg/ml and 2.5% glycine were added to 100 ml of 2YT medium free of glucose, and the seed culture was inoculated into the 2YT medium to reach an OD₆₁₀ value of 0.3, and then cultured at 18° C. and 180 rpm for 12 to 16 hours so that the OD₆₁₀ value reached 1.2 to 1.4. The culture was kept on ice for 30 minutes, and then centrifuged at 4,000 rpm at 4° C. for 15 minutes. Thereafter, the supernatant was discarded and the precipitated U3 strain was washed 4 times with a 10% glycerol solution and finally re-suspended in 0.5 ml of a 10% glycerol solution, thereby preparing competent cells. Electroporation was performed using a Bio-Rad electroporator. The prepared competent cells and the pK_pnuC (A163T) vector were placed in an electroporation cuvette (0.2 mm), and then subjected to electroporation under conditions of 2.5 kV, 200 Ω and 12.5 μF. Immediately after completion of the electroporation, 1 ml of a regeneration medium (containing 18.5 g/l brain heart infusion and 0.5 M sorbitol) was added to the cells which were then heat-treated at 46° C. for 6 minutes. Next, the cells were cooled at room temperature, transferred into a 15-ml cap tube, incubated at 30° C. for 2 hours, and plated on a selection medium (containing 5 g/l tryptone, 5 g/l NaCl, 2.5 g/l yeast extract, 18.5 g/l brain heart infusion powder, 15 g/l agar, 91 g/l sorbitol, and 20 μg/l kanamycin). The cells were cultured at 30° C. for 72 hours, and the generated colonies were cultured in BHI medium until the stationary phase to induce secondary recombination. Then, the cells were diluted to 10⁻⁵ to 10⁻⁷, and plated on an antibiotic-free 2YT plate medium (containing 10% sucrose), and a strain having no kanamycin resistance and grown on the medium containing 10% sucrose was selected and named PNU1.

Experimental Example 1. Evaluation of L-Glutamic Acid Productivity of Mutant Strain into Which Nicotinamide Mononucleotide Transporter Variant Has Been Introduced

[0061] L-glutamic acid productivity was compared between the parent strain U3 and the mutant strain PNU1 into which the nicotinamide mononucleotide transporter variant has been introduced.

[0062] Each strain (parent strain or mutant strain) was inoculated at 1% by volume into a 100-mL flask containing 10 mb of the medium for glutamic acid production shown in Table? below, and cultured with shaking at 200 rpm at 30° C. for 48 hours. After completion of the culturing, the concentration of L-glutamic acid in the medium was measured 5 using HPLC (Agilent), and the results are shown in Table 3 below.

TABLE 2

Component	Content
Glucose	70 g/L
(NH ₄) ₂ SO ₄	5 g/L
MgSO ₄	0.4 g/L
Urea	2 g/L
Soybean hydrolyzate	1.5% v/v
KH ₂ PO ₄	1.0 g/L
FeSO ₄	10 mg/L
MnSO ₄	10 mg/L
Thiamine_HCl	200 ug/L
biotin	2 ug/L
CaCO ₃	5%

TABLE 3

Strain	L-glutamic acid production (g/L)
U3	10.8
PNU1	11.9

[0063] As shown in Table 3 above, it was confirmed that the amount of L-glutamic acid produced by the mutant strain into which the nicotinamide mononucleotide transporter variant has been introduced increased by about 9.6% compared to that produced by the parent strain. These results suggest that the energy flux in the glutamic acid biosynthetic pathway is enhanced by the nicotinamide mononucleotide transporter variant, thereby increasing L-glutamic acid productivity.

[0064] So far, the present invention has been described with reference to the preferred embodiments. Those of ordinary skill in the art to which the present invention pertains will appreciate that the present invention may be embodied in modified forms without departing from the essential characteristics of the present invention. Therefore, the disclosed embodiments should be considered from an illustrative point of view, not from a restrictive point of view. The scope of the present invention is defined by the claims rather than the foregoing description, and all differences within the scope equivalent thereto should be construed as being included in the present invention

[Accession Number]

[0065] Depository Authority: Korean Culture Center of Microorganisms (KCCM)

[0066] Accession Number: KCCM13218P

[0067] Deposit Date: Jun. 29, 2022

SEQUENCE LISTING

Sequence total quantity: 9

SEQ ID NO: 1 moltype = DNA length = 693
 FEATURE Location/Qualifiers
 source 1..693
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 1
 atgaatccta taaccgaatt attagacgca acattatgga tcggcggagt tccgattctg 60
 tggcgcgaaa tcacggcga cgttttcgga ttatttagcg cgtgggcagg aatgcgacgc 120
 atcgtgtggg catggcccat cggcatcata ggcaacgcgc tgctgttcac agtatattatg 180
 ggcggccttt tccacactcc acaaaacctc gatctctacg gccaaagcggg tcgccagatc 240
 atgttcatca tcgtcagtggt ttatggctgg taccaatggt cggccgcaaa acgtcgcgca 300
 ctcaccccag aaaatgcagt agcagtggtt cctcgctggg caagcaccaa agaacgcgcc 360
 ggcatgtgga tcgcagcgggt tgcggaaca ctcagcttcg cctggatttt ccaagcactt 420
 ggctcctggg gaccatgggc cgacgcgtgg attttcgtcg gctcaatcct ggctacctac 480
 ggaatgactc gccgatggac agagttctcg ctgatctgga tcgcgcgcga catagttggc 540
 gttcctctac ttttgactgc tggctactac ccatccgcgg tgctttacct ggtgtatggc 600
 gcgtttgtca gctggggatt tgcgtgtgg ctgcgggtgc aaaaagcaga caaggtcgt 660
 gcgctggaag ctcaagagtc tgtgacagtc tga 693

SEQ ID NO: 2 moltype = AA length = 230
 FEATURE Location/Qualifiers
 source 1..230
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 2
 MNPITELLDA TLWIGGVPII WREIIGNVFG LFSAWAGMRR IVWAWPIGII GNALLPTVFM 60
 GGLFHTPQNL DLYGQAGRQI MFIIVSGYGW YQWSAAKRRA LTPENAVAVV PRWASTKERA 120
 GVIAAVVGT LSPAWIFQAL GSWGPWADAW IFVGSILATY GMTRGWTEFW LIWIAVDIVG 180
 VPLLLTAGIY PSAVLYLVYG AFVSWGFVVW LRVQKADKAR ALEAQESVTV 230

SEQ ID NO: 3 moltype = DNA length = 693
 FEATURE Location/Qualifiers
 source 1..693
 mol_type = genomic DNA
 organism = Corynebacterium sp.

SEQUENCE: 3
 atgaatccta taaccgaatt attagacgca acattatgga tcggcggagt tccgattctg 60
 tggcgcgaaa tcacggcga cgttttcgga ttatttagcg cgtgggcagg aatgcgacgc 120
 atcgtgtggg catggcccat cggcatcata ggcaacgcgc tgctgttcac agtatattatg 180
 ggcggccttt tccacactcc acaaaacctc gatctctacg gccaaagcggg tcgccagatc 240
 atgttcatca tcgtcagtggt ttatggctgg taccaatggt cggccgcaaa acgtcgcgca 300
 ctcaccccag aaaatgcagt agcagtggtt cctcgctggg caagcaccaa agaacgcgcc 360
 ggcatgtgga tcgcagcgggt tgcggaaca ctcagcttcg cctggatttt ccaagcactt 420
 ggctcctggg gaccatgggc cgacgcgtgg attttcgtcg gctcaatcct ggctacctac 480
 ggaatggctc gccgatggac agagttctcg ctgatctgga tcgcgcgcga catagttggc 540
 gttcctctac ttttgactgc tggctactac ccatccgcgg tgctttacct ggtgtatggc 600
 gcgtttgtca gctggggatt tgcgtgtgg ctgcgggtgc aaaaagcaga caaggtcgt 660
 gcgctggaag ctcaagagtc tgtgacagtc tga 693

SEQ ID NO: 4 moltype = AA length = 230
 FEATURE Location/Qualifiers
 source 1..230
 mol_type = protein
 organism = Corynebacterium sp.

SEQUENCE: 4
 MNPITELLDA TLWIGGVPII WREIIGNVFG LFSAWAGMRR IVWAWPIGII GNALLPTVFM 60
 GGLFHTPQNL DLYGQAGRQI MFIIVSGYGW YQWSAAKRRA LTPENAVAVV PRWASTKERA 120
 GVIAAVVGT LSPAWIFQAL GSWGPWADAW IFVGSILATY GMARGWTEFW LIWIAVDIVG 180
 VPLLLTAGIY PSAVLYLVYG AFVSWGFVVW LRVQKADKAR ALEAQESVTV 230

SEQ ID NO: 5 moltype = DNA length = 5719
 FEATURE Location/Qualifiers
 source 1..5719
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 5
 tgccgcaagc actcaggcgg caagggtcgc taaaggaagc ggaacacgta gaaagccagt 60
 ccgcagaaac ggtgctgacc cgggatgaat gtcagctact gggctatctg gacaagggaa 120
 aacgcaagcg caaagagaaa gcaggtagct tgcagtgggc ttacatggcg atagctagac 180
 tggcgcggtt tatggacagc aagcgaaccg gaattgccag ctggggcgcc ctctggttaag 240
 gttgggaagc cctgcaaaagt aaactggatg gctttcttcg cgccaaggat ctgatggcgc 300
 aggggatcaa gatctgatca agagacagga tgaggatcgt ttccgatgat tgaacaagat 360
 ggattgcacg caggttctcc ggccgcttgg gtggagaggc tattcggcta tgactgggca 420
 caacagacaa tcggctgctc tgatgccgcc gtgttcgggc tgtcagcgca ggggcgccc 480

-continued

gttctttttg	tcaagaccga	cctgtccggg	gccctgaatg	aactccaaga	cgaggcagcg	540
cggttatcgt	ggctggccac	gacgggcggt	ccttgccgag	ctgtgctcga	cgttgtcact	600
gaagcgggaa	gggactggct	gctattgggc	gaagtgccgg	ggcaggatct	cctgtcatct	660
caacctgtgc	ctgcgagaaa	agtatccatc	atggctgatg	caatgcggcg	gctgcatacg	720
cttgatccgg	ctacctgcc	attcgaccac	caagcgaaac	atcgcatcga	gcgagcacgt	780
actcggatgg	aagccggctc	tgtcgatcag	gatgatctgg	acgaagagca	tcaggggctc	840
gcgcagccg	aactgttcgc	caggctcaag	gcgcggatgc	ccgacggcga	ggatctcgtc	900
gtgacccatg	gcgatgcctg	cttgccgaat	atcatgggtg	aaaatggccg	cttttctgga	960
ttcatcgact	gtggccggct	gggtgtggcg	gaccgctatc	aggacatagc	gttggtctacc	1020
cgtgatattg	ctgaagagct	tggcggcgaa	tgggctgacc	gcttctcgt	gctttacggg	1080
atcgccgctc	ccgattcgca	gcgcacgcgc	ttctatcgcc	ttcttgacga	gttctctga	1140
gcgggactct	ggggttcgct	agaggatcga	tcctttttaa	cccatcacat	atacctgccg	1200
ttcactatta	tttagtgaaa	tgagatatta	tgatattttc	tgaattgtga	ttaaaaaggc	1260
aactttatgc	ccatgcacaa	gaaactataa	aaaatcacga	gaatgaaaag	aaacagatag	1320
atttttttagt	tccttagggc	cgtagtctgc	aaatcctttt	atgattttct	atcaaacaaa	1380
agaggaaaaa	agaccagttg	caatccaaac	gagagtctaa	tagaatgagg	tcgaaaaagta	1440
aatcgccggg	gtttgttact	gataaaagcag	gcaagaccta	aaatgtgtaa	agggcacaagt	1500
gtatactttg	gcgtcacccc	ttacatattt	taggtctttt	tttatgtgtc	gtaactaact	1560
tgccatcttc	aaacaggagg	gctggaagaa	gcagaccgct	aacacagtac	ataaaaaagg	1620
agacatgaac	gatgaacatc	aaaaagtttg	caaaacaagc	aacagtatta	acctttacta	1680
ccgcaactgc	ggcaggaggc	gcaactcaag	cgtttgcgaa	agaaacgaac	caaaagccat	1740
ataaggaaaac	atacggcatc	tcccatatta	cacgccatga	tatgctgcaa	atcctggaac	1800
agcaaaaaaa	tgaaaaatat	caagtttctg	aatttgatcc	gtccacaatt	aaaaatatct	1860
cttctgcaaa	aggcctggac	gcttgggaca	gctggccatt	acaaaacgct	gacggcactg	1920
tcgcaaaacta	tacgggctac	ccatcgtct	ttgcattagc	cggagatcct	aaaaatgcgg	1980
atgacacatc	gatttacatg	ttctatcaaa	aagtcggcga	aacttctatt	gacagctgga	2040
aaaaacgctg	ccgcgtcttt	aaagacagcg	acaaattcga	tgcaaatgat	ttctatcctaa	2100
aagaccaaaac	acaagaatgg	tcaggttcag	ccacattttc	atctgacgga	aaaatccggt	2160
tattctacac	tgattttctc	ggtaaacatt	acggcaaaac	aacactgaca	actgcacaa	2220
ttaacgctac	agcatcagac	agctctttga	acatcaacgg	tgtagaggat	tataaatcaa	2280
tccttgacgg	tgacggaaaa	acgtatcaaa	atgtacagca	gttcatcgat	gaaggcaact	2340
acagctcagg	cgacaacccat	acgctgagag	atcctcacta	cgtagaagat	aaaggccaca	2400
aatacttagt	atttgaagca	aacactggaa	ctgaagatgg	ctaccaaggc	gaagaatctt	2460
tatttaacaa	agcatactat	ggcaaaaagca	catcattctt	ccgtcaagaa	agtcacaaa	2520
ttctgcaaa	cgataaaaaa	cgcacggctg	agttagcaaa	cggcgctctc	gggtatgattg	2580
agctaaacga	tgatttacaca	ctgaaaaaag	tgatgaaacc	gctgattgca	ttcaacacag	2640
taacagatga	aattgaacgc	gcgaacgtct	ttaaaatgaa	cggcaaatgg	tacctgttca	2700
ctgactcccg	cggatcaaaa	atgacgattg	acggcattac	gtctaacgat	atttacatgc	2760
ttgggttatg	ttctaatctc	ttactggcc	catacaagcc	gctgaacaaa	actggccttg	2820
tggttaaaaat	ggatcttgat	cctaaccgatg	taacctttac	ttactcacac	ttcgtgtgac	2880
ctcaagcgaa	aggaacaacat	gtcgtgatta	caagctatat	gacaaacaga	ggattctacg	2940
cagacaaaac	atcaacgctt	gcgcggagct	tcctgctgaa	catcaaaagg	aagaaaaacat	3000
ctgttgctcaa	agcagcatc	cttgaacaag	gacaattaac	agttacacaa	taaaaacgca	3060
aaagaaaaatg	ccgatgggta	ccgagcgaaa	tgaccgacca	agcgacgccc	aaacctgccat	3120
cacgagattt	cgattccacc	gcgcgcttct	atgaaagggt	gggcttcgga	atcgttttcc	3180
gggacgcctc	cgcgagcggt	ctcatagtcc	acgacgccc	tgattttgta	gccctggccg	3240
acggccagca	ggtagggcga	caggctcatg	ccggccgccc	ccgccttttc	ctcaatcgct	3300
cttcggtcgt	ctggaaggca	gtacaccttg	ataggtgggc	tgcccttcc	gggtggcttg	3360
gtttcatcag	ccatccgctt	gccctcatct	gttacgccc	cggtagccgg	ccagcctcgc	3420
agagcaggat	tcccggtgag	acccgccagg	tgcaataaag	ggacagtga	gaagggaacac	3480
ccgctccggg	gtgggcccac	ttcacctatc	ctgcccggct	gacgcgcttg	gatacaccac	3540
ggaaagtcta	cacgaaccct	ttggcaaaaat	cctgtatatc	gtgcgaaaaa	ggatggatat	3600
accgaaaaaa	tcgctataat	gaccccgaa	cagggttatg	cagcgaaaaa	gcgctgcttc	3660
cctgctgttt	tggtgaatat	ctaccgactg	gaaacaggca	aatcgaggaa	attactgaac	3720
tgaggggaca	ggcgagagac	agctcctgaa	aatctcgata	actcaaaaaa		3780
tacgcccggg	agtgatctta	tttcattatg	gtgaaagtgg	gaacctctta	cgtgccgatc	3840
aacgtctcat	tttcgcaaaa	agttggccca	gggcttcccg	gtatcaacag	ggacaccagg	3900
attttattat	ctcggaagt	gatcttccgt	cacaggattt	tattcgccgc	aaagtgcgtc	3960
gggtgatgct	gccaaactac	tgatttagtg	tatgatgggt	tttttgaggt	gctccagtg	4020
cttctgtttc	tatcagctcc	tgaaaatctc	gataaactca	aaaatacgcc	cggtagtgat	4080
cttatttcat	tatggtgaaa	gttggaaact	cttacgtgcc	gatcaacgtc	tcattttcgc	4140
caaaagtgtg	cccagggtct	cccggtatca	acagggacac	caggatttat	ttattctgcg	4200
aagtgatctt	ccgtcacagg	tatttattcg	gcgcaaaagt	cgtcgggtga	tgctgccaac	4260
ttactgattt	agtgtatgat	ggtgtttttg	aggtgctcca	gtggcttctg	ttctatcag	4320
ggctggatga	tcctccagcg	cggggatctc	atgctggagt	ttctgcacca	ccccaaaagg	4380
atctagggtga	agatcctttt	tgataatctc	atgacaaaaa	tccttaaacg	tgagttttcg	4440
ttccactgag	cgtcagaccc	cgtagaaaaa	atcaaaaggat	cttcttgaga	tccttttttt	4500
ctgcgcgtaa	ctcgtcgttt	gcaaacacaa	aaaccaccgc	taccagcggg	ggtttgtttg	4560
ccggatcaag	agctaccaac	tcctttttcc	aaggtaactg	gcttcagcag	agcgcagata	4620
ccaaatactg	ttcttctagt	gtagccgtag	ttaggccacc	acttcaagaa	ctctgtagca	4680
ccgctacat	acctcgctct	gctaactcgt	ttaccagtgg	ctgctgccag	tgccgataag	4740
tcgtgtctta	ccgggttgga	ctcaagacga	tagttaccgg	ataaggcgca	gcggctcggc	4800
tgaaacgggg	gttcgtgcac	acagcccgac	ttggagcgaa	cgacctacac	cgaactgaga	4860
tacctacagc	gtgagctatg	agaaagcgcc	acgcttcccg	aaggagagaa	ggcggacagg	4920
tatccggtaa	ccggcagggt	cggaacaggga	gagcgcacga	gggagcttcc	aggggaaac	4980
gcctggtatc	tttatagttc	tgtcgggttt	cgccacctct	gacttgagcg	tcgatttttg	5040

-continued

```

tgatgctcgt caggggggag gagcctatgg aaaaacgcca gcaacgcggc ctttttacgg 5100
ttcctggcct tttgtggcc ttttgctcac atgttctttc ctgcgttatc cctgattctt 5160
gtggataaac gtattaccgc ctttgagtga gctgataccg ctgcgcgcag ccgaacgacc 5220
gagcgacagc agtcagtgag cgaggaagcg gaagagcgcc caatacgcaa accgcctctc 5280
ccgcgcggtt ggccgattca ttaatgcagc tggcacgaca ggtttccgga ctggaaagcg 5340
ggcagtgagc gcaacgcaat taatgtgagt tagctcactc attaggcacc ccaggcttta 5400
cactttatgc ttccggctcg tatgttgtgt ggaattgtga gcggataaca atttcacaca 5460
ggaaacacgt atgacatgat tacgccaaagc ttgcatgcct gcaggctgac tctagaggat 5520
ccccgggtac cgagctcgaa ttcaactggc gtctgtttac aacgtcgtga ctgggaaaac 5580
cctggcggtt cccaacttaa tcgccttgca gcacatcccc ctttcgccag ctggcgtaat 5640
agcgaagagg cccgcaccga tcgcccttcc caacagttgc gcagcctgaa tggcgaaatgg 5700
cgataagcta gcttcacgc 5719

```

```

SEQ ID NO: 6      moltype = DNA length = 23
FEATURE          Location/Qualifiers
source           1..23
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 6
cagtttcaac cagggcggtt cat 23

```

```

SEQ ID NO: 7      moltype = DNA length = 22
FEATURE          Location/Qualifiers
source           1..22
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 7
catccgcgag tcattccgta gg 22

```

```

SEQ ID NO: 8      moltype = DNA length = 22
FEATURE          Location/Qualifiers
source           1..22
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 8
cctacggaat gactcgcgga tg 22

```

```

SEQ ID NO: 9      moltype = DNA length = 19
FEATURE          Location/Qualifiers
source           1..19
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 9
gcctcgccca aggggtgtga 19

```

1. (canceled)

2. (canceled)

3. A composition for producing L-glutamic acid comprising a *Corynebacterium* sp. microorganism expressing a nicotinamide mononucleotide transporter variant consisting of the amino acid sequence of SEQ ID NO: 2 or a polynucleotide encoding the variant.

4. The composition of claim 3, wherein the polynucleotide comprises the nucleotide sequence represented by SEQ ID NO: 1.

5. (canceled)

6. A method for producing L-glutamic acid, comprising steps of:

culturing a transformant, which is a *Corynebacterium* sp. microorganism expressing a nicotinamide mononucleotide transporter variant consisting of the amino acid sequence of SEQ ID NO: 2 or a polynucleotide encoding the variant, in a medium; and recovering L-glutamic acid from the transformant or the medium in which the transformant has been cultured.

* * * * *