

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent Application Publication

20250255299

Kind Code

A1

Publication Date

August 14, 2025

Inventor(s)

YAO; CHENGLIN et al.

SYNERGISTIC MIXTURES FOR FUNGAL CONTROL IN CEREALS

Abstract

A fungicidal composition containing a fungicidally effective amount of the compound of Formula I, (S)-1,1-bis(4-fluorophenyl) propan-2-yl (3-acetoxy-4-methoxypicolinoyl)-L-alaninate, and at least one fungicide selected from the group consisting of tebuconazole, prothioconazole, difenconazole, epoxiconazole, mefentrifluconazole, benzovindiflupyr, penthiopyrad, fluxapyroxad, bixafen, fluopyram, picoxystrobin, pyraclostrobin, azoxystrobin, mancozeb and chlorothalonil, provides synergistic control of selected fungi.

##STR00001##

Inventors: YAO; CHENGLIN (INDIANAPOLIS, IN), MATHIESON; JOHN T. (CARSON CITY, NV)

Applicant: CORTEVA AGRISCIENCE LLC (INDIANAPOLIS, IN)

Family ID: 1000008575219

Assignee: CORTEVA AGRISCIENCE LLC (INDIANAPOLIS, IN)

Appl. No.: 19/027014

Filed: January 17, 2025

Related U.S. Application Data

parent US continuation 18452923 20230821 PENDING child US 19027014

parent US continuation 16610116 20191101 parent-grant-document US 11771085 US continuation PCT/US2018/030558 20180502 child US 18452923

us-provisional-application US 62500183 20170502

Publication Classification

Int. Cl.: A01N43/40 (20060101)

U.S. Cl.:

CPC A01N43/40 (20130101);

Background/Summary

CROSS REFERENCE TO RELATED APPLICATIONS [0001] This is a national phase entry under 35 U.S.C. § 371 of international patent application PCT/US18/030558, filed on May 2, 2018 and published in English as international patent publication WO2018204435 on Nov. 8, 2018, which claims the benefit of U.S. Provisional Patent Application Ser. No. 62/500,183 filed May 2, 2017, which is expressly incorporated by reference herein.

FIELD

[0002] This disclosure concerns a synergistic fungicidal composition containing (a) the compound of Formula I and (b) at least one fungicide selected from the group consisting of a sterol biosynthesis-inhibitor, for example prothioconazole, epoxiconazole, cyproconazole, myclobutanil, metconazole, mefentrifluconazole, difenoconazole, tebuconazole, tetraconazole, fenbuconazole, propiconazole, fluquinconazole, flusilazole, flutriafol, fenpropimorph, and prochloraz; a succinate dehydrogenase-inhibitor, for example fluxapyroxad, benzovindiflupyr, penthiopyrad, isopyrazam, bixafen, boscalid, penflufen, and fluopyram; a strobilurin, for example pyraclostrobin, fluoxastrobin, azoxystrobin, trifloxystrobin, picoxystrobin, and kresoxim methyl; and a multi-site-inhibitor, for example mancozeb and chlorothalonil, or other commercial fungicides to provide control of any plant fungal pathogen.

BACKGROUND AND SUMMARY

[0003] Fungicides are compounds, of natural or synthetic origin, which act to protect plants against damage caused by fungi. Current methods of agriculture rely heavily on the use of fungicides. In fact, some crops cannot be grown usefully without the use of fungicides. Using fungicides allows a grower to increase the yield and the quality of the crop, and consequently, increase the value of the crop. In most situations, the increase in value of the crop is worth at least three times the cost of the use of the fungicide.

[0004] However, no one fungicide is useful in all situations and repeated usage of a single fungicide frequently leads to the development of resistance to that and related fungicides. Consequently, research is being conducted to produce fungicides and combinations of fungicides that are safer, that have better performance, that require lower dosages, that are easier to use, and that cost less.

[0005] Synergism occurs when the activity of two or more compounds exceeds the activities of the compounds when used alone.

[0006] It is an object of this disclosure to provide synergistic compositions comprising fungicidal compounds. It is a further object of this disclosure to provide processes that use these synergistic compositions. The synergistic compositions are capable of preventing or curing, or both, diseases caused by fungi of the classes Ascomycetes and Basidiomycetes. In addition, the synergistic compositions have improved efficacy against the Ascomycete and Basidiomycete pathogens, including leaf blotch and brown rust of wheat. In accordance with this disclosure, synergistic compositions are provided along with methods for their use.

Description

DETAILED DESCRIPTION

[0007] The present disclosure concerns a synergistic fungicidal mixture comprising a fungicidally effective amount of (a) the compound of Formula I and (b) at least one fungicide selected from the compounds of the following groups A.1, B.1 and C.1:

[0008] A.1 Sterol biosynthesis inhibitors (SBI fungicides) selected from the following groups a), b) and c): [0009] a) C14 demethylase inhibitors (DMI fungicides), for example prothioconazole, epoxiconazole, cyproconazole, myclobutanil, metconazole, mefentrifluconazole, difenoconazole, tebuconazole, tetraconazole, fenbuconazole, propiconazole, fluquinconazole, flusilazole, flutriafol and prochloraz; [0010] b) Delta 14-reductase inhibitors, for example, fenpropimorph and aldimorph; [0011] c) Inhibitors of 3-keto reductase such as fenhexamid;

[0012] B.1 Respiration inhibitors selected from the following groups a) and b): [0013] a) inhibitors of complex II (SDHI fungicides, e.g. carboxamides), for example fluxapyroxad, benzovindiflupyr, penthiopyrad, isopyrazam, bixafen, boscalid, penflufen, and fluopyram; [0014] b) inhibitors of complex III at the Q.sub.o site (e.g. strobilurins), for example pyraclostrobin, fluoxastrobin, azoxystrobin, trifloxystrobin, picoxystrobin, and kresoxim methyl;

[0015] C.1 Inhibitors with multi-site action selected from the following groups a) and b): [0016] a) thio- and dithiocarbamates, such as mancozeb; [0017] b) organochlorine compounds (e.g. phthalimides, sulfamides, chloronitriles) such as chlorothalonil; [0018] or other commercial fungicides to provide control of any plant fungal pathogen.

##STR00002##

[0019] As used herein, the compound of Formula I is (S)-1,1-bis(4-fluorophenyl) propan-2-yl (3-acetoxy-4-methoxypicolinoyl)-L-alaninate. The compound of Formula I provides control of a variety of pathogens in economically important crops including, but not limited to, the causal agent of leaf blotch in wheat, *Zymoseptoria tritici* (SEPTTR).

[0020] As used herein, epoxiconazole is the common name for (2RS,3SR)-1-[3-(2-chlorophenyl)-2,3-epoxy-2-(4-fluorophenyl) propyl]-1H-1,2,4-triazole and possesses the following structure:

##STR00003##

[0021] Its fungicidal activity is described in The Pesticide Manual, Fifteenth Edition, 2009. Epoxiconazole provides broad spectrum control, with preventive and curative action, of diseases caused by Ascomycetes, Basidiomycetes and Deuteromycetes in bananas, cereals, coffee, rice and sugar beet.

[0022] As used herein, prothioconazole is the common name for 2-[(2RS)-2-(1-chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl]-2H-1,2,4-triazole-3 (4H)-thione and possesses the following structure:

##STR00004##

[0023] Its fungicidal activity is described in The Pesticide Manual, Fifteenth Edition, 2009. Prothioconazole provides control of diseases such as eyespot (*Pseudocercospora herpotrichoides*), *Fusarium* ear blight (*Fusarium* spp., *Microdochium nivale*), leaf blotch diseases (*Zymoseptoria tritici*, *Parastagonospora nodorum*, *Pyrenophora* spp., *Rhynchosporium secalis*, etc.), rust (*Puccinia* spp.) and powdery mildew (*Blumeria graminis*), by foliar application, in wheat, barley and other crops.

[0024] As used herein, difenoconazole is the common name for 3-chloro-4-[(2RS, 4RS, 2RS, 4RS)-4-methyl-2-(1H-1,2,4-triazol-1-ylmethyl)-1,3-dioxolan-2-yl]phenyl 4-chlorophenyl ether and possesses the following structure:

##STR00005##

[0025] Its fungicidal activity is described in BCPC Online Pesticide Manual-Latest Version. Difenoconazole provides broad spectrum control, with preventive and curative action, of diseases caused by Ascomycetes, Basidiomycetes and Deuteromycetes in grapes, pome fruit, stone fruit,

potatoes, sugar beet, oilseed rape, bananas, cereals, rice, soybeans, ornamentals and various vegetable crops.

[0026] As used herein, tebuconazole is the common name for (RS)-1-p-chlorophenyl-4,4-dimethyl-3-(1H-1,2,4-triazol-2-ylmethyl) pentan-3-ol and possesses the following structure:

##STR00006##

[0027] Its fungicidal activity is described in BCPC Online Pesticide Manual—Latest Version. Tebuconazole provides control of a broad range of diseases on a variety of crops. For example, on cereals it controls diseases caused by *Puccinia* spp., *Erysiphe graminis*, *Rhynchosporium secalis*, *Septoria* spp., *Pyrenophora* spp., *Cochliobolus sativus* and *Fusarium* spp., and in peanuts it controls diseases caused by *Mycosphaerella* spp., *Puccinia arachidis* and *Sclerotium rolfsii*. Other uses are in bananas, against *Mycosphaerella fijiensis*; in oilseed rape, against *Sclerotinia sclerotiorum*, *Alternaria* spp., *Leptosphaeria maculans*, and *Pyrenopeziza brassicae*; in tea, against *Exobasidium vexans*; in soybeans, against *Phakopsora pachyrhizi*; in pome and stone fruit, against *Monilinia* spp., *Podosphaera leucotricha*, *Sphaerotheca pannosa*, and *Venturia* spp.; in grapevines, against *Uncinula necator*; in coffee, against *Hemileia vastatrix*, *Cercospora coffeicola* and *Mycena citricolor*; in bulb vegetables, against *Sclerotium cepivorum* and *Alternaria porri*; in beans, against *Phaeoisariopsis griseola*; and in tomatoes and potatoes, against *Alternaria solani*.

[0028] As used herein, picoxystrobin is the common name for methyl (E)-3-methoxy-2-[2-(6-trifluoromethyl-2-pyridyloxymethyl)phenyl]acrylate and possesses the following structure:

##STR00007##

[0029] Its fungicidal activity is described in The e-Pesticide Manual, Version 5.2, 2011. Exemplary uses of picoxystrobin include, but are not limited to, broad-spectrum disease control in cereals, including *Mycosphaerella graminicola*, *Phaeosphaeria nodorum*, *Puccinia recondita* (brown rust), *Helminthosporium tritici-repentis* (tan spot) and *Blumeria graminis* f.sp. *tritici* (strobilurin-sensitive powdery mildew) in wheat; *Helminthosporium teres* (net blotch), *Rhynchosporium secalis*, *Puccinia hordei* (brown rust) and *Erysiphe graminis* f.sp. *hordei* (strobilurin-sensitive powdery mildew) in barley; *Puccinia coronata* and *Helminthosporium avenae* in oats; and *Puccinia recondita* and *Rhynchosporium secalis* in rye.

[0030] As used herein, azoxystrobin is the common name for (E)-2-{2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl}-3-methoxyacrylate and possesses the following structure:

##STR00008##

[0031] Its fungicidal activity is exemplified in The e-Pesticide Manual, Version 5.2, 2011. Exemplary uses of azoxystrobin include, but are not limited to, control of the following pathogens: *Erysiphe graminis*, *Puccinia* spp., *Parastagonospora nodorum*, *Zymoseptoria tritici* and *Pyrenophora teres* on temperate cereals; *Pyricularia oryzae* and *Rhizoctonia solani* on rice; *Plasmopara viticola* and *Uncinula necator* on vines; *Sphaerotheca fuliginea* and *Pseudoperonospora cubensis* on cucurbitaceae; *Phytophthora infestans* and *Alternaria solani* on potato and tomato; *Mycosphaerella arachidis*, *Rhizoctonia solani* and *Sclerotium rolfsii* on peanut; *Monilinia* spp. and *Cladosporium carpophilum* on peach; *Pythium* spp. and *Rhizoctonia solani* on turf; *Mycosphaerella* spp. on banana; *Cladosporium caryigenum* on pecan; *Elsinoë fawcettii*, *Colletotrichum* spp. and *Guignardia citricarpa* on citrus; *Colletotrichum* spp. and *Hemileia vastatrix* on coffee.

[0032] As used herein, pyraclostrobin is the common name for methyl N-[2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy]methyl]phenyl]-N-methoxycarbamate and possesses the following structure:

##STR00009##

[0033] Its fungicidal activity is described in BCPC Online Pesticide Manual-Latest Version. Exemplary uses of pyraclostrobin include, but are not limited to, broad spectrum disease control of

major plant pathogens, including *Zymoseptoria tritici*, *Puccinia* spp., *Drechslera tritici-repentis*, *Pyrenophora teres*, *Rhynchosporium secalis* and *Septoria nodorum* in cereals; *Mycosphaerella* spp. in peanuts; *Septoria glycines*, *Cercospora kikuchii* and *Phakopsora pachyrhizi* in soybeans; *Plasmopara viticola* and *Erysiphe necator* in grapes; *Phytophthora infestans* and *Alternaria solani* in potatoes and tomatoes; *Sphaerotheca fuliginea* and *Pseudoperonospora cubensis* in cucumber; *Mycosphaerella fijiensis* in bananas; *Elsinoë fawcettii* and *Guignardia citricarpa* in citrus and *Rhizoctonia solani* and *Pythium aphanidermatum* in turf.

[0034] As used herein, fluxapyroxad is the common name for 3-(difluoromethyl)-1-methyl-N-(3',4',5'-trifluorobiphenyl-2-yl) pyrazole-4-carboxamide and possesses the following structure:
##STR00010##

[0035] Its fungicidal activity is exemplified in Agrow Intelligence (<https://www.agrow-intelligence.com/agrow/agrow/databases/agrow-intelligence/>). Exemplary uses of fluxapyroxad include, but are not limited to, the control of plant pathogens, such as *Helminthosporium teres* (net blotch), *Rhynchosporium secalis* (leaf scald), *Puccinia hordei* (brown rust), and *Erysiphe graminis* f.sp. *hordei* (powdery mildew) in a range of crops, such as barley, maize, and soybeans.

[0036] As used herein, penthiopyrad is the common name for N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide and possesses the following structure:

##STR00011##

[0037] Its fungicidal activity is described in The Pesticide Manual, Fourteenth Edition, 2006. Penthiopyrad provides control of rust and *Rhizoctonia* diseases, as well as grey mold, powdery mildew and apple scab.

[0038] As used herein, benzovindiflupyr is the common name for N-[(1RS,4SR)-9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methylpyrazole-4-carboxamide and possesses the following structure:

##STR00012##

[0039] Its fungicidal activity is exemplified in Agrow Intelligence (<https://www.agrow-intelligence.com/agrow/agrow/databases/agrow-intelligence/>). Exemplary uses of benzovindiflupyr include, but are not limited to, controlling a variety of pathogens such as *Botrytis* spp., *Erysiphe* spp., *Rhizoctonia* spp., *Septoria* spp., *Phytophthora* spp., *Pythium* spp., *Phakopsora pachyrhizi*, and *Puccinia recondita*, in a range of crops including vines, cereals, soybeans, cotton, and fruit and vegetable crops.

[0040] As used herein, bixafen is the common name for N-(3',4'-dichloro-5-fluoro[1,1'-biphenyl]-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide and possesses the following structure:

##STR00013##

[0041] Its fungicidal activity is described in BCPC Online Pesticide Manual-Latest Version. Exemplary uses of bixafen include, but are not limited to, broad-spectrum disease control in cereals, including *Zymoseptoria tritici*, *Puccinia triticina*, *Puccinia striiformis*, *Oculimacula* spp. and *Pyrenophora tritici-repentis* in wheat and against *Pyrenophora teres*, *Ramularia collo-cygni*, *Rhynchosporium secalis* and *Puccinia hordei* in barley.

[0042] As used herein, fluopyram is the common name for N-[2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl]-2-(trifluoromethyl)benzamide and possesses the following structure:

##STR00014##

[0043] Its fungicidal activity is described in BCPC Online Pesticide Manual-Latest Version. Fluopyram provides control of grey mold, powdery mildew and *Sclerotinia* and *Monilinia* diseases in vines, table grapes, pome fruit, stone fruit, vegetables and field crops and control of Sigatoka in bananas. It also provides control of nematodes when used as a seed treatment.

[0044] As used herein, chlorothalonil is the common name for tetrachloroisophthal-onitrile and possesses the following structure:

##STR0015##

[0045] Its fungicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Chlorothalonil provides control of many fungal diseases in a wide range of crops, including pome fruit, stone fruit, almonds, citrus fruit, bush and cane fruit, cranberries, strawberries, pawpaws, bananas, mangoes, coconut palms, oil palms, rubber, pepper, vines, hops, vegetables, cucurbits, tobacco, coffee, tea, rice, soya beans, peanuts, potatoes, sugar beet, cotton, maize, ornamentals, mushrooms, and turf.

[0046] As used herein, mancozeb is the common name for [[2-[(dithiocarboxy)amino]ethyl]carbamodithioato (2-)-κS,κS']manganese mixture with [[2-[(dithiocarboxy)amino]ethyl]carbamodithioato (2-)-κS,κS']zinc and possesses the following structure:

##STR00016##

[0047] Its fungicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Mancozeb provides control of a wide range of fungal pathogens on a variety of fruits, vegetables and field crops.

[0048] In the compositions described herein, the concentration ratio of the mixture of the compound of Formula I to other fungicides at which the fungicidal effect is synergistic against wheat leaf blotch caused by *Zymoseptoria tritici* (SEPTTR) in protectant and curative applications lies within the range from about 7:1 to about 1:3,200. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to other fungicides at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 7:1 to about 1:3,200. In another embodiment, the concentration ratio of the mixture of the compound of Formula I to other fungicides at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 4:1 to about 1:2,500.

[0049] In the compositions described herein, the concentration ratio of the mixture of the compound of Formula I to a sterol biosynthesis-inhibitor at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 4:1 to about 1:52. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to a sterol biosynthesis-inhibitor at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 4:1 to about 1:52. In another embodiment, the concentration ratio of the mixture of the compound of Formula I to a sterol biosynthesis-inhibitor at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 4:1 to about 1:52. In some embodiments, the concentration ratio of the mixture of the compound of Formula I to tebuconazole at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 2:1 to about 1:4. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to tebuconazole at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 2:1 to about 1:2, and in another embodiment, the concentration ratio of the mixture of the compound of Formula I to tebuconazole at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 1:1 to about 1:4. In some embodiments, the concentration ratio of the mixture of the compound of Formula I to prothioconazole at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 1:6.5 to about 1:52. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to prothioconazole at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 1:6.5 to about 1:52, and in another embodiment, the concentration ratio of the mixture of the compound of Formula I to prothioconazole at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 1:6.5 to about 1:52. In some embodiments, the concentration ratio of the mixture of the compound of Formula I to difenoconazole at which the fungicidal effect is synergistic against

SEPTTR in protectant and curative applications lies within the range from about 2:1 to about 1:2. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to difenoconazole at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 2:1 to about 1:2, and in another embodiment the concentration ratio of the mixture of the compound of Formula I to difenoconazole at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 2:1 to about 1:2. In some embodiments, the concentration ratio of the mixture of the compound of Formula I to epoxiconazole at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 4:1 to about 1:2. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to epoxiconazole at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 4:1 to about 1:2, and in another embodiment the concentration ratio of the mixture of the compound of Formula I to epoxiconazole at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 4:1 to about 1:1. In some embodiments, the concentration ratio of the mixture of the compound of Formula I to mefentrifluconazole at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 4:1 to about 1:2. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to mefentrifluconazole at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 2:1 to about 1:2, and in another embodiment the concentration ratio of the mixture of the compound of Formula I to mefentrifluconazole at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 4:1 to about 1:2.

[0050] In the compositions described herein, the concentration ratio of the mixture of the compound of Formula I to a succinate dehydrogenase-inhibitor at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 1:1.2 to about 1:160. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to a succinate dehydrogenase-inhibitor at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 1:1.2 to about 1:80. In another embodiment, the concentration ratio of the mixture of the compound of Formula I to a succinate dehydrogenase-inhibitor at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 1:2.4 to about 1:160. In some embodiments, the concentration ratio of the mixture of the compound of Formula I to benzovindiflupyr at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 1:1.2 to about 1:5. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to benzovindiflupyr at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 1:1.2 to about 1:5, and in another embodiment, the concentration ratio of the mixture of the compound of Formula I to benzovindiflupyr at which the fungicidal effect is synergistic against SEPTTR in curative applications is about 1:2.4. In some embodiments, the concentration ratio of the mixture of the compound of Formula I to penthiopyrad at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 1:10 to about 1:160. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to penthiopyrad at which the fungicidal effect is synergistic against SEPTTR in protectant applications is about 1:80, and in another embodiment, the concentration ratio of the mixture of the compound of Formula I to penthiopyrad at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 1:10 to about 1:160. In some embodiments, the concentration ratio of the mixture of the compound of Formula I to fluxapyroxad at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 1:2.5 to about 1:40. In one embodiment, the concentration ratio of

the mixture of the compound of Formula I to fluxapyroxad at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 1:2.5 to about 1:20, and in another embodiment, the concentration ratio of the mixture of the compound of Formula I to fluxapyroxad at which the fungicidal effect is synergistic against SEPTTR in curative applications is about 1:40. In another embodiment, the concentration ratio of the mixture of the compound of Formula I to bixafen at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 1:2.5 to about 1:10. In some embodiments, the concentration ratio of the mixture of the compound of Formula I to fluopyram at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 1:1 to about 1:32. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to fluopyram at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 1:1 to about 1:32, and in another embodiment the concentration ratio of the mixture of the compound of Formula I to fluopyram at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 1:1 to about 1:16.

[0051] In the compositions described herein, the concentration ratio of the mixture of the compound of Formula I to a strobilurin fungicide at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 7:1 to about 1:80. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to a strobilurin fungicide at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 7:1 to about 1:80. In another embodiment, the concentration ratio of the mixture of the compound of Formula I to a strobilurin fungicide at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 1:1 to about 1:80. In some embodiments, the concentration ratio of the mixture of the compound of Formula I to picoxystrobin at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 1:5 to about 1:80. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to picoxystrobin at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 1:10 to about 1:80, and in another embodiment, the concentration ratio of the mixture of the compound of Formula I to picoxystrobin at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 1:5 to about 1:80. In some embodiments, the concentration ratio of the mixture of the compound of Formula I to azoxystrobin at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 2.5:1 to about 1:4. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to azoxystrobin at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 2.5:1 to about 1:3.2, and in another embodiment, the concentration ratio of the mixture of the compound of Formula I to azoxystrobin at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 1:1 to about 1:4. In some embodiments, the concentration ratio of the mixture of the compound of Formula I to pyraclostrobin at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 7:1 to about 1:4. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to pyraclostrobin at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 7:1 to about 1:4, and in another embodiment, the concentration ratio of the mixture of the compound of Formula I to pyraclostrobin at which the fungicidal effect is synergistic against SEPTTR in curative applications is about 1:1.

[0052] In the compositions described herein, the concentration ratio of the mixture of the compound of Formula I to a multi-site inhibitor at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 1:325 to about

1:3,200. In some embodiments, the concentration ratio of the compound of Formula I to a multi-site inhibitor at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 1:325 to about 1:3,200. In another embodiment, the concentration ratio of the compound of Formula I to a multi-site inhibitor at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 1:1,250 to about 1:2,600. In some embodiments, the concentration ratio of the mixture of the compound of Formula I to chlorothalonil at which the fungicidal effect is synergistic against SEPTTR in protectant and curative applications lies within the range from about 1:325 to about 1:2,500. In one embodiment, the concentration ratio of the mixture of the compound of Formula I to chlorothalonil at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 1:325 to about 1:2,600, and in another embodiment, the concentration ratio of the mixture of the compound of Formula I to chlorothalonil at which the fungicidal effect is synergistic against SEPTTR in curative applications lies within the range from about 1:1,250 to about 1:2,500. In another embodiment, the concentration ratio of the mixture of the compound of Formula I to mancozeb at which the fungicidal effect is synergistic against SEPTTR in protectant applications lies within the range from about 1:325 to about 1:3,200.

[0053] The rate at which the synergistic composition is applied will depend upon the particular type of fungus to be controlled, the degree of control required and the timing and method of application. In general, the compositions described herein can be applied at an application rate of between about 35 grams per hectare (g/ha) and about 2600 g/ha based on the total amount of active ingredients in the composition.

[0054] The compositions comprising the compound of Formula I and a sterol biosynthesis-inhibitor can be applied at an application rate of between about 40 g/ha and about 350 g/ha based on the total amount of active ingredients in the composition. Epoxiconazole is applied at a rate of between about 50 g/ha and about 250 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha. Prothioconazole is applied at a rate of between about 50 g/ha and about 250 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha. Difenconazole is applied at a rate of between about 30 g/ha and about 125 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha. Tebuconazole is applied at a rate of between about 50 g/ha and about 300 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha. Mefentrifluconazole is applied at a rate of between about 10 g/ha and about 200 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha.

[0055] The compositions comprising the compound of Formula I and a succinate dehydrogenase-inhibitor can be applied at an application rate of between about 35 g/ha and about 500 g/ha based on the total amount of active ingredients in the composition. Benzovindiflupyr is applied at a rate of between about 25 g/ha and about 300 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha. Penthiopyrad is applied at a rate of between about 100 g/ha and about 400 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha. Fluxapyroxad is applied at a rate of between about 45 g/ha and about 200 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha. Bixafen is applied at a rate of between about 30 g/ha and about 200 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha. Fluopyram is applied at a rate of between about 50 g/ha and about 300 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha.

[0056] The compositions comprising the compound of Formula I and a strobilurin can be applied at an application rate of between about 60 g/ha and about 475 g/ha based on the total amount of active ingredients in the composition. Picoxystrobin is applied at a rate of between about 50 g/ha and about 250 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha. Azoxystrobin is applied at a rate of between about 100 g/ha and about 375 g/ha and the

compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha. Pyraclostrobin is applied at a rate of between about 50 g/ha and about 250 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha.

[0057] The compositions comprising the compound of Formula I and a multi-site inhibitor can be applied at an application rate of between about 1010 g/ha and about 2600 g/ha based on the total amount of active ingredients in the composition. Chlorothalonil is applied at a rate of between about 1000 g/ha and about 2500 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha. Mancozeb is applied at a rate of between about 1500 g/ha and about 2000 g/ha and the compound of Formula I is applied at a rate between about 10 g/ha and about 100 g/ha.

[0058] The components of the synergistic mixture described herein can be applied either separately or as part of a multipart fungicidal system.

[0059] The synergistic mixture of the present disclosure can be applied in conjunction with one or more other fungicides to control a wider variety of undesirable diseases. When used in conjunction with other fungicide(s), the presently claimed compounds may be formulated with the other fungicide(s), tank mixed with the other fungicide(s) or applied sequentially with the other fungicide(s). Such other fungicides may include 2-(thiocyanatomethylthio)-benzothiazole, 2-phenylphenol, 8-hydroxyquinoline sulfate, ametoctradin, amisulbrom, antimycin, *Ampelomyces quisqualis*, azaconazole, *Bacillus subtilis*, *Bacillus subtilis* strain QST713, benalaxyl, benomyl, benthiavalicarb-isopropyl, benzylaminobenzene-sulfonate (BABS) salt, bicarbonates, biphenyl, bismethiazol, bitertanol, blasticidin-S, borax, Bordeaux mixture, boscalid, bromuconazole, bupirimate, calcium polysulfide, captafol, captan, carbendazim, carboxin, carpropamid, carvone, chlazafenone, chloroneb, chlozolate, *Coniothyrium minitans*, copper hydroxide, copper octanoate, copper oxychloride, copper sulfate, copper sulfate (tribasic), cuprous oxide, cyazofamid, cyflufenamid, cymoxanil, cyproconazole, cyprodinil, dazomet, debacarb, diammonium ethylenebis-(dithiocarbamate), dichlofluanid, dichlorophen, diclocymet, diclomezine, dichloran, diethofencarb, difenzoquat ion, diflumetorim, dimethomorph, dimoxystrobin, diniconazole, diniconazole-M, dinobuton, dinocap, diphenylamine, dipymetitrone, dithianon, dodemorph, dodemorph acetate, dodine, dodine free base, edifenphos, enestrobin, enestroburin, ethaboxam, ethoxyquin, etridiazole, famoxadone, fenamidone, fenarimol, fenbuconazole, fenfuram, fenhexamid, fenoxanil, fenpiclonil, fenpropidin, fenpropimorph, fenpyrazamine, fentin, fentin acetate, fentin hydroxide, ferbam, ferimzone, fluazinam, fludioxonil, fluindapyr, flumorph, fluopicolide, fluopyram, fluoroimide, fluoxastrobin, fluquinconazole, flusilazole, flusulfamide, flutianil, flutolanil, flutriafol, folpet, formaldehyde, fosetyl, fosetyl-aluminium, fuberidazole, furalaxyl, furametpyr, guazatine, guazatine acetates, GY-81, hexachlorobenzene, hexaconazole, hymexazol, imazalil, imazalil sulfate, imibenconazole, iminoctadine, iminoctadine triacetate, iminoctadine tris(albesilate), iodocarb, ipconazole, ipfenpyrazolone, iprobenfos, iprodione, iprovalicarb, isofetamide, isoprothiolane, isopyrazam, isotianil, kasugamycin, kasugamycin hydrochloride hydrate, kresoxium-methyl, laminarin, mancopper, mandipropamid, maneb, mefenoxam, mepanipyrim, mepronil, meptyl-dinocap, mercuric chloride, mercuric oxide, mercurous chloride, metalaxyl, metalaxyl-M, metam, metam-ammonium, metam-potassium, metam-sodium, metconazole, methasulfocarb, methyl iodide, methyl isothiocyanate, metiram, metominostrobin, metrafenone, mildiomyacin, myclobutanil, nabam, nitrothal-isopropyl, nuarimol, octhilinone, ofurace, oleic acid (fatty acids), orysastrobin, oxadixyl, oxathiapiprolin, oxine-copper, oxpoconazole fumarate, oxycarboxin, pefurazoate, penconazole, pencycuron, penflufen, pentachlorophenol, pentachlorophenyl laurate, phenylmercury acetate, phosphonic acid, phthalide, polyoxin B, polyoxins, polyoxorim, potassium bicarbonate, potassium hydroxyquinoline sulfate, probenazole, prochloraz, procymidone, propamocarb, propamocarb hydrochloride, propiconazole, propineb, proquinazid, pydiflumetofen, pyrametostrobin, pyraoxystrobin, pyraziflumid, pyrazophos, pyribencarb, pyributicarb, pyrifenoxy, pyrimethanil, pyriofenone, pyroquilon,

quinoclamine, quinoxifen, quinoxaline, *Reynoutria sachalinensis* extract, sedaxane, silthiofam, simeconazole, sodium 2-phenylphenoxide, sodium bicarbonate, sodium pentachlorophenoxide, spiroxamine, sulfur, SYP-Z048, tar oils, tebufloquin, tecnazene, tetraconazole, thiabendazole, thifluzamide, thiophanate-methyl, thiram, tiadinil, tolclofos-methyl, tolylfluanid, triadimefon, triadimenol, triazoxide, tricyclazole, tridemorph, trifloxystrobin, triflumizole, triforine, triticonazole, validamycin, valifenalate, valiphenal, vinclozolin, zineb, ziram, zoxamide, *Candida oleophila*, *Fusarium oxysporum*, *Gliocladium* spp., *Phlebiopsis gigantea*, *Streptomyces griseoviridis*, *Trichoderma* spp., (RS)—N-(3,5-dichlorophenyl)-2-(methoxymethyl)-succinimide, 1,2-dichloropropane, 1,3-dichloro-1,1,3,3-tetrafluoroacetone hydrate, 1-chloro-2,4-dinitronaphthalene, 1-chloro-2-nitropropane, 2-(2-heptadecyl-2-imidazolin-1-yl) ethanol, 2,3-dihydro-5-phenyl-1,4-dithiane 1,1,4,4-tetraoxide, 2-methoxyethylmercury acetate, 2-methoxyethylmercury chloride, 2-methoxyethylmercury silicate, 3-(4-chlorophenyl)-5-methylrhodanine, 4-(2-nitroprop-1-enyl)phenyl thiocyanate, aminopyrifin, ampropylfos, anilazine, azithiram, barium polysulfide, Bayer 32394, benodanil, benquinox, bentazon, benzamocryl; benzamocryl-isobutyl, benzamocryl, binapacryl, bis(methylmercury) sulfate, bis(tributyltin) oxide, buthiolate, cadmium calcium copper zinc chromate sulfate, carbamothion, CECA, chlorobenzothiazole, chloranilformetan, chlorfenazole, chlorquinox, climbazole, copper bis(3-phenylsalicylate), copper zinc chromate, cufraneb, cupric hydrazinium sulfate, cuprothion, cyclofuramid, cypendazole, cyproflumuron, decafenin, dichlobenzothiazole, dichlorodimethylpate, diclobutrazol, dimethirimol, dinocroton, dinosulfon, dinoterbon, dipyrithione, ditalimfos, dodicil, drazoxolon, EBP, ESBP, etaconazole, etem, ethirim, fenaminosulf, fenapanil, fenitropan, fluindapyr, fluopimomide, fluotrimazole, furcarbanil, furconazole, furconazole-cis, furmecycloz, furophanate, glyodine, griseofulvin, halacrinat, Hercules 3944, hexylthiofos, ICIA0858, inpyrfluxam, ipfentrifluconazole, ipflufenquin, isoflucypram, isopamphos, isovaldione, mandestrobin, mebenil, mecarbinid, metazoxolon, methfuroxam, methylmercury dicyandiamide, metsulfovax, metyltetraprole, milneb, mucochloric anhydride, myclozolin, N-3,5-dichlorophenylsuccinimide, N-3-nitrophenylitaconimide, natamycin, N-ethylmercurio-4-toluenesulfonanilide, nickel bis(dimethyldithiocarbamate), OCH, phenylmercury dimethyldithiocarbamate, phenylmercury nitrate, phosdiphen, prothiocarb; prothiocarb hydrochloride, pydiflumetofen, pyracarbolid, pyrapropoyne, pyridachlometyl, pyridinitril, pyroxychlor, pyroxyfur, quinacetol; quinacetol sulfate, quinazamid, quinconazole, quinofumelin, rabenzazole, salicylanilide, SSF-109, sultropan, tecoram, thiadiflour, thicyofen, thiochlorfenphim, thiophanate, thioquinox, tioxyimid, triamphos, triarimol, triazbutyl, trichlamide, urbacid, zarilamid, and any combinations thereof.

[0060] The compositions of the present disclosure are preferably applied in the form of a formulation comprising a composition of (a) a compound of Formula I and (b) at least one fungicide selected from the group consisting of tebuconazole, prothioconazole, difenconazole, epoxiconazole, benzovindiflupyr, penthiopyrad, fluxapyroxad, bixafen, picoxystrobin, pyraclostrobin, azoxystrobin, mefentrifluconazole, fluopyram, mancozeb and chlorothalonil, together with a phytologically acceptable carrier.

[0061] Concentrated formulations can be dispersed in water, or another liquid, for application, or formulations can be dust-like or granular, which can then be applied without further treatment. The formulations are prepared according to procedures which are conventional in the agricultural chemical art, but which are novel and important because of the presence therein of a synergistic composition.

[0062] The formulations that are applied most often are aqueous suspensions or emulsions. Either such water-soluble, water-suspendable, or emulsifiable formulations are solids, usually known as wettable powders, or liquids, usually known as emulsifiable concentrates, aqueous suspensions, or suspension concentrates. The present disclosure contemplates all vehicles by which the synergistic compositions can be formulated for delivery and use as a fungicide.

[0063] As will be readily appreciated, any material to which these synergistic compositions can be

added may be used, provided they yield the desired utility without significant interference with the activity of these synergistic compositions as antifungal agents.

[0064] Wettable powders, which may be compacted to form water-dispersible granules, comprise an intimate mixture of the synergistic composition, a carrier and agriculturally acceptable surfactants. The concentration of the synergistic composition in the wettable powder is usually from about 10% to about 90% by weight, more preferably about 25% to about 75% by weight, based on the total weight of the formulation. In the preparation of wettable powder formulations, the synergistic composition can be compounded with any of the finely divided solids, such as prophyllite, talc, chalk, gypsum, Fuller's earth, bentonite, attapulgite, starch, casein, gluten, montmorillonite clays, diatomaceous earths, purified silicates or the like. In such operations, the finely divided carrier is ground or mixed with the synergistic composition in a volatile organic solvent. Effective surfactants, comprising from about 0.5% to about 10% by weight of the wettable powder, include sulfonated lignins, naphthalenesulfonates, alkylbenzenesulfonates, alkyl sulfates, and non-ionic surfactants, such as ethylene oxide adducts of alkyl phenols.

[0065] Emulsifiable concentrates of the synergistic composition comprise a convenient concentration, such as from about 10% to about 50% by weight, in a suitable liquid, based on the total weight of the emulsifiable concentrate formulation. The components of the synergistic compositions, jointly or separately, are dissolved in a carrier, which is either a water-miscible solvent or a mixture of water-immiscible organic solvents, and emulsifiers. The concentrates may be diluted with water and oil to form spray mixtures in the form of oil-in-water emulsions. Useful organic solvents include aromatics, especially the high-boiling naphthalenic and olefinic portions of petroleum such as heavy aromatic naphtha. Other organic solvents may also be used, such as, for example, terpenic solvents, including rosin derivatives, aliphatic ketones, such as cyclohexanone, and complex alcohols, such as 2-ethoxyethanol.

[0066] Emulsifiers which can be advantageously employed herein can be readily determined by those skilled in the art and include various nonionic, anionic, cationic and amphoteric emulsifiers, or a blend of two or more emulsifiers. Examples of nonionic emulsifiers useful in preparing the emulsifiable concentrates include the polyalkylene glycol ethers and condensation products of alkyl and aryl phenols, aliphatic alcohols, aliphatic amines or fatty acids with ethylene oxide, propylene oxides such as the ethoxylated alkyl phenols and carboxylic esters solubilized with the polyol or polyoxyalkylene. Cationic emulsifiers include quaternary ammonium compounds and fatty amine salts. Anionic emulsifiers include the oil-soluble salts (e.g., calcium) of alkylaryl sulfonic acids, oil-soluble salts or sulfated polyglycol ethers and appropriate salts of phosphated polyglycol ether.

[0067] Representative organic liquids which can be employed in preparing the emulsifiable concentrates of the present disclosure are the aromatic liquids such as xylene, propyl benzene fractions, or mixed naphthalene fractions, mineral oils, substituted aromatic organic liquids such as dioctyl phthalate, kerosene, dialkyl amides of various fatty acids, particularly the dimethyl amides of fatty glycols and glycol derivatives such as the n-butyl ether, ethyl ether or methyl ether of diethylene glycol, and the methyl ether of triethylene glycol. Mixtures of two or more organic liquids are also often suitably employed in the preparation of the emulsifiable concentrate. The preferred organic liquids are xylene, and propyl benzene fractions, with xylene being most preferred. The surface-active dispersing agents are usually employed in liquid formulations and in the amount of from 0.1 to 20 percent by weight of the combined weight of the dispersing agent with the synergistic compositions. The formulations can also contain other compatible additives, for example, plant growth regulators and other biologically active compounds used in agriculture.

[0068] Aqueous suspensions comprise suspensions of one or more water-insoluble compounds, dispersed in an aqueous vehicle at a concentration in the range from about 5% to about 70% by weight, based on the total weight of the aqueous suspension formulation. Suspensions are prepared by finely grinding the components of the synergistic combination either together or separately, and vigorously mixing the ground material into a vehicle comprised of water and surfactants chosen

from the same types discussed above. Other ingredients, such as inorganic salts and synthetic or natural gums, may also be added to increase the density and viscosity of the aqueous vehicle. It is often most effective to grind and mix at the same time by preparing the aqueous mixture and homogenizing it in an implement such as a sand mill, ball mill, or piston-type homogenizer.

[0069] The synergistic composition may also be applied as a granular formulation, which is particularly useful for applications to the soil. Granular formulations usually contain from about 0.5% to about 10% by weight of the compounds, based on the total weight of the granular formulation, dispersed in a carrier which consists entirely or in large part of coarsely divided attapulgite, bentonite, diatomite, clay or a similar inexpensive substance. Such formulations are usually prepared by dissolving the synergistic composition in a suitable solvent and applying it to a granular carrier which has been preformed to the appropriate particle size, in the range of from about 0.5 to about 3 millimeters (mm). Such formulations may also be prepared by making a dough or paste of the carrier and the synergistic composition, and crushing and drying to obtain the desired granular particle.

[0070] Dusts containing the synergistic composition are prepared simply by intimately mixing the synergistic composition in powdered form with a suitable dusty agricultural carrier, such as, for example, kaolin clay, ground volcanic rock, and the like. Dusts can suitably contain from about 1% to about 10% by weight of the synergistic composition/carrier combination.

[0071] The formulations may contain agriculturally acceptable adjuvant surfactants to enhance deposition, wetting and penetration of the synergistic composition onto the target crop and organism. These adjuvant surfactants may optionally be employed as a component of the formulation or as a tank mix. The amount of adjuvant surfactant will vary from 0.01 percent to 1.0 percent volume/volume (v/v) based on a spray-volume of water, preferably 0.05 to 0.5 percent. Suitable adjuvant surfactants include ethoxylated nonyl phenols, ethoxylated synthetic or natural alcohols, salts of the esters or sulfosuccinic acids, ethoxylated organosilicones, ethoxylated fatty amines and blends of surfactants with mineral or vegetable oils.

[0072] The formulations may optionally include combinations that can comprise at least 1% by weight of one or more of the synergistic compositions with another pesticidal compound. Such additional pesticidal compounds may be fungicides, insecticides, nematocides, miticides, arthropodicides, bactericides or combinations thereof that are compatible with the synergistic compositions of the present disclosure in the medium selected for application, and not antagonistic to the activity of the present compounds. Accordingly, in such embodiments the other pesticidal compound is employed as a supplemental toxicant for the same or for a different pesticidal use. The pesticidal compound and the synergistic composition can generally be mixed together in a weight ratio of from 1:100 to 100:1.

[0073] The present disclosure includes within its scope methods for the control or prevention of fungal attack. These methods comprise applying to the locus of the fungus, or to a locus in which the infestation is to be prevented (for example applying to wheat or barley plants), a fungicidally effective amount of the synergistic composition. The synergistic composition is suitable for treatment of various plants at fungicidal levels, while exhibiting low phytotoxicity. The synergistic composition is useful in a protectant or eradicator fashion. The synergistic composition is applied by any of a variety of known techniques, either as the synergistic composition or as a formulation comprising the synergistic composition. For example, the synergistic compositions may be applied to the roots, seeds or foliage of plants for the control of various fungi, without damaging the commercial value of the plants. The synergistic composition is applied in the form of any of the generally used formulation types, for example, as solutions, dusts, wettable powders, flowable concentrates, or emulsifiable concentrates. These materials are conveniently applied in various known fashions.

[0074] The synergistic composition has been found to have significant fungicidal effect, particularly for agricultural use. The synergistic composition is particularly effective for use with

agricultural crops and horticultural plants, or with wood, paint, leather or carpet backing.

[0075] In particular, the synergistic composition is effective in controlling a variety of undesirable fungi that infect useful plant crops. The synergistic composition may be used against a variety of Ascomycete and Basidiomycete fungi, including for example the following representative fungi species: barley leaf scald (*Rhynchosporium secalis*); barley *Ramularia* leaf spot (*Ramularia collo-cygni*); barley net blotch (*Pyrenophora teres*); barley powdery mildew (*Blumeria graminis* f. sp. *hordei*); wheat powdery mildew (*Blumeria graminis* f. sp. *tritici*); wheat brown rust (*Puccinia triticina*); stripe rust of wheat (*Puccinia striiformis*); leaf blotch of wheat (*Zymoseptoria tritici*); glume blotch of wheat (*Parastagonospora nodorum*); leaf spot of sugar beets (*Cercospora beticola*); leaf spot of peanut (*Mycosphaerella arachidis*); cucumber anthracnose (*Colletotrichum lagenarium*); cucumber powdery mildew (*Erysiphe cichoracearum*); watermelon stem gummy blight (*Didymella bryoniae*); apple scab (*Venturia inaequalis*); apple powdery mildew (*Podosphaera leucotricha*); grey mold (*Botrytis cinerea*); *Sclerotinia* white mold (*Sclerotinia sclerotiorum*); grape powdery mildew (*Erysiphe necator*); early blight of tomato (*Alternaria solani*); rice blast (*Pyricularia oryzae*); brown rot of stone fruits (*Monilinia fructicola*) and black sigatoka disease of banana (*Mycosphaerella fijiensis*). It will be understood by those in the art that the efficacy of the synergistic compositions for one or more of the foregoing fungi establishes the general utility of the synergistic compositions as fungicides.

[0076] The synergistic compositions have a broad range of efficacy as a fungicide. The exact amount of the synergistic composition to be applied is dependent not only on the relative amounts of the components, but also on the particular action desired, the fungal species to be controlled, and the stage of growth thereof, as well as the part of the plant or other product to be contacted with the synergistic composition. Thus, formulations containing the synergistic composition may not be equally effective at similar concentrations or against the same fungal species.

[0077] The synergistic compositions are effective in use with plants in a disease-inhibiting and phytologically acceptable amount. The term “disease-inhibiting and phytologically acceptable amount” refers to an amount of the synergistic composition that kills or inhibits the plant disease for which control is desired, but is not significantly toxic to the plant. The exact concentration of synergistic composition required varies with the fungal disease to be controlled, the type of formulation employed, the method of application, the particular plant species, climate conditions, and the like.

[0078] The present compositions can be applied to fungi or their locus by the use of conventional ground sprayers, granule applicators, and by other conventional means known to those skilled in the art.

[0079] The following examples are provided for illustrative purposes and should not be construed as limitations to the disclosure.

EXAMPLES

[0080] Evaluation of Curative and Protectant Activity of Fungicide Mixtures vs. Leaf Blotch of Wheat (*Zymoseptoria tritici*; Bayer code: SEPTTR):

[0081] Wheat plants (variety Yuma) were grown from seed in a greenhouse in plastic pots with a surface area of 27.5 square centimeters (cm.sup.2) containing 50% mineral soil/50% soil-less Metro mix, with 8-12 seedlings per pot. The plants were employed for testing when the first leaf was fully emerged, which typically took 7 to 8 days after planting.

[0082] Treatments consisted of fungicide compounds difenoconazole, epoxiconazole, prothioconazole, tebuconazole, mefentrifluconazole, azoxystrobin, picoxystrobin, pyraclostrobin, penthiopyrad, fluxapyroxad, benzovindiflupyr, bixafen, fluopyram, mancozeb and chlorothalonil, either using individually or as two-way mixture with the compound of Formula I.

[0083] The compounds were tested as technical grade material formulated in acetone, and spray solutions contained 10% acetone and 100 ppm Triton X-100. Fungicide solutions were applied onto plants using an automated booth sprayer, which utilized two 6218-1/4 JAUPM spray nozzles

operating at 20 pounds per square inch (psi) set at opposing angles to cover both leaf surfaces. All sprayed plants were allowed to air dry prior to further handling. Control plants were sprayed in the same manner with the solvent blank.

[0084] Test plants were inoculated with an aqueous spore suspension of *Zymoseptoria tritici* either 3 days prior to (3-day curative test) or 1 day after fungicide treatments (1-day protectant test). After inoculation the plants were kept in 100% relative humidity (one day in a dark dew chamber followed by two days in a lighted mist chamber) to permit spores to germinate and infect the leaf. The plants were then transferred to a greenhouse for disease to develop. When disease fully developed on untreated plants, disease severity on the first leaf of the seedlings was assessed and activity was represented by percent of leaf area free of SEPTTR infection relative to the untreated plants.

[0085] Colby's equation was used to determine the fungicidal effects expected from the mixtures. (See Colby, S. R. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 1967, 15, 20-22.)

[0086] The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

Expected=A+B-(A×B/100) [0087] A=observed efficacy of active component A at the same concentration as used in the mixture; [0088] B=observed efficacy of active component B at the same concentration as used in the mixture.

[0089] Synergistic interactions between compound I and other fungicides were detected in curative and/or protectant assays vs. SEPTTR (Tables 1-7).

TABLE-US-00001 TABLE 1 Synergistic Interactions of the Compound of Formula I and Other Fungicides In a 1-Day Protectant (1DP) *Zymoseptoria tritici* (SEPTTR) Assay-Test 1. Rates SEPTTR* Synergism Composition (ppm)* Observed* Expected* Factor* Epoxiconazole + Compound I 0.05 + 0.1 83 76 1.09 Epoxiconazole + Compound I 0.025 + 0.1 91 72 1.26 Epoxiconazole + Compound I 0.1 + 0.05 98 68 1.43 Epoxiconazole + Compound I 0.05 + 0.05 69 44 1.57 Epoxiconazole + Compound I 0.025 + 0.05 62 34 1.79 Prothioconazole + Compound I 1.28 + 0.1 83 73 1.13 Prothioconazole + Compound I 0.64 + 0.1 90 81 1.11 Prothioconazole + Compound I 1.28 + 0.05 93 37 2.53 Prothioconazole + Compound I 0.64 + 0.05 81 55 1.46 Prothioconazole + Compound I 0.32 + 0.05 62 51 1.21 Benzovindiflupyr + Compound I 0.3 + 0.1 89 83 1.06 Benzovindiflupyr + Compound I 0.075 + 0.05 45 39 1.13 Bixafen + Compound I 0.5 + 0.1 78 73 1.08 Pyraclostrobin + Compound I 0.014 + 0.1 97 89 1.09 Pyraclostrobin + Compound I 0.028 + 0.05 85 47 1.80 Pyraclostrobin + Compound I 0.014 + 0.05 98 75 1.31 Chlorothalonil + Compound I 129 + 0.05 94 82 1.15 Mancozeb + Compound I 160 + 0.05 95 89 1.07 *SEPTTR = Leaf Blotch of Wheat; *Zymoseptoria tritici* *Observed = Observed percent disease control at the test rates *Expected = Percent disease control expected as predicted by the Colby equation *ppm = Parts per million *Synergism factor = Observed/Expected

TABLE-US-00002 TABLE 2 Synergistic Interactions of the Compound of Formula I and Other Fungicides In a 1-Day Protectant (1DP) *Zymoseptoria tritici* (SEPTTR) Assay-Test 2. Rates SEPTTR* Synergism Composition (ppm)* Observed* Expected* Factor* Epoxiconazole + Compound I 0.1 + 0.1 100 67 1.48 Epoxiconazole + Compound I 0.05 + 0.1 89 64 1.39 Epoxiconazole + Compound I 0.025 + 0.1 91 58 1.58 Prothioconazole + Compound I 2.6 + 0.1 99 91 1.09 Prothioconazole + Compound I 1.3 + 0.1 98 84 1.16 Prothioconazole + Compound I 0.65 + 0.1 99 82 1.21 Prothioconazole + Compound I 2.6 + 0.05 99 84 1.18 Prothioconazole + Compound I 1.3 + 0.05 94 72 1.30 Prothioconazole + Compound I 0.65 + 0.05 79 68 1.16 Difenoconazole + Compound I 0.1 + 0.1 100 87 1.15 Difenoconazole + Compound I 0.05 + 0.1 100 73 1.36 Difenoconazole + Compound I 0.1 + 0.05 96 76 1.27 Difenoconazole + Compound I 0.05 + 0.05 91 52 1.74 Tebuconazole + Compound I 0.2 + 0.1 96 62 1.56 Tebuconazole + Compound I 0.1 + 0.1 89 60 1.49 Tebuconazole + Compound I 0.05 + 0.1 82 59 1.39

Benzovindiflupyr + Compound I 0.5 + 0.1 75 70 1.08 Benzovindiflupyr + Compound I 0.25 + 0.1
 86 59 1.45 Benzovindiflupyr + Compound I 0.12 + 0.1 91 59 1.56 Benzovindiflupyr + Compound
 I 0.25 + 0.05 40 27 1.48 Benzovindiflupyr + Compound I 0.12 + 0.05 32 26 1.22 Fluxapyroxad +
 Compound I 1 + 0.1 93 63 1.47 Fluxapyroxad + Compound I 0.5 + 0.1 98 59 1.68 Fluxapyroxad
 + Compound I 0.25 + 0.1 99 58 1.72 Fluxapyroxad + Compound I 1 + 0.05 51 34 1.50
 Fluxapyroxad + Compound I 0.5 + 0.05 30 26 1.15 Fluxapyroxad + Compound I 0.25 + 0.05 26
 25 1.07 Bixafen + Compound I 1 + 0.1 86 65 1.33 Bixafen + Compound I 0.5 + 0.1 77 65 1.20
 Bixafen + Compound I 0.25 + 0.1 77 65 1.20 Bixafen + Compound I 0.5 + 0.05 40 36 1.11
 Bixafen + Compound I 0.25 + 0.05 46 36 1.25 Azoxystrobin + Compound I 0.16 + 0.1 79 62 1.28
 Azoxystrobin + Compound I 0.08 + 0.1 96 62 1.54 Azoxystrobin + Compound I 0.04 + 0.1 98 59
 1.65 Azoxystrobin + Compound I 0.16 + 0.05 49 31 1.58 Azoxystrobin + Compound I 0.08 + 0.05
 47 33 1.46 Pyraclostrobin + Compound I 0.2 + 0.1 100 84 1.19 Pyraclostrobin + Compound I 0.1 +
 0.1 95 6 1.43 Pyraclostrobin + Compound I 0.05 + 0.1 96 61 1.59 Pyraclostrobin + Compound I
 0.2 + 0.05 96 71 1.36 Pyraclostrobin + Compound I 0.1 + 0.05 51 39 1.30 Pyraclostrobin +
 Compound I 0.05 + 0.05 33 30 1.12 Chlorothalonil + Compound I 130 + 0.1 100 67 1.50
 Chlorothalonil + Compound I 65 + 0.1 100 66 1.51 Chlorothalonil + Compound I 32.5 + 0.1 100
 69 1.44 Chlorothalonil + Compound I 130 + 0.05 92 40 2.26 Chlorothalonil + Compound I 65 +
 0.05 79 39 2.02 Chlorothalonil + Compound I 32.5 + 0.05 61 44 1.38 Mancozeb + Compound I
 130 + 0.1 100 66 1.51 Mancozeb + Compound I 65 + 0.1 100 79 1.26 Mancozeb + Compound I
 32.5 + 0.1 99 61 1.63 Mancozeb + Compound I 130 + 0.05 95 39 2.43 Mancozeb + Compound I
 65 + 0.05 89 63 1.41 Mancozeb + Compound I 32.5 + 0.05 61 30 2.06 *SEPTTR = Leaf Blotch
 of Wheat; *Zymoseptoria tritici* *Observed = Observed percent disease control at the test rates
 *Expected = Percent disease control expected as predicted by the Colby equation *ppm = Parts per
 million *Synergism factor = Observed/Expected

TABLE-US-00003 TABLE 3 Synergistic Interactions of the Compound of Formula I and Other
 Fungicides In a 1-Day Protectant (1DP) *Zymoseptoria tritici* (SEPTTR) Assay-Test 3. Rates
 SEPTTR* Synergism Composition (ppm)* Observed* Expected* Factor* Penthiopyrad +
 Compound I 2 + 0.025 39 26 1.49 Picoxystrobin + Compound I 4 + 0.05 70 60 1.17
 Picoxystrobin + Compound I 2 + 0.05 58 36 1.59 Picoxystrobin + Compound I 2 + 0.025 36
 27 1.31 Picoxystrobin + Compound I 1 + 0.05 39 18 2.23 Picoxystrobin + Compound I 1 +
 0.025 22 6 3.67 Picoxystrobin + Compound I 0.5 + 0.05 32 26 1.22 Picoxystrobin + Compound I
 0.5 + 0.025 18 16 1.12 *SEPTTR = Leaf Blotch of Wheat; *Zymoseptoria tritici* *Observed =
 Observed percent disease control at the test rates *Expected = Percent disease control expected as
 predicted by the Colby equation *ppm = Parts per million *Synergism factor = Observed/Expected
 TABLE-US-00004 TABLE 4 Synergistic Interactions of the Compound of Formula I and Other
 Fungicides In a 3-Day Curative (3DC) *Zymoseptoria tritici* (SEPTTR) Assay-Test 4. Rates
 SEPTTR* Synergism Composition (ppm)* Observed* Expected* Factor* Prothioconazole +
 Compound I 0.65 + 0.05 77 63 1.22 Tebuconazole + Compound I 0.2 + 0.1 97 81 1.20
 Tebuconazole + Compound I 0.2 + 0.05 43 25 1.74 Tebuconazole + Compound I 0.1 + 0.05 43
 13 3.25 Fluxapyroxad + Compound I 2 + 0.05 100 95 1.05 Benzovindiflupyr + Compound I
 0.12 + 0.05 41 25 1.60 Azoxystrobin + Compound I 0.2 + 0.05 48 23 2.03 Azoxystrobin +
 Compound I 0.05 + 0.05 21 17 1.27 Pyraclostrobin + Compound I 0.05 + 0.05 96 63 1.52
 Chlorothalonil + Compound I 125 + 0.05 48 15 3.11 Chlorothalonil + Compound I 62.5 + 0.05 39
 11 3.64 *SEPTTR = Leaf Blotch of Wheat; *Zymoseptoria tritici* *Observed = Observed percent
 disease control at the test rates *Expected = Percent disease control expected as predicted by the
 Colby equation *ppm = Parts per million *Synergism factor = Observed/Expected
 TABLE-US-00005 TABLE 5 Synergistic Interactions of the Compound of Formula I and Other
 Fungicides In a 3-Day Curative (3DC) *Zymoseptoria tritici* (SEPTTR) Assay-Test 5. Rates
 SEPTTR* Synergism Composition (ppm)* Observed* Expected* Factor* Epoxiconazole +
 Compound I 0.1 + 0.1 100 67 1.48 Epoxiconazole + Compound I 0.05 + 0.1 89 64 1.39

Epoxiconazole + Compound I 0.025 + 0.1 91 58 1.58 Prothioconazole + Compound I 2.6 + 0.1 99 91 1.09 Prothioconazole + Compound I 1.3 + 0.1 98 84 1.16 Prothioconazole + Compound I 0.65 + 0.1 99 82 1.21 Prothioconazole + Compound I 2.6 + 0.05 99 84 1.18 Prothioconazole + Compound I 1.3 + 0.05 94 72 1.30 Prothioconazole + Compound I 0.65 + 0.05 79 68 1.16 Difenoconazole + Compound I 0.1 + 0.1 100 87 1.15 Difenoconazole + Compound I 0.05 + 0.1 100 73 1.36 Difenoconazole + Compound I 0.1 + 0.05 96 76 1.27 Difenoconazole + Compound I 0.05 + 0.05 91 52 1.74 Tebuconazole + Compound I 0.2 + 0.1 96 62 1.56 Tebuconazole + Compound I 0.1 + 0.1 89 60 1.49 Penthiopyrad + Compound I 4 + 0.1 94 64 1.47 Penthiopyrad + Compound I 4 + 0.05 93 54 1.71 Penthiopyrad + Compound I 4 + 0.025 93 42 2.22 Penthiopyrad + Compound I 2 + 0.05 50 45 1.10 Penthiopyrad + Compound I 2 + 0.025 34 31 1.11 Penthiopyrad + Compound I 1 + 0.1 46 39 1.17 Penthiopyrad + Compound I 0.5 + 0.05 29 21 1.39 Picoxystrobin + Compound I 4 + 0.1 85 53 1.60 Picoxystrobin + Compound I 4 + 0.05 68 41 1.67 Picoxystrobin + Compound I 2 + 0.1 89 48 1.86 Picoxystrobin + Compound I 2 + 0.05 59 34 1.74 Picoxystrobin + Compound I 2 + 0.025 21 17 1.25 Picoxystrobin + Compound I 1 + 0.1 83 40 2.08 Picoxystrobin + Compound I 1 + 0.05 42 24 1.73 Picoxystrobin + Compound I 1 + 0.025 25 4 6.25 Picoxystrobin + Compound I 0.5 + 0.1 77 38 2.03 Picoxystrobin + Compound I 0.5 + 0.05 54 21 2.58 *SEPTTR = Leaf Blotch of Wheat; *Zymoseptoria tritici* *Observed = Observed percent disease control at the test rates *Expected = Percent disease control expected as predicted by the Colby equation *ppm = Parts per million *Synergism factor = Observed/Expected

TABLE-US-00006 TABLE 6 Synergistic Interactions of the Compound of Formula I and Other Fungicides In a 1-Day Protectant (1DP) *Zymoseptoria tritici* (SEPTTR) Assay-Test 6. Rates

SEPTTR Synergism Composition (ppm) Observed* Expected* Factor* Mefentrifluconazole + Compound I 0.05 + 0.1 91 74 1.23 Mefentrifluconazole + Compound I 0.1 + 0.05 95 82 1.16 Fluopyram + Compound I 3.2 + 0.1 97 86 1.13 Fluopyram + Compound I 1.6 + 0.1 95 70 1.36 Fluopyram + Compound I 0.8 + 0.1 96 69 1.38 Fluopyram + Compound I 0.4 + 0.1 99 68 1.45 Fluopyram + Compound I 0.2 + 0.1 91 68 1.33 Fluopyram + Compound I 0.1 + 0.1 82 68 1.21 Fluopyram + Compound I 0.8 + 0.05 57 51 1.13 Fluopyram + Compound I 0.4 + 0.05 56 50 1.12

*SEPTTR = Leaf Blotch of Wheat; *Zymoseptoria tritici* *Observed = Observed percent disease control at the test rates *Expected = Percent disease control expected as predicted by the Colby equation *ppm = Parts per million *Synergism factor = Observed/Expected

TABLE-US-00007 TABLE 7 Synergistic Interactions of the Compound of Formula I and Other Fungicides In a 3-Day Curative (3DC) *Zymoseptoria tritici* (SEPTTR) Assay-Test 7. Rates

SEPTTR Synergism Composition (ppm) Observed* Expected* Factor* Mefentrifluconazole + Compound I 0.05 + 0.2 82 74 1.11 Mefentrifluconazole + Compound I 0.05 + 0.1 45 35 1.30 Fluopyram + Compound I 3.2 + 0.2 99 83 1.20 Fluopyram + Compound I 0.8 + 0.2 97 69 1.42 Fluopyram + Compound I 0.4 + 0.2 93 68 1.37 Fluopyram + Compound I 0.2 + 0.2 85 68 1.25 Fluopyram + Compound I 1.6 + 0.1 53 39 1.38 Fluopyram + Compound I 0.8 + 0.1 62 21 2.89 Fluopyram + Compound I 0.4 + 0.1 57 20 2.83 Fluopyram + Compound I 0.2 + 0.1 33 20 1.67 Fluopyram + Compound I 1.6 + 0.05 40 31 1.29 Fluopyram + Compound I 0.8 + 0.05 23 12 2.03 Fluopyram + Compound I 0.4 + 0.05 22 10 2.17 *SEPTTR = Leaf Blotch of Wheat; *Zymoseptoria tritici* *Observed = Observed percent disease control at the test rates *Expected = Percent disease control expected as predicted by the Colby equation *ppm = Parts per million *Synergism factor = Observed/Expected

Claims

1. A synergistic fungicidal mixture, comprising: a fungicidally effective amount of the compound of Formula I, (S)-1,1-bis(4-fluorophenyl) propan-2-yl (3-acetoxy-4-methoxypicolinoyl)-L-alaninate: ##STR00017## and at least one additional fungicide selected from the group consisting

of sterol biosynthesis inhibitors, respiration inhibitors, and multi-site action inhibitors.

2. The mixture of claim 1 wherein the sterol biosynthesis inhibitor is selected from the group consisting of tebuconazole, prothioconazole, difenoconazole, epoxiconazole, and mefentrifluconazole.
 3. The mixture of claim 2 wherein a concentration ratio of the compound of Formula I to tebuconazole is from about 2:1 to about 1:4.
 4. The mixture of claim 2 wherein a concentration ratio of the compound of Formula I to prothioconazole is from about 1:6.4 to about 1:52.
 5. The mixture of claim 2 wherein a concentration ratio of the compound of Formula I to difenoconazole is from about 2:1 to about 1:2.
 6. The mixture of claim 2 wherein a concentration ratio of the compound of Formula I to epoxiconazole is from about 4:1 to about 1:2.
 7. The mixture of claim 2 wherein a concentration ratio of the compound of Formula I to mefentrifluconazole is from about 4:1 to about 1:2.
 8. The mixture of claim 1 wherein respiration inhibitor is selected from the group consisting of benzovindiflupyr, penthiopyrad, fluxapyroxad, bixafen, and fluopyram.
 9. The mixture of claim 8 wherein a concentration ratio of the compound of Formula I to benzovindiflupyr is from about 1:1.2 to about 1:5.
 10. The mixture of claim 8 wherein a concentration ratio of the compound of Formula I to penthiopyrad is from about 1:10 to about 1:160.
 11. The mixture of claim 8 wherein a concentration ratio of the compound of Formula I to fluxapyroxad is from about 1:2.5 to about 1:40.
 12. The mixture of claim 8 wherein a concentration ratio of the compound of Formula I to bixafen is from about 1:2.5 to about 1:10.
 13. The mixture of claim 8 wherein a concentration ratio of the compound of Formula I to fluopyram is from about 1:1 to about 1:32.
 14. The mixture of claim 1 wherein the respiration inhibitor is selected from the group consisting of picoxystrobin, azoxystrobin, and pyraclostrobin.
 15. The mixture of claim 14 wherein a concentration ratio of the compound of Formula I to picoxystrobin is from about 1:5 to about 1:80.
 16. The mixture of claim 14 wherein a concentration ratio of the compound of Formula I to azoxystrobin is from about 2.5:1 to about 1:4.
 17. The mixture of claim 14 wherein a concentration ratio of the compound of Formula I to pyraclostrobin is from about 7:1 to about 1:4.
 18. The mixture of claim 1 wherein the multi-site inhibitor is selected from the group consisting of chlorothalonil and mancozeb.
 19. The mixture of claim 18 wherein a concentration ratio of the compound of Formula I to chlorothalonil is from about 1:325 to about 1:2,600.
 20. The mixture of claim 18 wherein a concentration ratio of the compound of Formula I to mancozeb is from about 1:325 to about 1:3,200.
-