

# US Patent & Trademark Office

## Patent Public Search | Text View

---

United States Patent	12384828
Kind Code	B2
Date of Patent	August 12, 2025
Inventor(s)	Pentecost; Mickey et al.

---

### Engineered extracellular vesicles comprising fusion proteins

---

#### Abstract

Described herein are compositions and techniques related to generation and therapeutic application of artificial synapses. Artificial synapses are engineered extracellular vesicles, including exosomes, which incorporate sticky binders on their surface to anchor signaling domains against biological targets, such as receptors. These engineered additives can be organized in genetic vector constructs, expressed in mammalian cells, wherein the sticky binders attach to extracellular vesicles such as exosomes, thereby presenting their joined signaling domains which are rapidly taken up by recipient cells. Artificial synapses adopt the hallmark biophysical and biochemical features of extracellular vesicles, allowing for rapid deployment and scale-up. Importantly, this strategy can allow for kinetically favorable signal generation and signal propagation. This includes, for example, increasing density of agonist presentation to support receptor clustering—an onerous barrier for traditional receptor targeting strategies.

---

**Inventors:** Pentecost; Mickey (West Hollywood, CA), Bartkowski; Wojciech (La Crescenta, CA)

**Applicant:** DIADEM BIOTHERAPEUTICS, INC. (Torrance, CA)

**Family ID:** 1000008748815

**Assignee:** Diadem Biotherapeutics, Inc. (Torrance, CA)

**Appl. No.:** 18/056029

**Filed:** November 16, 2022

#### Prior Publication Data

<b>Document Identifier</b>	<b>Publication Date</b>
US 20230312678 A1	Oct. 05, 2023

#### Related U.S. Application Data

## Publication Classification

**Int. Cl.:** **C07K14/705** (20060101); **A61K35/17** (20250101); **A61K47/54** (20170101); **A61K47/58** (20170101); **A61K47/62** (20170101); **A61K47/69** (20170101); **A61P27/02** (20060101); **A61P37/06** (20060101); **C07K14/75** (20060101); **C12N9/14** (20060101); A61K38/00 (20060101)

**U.S. Cl.:**

**CPC** **C07K14/70532** (20130101); **A61K35/17** (20130101); **A61K47/544** (20170801); **A61K47/605** (20170801); **A61K47/62** (20170801); **A61K47/69** (20170801); **A61K47/6901** (20170801); **A61K47/6911** (20170801); **A61K47/6917** (20170801); **A61P27/02** (20180101); **A61P37/06** (20180101); **C07K14/705** (20130101); **C07K14/70521** (20130101); **C07K14/7056** (20130101); **C07K14/70596** (20130101); **C07K14/75** (20130101); **C12N9/14** (20130101); A61K38/00 (20130101); C07K14/70503 (20130101); C07K2319/01 (20130101); C07K2319/02 (20130101); C07K2319/035 (20130101); C07K2319/055 (20130101); C07K2319/30 (20130101); C07K2319/74 (20130101); C07K2319/75 (20130101); C07K2319/912 (20130101); C07K2319/915 (20130101); C12Y304/24081 (20130101)

## Field of Classification Search

**CPC:** C07K (14/70532); C07K (14/70521); C07K (14/70596); C07K (2319/035); C07K (2319/055); C07K (2319/30); C07K (2319/75); C07K (2319/92); C07K (2319/915); A61K (47/6917); A61K (38/00); C12Y (304/24081)

---

## References Cited

### U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
3773919	12/1972	Boswell et al.	N/A	N/A
4186183	12/1979	Alving et al.	N/A	N/A
4217344	12/1979	Handjani et al.	N/A	N/A
4235871	12/1979	Papahadjopoulos et al.	N/A	N/A
4261975	12/1980	Fullerton et al.	N/A	N/A
4485054	12/1983	Mezei et al.	N/A	N/A
4501728	12/1984	Geho et al.	N/A	N/A
4774085	12/1987	Fidler	N/A	N/A
4816567	12/1988	Cabilly et al.	N/A	N/A
4837028	12/1988	Allen	N/A	N/A
4873316	12/1988	Meade et al.	N/A	N/A
4897355	12/1989	Eppstein et al.	N/A	N/A
4946787	12/1989	Eppstein et al.	N/A	N/A

5049386	12/1990	Eppstein et al.	N/A	N/A
5585089	12/1995	Queen et al.	N/A	N/A
5693780	12/1996	Newman et al.	N/A	N/A
7704964	12/2009	Delcayre et al.	N/A	N/A
9546371	12/2016	Mamoun et al.	N/A	N/A
9611481	12/2016	Mamoun	N/A	N/A
10195290	12/2018	Dooley et al.	N/A	N/A
10370663	12/2018	Lotvall et al.	N/A	N/A
10617768	12/2019	Lu et al.	N/A	N/A
10695443	12/2019	Lotvall et al.	N/A	N/A
10723782	12/2019	Lewis et al.	N/A	N/A
11260076	12/2021	Copik	N/A	N/A
11578116	12/2022	Pentecost et al.	N/A	N/A
11746138	12/2022	Pentecost et al.	N/A	N/A
11851470	12/2022	Riazifar	N/A	C07K 14/70521
2016/0137716	12/2015	El Andaloussi et al.	N/A	N/A
2017/0087087	12/2016	Leonard et al.	N/A	N/A
2017/0258938	12/2016	Lotvall et al.	N/A	N/A
2017/0333479	12/2016	Copik et al.	N/A	N/A
2018/0015182	12/2017	Lu et al.	N/A	N/A
2018/0117117	12/2017	Choi et al.	N/A	N/A
2018/0135056	12/2017	Lotvall et al.	N/A	N/A
2018/0236104	12/2017	Lotvall et al.	N/A	N/A
2019/0015333	12/2018	Lu et al.	N/A	N/A
2019/0060483	12/2018	Dooley et al.	N/A	N/A
2019/0117792	12/2018	Dooley et al.	N/A	N/A
2019/0151456	12/2018	McConnell et al.	N/A	N/A
2019/0167810	12/2018	Hean et al.	N/A	N/A
2019/0202892	12/2018	Lewis et al.	N/A	N/A
2019/0224331	12/2018	Wiklander	N/A	N/A
2019/0290585	12/2018	Wiklander	N/A	N/A
2019/0388347	12/2018	Wiklander et al.	N/A	N/A
2020/0054686	12/2019	Rodriguez-Borlado et al.	N/A	N/A
2020/0062813	12/2019	Nordin et al.	N/A	N/A
2020/0109183	12/2019	Wiklander et al.	N/A	N/A
2020/0155703	12/2019	Lotvall et al.	N/A	N/A
2020/0163998	12/2019	Park et al.	N/A	N/A
2020/0206360	12/2019	Choi et al.	N/A	N/A
2020/0207833	12/2019	El Andaloussi et al.	N/A	N/A
2020/0222556	12/2019	Dooley et al.	N/A	N/A
2020/0347112	12/2019	McConnell et al.	N/A	N/A
2020/0407419	12/2019	Lewis et al.	N/A	N/A
2021/0030850	12/2020	Leonard et al.	N/A	N/A
2021/0371497	12/2020	Pentecost et al.	N/A	N/A
2021/0379198	12/2020	Pentecost et al.	N/A	N/A
2022/0411481	12/2021	Pentecost et al.	N/A	N/A

## FOREIGN PATENT DOCUMENTS

<b>Patent No.</b>	<b>Application Date</b>	<b>Country</b>	<b>CPC</b>
3073162	12/2018	CA	N/A
0264166	12/1995	EP	N/A
2021508691	12/2020	JP	N/A
2021515570	12/2020	JP	N/A
2021532736	12/2020	JP	N/A
WO-9116024	12/1990	WO	N/A
WO-9117424	12/1990	WO	N/A
WO-9614057	12/1995	WO	N/A
2017081082	12/2016	WO	N/A
2018015535	12/2017	WO	N/A
2018129207	12/2017	WO	N/A
2019027847	12/2018	WO	N/A
WO-2019035057	12/2018	WO	N/A
2019178334	12/2018	WO	N/A
WO-2019236082	12/2018	WO	N/A
2020154746	12/2019	WO	N/A
2020257710	12/2019	WO	N/A
2021159016	12/2020	WO	N/A

## **OTHER PUBLICATIONS**

Audagnotto M and Dal Pararo M. (2017) Computational and Structural Biotechnology Journal 15:307-319. (doi.org/10.1016/j.csbj.2017.03.004). cited by examiner

Geneovese S, et al. (2009) Phytochemistry. 70:1082-1091. (doi: 10.1016/j.phytochem.2009.06.016). cited by examiner

Clarke CF and Allan CM. (Jun. 25, 2015) Nature. 522:427. cited by examiner

Resh MD. (Jul. 2016) Prog Lipid Res. 63:120-131. (doi:10.1016/j.plipres.2016.05.002). cited by examiner

Trinh R, et al. (2004) Molecular Immunology. 40:717-722. (doi: 10.1016/j.molimm.2003.08.006). cited by examiner

Certified Copy of U.S. Appl. No. 62/864,566, filed Jun. 21, 2019, 6 pages. cited by applicant

Certified Copy of U.S. Appl. No. 62/875,001, filed Jul. 17, 2019, 9 pages. cited by applicant

Riazifar et al., Stem Cell Extracellular Vesicles: Extended Messages of Regeneration; Annu Rev Pharmacol Toxicol., 2017, 57:125-154. cited by applicant

Riazifar et al., Stem Cell-Derived Exosomes as Nanotherapeutics for Autoimmune and Neurodegenerative Disorders, ACS Nano. 2019, 13(6):6670-6688. cited by applicant

Yáñez-Mó, et al., Biological properties of extracellular vesicles and their physiological functions, J Extracell Vesicles, 2015(4):27066. Published May 14, 2015. doi:10.3402/jev.v4.27066. cited by applicant

Yadid et al., Endothelial extracellular vesicles contain protective proteins and rescue ischemia-reperfusion injury in a human heart-on-chip, 2020, Science Translation Medicine 12, 17 pages. cited by applicant

de Abreu et al., Native and bioengineered extracellular vesicles for cardiovascular therapeutics, Nature Reviews Cardiology, 2020, 17(11), 685-697. cited by applicant

Zhang et al., Characterization of Protein Profiling and mRNA Expression of LLC Exosomes, 2019, Protein J 38:586-597 (2019). cited by applicant

Zha et al., Extracellular vesicles: An overview of biogenesis, function, and role in breast cancer, Tumor Biology, 2017, pp. 1-7. cited by applicant

Tan et al., Thrombin stimulated platelet-derived exosomes inhibit platelet-derived growth factor

receptor-beta expression in vascular smooth muscle cells, *Cellular Physiology and Biochemistry*, 2016, 38:2348-2365. cited by applicant

Kalluri et al., The biology, function, and biomedical applications of exosomes, *Science* 2020, 367, 17 pages. cited by applicant

Resh, Fatty acylation of proteins: new insights into membrane targeting of myristoylated and palmitoylated proteins, *Biochimica et Biophysica Acta (BBA)—Molecular Cell Research*, 1999, vol. 1451, Issue 1, pp. 1-17, doi.org/10.1016/S0167-4889(99)00075-0. cited by applicant

Alberts et al., *Membrane Proteins*, *Molecular Biology of the Cell*, 2020, 4th edition, New York: Garland Science, 2002, 16 pages, <https://www.ncbi.nlm.nih.gov/books/NBK26878/>. cited by applicant

Apolloni et. al., H-ras but Not K-ras Traffics to the Plasma Membrane through the Exocytic Pathway, *Molecular and Cellular Biology*, Apr. 2000, 20 (7) 2475-2487, DOI: 10.1128/MCB.20.7.2475-2487.2000. cited by applicant

Dawaliby et. al., Phosphatidylethanolamine Is a Key Regulator of Membrane Fluidity in Eukaryotic Cells, *Membrane Biology*, 2016, vol. 291(7), 10 pages, doi.org/10.1074/jbc.M115.706523. cited by applicant

Deschenes, *Protein Palmitoylation*, *Encyclopedia of Biological Chemistry (Second Edition)*, Academic Press, 2013, pp. 645-647, ISBN 9780123786319, <https://doi.org/10.1016/B978-0-12-378630-2.00022-0>. cited by applicant

Palsuledesai et al., Protein Prenylation: Enzymes, Therapeutics, and Biotechnology Applications, *ACS Chemical Biology* 2015 10 (1), 51-62, DOI: 10.1021/cb500791f. cited by applicant

Hung et al., Stabilization of exosome-targeting peptides via engineered glycosylation, *J Biol Chem*, Mar. 27, 2015;290(13):8166-72, doi: 10.1074/jbc.M114.621383. cited by applicant

O'Shea et al., Peptide 'Velcro': design of a heterodimeric coiled coil, *Curr Biol*. Oct. 1, 1993;3(10):658-67. doi: 10.1016/0960-9822(93)90063-t. PMID: 15335856. cited by applicant

Udenwobe, et al., Myristoylation: An Important Protein Modification in the Immune Response, *Frontiers in Immunology*, vol. 8, 2017, DOI=10.3389/fimmu.2017.00751. cited by applicant

Kinoshita, Biosynthesis and biology of mammalian GPI-anchored proteins *Open Biol*. 2020, 16 pages, 10190290, <http://doi.org/10.1098/rsob.190290>. cited by applicant

Chen et al., Fusion protein linkers: property, design and functionality, *Adv Drug Deliv Rev*. 2013, 65(10):1357-1369 doi:10.1016/j.addr.2012.09.039. cited by applicant

Müller et al., Protein fusions to coiled-coil domains, *Methods Enzymol*. 2000; 328:261-82. doi: 10.1016/s0076-6879(00)28402-4. PMID: 11075350. cited by applicant

Whitford et al., Exosome manufacturing status, *Future Med Chem*. May 2019; 11(10):1225-1236. doi: 10.4155/mc-2018-0417. PMID: 31280675. cited by applicant

Patel et al., Towards rationally designed biomanufacturing of therapeutic extracellular vesicles: impact of the bioproduction microenvironment, *Biotechnol Adv*. Dec. 2018; 36(8):2051-2059. doi: 10.1016/j.biotechadv.2018.09.001. Epub Sep. 12, 2018. PMID: 30218694; PMCID: PMC6250573. cited by applicant

Ng et al., Bioprocess decision support tool for scalable manufacture of extracellular vesicles, *Biotechnol Bioeng*. Feb. 2019; 116(2):307-319. doi: 10.1002/bit.26809. Epub Nov. 8, 2018. PMID: 30063243; PMCID: PMC6322973. cited by applicant

Paganini et al., Scalable Production and Isolation of Extracellular Vesicles: Available Sources and Lessons from Current Industrial Bioprocesses, *Biotechnol J*. Oct. 2019; 14 (10):e1800528. doi: 10.1002/biot.201800528. Epub Jul. 8, 2019. PMID: 31140717. cited by applicant

Zhang et al., Exosome: A Review of Its Classification, Isolation Techniques, Storage, Diagnostic and Targeted Therapy Applications, *Int J Nanomedicine*, Sep. 22, 2020;15:6917-6934. doi: 10.2147/IJN.S264498. PMID: 33061359; PMCID: PMC7519827. cited by applicant

Kluszczynska et al., Methods for the Determination of the Purity of Exosomes, *Curr Pharm Des*. 2019; 25(42):4464-4485. doi: 10.2174/1381612825666191206162712. PMID: 31808383. cited by

applicant

Nolan et al. Analysis of Individual Extracellular Vesicles by Flow Cytometry, *Methods Mol Biol.* 2018; 1678:79-92. doi: 10.1007/978-1-4939-7346-0\_5. PMID: 29071676. cited by applicant

Doyle et al., Overview of Extracellular Vesicles, Their Origin, Composition, Purpose, and Methods for Exosome Isolation and Analysis, *Cells*, Jul. 15, 2019; 8(7):727. doi: 10.3390/cells8070727. PMID: 31311206; PMCID: PMC6678302. cited by applicant

Pugholm et al., Antibody-Based Assays for Phenotyping of Extracellular Vesicles, *Biomed Res Int.* 2015; 2015:524817. doi: 10.1155/2015/524817. Epub Dec. 3, 2015. PMID: 26770974; PMCID: PMC4681819. cited by applicant

Shao et al., New Technologies for Analysis of Extracellular Vesicles, *Chem Rev.* Feb. 28, 2018; 118(4):1917-1950. doi: 10.1021/acs.chemrev.7b00534. Epub Jan. 31, 2018. PMID: 29384376; PMCID: PMC6029891. cited by applicant

Elshaer et al., Adipose stem cells and their paracrine factors are therapeutic for early retinal complications of diabetes In the Ins2Akita mouse, *Stem Cell Research & Therapy*, 2018, 9:322, 18 pages. cited by applicant

Jha et al., TSG-6 in conditioned media from adipose mesenchymal stem cells protects against visual deficits in mild traumatic brain injury model through neurovascular modulation, *Stem Cell Research & Therapy*, 2019, 10:318, 15 pages. cited by applicant

Jha et al., Concentrated conditioned media from adipose tissue derived mesenchymal stem cells mitigates visual deficits and retinal inflammation following mild traumatic brain injury, *International Journal of Molecular Sciences*, 2018. 19:1-22. cited by applicant

ISR and WO for PCT/US2021/016949 dated Jul. 22, 2021, 12 pages. cited by applicant

Andreu et al., Tetraspanins in Extracellular Vesicle Formation and Function, *Frontiers in Immunology*, 2014, vol. 5 (442), pp. 1-12. cited by applicant

Armstrong et al., Re-Engineering Extracellular Vesicles as Smart Nanoscale Therapeutics, *ACS Nano*, 2017, vol. 11(1), pp. 69-83. cited by applicant

“Engineered”, Lexico, available online at <https://www.lexico.com/en/definition/engineered>, 4 pages (accessed on Mar. 7, 2022) (Year: 2022). cited by applicant

“Homology”, Encyclopedia Britannica, available online at [www.britannica.com/science/homology-evolution](http://www.britannica.com/science/homology-evolution), 3 pages at p. 1, 1st paragraph (accessed on Mar. 8, 2022) (Year: 2022). cited by applicant

Kanduc et al., Homology, Similarity and Identity in Peptide Epitope Immunodefinition, *Journal of Peptide Science*, 2012, vol. 18, pp. 487-494. cited by applicant

Pearson et al., An Introduction to Sequence Similarity (“Homology”) Searching, *Curr Protoc Bioinformatics*, 2013, 9 pages. cited by applicant

Samudrala et al., Difference between Homology, Identity and Similarity, available on line at <http://www.bio.net/mm/proteins/1998-July/006538.html>, 1 page (1998) (Year: 1998). cited by applicant

Yang et al., Engineering of Fc Fragments with Optimized Physicochemical Properties Implying Improvement of Clinical Potentials for Fc-Based Therapeutics, *Frontiers in Immunology*, 2018, vol. 8, 14 pages. cited by applicant

Barile et al., Exosomes: Therapy delivery tools and biomarkers of diseases *Pharmacology and Therapeutics*, 2017, 174:63-78 (<http://dx.doi.org/10.1016/j.pharmthera.2017-02-020>). cited by applicant

Ferguson et al., Exosomes as therapeutics: the implications of molecular composition and exosomal heterogeneity (2016) *Journal of Controlled Release*. 228:179-190 (<http://dx.doi.org/10.1016/j.conrel.2016.02.037>). cited by applicant

Murphy, et al., Extracellular vesicle-based therapeutics: natural versus engineered targeting and trafficking (2019) *Experimental & Molecular Medicine*. 51:32 (<https://doi.org/10.1038/s12276-019-0223-5>). cited by applicant

Liu et al., Design strategies and application progress of therapeutic exosomes (2019) *Theranostics*.

9(4):1015-1028 (doi: 10.7150/thno.30853). cited by applicant

Supp ESR for 21750784.7 dated Apr. 3, 2024, 11 pages. cited by applicant

ISR/WO for PCT/US2024/010433 dated Apr. 17, 2024, 15 pages. cited by applicant

Umeda et al., Structural insights into tetraspanin CD9 function,, supplemental information, Nat. Commun. 11:11 (2020). cited by applicant

Umeda et al., Structural insights into tetraspanin CD9 function, Nat. Commun. 11:152 (2020). cited by applicant

UniProt Accession No. A0A024RB05, 5 pages (2014). cited by applicant

Yang et al., Extracellular vesicles as a platform for membrane-associated therapeutic protein delivery, J. Extracellular Vesicles 2018, 7:15. cited by applicant

Adjei, Akwete, and Garren, Julie. Pulmonary delivery of peptide drugs: effect of particle size on bioavailability of leuprolide acetate in healthy male volunteers. Pharmaceutical Research 7(6):565-569 (1990). cited by applicant

Ahmad, Imran, and Allen, Theresa M. Antibody-mediated specific binding and cytotoxicity of liposome-entrapped doxorubicin to lung cancer cells in vitro. Cancer Research 52(17):4817-4820 (1992). cited by applicant

Allen, Loyd V. et al. Remington: The Science and Practice of Pharmacy, 22nd Edition. Pharmaceutical Press (2012). cited by applicant

Anderson, Paula J. et al. Effect of cystic fibrosis on inhaled aerosol boluses. 1-3. Am Rev Respir Dis 140(5):1317-1324 (1989). cited by applicant

Banerji, Julian. et al. A lymphocyte-specific cellular enhancer is located downstream of the joining region in immunoglobulin heavy chain genes. Cell 33(3):729-740 (1983). cited by applicant

Behr, Jean-Paul. Gene transfer with synthetic cationic amphiphiles: prospects for gene therapy. Bioconjugate Chemistry 5(5):382-389 (1994). cited by applicant

Bird, Robert E. et al. Single-chain Antigen-binding Proteins. Science 242(4877):423-426 (1988). cited by applicant

Blaese, M. et al. Vectors in cancer therapy: how will they deliver? Cancer Gene Therapy 2(4):291-297 (1995). cited by applicant

Brinkmann, Ulrich et al. The making of bispecific antibodies. MAbs 9(2):182-212 (2017). cited by applicant

Byrne, G. W., and Ruddle, F. H. Multiplex gene regulation: a two-tiered approach to transgene regulation in transgenic mice. Proceedings of the National Academy of Sciences 86(14):5473-5477 (1989). cited by applicant

Byron, Peter R. Determinants of drug and polypeptide bioavailability from aerosols delivered to the lung. Advanced Drug Delivery Reviews 5(1-2):107-132 (1990). cited by applicant

Calame, Kathryn, and Eaton, Suzanne. Transcriptional controlling elements in the immunoglobulin and T cell receptor loci. Advances in Immunology 43:235-275 (1988). cited by applicant

Camper, Sally A., and Tilghman, Shirley M. Postnatal repression of the alpha-fetoprotein gene is enhancer independent. Genes and Development 3(4):537-546 (1989). cited by applicant

Chothia, Cyrus et al. Canonical Structures for the Hypervariable Regions of Immunoglobulins. Journal of Molecular Biology 196(4):901-917 (1987). cited by applicant

Chothia, Cyrus et al. Conformations of Immunoglobulin Hypervariable Regions. Nature 342(6252):877-883 (1989). cited by applicant

Crystal, Ronald G. Transfer of genes to humans: early lessons and obstacles to success. Science 270(5235):404-410 (1995). cited by applicant

Damms, Bob; et al. The cost of delivering drugs without needles. Bio/Technology 13.12: 1438-1440. (1995). cited by applicant

Dordunoo, S. K. et al. Preformulation studies on solid dispersions containing triamterene or temazepam in polyethylene glycols or Gelucire 44/14 for liquid filling of hard gelatin capsules. Drug Development and Industrial Pharmacy 17(12):1685-1713 (1991). cited by applicant

Edlund, Thomas. et al. Cell-specific expression of the rat insulin gene: evidence for role of two distinct 5' flanking elements. *Science* 230(4728):912-916 (1985). cited by applicant

French, Donna L. et al. The influence of formulation on emission, deaggregation and deposition of dry powders for inhalation. *Journal of Aerosol Science* 27(5):769-783 (1996). cited by applicant

Gao, X, and HUANG, Long-bin. Cationic liposome-mediated gene transfer. *Gene Therapy* 2(10):710-722 (1995). cited by applicant

Goh, Wei Jiang. et al. Bioinspired cell-derived nanovesicles versus exosomes as drug delivery systems: a cost-effective alternative. *Scientific Reports* 7(1):14322, 1-10 (2017). cited by applicant

Gonda, Igor. Aerosols for delivery of therapeutic and diagnostic agents to the respiratory tract. *Critical Reviews in Therapeutic Drug Carrier Systems* 6(4):273-313 (1990). cited by applicant

Greenfield, Edward A. *Antibodies, A Laboratory Manual*. Second Edition. Cold Spring Harbor Press (2013). cited by applicant

Holliger, Philipp et al. "Diabodies": Small Bivalent and Bispecific Antibody Fragments. *Proceedings of the National Academy of Sciences of the United States of America* 90(14):6444-6448 (1993). cited by applicant

Hornyak, Gabor L. et al. *Introduction to Nanoscience and Nanotechnology*. CRC Press (2008). cited by applicant

Huston, James S. et al. Protein Engineering of Antibody Binding Sites: Recovery of Specific Activity in an Anti-digoxin Single-chain Fv Analogue Produced in *Escherichia coli*. *PNAS USA* 85(16):5879-5883 (1988). cited by applicant

Kabat, Elvin A. et al. *Sequences of proteins of immunological interest*. 5th Edition. Public Health Service, National Institutes of Health, Bethesda MD (1991). cited by applicant

Kaufman, Randal J. et al. Translational efficiency of polycistronic mRNAs and their utilization to express heterologous genes in mammalian cells. *The EMBO Journal* 6(1):187-193 (1987). cited by applicant

Kessel, Michael, and Peter Gruss. Murine developmental control genes. *Science* 249(4967):374-379 (1990). cited by applicant

Kobayashi, Shinzo et al. Pulmonary delivery of salmon calcitonin dry powders containing absorption enhancers in rats. *Pharmaceutical Research* 13(1):80-83 (1996). cited by applicant

Kohler, G., and Milstein, C. Derivation of specific antibody-producing tissue culture and tumor lines by cell fusion. *European Journal of Immunology* 6(7):511-519 (1976). cited by applicant

Kontermann, Roland, and Dubel, Stefan. *Antibody Engineering*. Springer Verlag (2001). cited by applicant

Lanzavecchia, Antonio, and Scheidegger, D. The use of hybrid hybridomas to target human cytotoxic T lymphocytes. *European journal of immunology* 17(1):105-111 (1987). cited by applicant

Lasic, D.D. *Liposomes: From Physics to Applications*. Elsevier (1993). cited by applicant

Morrison, Sherie L. et al. Chimeric Human Antibody Molecules: Mouse Antigen-Binding Domains With Human Constant Region Domains. *PNAS USA* 81(21):6851-6855 (1984). cited by applicant

New, R C C. *Liposomes. A practical approach*, IRL Press (1990). cited by applicant

Niven, Ralph W. et al. The pulmonary absorption of aerosolized and intratracheally instilled rhG-CSF and monoPEGylated rhG-CSF. *Pharmaceutical Research* 12(9):1343-1349 (1995). cited by applicant

Osol, Arthur. *Remington's Pharmaceutical Sciences*, 16th Edition. Mack Publishing Company (1980). cited by applicant

Patton, J. S. et al. Bioavailability of pulmonary delivered peptides and proteins: alpha-interferon, calcitonins and parathyroid hormones. *Journal of Controlled Release* 28(1-3):79-85 (1994). cited by applicant

Patton, John S, and Platz, Robert M. (D) Routes of delivery: Case studies:(2) Pulmonary delivery of peptides and proteins for systemic action. *Advanced Drug Delivery Reviews* 8(2-3): 179-196



(1992). cited by applicant

PCT/US2021/016949 International Preliminary Report on Patentability dated Aug. 18, 2022. cited by applicant

Pinkert, Carl A. et al. An albumin enhancer located 10 kb upstream functions along with its promoter to direct efficient, liver-specific expression in transgenic mice. *Genes and Development* 1(3):268-276 (1987). cited by applicant

Poljak, Roberto J. Production and Structure of Diabodies. *Structure* 2(12): 1121-1123 (1994). cited by applicant

Queen, Cary, and Baltimore, David. Immunoglobulin gene transcription is activated by downstream sequence elements. *Cell* 33(3):741-748 (1983). cited by applicant

Remy, Jean-Serge et al. Gene transfer with a series of lipophilic DNA-binding molecules. *Bioconjugate Chemistry* 5(6):647-654 (1994). cited by applicant

Riechmann, Lutz et al. Reshaping Human Antibodies for Therapy. *Nature* 332(6162):323-327 (1988). cited by applicant

Rudd, S., and R. H. Muller. In vitro phagocytosis assay of nano-and microparticles by chemiluminescence. I. Effect of analytical parameters, particle size and particle concentration. *Journal of Controlled Release* 22(3):263-271 (1992). cited by applicant

Sambrook, Joseph, and Green, Michael R. *Molecular Cloning: A Laboratory Manual*, 4th edition. Cold Spring Harbor Laboratory Press (2012). cited by applicant

Sambrook, Joseph et al. *Molecular Cloning: A Laboratory Manual*, 2nd Edition. Cold Spring Harbor Laboratory Press (1989). cited by applicant

Seed, Brian. An LFA-3 cDNA encodes a phospholipid-linked membrane protein homologous to its receptor CD2. *Nature* 329(6142):840-842 (1987). cited by applicant

Sheen, Pai-Chang et al. Bioavailability of a poorly water-soluble drug from tablet and solid dispersion in humans. *Journal of Pharmaceutical Sciences* 80(7):712-714 (1991). cited by applicant

Singleton, Paul, and Sainsbury, Diana. *Dictionary of Microbiology and Molecular Biology*, 3rd Edition revised. John Wiley and Sons (2006). cited by applicant

Singleton, Paul. *Dictionary of DNA and Genome Technology*, Third Edition. Wiley-Blackwell (2012). cited by applicant

Smith, Michael B. *March's Advanced Organic Chemistry, Reactions, Mechanisms, and Structure*. 7th Edition. John Wiley and Sons (2013). cited by applicant

Sonnhammer, Erik LL. et al. A hidden Markov model for predicting transmembrane helices in protein sequences. *International Conference on Intelligent Systems for Molecular Biology* 6:175-182 (1998). cited by applicant

Tabata, Yasuhiko, and Ikada, Yoshito. Macrophage phagocytosis of biodegradable microspheres composed of L-lactic acid/glycolic acid homo-and copolymers. *Journal of Biomedical Materials Research* 22(10):837-858 (1988). cited by applicant

Tan, Lina et al. Recent advances of exosomes in immune modulation and autoimmune diseases. *Autoimmunity* 49(6):357-365 (2016). cited by applicant

Tansey, Ian P. The challenges in the development of metered dose inhalation aerosols using ozone-friendly propellants. *Spray Technol and Market* 4:26-29 (1994). cited by applicant

Timsina, M. P. et al. Drug delivery to the respiratory tract using dry powder inhalers. *International Journal of Pharmaceutics* 101(1-2):1-13 (1994). cited by applicant

Visser, J. An Invited Review: Van der Waals and other cohesive forces affecting powder fluidization. *Powder Technology* 58(1):1-10 (1989). cited by applicant

Wall, Doris A. Pulmonary absorption of peptides and proteins. *Drug Delivery* 2(1):1-20 (1995). cited by applicant

Winkler, Johannes. Oligonucleotide conjugates for therapeutic applications. *Therapeutic Delivery* 4(7):791-809 (2013). cited by applicant

Winoto, Astar, and Baltimore, David. A novel, inducible and T cell-specific enhancer located at the

3' end of the T cell receptor alpha locus. The EMBO Journal 8(3):729-733 (1989). cited by applicant  
Zanen, Pieter et al. The optimal particle size for parasympatholytic aerosols in mild asthmatics. International Journal of Pharmaceutics 114(1):111-115 (1995). cited by applicant  
Zapata, Gerardo et al. Engineering Linear F(ab')<sub>2</sub> Fragments for Efficient Production in *Escherichia coli* and Enhanced Antiproliferative Activity. Protein Engineering 8(10):1057-1062 (1995). cited by applicant  
Zha, Quan Bin et al. Extracellular vesicles: An overview of biogenesis, function, and role in breast cancer. Tumor Biology 39(2):1-7 (2017). cited by applicant

---

*Primary Examiner:* Landsman; Robert S

*Attorney, Agent or Firm:* WILSON SONSINI GOODRICH & ROSATI

---

## **Background/Summary**

CROSS-REFERENCE TO RELATED APPLICATIONS (1) This application is a division of U.S. application Ser. No. 17/377,550, filed Jul. 16, 2021, now U.S. Pat. No. 11,578,116, which is a continuation of International Application No. PCT/US2021/016949, filed Feb. 5, 2021, which designated the U.S., and also claims the benefit of priority under 35 U.S.C. § 119(e) of U.S. Provisional Application No. 62/970,374, filed Feb. 5, 2020, the contents of each of which are incorporated herein by reference in their entirety.

### **SEQUENCE LISTING**

(1) The instant application contains a Sequence Listing which has been submitted electronically in XML format and is hereby incorporated by reference in its entirety. Said XML copy, created on Nov. 9, 2022, is named 085172\_000001USD1\_ST.26.xml and is 724,326 bytes in size.

### **FIELD OF THE INVENTION**

(2) This invention relates to the generation of artificial synapses or extracellular vesicles, including features of extracellular vesicles engineered to deliver signaling, for therapeutic use, including treatment of immune diseases and cancer.

### **BACKGROUND**

(3) Extracellular vesicles (EVs) play a critical role in intercellular communication by transferring microRNAs, lipids, and proteins to neighboring cells. The delivery of encapsulated molecules within EVs is a highly promising strategy as a therapeutic platform in many contexts, exploiting the unique biophysical and biochemical characteristics of extracellular vesicles (EVs). However, there remains a great need in the art for a flexible and dynamic platform, where specific biological signals can be reliably targeted without off-target effects and that provide a robust cellular response to achieve a therapeutic effect, such as modulating inflammation.

### **SUMMARY**

(4) The compositions and methods provided herein are based, in part, on the discovery that extracellular vesicles can be used to express engineered fusion polypeptides that can modulate biological signal generation. These engineered vesicles, also termed artificial synapses, adopt the hallmark biophysical and biochemical features of extracellular vesicles, but are further engineered with vesicle targeting domains (e.g., sticky binders) and signaling domains, optionally joined by a linker with specific functions. The fusion polypeptides provided herein are designed and produced as nucleic acid constructs (e.g., vectors) and expressed in cells, such as mammalian cells. In particular, the vesicle targeting domain of each fusion polypeptide anchors the polypeptide to the extracellular vesicle lipid membrane, thereby presenting the signaling domain(s) of the

polypeptide. The signaling domains on or within the vesicle membrane can make contact with recipient cells via target polypeptides (e.g., receptors on the extracellular surface of the recipient cell). Importantly, this strategy can allow for kinetically favorable signal generation and signal propagation. This includes, for example, increasing density of agonist presentation to support receptor clustering of a target receptor located on a target cell—an onerous barrier for traditional receptor targeting strategies.

(5) This strategy was applied to alter immune checkpoint signaling, by engineering artificial synapses through genetic constructs with lipid binding glycosylphosphatidylinositol (GPI) sticky binders joined with programmed death-ligand 1 (PD-L1) signaling domain, e.g., human programmed death-ligand 1 (hPD-L1), expressed in cells and capable of attachment to exosomes. Isolation, purification, and analysis of artificial synapses revealed a high density of signaling domains of the hPD-L1-GPI fusion polypeptide. The hPD-L1 artificial synapse exosomes further demonstrated enhanced agonist signaling than soluble PD-L1 ligand alone, supporting receptor clustering on a target cell. When applied to a model of experimental autoimmune uveoretinitis (EAU), a statistically significant reduction in EAU symptoms was observed.

(6) Thus, in one aspect, provided herein is an engineered extracellular vesicle or artificial vesicle comprising: at least one fusion polypeptide comprising: at least one protein of interest (POI) domain; and at least one vesicle targeting domain. In some embodiments of any of the aspects, the engineered extracellular vesicle is an exosome. In some embodiments, of any of the aspects, the fusion protein further comprises at least one linker. In some embodiments of any of the aspects, the POI domain can substantially bind to a target polypeptide.

(7) In another aspect, provided herein is an engineered extracellular vesicle comprising: at least one fusion polypeptide comprising: (i) at least one protein of interest (POI) domain or a fragment thereof; and (ii) at least one vesicle targeting domain, wherein the POI domain is in an extracellular position relative to a lipid membrane of the extracellular vesicle.

(8) In another aspect, provided herein is an engineered extracellular vesicle comprising: (a) a first fusion polypeptide comprising: (i) at least one protein of interest (POI) domain or a fragment thereof; and (ii) at least one vesicle targeting domain, wherein the at least one POI domain is in an extracellular position relative to a lipid membrane of the extracellular vesicle, (b) a second fusion polypeptide comprising: (i) at least one protein of interest (POI) domain or a fragment thereof; and (ii) at least one vesicle targeting domain, wherein the POI domain is in an extracellular position relative to a lipid membrane of the extracellular vesicle, and wherein the at least one vesicle targeting domain is within a lipid membrane of the extracellular vesicle.

(9) In another aspect, provided herein is an extracellular vesicle composition comprising: a plurality of artificial synapses, wherein each artificial synapse comprises (i) an extracellular vesicle; (ii) one or more sticky binders; and (iii) one or more signaling domains.

(10) In another aspect, provided herein is a composition comprising a plurality of the engineered extracellular vesicles provided herein.

(11) In another aspect, provided herein is a composition comprising two or more of the engineered extracellular vesicles provided herein.

(12) In another aspect, provided herein is a composition comprising three or more of the engineered extracellular vesicles provided herein.

(13) In another aspect, provided herein is a method of producing the engineered extracellular vesicle or the compositions provided herein, comprising: (a) providing a population of cells expressing a vector construct encoding one or more sticky binder and one or more signaling domains; and (b) isolating a plurality of artificial synapses from the population of cells.

(14) In another aspect, provided herein is a method of producing the engineered extracellular vesicle or the compositions provided herein, comprising: (a) providing a population of cells expressing a vector construct encoding one or more sticky binder and one or more signaling domains; and (b) isolating a plurality of artificial synapses from the population of cells; and (c)

purifying the plurality of artificial synapses from the population of cells.

(15) In another aspect, provided herein is a method of modulating inflammation in a subject, the method comprising: administering a composition comprising a plurality of engineered extracellular vesicles to a subject in need thereof, wherein the engineered extracellular vesicles comprise at least one fusion polypeptide comprising: (i) at least one protein of interest (POI) domain or a fragment thereof; and (ii) at least one vesicle targeting domain.

(16) In another aspect, provided herein is a use of a composition or engineered extracellular vesicle provided herein for the treatment of an inflammatory disease or condition.

(17) In another aspect, provided herein is a use of a composition or engineered extracellular vesicle provided herein for the treatment of an autoimmune disease or condition.

(18) In another aspect, provided herein is a use of a composition or engineered extracellular vesicle provided herein for the treatment of cancer.

(19) In one embodiment of any of the aspects, the engineered extracellular vesicle is an exosome.

(20) In another embodiment of any of the aspects, the protein of interest (POI) domain or a fragment thereof is a N-terminal domain of the fusion polypeptide. In another embodiment of any of the aspects, the POI domain is selected from the group consisting of: Table 1. In another embodiment of any of the aspects, the POI domain is PD-L1 or a fragment thereof. In another embodiment of any of the aspects, the POI domain is PD-L2 or a fragment thereof. In another embodiment of any of the aspects, the POI domain is FGL1 or a fragment thereof. In another embodiment of any of the aspects, the POI domain is 4-1BBL or a fragment thereof. In another embodiment of any of the aspects, the POI domain is CTLA-4 or a fragment thereof. In another embodiment of any of the aspects, the protein of interest (POI) domain is HVEM or a fragment thereof.

(21) In another embodiment of any of the aspects, the vesicle targeting domain is a C-terminal domain of the fusion polypeptide. In another embodiment of any of the aspects, the vesicle targeting domain is in a luminal position relative to the lipid membrane of the extracellular vesicle. In another embodiment of any of the aspects, the vesicle targeting domain is in an exterior position relative to the lipid membrane of the extracellular vesicle. In another embodiment of any of the aspects, the vesicle targeting domain is selected from the group consisting of: Table 3. In another embodiment of any of the aspects, the vesicle targeting domain is selected from the group consisting of: a Glycosylphosphatidylinositol (GPI) anchor, a fatty acylation site, and a prenylation site. In another embodiment of any of the aspects, the vesicle targeting domain is C1C2. In another embodiment of any of the aspects, the vesicle targeting domain is a GPI anchor.

(22) In another embodiment of any of the aspects, the fusion polypeptide comprises at least two POI domains and/or at least two exosome targeting domains.

(23) In another embodiment of any of the aspects, the POI domain substantially binds to one or more of a target polypeptide. In another embodiment of any of the aspects, the target polypeptide is selected from the group consisting of: Table 2.

(24) In another embodiment of any of the aspects, the fusion polypeptide further comprises a peptide linker. In another embodiment of any of the aspects, the fusion polypeptide further comprises a fragment crystallizable region (Fc) domain. In another embodiment of any of the aspects, the linker is in an exterior position relative to the lipid membrane of the extracellular vesicle. In another embodiment of any of the aspects, the linker is a transmembrane linker. In another embodiment of any of the aspects, the linker is in a luminal position relative to the lipid membrane of the extracellular vesicle.

(25) In another embodiment of any of the aspects, the engineered extracellular vesicle does not comprise an endogenous POI polypeptide.

(26) In another embodiment of any of the aspects, the composition further comprises a pharmaceutically acceptable carrier.

(27) In another embodiment of any of the aspects, the one or more sticky binders or the vesicle

targeting domain is selected from the group consisting of: a GPI anchor, a fatty acylation site, and a prenylation site.

(28) In another embodiment of any of the aspects, the signaling domain or the protein of interest comprises one or more of: PD-L1, PD-L2, CTLA-4 (CD152), 4-1BBL (CD137L), HVEM (CD270), FGL1, OX-2 (CD200), Galectin-9, PVR (CD155), Nectin-2 (CD112) isoform alpha, Nectin-2 (CD112) isoform beta, Nectin-2 (CD112) isoform delta, IL-10, TSG-6, B7-H3 (CD276), B7-H4 (VTCN1), B7-H5 (VISTA), B7-H7 (HHLA2), BTNL1, VSIG8, VSIG3 (IGSF11), VSIG4, TIM-3 (HAVCR2), TIM-4 (TIMD4), CEACAM1, BTN3A1, BTN3A2, BTN2A1, BTNL8, BTN2A2, BTN1A1, TIGIT, CD27L (CD70), CD30L (CD153), GITRL, CD40L (CD154), LIGHT (CD258), TL1, CD80, CD86, LFA-3 (CD58), SLAM (CD150), CD40, CD28, CD28H, CD2, LFA-3 (CD58), CD48, CD226, DR3, DcR3, FasL, TIM-1 (CD365), PD-1, or active fragment thereof.

(29) In another embodiment of any of the aspects, the isolating is via size exclusion chromatography. In another embodiment of any of the aspects, the purifying is via multimodal chromatography. In another embodiment of any of the aspects, the method further comprises performing an assay for POI binding to a target polypeptide.

(30) In another embodiment of any of the aspects, the vector construct further encodes a promoter. In another embodiment of any of the aspects, the promoter is a tissue-specific promoter or an inducible promoter.

(31) In one embodiment of any of the aspects, the method further comprises selecting a subject that has or is suspected of having an autoimmune disease or an inflammatory disease or condition. In another embodiment of any of the aspects, the inflammatory disease and/or condition is acute. In another embodiment of any of the aspects, the inflammatory related disease and/or condition is chronic.

(32) In another embodiment of any of the aspects, administering the composition provided herein comprises injection, topical administration, or inhalation.

---

## Description

### A BRIEF DESCRIPTION OF THE DRAWINGS

(1) FIG. 1 shows construct representation of fusion polypeptides for labeling an exosome surface with Type I membrane proteins.

(2) FIG. 2A shows nucleic acid and translated protein sequences of full-length Phosphatidylserine binding: Lactadherin (MFGE8) C1C2. Underlined nucleic acid sequence highlights the sequence translated to the C1C2 protein. Bold and underlined text highlights the C1C2 domain used to anchor signaling domains of interest (i.e., PD-L1 extracellular domain) onto the surface of the Inventors' artificial synapses. FIG. 2B shows nucleic acid and translated protein sequences of full length CD55 (DAF) Glycosylphosphatidylinositol (GPI) anchor. Bold and underlined text highlights the GPI anchor domain used to anchor signaling domains of interest (i.e., PD-L1 extracellular domain) onto the surface of the Inventors' artificial synapses engineered from exosomes. FIGS. 2A-2B disclose SEQ ID NOS: 199-200, and 196-197, respectively, in order of appearance.

(3) FIG. 3 demonstrates the nucleic acid and translated protein sequence for the Fc linker used in genetically engineered constructs is shown in bold and underlined. FIG. 3 discloses SEQ ID NOS: 219-220, respectively, in order of appearance.

(4) FIG. 4A demonstrates nucleic acid and translated protein sequence of human PD-L1 (CD274). Bold and underlined sequence highlights the PD-L1 extracellular domain used in the Inventors' artificial synapses engineered from exosomes. FIG. 4B demonstrates nucleic acid and protein sequence of human PD-L2. Bold and underlined sequence highlights the PD-L2 extracellular domain used in the Inventors' artificial synapses engineered from exosomes. FIG. 4C shows mRNA

and protein sequence of human CTLA-4 (CD152). **CD152** and underlined sequence highlights the CTLA-4 extracellular domain used in the Inventors' artificial synapses. FIGS. **4A-4C** disclose SEQ ID NOS: 1-2, 5-6, and 9-10, respectively, in order of appearance.

(5) FIG. 5A shows an exemplary embodiment of pcDNA5-FRT cloning vector with a gene sequence coding for a fusion polypeptide inserted into a multiple cloning site. FIG. 5B shows an exemplary embodiment of the Gateway® destination vector pEF5-FRT-V5-DEST with a gene sequence coding for a fusion polypeptide inserted into a multiple cloning site. The vectors were used for constitutive high-level expression of fusion polypeptide described herein in mammalian cells. FIG. 5C shows the nucleic acid and protein sequence for the hCTLA4-Fc-GPI fusion polypeptide wherein the text for the signaling domain is **bolded**, Fc linker is underlined, and sticky binder is *italicized*. FIG. 5D shows the nucleic acid and protein sequence for the hPDL1-GPI-P2A-hHVEM-GPI fusion polypeptide wherein the text for the signaling domain hPDL1 and hHVEM are **bolded**, P2A sequence is underlined, and sticky binder GPI is *italicized*. With P2A included, a self-cleaving peptide sequence, artificial synapses with this feature will have both hPDL1-GPI and hHVEM-GPI loaded onto the surface. FIG. 5E shows the nucleic acid and protein sequence for the hPDL1-GPI-P2A-hFGL1-GPI fusion polypeptide wherein the text for the signaling domain hPDL1 and hFGL1 are **bolded**, P2A sequence is underlined, and sticky binder GPI is *italicized*. With P2A included, a self-cleaving peptide sequence, artificial synapses with this feature will have both hPDL1-GPI and FGL1-GPI loaded onto the surface. FIG. 5F shows the nucleic acid and protein sequence for the hPDL1-GPI fusion polypeptide wherein the text for the signaling domain hPDL1 is **bolded** and sticky binder GPI is *italicized*. FIG. 5G shows the nucleic acid and protein sequence for the hPDL1-Fc-GPI fusion polypeptide wherein the text for the signaling domain hPDL1 is **bolded**, Fc is underlined, and sticky binder GPI is *italicized*. FIG. 5H shows the nucleic acid and protein sequence for the hPDL2-Fc-GPI fusion polypeptide wherein the text for the signaling domain hPDL2 is **bolded**, Fc is underlined, and sticky binder GPI is *italicized*. FIG. 5I shows the nucleic acid and protein sequence for the hPDL1-C1C2 fusion polypeptide wherein the text for the signaling domain hPDL1 is **bolded** and sticky binder C1C2 is *italicized*. FIG. 5J shows the nucleic acid and protein sequence for the hPDL2-C1C2 fusion polypeptide wherein the text for the signaling domain hPDL2 is **bolded** and sticky binder C1C2 is *italicized*. FIG. 5K shows the nucleic acid and protein sequence for the 4F2-h41BBL fusion polypeptide wherein the text for the signaling domain h41BBL is **bolded** and sticky binder 4F2 is *italicized*. FIG. 5L shows the nucleic acid and protein sequence for the hPDL1-4Fc-GPI fusion polypeptide wherein the text for the signaling domain hPDL1 is **bolded**, 4Fc is underlined, and sticky binder GPI is *italicized*. FIG. 5M shows the nucleic acid and protein sequence for the Myr-NanoLuc® Luciferase fusion polypeptide wherein the text for the signaling domain NanoLuc® Luciferase is **bolded**, and sticky binder Myr is *italicized*. FIG. 5N shows the nucleic acid and protein sequence for the Myr-mScarlet fusion polypeptide wherein the text for the signaling domain mScarlet is **bolded**, and sticky binder Myr is *italicized*. FIG. 5O shows the nucleic acid and protein sequence for the secreted isoform of hPDL1 (SecPDL1) fusion polypeptide hSecPDL1-GPI wherein the text for the signaling domain hSecPDL1 is **bolded** and sticky binder GPI is *italicized*. FIG. 5P shows the nucleic acid and protein sequence for the Tfr2-h41BBL fusion polypeptide wherein the text for the signaling domain h41BBL is **bolded** and sticky binder Tfr2 is *italicized*. FIG. 5Q shows the nucleic acid and protein sequence for the CD9tm3-h41BBL fusion polypeptide wherein the text for the signaling domain h41BBL is **bolded** and sticky binder CD9tm3 is *italicized*. FIG. 5R shows the nucleic acid and protein sequence for the Myr/Palm-4F2-h41BBL fusion polypeptide wherein the text for the signaling domain h41BBL is **bolded**, sticky binder Myr/Palm is underlined, and sticky binder 4F2 is *italicized*. FIG. 5S shows the nucleic acid and protein sequence for the Myr/Palm-Link-h41BBL fusion polypeptide wherein the text for the signaling domain h41BBL is **bolded**, sticky binder Myr/Palm is *italicized* and underlined, and sticky binder Link (in this embodiment a GSSG linker (SEQ ID NO: 319)) is in regular text (not underlined and not italicized). FIG. 5T shows the nucleic

acid and protein sequence for the hPDL1-Link-GPI fusion polypeptide wherein the text for the signaling domain hPDL1 is bolded, Link is underlined (in this embodiment a GSSG linker (SEQ ID NO: 319)), and sticky binder GPI is italicized. FIG. 5U shows the nucleic acid and protein sequence for the secreted isoform of hPDL1 (SecPDL1) fusion polypeptide hSecPDL1-CD9tm2 wherein the text for the signaling domain hSecPDL1 is bolded and sticky binder CD9tm2 is italicized. FIG. 5V shows the nucleic acid and protein sequence for the secreted isoform of hPDL1 (SecPDL1) fusion polypeptide hSecPDL1-CD9tm2-KRAS wherein the text for the signaling domain hSecPDL1 is bolded, sticky binder CD9tm2 is italicized, and sticky binder KRAS is italicized and underlined. FIG. 5W shows the nucleic acid and protein sequence for the secreted isoform of hPDL1 (SecPDL1) fusion polypeptide hSecPDL1-CD9tm4 wherein the text for the signaling domain hSecPDL1 is bolded and sticky binder CD9tm4 is italicized. FIG. 5X shows the nucleic acid and protein sequence for the secreted isoform of hPDL1 (SecPDL1) fusion polypeptide hSecPDL1-CD81 wherein the text for the signaling domain hSecPDL1 is bolded and sticky binder CD81 is italicized. FIG. 5Y shows the nucleic acid and protein sequence for the hCD200-Fc-GPI fusion polypeptide wherein the text for the signaling domain hCD200 is bolded, Fc is underlined, and sticky binder GPI is italicized, a spacer sequence domain (regular text, not underlined and not italicized) separates hCD200 sequence from the Fc domain, a spacer sequence domain (regular text, not underlined and not italicized) separates Fc sequence from the GPI. FIG. 5Z shows the nucleic acid and protein sequence for the hFGL1-GPI fusion polypeptide wherein the text for the signaling domain hFGL1 is bolded, and sticky binder GPI is italicized. FIG. 5AA shows the nucleic acid and protein sequence for the hGal9-Fc-GPI fusion polypeptide wherein the text for the signaling domain hGal9 is bolded, Fc is underlined, and sticky binder GPI is italicized. FIG. 5BB shows the nucleic acid and protein sequence for the hCD200-GPI fusion polypeptide wherein the text for the signaling domain hCD200 is bolded, and sticky binder GPI is italicized. FIG. 5CC shows the nucleic acid and protein sequence for the hGal9-GPI fusion polypeptide wherein the text for the signaling domain hGal9 is bolded, and sticky binder GPI is italicized. FIG. 5DD shows the nucleic acid and protein sequence for the hHVEM-GPI fusion polypeptide wherein the text for the signaling domain hHVEM is bolded, and sticky binder GPI is italicized. FIG. 5EE shows the nucleic acid and protein sequence for the hPDL2-GPI fusion polypeptide wherein the text for the signaling domain hPDL2 is bolded, and sticky binder GPI is italicized. FIG. 5FF shows the nucleic acid and protein sequence for the hTSG6-GPI fusion polypeptide wherein the text for the signaling domain hTSG6 is bolded, and sticky binder GPI is italicized. FIG. 5GG shows the nucleic acid and protein sequence for the hHVEM-Fc-GPI fusion polypeptide wherein the text for the signaling domain hHVEM is bolded, Fc is underlined, and sticky binder GPI is italicized. FIG. 5HH shows the nucleic acid and protein sequence for the mCTLA4-Fc-GPI fusion polypeptide wherein the text for the signaling domain mCTLA4 is bolded, Fc is underlined, and sticky binder GPI is italicized. FIG. 5II shows the nucleic acid and protein sequence for the mPDL1-C1C2 fusion polypeptide wherein the text for the signaling domain mPDL1 is bolded and sticky binder C1C2 is italicized. FIG. 5JJ shows the nucleic acid and protein sequence for the mPDL1-Fc-GPI fusion polypeptide wherein the text for the signaling domain mPDL1 is bolded, Fc is underlined, and sticky binder GPI is italicized. FIG. 5KK shows the nucleic acid and protein sequence for the mPDL1-GPI fusion polypeptide wherein the text for the signaling domain mPDL1 is bolded and sticky binder GPI is italicized. FIG. 5LL shows the nucleic acid and protein sequence for the mPDL2-C1C2 fusion polypeptide wherein the text for the signaling domain mPDL2 is bolded and sticky binder C1C2 is italicized. FIG. 5MM shows the nucleic acid and protein sequence for the mPDL2-Fc-GPI fusion polypeptide wherein the text for the signaling domain mPDL2 is bolded, Fc is underlined, and sticky binder GPI is italicized. FIG. 5NN shows the nucleic acid and protein sequence for the mPDL1-mFc-GPI fusion polypeptide wherein the text for the signaling domain mPDL2 is bolded, mFc is underlined, and sticky binder GPI is italicized. FIG. 5OO shows the nucleic acid and protein sequence for the mPDL2-GPI fusion polypeptide wherein the text for the signaling domain mPDL2

is bolded and sticky binder GPI is italicized. FIG. 5PP shows the nucleic acid and protein sequence for the mPDL1-GPI-P2A-mHVEM-GPI fusion polypeptide wherein the text for the signaling domain mPDL1 and mHVEM are bolded, P2A sequence is underlined, and sticky binder GPI is italicized. With P2A included, a self-cleaving peptide sequence, artificial synapses with this feature will have both mPDL1-GPI and mHVEM-GPI loaded onto the surface. FIG. 5QQ shows the nucleic acid and protein sequence for the hPDL1-ADAM10 fusion polypeptide wherein the text for the signaling domain mPDL1 is bolded and sticky binder ADAM10 is italicized. FIG. 5RR shows the nucleic acid and protein sequence for the hPDL1-4Fc-CD9tm2 fusion polypeptide wherein the text for the signaling domain hPDL1 is bolded, 4Fc is underlined, and sticky binder CD9tm2 is italicized. FIG. 5SS shows the nucleic acid and protein sequence for the fusion polypeptide hPDL1-4Fc-CD9tm2-KRAS wherein the text for the signaling domain hPDL1 is bolded, sticky binder 4Fc is underlined, sticky binder CD9tm2 is italicized, and sticky binder KRAS is italicized and underlined. FIG. 5TT shows the nucleic acid and protein sequence for the hPDL1-Fc-CD9tm2 fusion polypeptide wherein the text for the signaling domain hPDL1 is bolded, Fc is underlined, and sticky binder CD9tm2 is italicized. FIG. 5UU shows the nucleic acid and protein sequence for the fusion polypeptide hPDL1-Fc-CD9tm2-KRAS wherein the text for the signaling domain hPDL1 is bolded, sticky binder Fc is underlined, sticky binder CD9tm2 is italicized, and sticky binder KRAS is italicized and underlined. FIG. 5VV shows the nucleic acid and protein sequence for the mPDL1-mFc-CD9tm2 fusion polypeptide wherein the text for the signaling domain mouse PDL1 (mPDL1) is bolded, mouse mFc (mFc) is underlined, and sticky binder CD9tm2 is italicized. FIG. 5WW shows the nucleic acid and protein sequence for the fusion polypeptide mPDL1-mFc-CD9tm2-KRAS wherein the text for the signaling domain mPDL1 is bolded, sticky binder mFc is underlined, sticky binder CD9tm2 is italicized, and sticky binder KRAS is italicized and underlined. Wherein mPDL1 and mFc are mouse PDL1 and mouse Fc, respectively. FIGS. 5C-5WW disclose SEQ ID NOS: 223-224, 283-284, 239-240, 225-226, 229-230, 233-234, 227-228, 231-232, 235-238, 243-244, 241-242, 245-282, and 285-316, respectively, in order of appearance. (6) FIG. 6 shows hPD-L1-Fc-GPI artificial synapse purification via a multimodal resin marketed for exosome purification. Large MW artificial synapses elute in the first fraction as shown by the high hPD-L1 concentration and artificial synapse quantity (2.26E9 synapses/ml) in elution 1. Clean in place (CIP) fractions show bound and eliminated proteins from the Inventors' artificial synapse elution.

(7) FIG. 7 shows hPDL1-Fc-GPI exosome purification via size exclusion chromatography using a resin marketed for exosome purification. Artificial synapses engineered from exosomes eluted from via a multimodal resin may be further purified via size exclusion chromatography using a resin marketed for exosome purification as shown here. Using a size exclusion chromatography, artificial synapses elute in fractions 7-9. Total protein (determined by Qubit™) and hPD-L1 ng/ml (determined by ELISA) of each fraction is shown in the graph. Bars show exosome number per ml (i.e., 1E10 exosomes/ml etc.). Fractions 7-9 contain >99% purified artificial synapses. Fractions 7-9 are pooled and may be concentrated using a filtration device, for example a 10K MWCO Amicon® centrifugal filter. Final purified product may be filtered through a low protein binding filter, for example a 0.2 µm or 0.45 µm PES filter.

(8) FIG. 8 shows hPD-L1 Expression on exosomes, quantity and hPD-L1 concentration was determined in size exclusion chromatography fractions 7-9. Knowing the molecular weight of engineered hPD-L1, the Inventors can determine the number of hPD-L1 molecules per exosome to be approximately between 12 and 40 hPD-L1/exosome. This value is consistent between different purification runs and constructs.

(9) FIG. 9 shows the purification of hPD-L2-Fc-GPI artificial synapses engineered from exosomes via multimodal resin marketed for exosome purification. This graph shows Abs 280 of fractions and quantity of hPDL2 in indicated fractions. Exosomes eluted in Elution 1. Clean in place (CIP) fractions show bound and eliminated proteins from the Inventors' artificial synapse elution.



(10) FIG. 10 shows purification of hPD-L2-Fc-GPI labeled exosomes via size exclusion column as shown here using size exclusion resin marketed for exosome purification. Fractions containing large molecular weight exosomes (Fractions 7-9) showed high hPD-L2 concentration indicating that the purified exosomes contain hPD-L2-Fc-GPI. Total protein (determined by Qubit™) and hPD-L1 ng/ml (determined by ELISA) of each fraction is shown in the graph. Lower molecular weight unbound hPD-L2-Fc-GPI eluted at later fractions.

(11) FIG. 11 shows hCTLA4-Fc-GPI exosome purification via size exclusion column as shown here using size exclusion resin marketed for exosome purification. Using size exclusion chromatography, exosomes elute in fractions 7-9. Total protein (determined by Qubit™) and hCTLA4 ng/ml (determined by ELISA) of each fraction is shown in the graph. Fractions 7-9 are pooled and contain >99% purified exosomes. Pooled exosome fractions may then be concentrated using a filtration device, for example a 10K MWCO Amicon® centrifugal filter. Final purified product may be filtered through a low protein binding filter, for example a 0.2 µm or 0.45 µm PES filter. Knowing the molecular weight of engineered hCTLA-4, the Inventors can determine the number of hCTLA-4 molecules per exosome to be approximately 233 hCTLA-4/exosome.

(12) FIG. 12A shows PD-1 Signaling Bioassay Method. The Inventors established a method to validate that PD-L1 and PD-L2 artificial synapses engineered from exosomes can bind to cells expressing PD-1 ligand. To perform this validation method, the Inventors modified the PathHunter® PD-1 Signaling Bioassay from DISCOVERX® Briefly, the PathHunter® PD-1 Signaling Bioassay relies on the well-established PathHunter® Enzyme Fragment Complementation (EFC) technology to interrogate receptor activity. EFC consists of a split-galactosidase (β-gal) enzyme: the Enzyme Donor (ED) and Enzyme Acceptor (EA) fragments which independently have no β-gal activity. However, when forced to complement they form an active β-gal enzyme that will hydrolyze substrate to produce a chemiluminescent signal. The PathHunter® PD-1 Signaling Bioassay consists of human cells engineered to stably express an ED-tagged PD-1 receptor, while EA is fused to the phosphotyrosine-binding SH2 domain of the intracellular signaling protein, SHP1. Ligand or antibody-induced activation of the receptor results in phosphorylation of the receptor's cytosolic tail. Ligand engagement, through addition of ligand-presenting artificial synapses engineered from exosomes, results in phosphorylation of PD-1, leading to the recruitment of SHP1-EA. This forces complementation of the EFC components to create an active β-gal enzyme. This active enzyme hydrolyzes substrate to create chemiluminescence as a measure of receptor activity. Addition of an antagonist (e.g., antibody to PD-L1) blocks PD-1 signaling, and will prevent complementation, resulting in a loss of signal.

FIG. 12B shows that the Inventors obtained approximately 10,000× higher increase in Relative Light Units (RLU) in Jurkat signaling cells treated with PD-L1 or PD-L2 labeled artificial synapses when compared to soluble PD-L1-Fc or PD-L2-Fc ligand, respectively. Meaning, it took 10,000× less µg/ml of PD-L1 or PD-L2 on artificial synapses than solubilized PD-L1-Fc or PD-L2 ligand to achieve the same RLU signaling. Shown is a dose-response curve for the PD-L1 and PD-L2 artificial synapses engineered from exosomes vs soluble PD-L1 and PD-L2 signaling bioassay.

(13) FIG. 13A-13C shows experimental EAU outline Test Agent A—unmodified exosomes, Test Agent B—mPDL1-Fc-GPI artificial synapses engineered from exosomes 40 µg/ml, Test Agent C—mPDL1-Fc-GPI artificial synapses engineered from exosomes 400 µg/ml, IRBP-interphotoreceptor retinoid-binding protein (IRBP) peptide, BID-Bis in die (2× daily) p.o.—Per os (orally) (FIG. 13B) EAU symptoms appear at day 6. 1st intravitreal injection and 2nd intravenous injections are performed on Day 6. There is a statistically significant initial reduction in EAU in mouse PD-L1 (mPD-L1) artificial synapses engineered from exosomes treated rats via either the intravitreal and intravenous delivery modes. 2nd intravitreal and 3rd intravenous injections are performed on Day 12. There appears to be a more rapid rate of resolution in the 1× intravitreal and intravenous groups. (FIG. 13C) Weight of rats was monitored throughout the study. 3rd intravitreal and 4th intravenous injections are performed on Day 16. There does not appear to be any significant change

in EAU in any of the test groups. The aforementioned results provide proof of principle of successfully treating an autoimmune condition (i.e., EAU) with human cell derived artificial synapses with PD-L1.

(14) FIG. **14** shows 2 types of ligands displayed on the exosome surface (Type I and Type II membrane proteins). Type I membrane proteins wherein the N-Terminus is on the luminal (interior) side of the exosome membrane and the C-Terminus is on the exterior of the exosome.

(15) Type II membrane proteins wherein the N-Terminus is on the exterior while the C-Terminus is on the interior.

(16) FIG. **15** shows a schematic representation of several embodiments of Type I membrane protein constructs, which include but are not limited to: PD-L1, PD-L2, FGL1, OX40L. FIG. **15** discloses “GSSG link” as SEQ ID NO: 319.

(17) FIG. **16** shows a schematic representation of several embodiments of the surface of an extracellular vesicle engineered with a Type I membrane protein of interest (POI) with a variable membrane anchor. Vesicle targeting sequences such as select sequences from 4F2 (CD98), ADAM10, CD298, TFR2, transmembrane portions of CD9, MARCKS, KRAS, and GPI from CD55. Proteins engineered to include a targeting sequence domain may include one or more linkers between the sticky binder and signaling domain (e.g., an Fc linker or a bond sequence wherein the bond sequence may be dimerization or multimerization sequence).

(18) FIG. **17** shows a schematic representation of the surface of an exosome engineered with an extracellular portion of the Type II membrane protein of interest (POI) with transmembrane/exosome targeting domains.

(19) FIG. **18** shows a schematic representation of an exosome engineered with an extracellular portion of the Type II membrane protein 4-1BB.

(20) FIG. **19** demonstrates a construct design for labeling an exosome surface with Type II membrane proteins.

(21) FIG. **20** shows a schematic representation of a construct design for labeling an exosome surface with multiple POI domains operably linked by a cleavable (e.g., P2A) linker.

(22) FIG. **21** shows a flow chart of purification and analytical processes provided herein.

(23) FIG. **22** shows a PD-L1 labeled exosome constructs.

(24) FIG. **23** shows several embodiments of the surface of an exosome engineered with PD-L1. The PD-L1 can be the membrane-bound PD-L1 isotype or secreted PD-L1 (SecPD-L1).

(25) FIG. **24** demonstrates size exclusion chromatography for purifying human PD-L1-GPI (no Fc) exosomes. Left panel: Protein, RNA and DNA measurements in SEC fractions are shown.

Invitrogen Qubit™ fluorometric assays were used to measure biomolecules from unmodified concentrated cell media SEC fractions or hPD-L1-Exo-Tag concentrated cell media SEC fractions. PD-L1 was measured using an R&D systems PD-L1 ELISA kit. Right panel shows dot-blot immunoblot analysis of SEC fractions. A 96-well dot blot apparatus was used to immobilize 50 µl of each SEC fraction onto PVDF. Right bottom figures: Exosome size and concentration was measured in fraction 7 by tunable resistive pulse sensing (TRPS).

(26) FIG. **25** demonstrates that GPI anchors hPD-L1 on exosomes.

(27) FIG. **26** demonstrates that a multimodal resin marketed for exosome purification purifies and disaggregates exosomes.

(28) FIG. **27** shows the exosome decoration with hPD-L1-Fc-GPI.

(29) FIG. **28A** shows the exosome decoration with hPD-L1-Fc-GPI. Fraction 7 contained the purified hPD-L1-Fc-GPI vesicles. FIG. **28B** shows size exclusion chromatography (SEC) purification results of various embodiments of human PD-L1 displayed on the surface of extracellular vesicles.

(30) FIG. **29** shows that mouse PD-L1-Fc-GPI exosomes have higher valency than mPD-L1-GPI.

(31) FIG. **30A-30C** demonstrates comparison proteomics of transprotein expression and shows that surface labeling on the engineered extracellular vesicles provided herein do not affect the relative

expression of native and associated exosome proteins. FIG. 30A shows hPD-L1-Fc-GPI. FIG. 30B shows hPD-L2-FcGPI. FIG. 30C shows hCTLA4-Fc-GPI.

(32) FIG. 31 shows production of mPD-L1-Fc-GPI in STR Bioreactor.

(33) FIG. 32 shows purification of mPD-L1-Fc-GPI (STR) via SEC. Graph shows mPD-L1 ng/ml vs Total Protein µg/ml.

(34) FIG. 33 shows purification mPDL1-Fc-GPI (STR bioreactor).

(35) FIG. 34 shows a schematic representation of the 4-1BBL labeled exosomes. Top: Vector map showing the N-terminal cystolic domain, a transmembrane (TM) domain, and the POI domain at the C-terminus. Bottom: An embodiment of an engineered EV with a type-II membrane display of the fusion protein.

(36) FIG. 35 shows embodiments of a 4-1BBL display exosome.

(37) FIG. 36A-36B show the protein engineering and purification of 4F2-4-1BBL labeled exosomes. FIG. 36B confirms that h4-1BBL is displayed on the engineered exosomes.

(38) FIG. 37 shows internal fusion protein loading of exosomes.

(39) FIG. 38 shows internal loading of exosomes with mScarlet (RFP).

(40) FIG. 39A shows internal loading of exosomes with NanoLuc® luciferase. FIG. 39B shows tetraspanin characterization of exosomes internally loaded with NanoLuc® luciferase.

(41) FIG. 40A shows the mechanism of PD-L1 engineered extracellular vesicles induce membrane clustering and receptor agonism on a target cell. An exemplary model of proposed mechanism of extracellular vesicles with a Type I membrane protein signaling domain (PD-L1) promoting receptor clustering on a target cell, wherein receptor clustering promotes increased potency of signal transduction of the target receptor. Antagonist antibodies function well at blocking receptors. Antibodies are poor agonist modalities due to their general inability to cluster receptors. Ligands on a membrane surface are potent agonists, however the cost and cold chain logistics of cell therapies makes commercialization difficult and expensive. Extracellular vesicles engineered with Type I membrane protein are able to induce receptor clustering of target receptors and initiate and propagate a potent signal response on a target cell.

(42) FIG. 40B shows the mechanism of 4-1BBL engineered extracellular vesicles induce membrane clustering and receptor agonism on a target cell. An exemplary model of proposed mechanism of extracellular vesicles with a Type II membrane protein signaling domain (4-1BBL) promoting receptor clustering on a target cell, wherein receptor clustering promotes increased potency of signal transduction of the target receptor. Soluble ligands are often poor agonist modalities due to their general inability to cluster receptors. Ligands displayed on a membrane surface are potent agonists, however the cost and cold chain logistics of cell therapies makes commercialization difficult and expensive. Extracellular vesicles engineered with Type II membrane protein are able to induce receptor clustering of target receptors and initiate and propagate a potent signal response on a target cell.

#### DETAILED DESCRIPTION

(43) The compositions and methods provided herein are based, in part, on the discovery that engineered extracellular vesicles (e.g., exosomes) expressing an engineered fusion protein (e.g., PD-L1) reduces inflammation in an animal model of experimental autoimmune uveoretinitis (EAU), an autoimmune disorder. The compositions and methods provided herein are further based, in part, on the discovery that engineered extracellular vesicles produce enhanced signaling compared to an equal quantity of recombinant ligand. Since some cellular receptors, (e.g., PD-1) require clustering or super-clustering to promote a signaling response, it stands to reason that extracellular vesicles engineered to express ligands on their surface wherein the ligands may engage target receptors on target cells and promote clustering of said target receptors thereby promoting a signal response on said target cell.

(44) In one aspect, provided herein is an engineered extracellular vesicle comprising: at least one fusion polypeptide comprising: at least one protein of interest (POI) domain; and at least one

vesicle targeting domain. In some embodiments of any of the aspects, the engineered extracellular vesicle is an exosome. In some embodiments, of any of the aspects, the fusion protein further comprises at least one linker. In some embodiments of any of the aspects, the POI domain can substantially bind to a target polypeptide. In some embodiments of any of the aspects provided herein, the engineered extracellular vesicle is an artificial synapse.

(45) Generally, the extracellular vesicles (e.g., exosomes) provided herein are produced by contacting a population of cells with a nucleic acid construct encoding the fusion proteins provided herein and isolating a plurality of extracellular vesicles. The extracellular vesicles can then be purified by methods provided herein and are formulated for therapeutic use, including but not limited to, for the treatment of autoimmune diseases, cancer, or modulating inflammation in a subject.

(46) The compositions and methods provided herein are specifically designed to exploit the membrane trafficking mechanisms of extracellular vesicles and rely on the hallmark biophysical and biochemical properties of extracellular vesicles, such as exosomes. The vesicles/artificial synapses provided herein are specifically engineered to induce/agonize and propagate biological signaling via a target polypeptide (e.g., by activating a receptor or enzyme or agonizing said receptor or enzyme). Alternatively, the engineered extracellular vesicles provided herein can act as cellular decoys or to reduce or antagonize biological signaling, e.g., by blocking an endogenous ligand from binding to a target cellular receptor and preventing activation of the receptor.

(47) Engineering of the extracellular vesicles provided herein extends these capabilities significantly by incorporating sticky binders attaching to extracellular vesicles such as exosomes, further coupled with signaling domains of choice. For example, attachment of sticky binders to exosomes, along with their linked signaling domains, allows for receptor clustering for biological signal induction/agonism and propagation not otherwise possible. In this aspect, the aforementioned design achieves the aim of an engineered extracellular vesicle by inducing the desired biological signaling in a target recipient cell.

(48) Various aspects and embodiments of the compositions and methods are provided herein in detail below.

#### Engineered Extracellular Vesicle (EV) Compositions

(49) The compositions provided herein comprises at least one extracellular vesicle (also termed artificial synapse or abbrev: EV), wherein the extracellular vesicle comprises at least one fusion polypeptide or a plurality of fusion polypeptides comprising: at least one vesicle targeting domain (e.g., sticky binders); and at least one protein of interest domain or a fragment thereof (also termed signaling domains).

(50) Extracellular vesicles (EVs) are lipid particles that are released from various cell types that function to transfer “cargo” such as nucleic acids and proteins to other cells. EVs are not able to replicate but serve as cell messengers. EV-mediated signals can be transmitted by all the different biomolecule categories—protein, lipids, nucleic acids and sugars—and the unique package of this information provides both protection and the option of simultaneous delivery of multiple different messengers even to sites remote to the vesicular origin. See, e.g., Yáñez-Mó M, Siljander P R, Andreu Z, et al. Biological properties of extracellular vesicles and their physiological functions. *J Extracell Vesicles*. 2015; 4:27066. Published 2015 May 14. doi: 10.3402/jev.v4.27066, which is incorporated herein by reference in its entirety. Furthermore, there is an increasing amount of evidence that shows that EVs can modulate a milieu of cellular signaling processes. See, e.g., Yadi et al. *Science Translation Medicine* (2020); Cerqueira de Abreu et al. *Nature Reviews Cardiology* (2020); Zhang W. et al. *Protein J.* (2019); Zha Q B et al. *Tumor Biology*. February 2017; Tan et al. (2016) Recent advances of exosomes in immune modulation and autoimmune diseases, *Autoimmunity*, 49:6, 357-365; Kalluri R, LeBleu V S. et al. The biology, function, and biomedical applications of exosomes. *Science*. 2020 Feb. 7; 367(6478); which is incorporated herein by reference in its entirety.

(51) There are various types of extracellular vesicles that are named for their site of origin in a cell, size, and structural and/or functional properties. In some embodiments of any of the aspects provided herein, the extracellular vesicle is an exosome, ectosome, macrovesicle, microparticle, apoptotic body, vesicular organelle, oncosome, exosphere, exomeres, or cell derived nanovesicle (CDN) ((e.g., by genesis via grating or shearing cells), liposomes or the like known by one of ordinary skill in the art. In various embodiments, the extracellular vesicle comprises a phospholipid bilayer with an exterior phospholipid layer and an interior phospholipid layer, wherein the exterior phospholipid layer has an external surface and an internal surface, wherein the interior phospholipid layer has an internal surface and an external surface, and the internal surface of the exterior phospholipid layer faces the internal surface of the interior phospholipid layer, and the phospholipid bilayer encloses an internal space, wherein the external surface of the interior phospholipid layer faces the internal space and wherein the external surface of the exterior phospholipid layer faces an extracellular environment, and the external surface of the inner phospholipid layer is the internal surface of the extracellular vesicle.

(52) In various embodiments, the extracellular vesicles range in size from 30 nanometers (nm) to 300 nm. In various embodiments, the plurality of EVs range in size from about 30 nm to about 150 nm. In various embodiments, the plurality of EVs or artificial synapses includes one or more artificial synapses that are about 10 nm to about 250 nm in diameter, including those about 10 nm to about 15 nm, about 15 nm to about 20 nm, about 20 nm to about 25 nm, about 25 nm to about 30 nm, about 30 nm to about 35 nm, about 35 nm to about 40 nm, about 40 nm to about 50 nm, about 50 nm to about 60 nm, about 60 nm to about 70 nm, about 70 nm to about 80 nm, about 80 nm to about 90 nm, about 90 nm to about 95 nm, about 95 nm to about 100 nm, about 100 nm to about 105 nm, about 105 nm to about 110 nm, about 110 nm to about 115 nm, about 115 nm to about 120 nm, about 120 nm to about 125 nm, about 125 nm to about 130 nm, about 130 nm to about 135 nm, about 135 nm to about 140 nm, about 140 nm to about 145 nm, about 145 nm to about 150 nm, about 150 to about 200 nm, about 200 nm to about 250 nm, about 250 nm or more.

(53) In some embodiments of any of the aspects provided herein, the EV is an exosome. Exosomes are membrane-bound EVs that are produced in the endosomal compartment of most eukaryotic cells. As used herein, the term “exosome” refers to a species of extracellular vesicle between about 20 nm to about 400  $\mu$ m in diameter, e.g., about 30 nm-200 nm in diameter by inward invagination of a portion of a membrane of an endosome (for example an early or late endosome), wherein the endosome is within a cell comprising a plasma membrane, and the exosome is released from the cell upon fusion of another portion of the endosome membrane with the plasma membrane. An exosome may refer to a species of extracellular vesicle between 20 nm-400  $\mu$ m in diameter, more preferably 30 nm-200 nm in diameter, that originates by budding of a portion of a plasma membrane from a cell wherein the budded portion of the plasma membrane is released to the extracellular environment.

(54) The EVs (e.g., exosomes or cell derived vesicles) provided herein may comprise cargo, for example, peptides, proteins, nucleic acids, lipids, metabolites, carbohydrates, biomolecules, small molecules, large molecules, vesicles, organelles, or fragments thereof. Exosome cargo may be located within the internal space of the exosome. EV cargo may be membrane bound spanning one or both layers of the exosome phospholipid bilayer (for example a transmembrane protein). EV cargo may be in contact with the exterior or interior surface of the exosome, for example through a covalent bond or a non-covalent bond. The phospholipid bilayer of the EV or exosome provided herein may comprise one or more transmembrane proteins, wherein a portion of the one or more transmembrane membrane proteins is located within the internal space of the exosome. The phospholipid bilayer of the EV or exosome provided herein may comprise one or more transmembrane proteins, wherein a portion of the one or more transmembrane membrane proteins traverses the EV phospholipid bilayer. The phospholipid bilayer of the EV may comprise one or more transmembrane proteins, wherein the one or more transmembrane membrane proteins

comprises a domain on the exterior of the exosome.

(55) In some embodiments of any of the aspects, the extracellular vesicles or exosomes provided herein endogenously express CD81+, CD82+, CD37+, CD63+, CD9+, CD151+, CD105+, or any combination thereof. In various embodiments, the plurality of artificial synapses includes one or more artificial synapses expressing a biomarker. In certain embodiments, the biomarkers are tetraspanins. In other embodiments, the tetraspanins are one or more selected from the group including CD63, CD81, CD82, CD53, CD151, and CD37. In other embodiments, the artificial synapses express one or more lipid raft associated proteins (e.g., glycosylphosphatidylinositol-anchored proteins and flotillin), cholesterol, sphingolipids such as sphingomyelin, and/or hexosylceramides.

(56) In other embodiments, the biological protein is related to exosome formation and packaging of cytosolic proteins, e.g., Hsp70, Hsp90, 14-3-3 epsilon, PKM2, GW182 and AGO2. In certain embodiments, the artificial synapses express CD63, HSP70, CD105 or combinations thereof. In other embodiments, the artificial synapses do not express CD9 or CD81, or express neither. For example, plurality of artificial synapses can include one or more artificial synapses that are CD63+, HSP+, CD105+, CD9-, and CD81-.

(57) The EVs provided herein are specifically engineered to express fusion polypeptides that elicit biological signaling via a target cell. In some embodiments, the fusion polypeptide is overexpressed to elicit a biological response on a target cell or target polypeptide. The engineered EV comprises at least one fusion polypeptide and can comprise a plurality of the same or different fusion polypeptides provided herein. The fusion polypeptides provided herein comprise a protein of interest domain, also termed the signaling domain.

(58) The fusion polypeptides provided herein can comprise one or more of a protein of interest domain, such that expression of said fusion polypeptide is permitted and that the number of POI domains does not impede protein expression or folding. Furthermore, the EVs provided herein can express more than one fusion protein (e.g., encoded by multiple different nucleic acid constructs). One of skill in the art can appreciate that an engineered EV can include one or more combinations of different signaling domains and/or vesicle targeting domains, or that one can use a plurality of engineered EVs, each including one or more vesicle targeting domains and one or more signaling domains.

(59) In some embodiments, the EVs provided herein comprise one or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, or ten or more fusion proteins. The fusion proteins can be encoded by the same vector or separate vectors. In some embodiments of any of the aspects, the engineered extracellular vesicle comprises at least two POI domains and/or at least two vesicle targeting domains.

(60) In some embodiments, the fusion polypeptide comprises one or more, two or more, three or more, four or more, five or more, or six or more POI domains on the same polypeptide or nucleic acid construct encoding said polypeptide. For example, the fusion polypeptides provided herein can express a fusion polypeptide encoding one or more, two or more, three or more, four or more, five or more, or six or more signaling domains. In another example, the fusion polypeptides provided herein can express a fusion polypeptide encoding an immune checkpoint protein or a protein involved in immune or cell synapse or any combination or fragment thereof.

(61) In some embodiments, the EV comprises one or more, two or more, three or more, four or more, five or more, or six or more fusion polypeptides on the same EV. For example, EVs comprising one or more, two or more, three or more, four or more, five or more, or six or more fusion polypeptides wherein the fusion polypeptides encode a signaling domain. In another example, EVs comprising one or more, two or more, three or more, four or more, five or more, or six or more fusion polypeptides wherein the fusion polypeptides encode for one or more immune checkpoint proteins or proteins involved in immune or cell synapse, or any combination or fragment thereof.

(62) In various embodiments, the signaling domain is a protein or peptide of interest, or a fragment thereof. In various embodiments, the protein of interest (signaling domain) is an immune checkpoint protein. The terms “immune checkpoint protein” or “protein involved in immune or cell synapse” can include but are not limited to adenosine A2A receptor (A2AR), Galectin 9, fibrinogen-like protein 1 (FGL-1), platelet endothelial adhesion factor-1 (PECAM-1), tumor necrosis factor gene 6 protein (TSG-6), Stabilin-1 (STAB-1) also known as Clever-1, Neuropilin 1 (NRP1), Neuropilin 2 (NRP2), semaphorin-3A (SEMA3A), semaphorin-3F (SEMA3F), repulsive guidance molecule B (RGMB) also known as DRG11, T-cell immunoglobulin and mucin domain 3 (TIM-3), T cell immunoreceptor with Ig and ITIM domains (TIGIT), human leukocyte antigen (HLA) class I, HLA class II, high mobility group protein B1 (HMGB1), phosphatidylserine, carcinoembryonic antigen-related cell adhesion molecule 1 (CEACAM-1), T-cell receptor (TCR), Src homology 2 domain-containing protein tyrosine phosphatase 1 (SHP-1), SHP-2, F-Box protein 38 (FBXO38), signaling lymphocytic activation molecule (SLAM)-associated protein (SAP) also known as SH2D1A, B7RP1, indoleamine 2,3-dioxygenase (IDO), NADH oxidase 2 (NOX2), tumor necrosis factor receptor (TNFR) superfamily member 18 (TNFRSF18) (also known as activation inducible TNFR family receptor (AITR), glucocorticoid-induced TNFR related (GITR) protein, and CD357), B7-H4 also known as V-set domain containing T-cell activator inhibitor (VTCN1), B7-H5 (also known as V-domain Ig suppressor of T-cell activation (VISTA), platelet receptor Gi24, and stress induced secreted protein 1 (SISP1), B7-H6 (also known as NCR3LG1), B7-H7 (also known as human endogenous retrovirus-H (HERV-H) long terminal repeat-associating protein 2 (HHLA2), apelin receptor (APLNR), interferon gamma (IFN  $\gamma$ ) receptor, programmed cell death-1 (PD-1), Protein Wnt-5a (WNT5A), serine/threonine-protein kinase PAK4, interleukin 6 (IL-6), interleukin-10 (IL-10), NKG2 family of C-type lectin receptors (for example NKG2A, B, C, D, E, F and H), ligands of NKG2 family, killer cell immunoglobulin-like receptors, CD-2, cluster of differentiation 4 (CD4), CD8, CD27, CD27 ligand (CD27L, also known as CD70), CD28, CD28H (also known as transmembrane and immunoglobulin domain containing 2 (TMIGD2) and Ig containing and proline-rich receptor-1 (IGPR1)), CD39, CD40, CD44, integrin associated protein (CD47), carcinoembryonic antigen related cell adhesion molecule 1 (CEACAM1 also known as CD66a), CD73, B7-1 (also known as CD80), B7-2 (also known as CD86), CD94, CD96, immunoglobulin superfamily member 2 (IGSF2) also known as CD101, nectin cell adhesion molecule 2 (NECTIN2) (also known as herpesvirus entry mediator B (HVEB), poliovirus receptor related 2 (PRR2, PVRL2 and PVRR2) and CD112), poliovirus receptor related immunoglobulin domain containing protein (PVIRG) also known as CD112R, CD122 (also known as IL5RB and P70-75), OX40 (also known as tumor necrosis factor receptor superfamily member 4 (TNFRSF4) and CD134), OX40 ligand (OX40L), 4-1BB (also known as CD137), CD134 (also known as 4-1BB ligand (4-1BBL) and as tumor necrosis factor ligand superfamily member 9 (TNFSF9) and CD137L), cytotoxic T-lymphocyte-associated protein 4 (CTLA-4) also known as CD152, CD154 (also known as CD40L), poliovirus receptor (PVR) also known as CD155, killer-cell immunoglobulin-like receptors (KIRs) (for example but not limited to CD158 family, CD158a, CD158g, CD158h, KIR2DL1, KIR2DS1, KIRDS3, and KIR2DS5), CD160, signal-regulatory protein alpha (SIRP $\alpha$ ) also known as CD172a, OX-2 also known as CD200, CD200R, lymphocyte-activation gene 3 (LAG-3) also known as CD223, CD226, OX40L also known as CD252, herpes virus entry mediator (HVEM) also known as tumor necrosis factor receptor superfamily member 14 (TNFRSF14) and CD270, B- and T-lymphocyte attenuator (BTLA) also known as CD272, programmed cell death ligand-2 (PD-L2) (also known as B7-DC, PDCDILG2, and CD273), programmed cell death-ligand 1 (PD-L1) (also known as B7-H1 and CD274), B7-H2 (also known as inducible T-cell co-stimulator ligand (ICOSLG), B7RP1, and CD275), B7-H3 also known as CD276, inducible T-cell co-stimulator (ICOS) also known as CD278, programmed cell death protein 1 (PD-1) also known as CD279, leukocyte-associated Ig-like receptor-1 (LAIR-1) also known as CD305, collagen family of proteins (for example but not limited to collagen I, collagen II, collagen

III alpha 1, collagen IV, collagen XXIII alpha 1, collagen XXV alpha 1), sialic acid-binding immunoglobulin-type lectin 7 (SIGLEC7) also known as CD328, sialic acid-binding immunoglobulin-type lectin 7 (SIGLEC9) also known as CD329, natural cytotoxicity triggering receptor 3 (NKp30) also known as CD337, or any isoform, fragment, variation thereof, or a ligand to the aforementioned proteins thereof, or the like known by one of ordinary skill in the art. All variants are encompassed by the present invention.

(63) In some embodiments of any of the aspects provided herein, the protein of interest domain (POI domain) comprises a polypeptide or a fragment thereof or a nucleic acid encoding said polypeptide or fragment thereof selected from the group consisting of: Table 1 (below). Non-limiting examples of nucleic acid sequences that encode the POI domains provided herein are also provided in

(64) TABLE-US-00001 TABLE 1 Type I Proteins of Interest Amino Acid Sequence  
Protein of Transcript Sequence (SEQ ID NO:) Interest Amino Acid Sequence  
(SEQ ID NO:) Human >NM\_014143.4 *Homo sapiens* CD274 molecule (CD274),  
Programmed transcript variant 1, mRNA death-ligand 1

AGTTCTGCGCAGCTTCCCGAGGCTCCGCACCAGCCGCGCTTCTGTCCGCCTGCAGGG  
(PD-L1)  
CATTCCAGAAAGATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTG  
CTGAACGCATTTACTGTACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGC  
AATATGACAATTGAATGCAAATTCAGTAGAAAAACAATTAGACCTGGCTGCACTA  
ATTGTCTATTGGGAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAA  
GACCTGAAGGTTTCAGCATAGTAGCTACAGACAGAGGGCCCGGCTGTTGAAGGACCAG  
CTCTCCCTGGGAAATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGG  
GTGTACCGCTGCATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAA  
GTCAATGCCCCATACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACC  
TCTGAACATGAACTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGG  
ACAAGCAGTGACCATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGA  
GAGGAGAAGCTTTTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAG  
ATTTTCTACTGCACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTG  
GTCATCCCAGAACTACCTCTGGCACATCCTCCAAATGAAAGGACTCACTTGGTAATT  
CTGGGAGCCATCTTATTATGCCTTGGTGTAGCACTGACATTCATCTTCCGTTTAAGA  
AAAGGGAGAATGATGGATGTGAAAAAATGTGGCATCCAAGATACAAACTCAAAGAAG  
CAAAGTGATACACATTTGGAGGAGACGTAATCCAGCATTGGAACCTTCTGATCTTCAA  
GCAGGGATTCTCAACCTGTGGTTTAGGGGTTTCATCGGGGCTGAGCGTGACAAGAGGA  
AGGAATGGGCCCCGTGGGATGCAGGCAATGTGGGACTTAAAAGGCCCAAGCACTGAAA  
ATGGAACCTGGCGAAAGCAGAGGAGGAGAATGAAGAAAGATGGAGTCAAACAGGGAG  
CCTGGAGGGAGACCTTGATACTTTCAAATGCCTGAGGGGCTCATCGACGCCTGTGAC  
AGGGAGAAAGGATACTTCTGAACAAGGAGCCTCCAAGCAAATCATCCATTGCTCATC  
CTAGGAAGACGGGTTGAGAATCCCTAATTTGAGGGTCAGTTCCTGCAGAAGTGCCCT  
TTGCCTCCACTCAATGCCTCAATTTGTTTTCTGCATGACTGAGAGTCTCAGTGTTGG  
AACGGGACAGTATTTATGTATGAGTTTTTCCTATTTATTTTGAAGTCTGTGAGGTCTT  
CTTGTGTCATGTGAGTGTGGTTGTGAATGATTTCTTTTGAAGATATATTGTAGTAGATG  
TTACAATTTTGTGCCCCAACTAACTTGCTGCTTAATGATTTGCTCACATCTAGTAA  
AACATGGAGTATTTGTAAAGGTGCTTGGTCTCCTCTATAACTACAAGTATACATTGGA  
AGCATAAAGATCAAACCGTTGGTTGCATAGGATGTCACCTTTATTTAACCCTTAAT  
ACTCTGGTTGACCTAATCTTATTCTCAGACCTCAAGTGTCTGTGCAGTATCTGTTCC  
ATTTAAATATCAGCTTTACAATTATGTGGTAGCCTACACACATAATCTCATTTTCATC  
GCTGTAACCAACCCTGTTGTGATAACCACTATTATTTTACCCATCGTACAGCTGAGGA  
AGCAAACAGATTAAGTAACTTGCCCAAACCAGTAAATAGCAGACCTCAGACTGCCAC  
CCACTGTCCTTTTATAATACAATTTACAGCTATATTTTACTTTAAGCAATTCTTTTA



TTCAAACCAATTTATTAAAGTGGCCCTTGCAATATCAATCGCTGTGCCAGGCATTGAA  
TCTACAGATGTGAGCAAGACAAAGTACCTGTCCTCAAGGAGCTCATAGTATAATGAG  
GAGATTAACAAGAAAATGTATTATTACAATTTAGTCCAGTGTCATAGCATAAGGATG  
ATGCGAGGGGAAAACCCGAGCAGTGTTGCCAAGAGGAGGAAATAGGCCAATGTGGTC  
TGGGACGGTTGGATATACTTAAACATCTTAATAATCAGAGTAATTTTCATTTACAAA  
GAGAGGTCGGTACTTAAAATAACCCCTGAAAAATAACACTGGAATTCCTTTTCTAGCA  
TTATATTTATTCCTGATTTGCCTTTGCCATATAATCTAATGCTTGTTTATATAGTGT  
CTGGTATTGTTTAAACAGTTCTGTCTTTTCTATTTAAATGCCACTAAATTTTAAATTC  
ATACCTTTCCATGATTCAAAATTCAAAAGATCCCATGGGAGATGGTTGGAAAATCTC  
CACTTCATCCTCCAAGCCATTCAAGTTTCTTTTCCAGAAGCAACTGCTACTGCCTTT  
CATTCATATGTTCTTCTAAAGATAGTCTACATTTGGAAATGTATGTTAAAAGCACGT  
ATTTTTTAAAATTTTTTTTCTAAATAGTAACACATTGTATGTCTGCTGTGTACTTTGC  
TATTTTTATTTATTTTAGTGTTTCTTATATAGCAGATGGAATGAATTTGAAGTTCCC  
AGGGCTGAGGATCCATGCCTTCTTTGTTTCTAAGTTATCTTTCCCATAGCTTTTCAT  
TATCTTTCATATGATCCAGTATATGTTAAATATGTCCTACATATACATTTAGACAAC  
CACCATTTGTTAAGTATTTGCTCTAGGACAGAGTTTGGATTTGTTTATGTTTGCTCA  
AAAGGAGACCCATGGGCTCTCCAGGGTGCCTGAGTCAATCTAGTCCTAAAAAGCAA  
TCTTATTATTAACCTCTGTATGACAGAATCATGTCTGGAACTTTTGTTTTCTGCTTTC  
TGTC AAGTATAAACTTCACTTTGATGCTGTACTTGCAAATCACATTTTCTTTCTGG  
AAATTCCGGCAGTGTACCTTGACTGCTAGCTACCCTGTGCCAGAAAAGCCTCATTG  
TTGTGCTTGAACCCTTGAATGCCACCAGCTGTCATCACTACACAGCCCTCCTAAGAG  
GCTTCCTGGAGGTTTCGAGATTCAGATGCCCTGGGAGATCCCAGAGTTTCCTTTCCC  
TCTTGGCCATATTCTGGTGTCAATGACAAGGAGTACCTTGGCTTTGCCACATGTCAA  
GGCTGAAGAAACAGTGTCTCCAACAGAGCTCCTTGTGTTATCTGTTTGTACATGTGC  
ATTTGTACAGTAATTGGTGTGACAGTGTTCTTTGTGTGAATTACAGGCAAGAATTGT  
GGCTGAGCAAGGCACATAGTCTACTCAGTCTATTCTTAAGTCCTAACTCCTCCTTGT  
GGTGTGATTGTAAGGCACCTTATCCCTTTTGTCTCATGTTTCATCGTAAATGGC  
ATAGGCAGAGATGATACCTAATTCTGCATTTGATTGTCACTTTTTGTACCTGCATTA  
ATTTAATAAAATATTCTTATTTATTTTGTACTTGGTACACCAGCATGTCCATTTTC  
TTGTTTATTTTGTGTTTAAATAAAATGTTTCAGTTTAAACATCCCA (SEQ ID NO: 1)

>NP\_054862.1 programmed cell death 1 ligand 1 isoform a precursor [*Homo sapiens*]

MRIFAVFIFMTYWHL LNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVYW  
EMEDKNIIQFVHGEEDLKVQHSSYRQRRARLLKDQLSLGNAALQITDVKLQDAGVYRC  
MISYGGADYKRITVKVNAPYNKINQRILVVDPTSEHELTCQAEGYPKAEVIWTSSD  
HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPE  
LPLAHPPNERTHLVILGAILLCLGVALTFIFRLRKGRMMDVKKCGIQDTNSKKQSDT  
HLEET (SEQ ID NO: 2) Mouse PD-L1 >NM\_021893.3 *Mus musculus* CD274  
antigen (Cd274), mRNA

GAAATCGTGGTCCCCAAGCCTCATGCCAGGCTGCACTTGACGTCGCGGGGCCAGTCT  
CCTCGCCTGCAGATAGTTCCCAAAACATGAGGATATTTGCTGGCATTATATTCACAG  
CCTGCTGTCACTTGCTACGGGCGTTTACTATCACGGCTCCAAAGGACTTGTACGTGG  
TGGAGTATGGCAGCAACGTCACGATGGAGTGCAGATTCCCTGTAGAACGGGAGCTGG  
ACCTGCTTGCGTTAGTGGTGTACTGGGAAAAGGAAGATGAGCAAGTGATTCAAGTTG  
TGGCAGGAGAGGAGGACCTTAAGCCTCAGCACAGCAACTTCAGGGGGAGAGCCTCGC  
TGCCAAAGGACCAGCTTTTGAAGGGAAATGCTGCCCTTCAGATCACAGACGTCAAGC  
TGCAGGACGCAGGCGTTTACTGCTGCATAATCAGCTACGGTGGTGCGGACTACAAGC  
GAATCACGCTGAAAGTCAATGCCCCATACCGCAAAATCAACCAGAGAATTTCCGTGG  
ATCCAGCCACTTCTGAGCATGAACTAATATGTCAGGCCGAGGGTTATCCAGAAGCTG  
AGGTAATCTGGACAAACAGTGACCACCAACCCGTGAGTGGGAAGAGAAGTGTCACCA

CTTCCCGGACAGAGAGGGGATGCTTCTTCTCAATGTGACACAGCAGTCTGAGGGGTCAACGCCA  
CAGCGAATGATGTTTTCTACTGTACGTTTTTGAGATCACAGCCAGGGCAAACCACA  
CAGCGGAGCTGATCATCCCAGAACTGCCTGCAACACATCCTCCACAGAACAGGACTC  
ACTGGGTGCTTCTGGGATCCATCCTGTTGTTCTCATTGTAGTGTCCACGGTCCTCC  
TCTTCTTGAGAAAACAAGTGAGAATGCTAGATGTGGAGAAATGTGGCGTTGAAGATA  
CAAGCTCAAAAAACCGAAATGATACACAATTCGAGGAGACGTAAGCAGTGTTGAACC  
CTCTGATCGTCGATTGGCAGCTTGTGGTCTGTGAAAGAAAGGGGCCCATGGGACATGA  
GTCCAAAGACTCAAGATGGAACCTGAGGGAGAGAACCAAGAAAGTGTTGGGAGAGGA  
GCCTGGAACAACGGACATTTTTTCCAGGGAGACACTGCTAAGCAAGTTGCCCATCAG  
TCGTCTTGGGAAATGGATTGAGGGTTCTGGCTTAGCAGCTGGTCCTTGCACAGTGA  
CCTTTTCTCTGCTCAGTGCCGGGATGAGAGATGGAGTCATGAGTGTTGAAGAATAA  
GTGCCTTCTATTTATTTTGAAGTCTGTGTGTTCTCACTTTGGGCATGTAATTATGACT  
GGTGAATTCTGACGACATGATAGATCTTAAGATGTAGTCACCAAACCTCAACTGCTGC  
TTAGCATCCTCCGTAACTACTGATACAAGCAGGGAACACAGAGGTCACCTGCTTGGT  
TTGACAGGCTCTTGCTGTCTGACTCAAATAATCTTTATTTTTTCAGTCCTCAAGGCTC  
TTCGATAGCAGTTGTTCTGTATCAGCCTTATAGGTGTCAGGTATAGCACTCAACATC  
TCATCTCATTACAATAGCAACCCTCATCACCATAGCAACAGCTAACCTCTGTTATCC  
TCACTTCATAGCCAGGAAGCTGAGCGACTAAGTCACTTGCCACACAGAGTATCAGCTC  
TCAGATTTCTGTTCTTCAGCCACTGTCCTTTCAGGATAGAATTTGTTCGTTAAGAAAT  
TAATTTAAAAACTGATTATTGAGTAGCATTGTATATCAATCACAACATGCCTTGTGC  
ACTGTGCTGGCCTCTGAGCATAAAGATGTACGCCGGAGTACCGGTCGGACATGTTTA  
TGTGTGTTAAATACTCAGAGAAATGTTCAATTAACAAGGAGCTTGCATTTTAGAGACA  
CTGGAAAGTAACTCCAGTTCATTGTCTAGCATTACATTTACCTCATTTGCTATCCTT  
GCCATACAGTCTCTTGTTCTCCATGAAGTGTCAATGAATCTTGTTGAATAGTTCTTTT  
ATTTTTTAAATGTTTCTATTTAAATGATATTGACATCTGAGGCGATAGCTCAGTTGG  
TAAAACCTTTTCTCACAAGTGTGAAACCTGAGTCTTATCCCTAGAACCCACATAA  
AAAACAGTTGCGTATGTTTGTGCATGCTTTTGATCCCAGCACTAGGGAGGCAGAGGC  
AGGCAGATCCTGAGCTCTCATTGACCACCCAGCCTAGCCTACATGGTTAGCTCCAGG  
CCTACAGGAGCTGGCAGAGCCTGAAAAACGATGCCTAGACACACACACACACACA  
CACACACACACACACACACACACACACACCATGTACTCATAGACCTAAGTGCACCCTCCTA  
CACATGCACACACATACAATTCAAACACAAATCAACAGGGAATTGTCTCAGAATGGT  
CCCCAAGACAAAGAAGAAGAAAAACACCAAACCAGCTCTATTCCCTCAGCCTATCCT  
CTCTACTCCTTCCTAGAAGCAACTACTATTGTTTTTGTATATAAATTTACCCAACGA  
CAGTTAATATGTAGAATATATATTAAGTGTCTGTCAATATATATTATCTCTTTCTT  
TCTTTCTTCCTTTCTTTCTTTCTTTCTTTCTTTCTTTCTTTCTTTCTTTCTTT  
CTTCCTTCCTTCCTTCCTTCCTTCCTTCCTTCCTTCCTTCCTTCCTTCCTTCCTTT  
TTTCTGTCTATCTGTACCTAAATGGTTGCTCACTATGCATTTTCTGTGCTCTTCGCC  
CTTTTTATTTAATGTATGGATATTTATGCTGCTTCCAGAATGGATCTAAAGCTCTTT  
GTTTCTAGGTTTTCTCCCCCATCCTTCTAGGCATCTCTCACACTGTCTAGGCCAGAC  
ACCATGTCTGCTGCCTGAATCTGTAGACACCATTATATAAAGCACGTACTACCGAGT  
TTGTATTTGGCTTGTTCTGTGTCTGATTAAAGGGAGACCATGAGTCCCCAGGGTACA  
CTGAGTTACCCCAGTACCAAGGGGGAGCCTTGTTTGTGTCTCCATGGCAGAAGCAGG  
CCTGGAGCCATTTTGGTTTCTTCCTTGACTTCTCTCAAACACAGACGCCTCACTTGC  
TCATTACAGGTTCTCCTTTGGGAATGTCAGCATTGCTCCTTGACTGCTGGCTGCCCT  
GGAAGGAGCCCATTAGCTCTGTGTGAGCCCTTGACAGCTACTGCCTCTCCTTACCAC  
AGGGGCCTCTAAGATACTGTTACCTAGAGGTCTTGAGGATCTGTGTTCTCTGGGGGG  
AGGAAAGGAGGAGGAACCCAGAACTTTCTTACAGTTTTCTTGTTCTGTACATGTC  
AAGACTGAAGGAACAGGCTGGGCTACGTAGTGAGATCCTGTCTCAAAGGAAAGACGA  
GCATAGCCGAACCCCCGGTGGAACCCCCTCTGTTACCTGTTACACAAGCTTATTGA  
TGAGTCTCATGTTAATGTCTTGTTTGTATGAAGTTTAAGAAAATATCGGGTTGGGCA

ACACATTCTATTCTTTATTTGAAATCCTTAATGCCATTCTCATGGTGTGGA  
TGGTGTGGCACTTTATTCTTTTGTGTTGTGTATAACCATAAATTTTATTTTGCATCA  
GATTGTCAATGTATTGCATTAATTTAATAAATAATTTTATTTATTAATAAAAAAAAAA  
AAAAA (SEQ ID NO: 3)>NP\_068693.1 programmed cell death 1 ligand 1  
precursor [*Mus musculus*]

MRIFAGIIFTACCHLLRAFTITAPKDLVVEYGSNVTMECRFPVERELDLLALVVYW  
EKEDEQVIQFVAGEEDLK PQHSNFRGRASLPKDQLLKGNALQITDVKLQDAGVYCC  
IISYGGADYKRITLKVNPYRKINQRISVDPATSEHELICQAEGYPEAEVIWTNSDH  
QPVSGKRSVTTSRTEGMLLNVTSSLRVNATANDVFYCTFWRSQPGQNHTAELIPEL  
PATHPPQNRTHWVLLGSILLFLIVVSTVLLFLRKQVRMLDVEKCGVEDTSSKNRNDT  
QFEET (SEQ ID NO: 4) Human PD-L2 >NM\_025239.4 *Homo sapiens*  
programmed cell death 1 ligand 2 (PCD1LG2), mRNA

ACTCTCATGTTACGGCAAACCTTAAGCTGAATGAACAACCTTTTCTTCTCTTGAATAT  
ATCTTAACGCCAAATTTTGAGTGCTTTTTTGTACCCATCCTCATATGTCCCAGCTA  
GAAAGAATCCTGGGTTGGAGCTACTGCATGTTGATTGTTTTGTTTTCTTTTGGCT  
GTTCATTTTGGTGGCTACTATAAGGAAATCTAACACAAACAGCAACTGTTTTTTGTT  
GTTTACTTTTGCATCTTTACTTGTGGAGCTGTGGCAAGTCCTCATATCAAATACAGA  
ACATGATCTTCCTCCTGCTAATGTTGAGCCTGGAATTGCAGCTTCACCAGATAGCAG  
CTTTATTCACAGTGACAGTCCCTAAGGAAGTGTACATAATAGAGCATGGCAGCAATG  
TGACCCTGGAATGCAACTTTGACACTGGAAGTCATGTGAACCTTGGAGCAATAACAG  
CCAGTTTGCAAAAGGTGGAAAATGATACATCCCCACACCGTGAAAGAGCCACTTTGC  
TGGAGGAGCAGCTGCCCCTAGGGAAGGCCTCGTTCCACATACCTCAAGTCCAAGTGA  
GGGACGAAGGACAGTACCAATGCATAATCATCTATGGGGTCGCCTGGGACTACAAGT  
ACCTGACTCTGAAAGTCAAAGCTTCCTACAGGAAAATAAACACTCACATCCTAAAGG  
TTCCAGAAACAGATGAGGTAGAGCTCACCTGCCAGGCTACAGGTTATCCTCTGGCAG  
AAGTATCCTGGCCAAACGTCAGCGTTCCTGCCAACACCAGCCACTCCAGGACCCCTG  
AAGGCCTCTACCAGGTCACCAGTGTTCTGCGCCTAAAGCCACCCCTGGCAGAACT  
TCAGCTGTGTGTTCTGGAATACTCACGTGAGGGAAGTACTTTGGCCAGCATTGACC  
TTCAAAGTCAGATGGAACCCAGGACCCATCCAAGTGGCTGCTTCACATTTTCATCC  
CCTTCTGCATCATTGCTTTTCATTTTCATAGCCACAGTGATAGCCCTAAGAAAACAAC  
TCTGTCAAAAGCTGTATTCTTCAAAGACACAACAAAAAGACCTGTCACCACAACAA  
AGAGGGAAGTGAACAGTGCTATCTGAACCTGTGGTCTTGGGAGCCAGGGTGACCTGA  
TATGACATCTAAAGAAGCTTCTGGACTCTGAACAAGAATTCGGTGGCCTGCAGAGCT  
TGCCATTTGCACTTTTCAAATGCCTTTGGATGACCCAGCACTTTAATCTGAAACCTG  
CAACAAGACTAGCCAACACCTGGCCATGAAACTTGCCCCTTCACTGATCTGGACTCA  
CCTCTGGAGCCTATGGCTTTAAGCAAGCACTACTGCACCTTACAGAATTACCCCACT  
GGATCCTGGACCCACAGAATTCCTTCAGGATCCTTCTTGCTGCCAGACTGAAAGCAA  
AAGGAATTATTTCCCCTCAAGTTTCTAAGTGATTTCCAAAAGCAGAGGTGTGTGGA  
AATTTCCAGTAACAGAAACAGATGGGTTGCCAATAGAGTTATTTTTTATCTATAGCT  
TCCTCTGGGTACTAGAAGAGGCTATTGAGACTATGAGCTCACAGACAGGGCTTCGCA  
CAAAGTCAAATCATAATTGACATGTTTTATGGATTACTGGAATCTTGATAGCATAAT  
GAAGTTGTTCTAATTAACAGAGAGCATTTAAATATACACTAAGTGCACAAATTGTGG  
AGTAAAGTCATCAAGCTCTGTTTTTGGAGTCTAAGTCACAAAGCATTGTGTTTAACC  
TGTAATGGCACCATGTTTAATGGTGGTTTTTTTTTTTGAAGTACATCTTTCCTTTAAA  
AATTATTGGTTTCTTTTTTATTTGTTTTTACCTTAGAAATCAATTATATACAGTCAAA  
AATATTTGATATGCTCATAACGTTGTATCTGCAGCAATTCAGATAAGTAGCTAAAAT  
GGCCAAAGCCCCAACTAAGCCTCCTTTTCTGGCCCTCAATATGACTTTAAATTTGA  
CTTTTCAGTGCCTCAGTTTGCACATCTGTAATACAGCAATGCTAAGTAGTCAAGGCC  
TTTGATAATTGGCACTATGGAAATCCTGCAAGATCCCCTACATATGTGTGGAGCAG  
AAGGGTAACTCGGCTACAGTAACAGCTTAATTTTGTAAATTTGTTCTTTATACTGG

AGCCATGACATGACAGCATTAGCTTACCCTTGAACCTATTCAAATGGGGCACATTAGC  
TAGTATAACAGACTTACATAGGTGGGCCTAAAGCAAGCTCCTTAACTGAGCAAAATT  
TGGGGCTTATGAGAATGAAAGGGTGTGAAATTGACTAACAGACAAATCATACATCTC  
AGTTTCTCAATTCTCATGTAAATCAGAGAATGCCTTTAAAGAATAAAACTCAATTGT  
TATTCTTCAACGTTCTTTATATATTCTACTTTTGGGTA (SEQ ID NO: 5)

>NP\_079515.2 programmed cell death 1 ligand 2 precursor [*Homo sapiens*]

MIFLLMLSLQLHQAALFTVTVPKELYIIEHGSNVTLECNFDTGSHVNLGAITA  
SLQKVENDTSPHRERATLLEEQLPLGKASFHIPQVQVRDEGQYQCIIYGVAWDYKY  
LTLKVKASYRKINTHILKVPETDEVELTCQATGYPLAEVSWPNVSVPANTSHSRTPE  
GLYQVTSVLRLKPPPGRNFSVFWNTHVRELTLASIDLQSQMEPRTHPTWLLHIFIP  
FCIIAFIFIATVIALRKQLCQKLYSSKDTTKRPVTTTKREVNSAI (SEQ ID NO: 6)

Mouse PD-L2 >NM\_021396.2 *Mus musculus* programmed cell death 1 ligand 2  
(Pcdcllg2), mRNA

GACCACATCATTTTTGTTCCTTTGTTGGATATATCCTAATGTCAAATGTGGCATAT  
CTTTGTTGTCTCCTTCTGTCTCCCAACTAGAGAGAACACACTTACGGCTCCTGTCCC  
GGGCAGGTTTGGTTGTCTGGTGTGATTGGCTTCCAGGGAACCTGATACAAGGAGCAAC  
TGTGTGCTGCCTTTTCTGTGTCTTTGCTTGAGGAGCTGTGCTGGGTGCTGATATTGA  
CACAGACCATGCTGCTCCTGCTGCCGATACTGAACCTGAGCTTACAACCTTCATCCTG  
TAGCAGCTTTATTCACCGTGACAGCCCCTAAAGAAGTGACACCGTAGACGTCGGCA  
GCAGTGTGAGCCTGGAGTGCGATTTTGACCGCAGAGAATGCACTGAACTGGAAGGGA  
TAAGAGCCAGTTTGCAGAAGGTAGAAAATGATACGTCTCTGCAAAGTGAAAGAGCCA  
CCCTGCTGGAGGAGCAGCTGCCCCCTGGGAAAGGCTTTGTTCCACATCCCTAGTGTCC  
AAGTGAGAGATTCCGGGCAGTACCGTTGCCTGGTCATCTGCGGGGCGCCTGGGACT  
ACAAGTACCTGACGGTGAAAGTCAAAGCTTCTTACATGAGGATAGACACTAGGATCC  
TGGAGGTTCCAGGTACAGGGGAGGTGCAGCTTACCTGCCAGGCTAGAGGTTATCCCC  
TAGCAGAAGTGTCCTGGCAAATGTCAGTGTTCTGCCAACACCAGCCACATCAGGA  
CCCCGAAGGCCTCTACCAGGTCACCAGTGTTCTGCGCCTCAAGCCTCAGCCTAGCA  
GAACTTCAGCTGCATGTTCTGGAATGCTCACATGAAGGAGCTGACTTCAGCCATCA  
TTGACCCTCTGAGTCGGATGGAACCCAAAGTCCCCAGAACGTGGCCACTTCATGTTT  
TCATCCCGGCCTGCACCATCGCTTTGATCTTCCTGGCCATAGTGATAATCCAGAGAA  
AGAGGATCTAGGGGAAGCTGTATTACGGAAGAAGATCTGGACCTGCGGTCTTGGGAG  
TTGGAAGGATCTGATGGGAAACCCTCAAGAGACTTCTGGACTCAAAGTGAGAATCTT  
GCAGGACCTGCCATTTGCACTTTTGAACCCTTTGGACGGTGACCCAGGGCTCCGAAG  
AGGAGCTTGTAAGACTGACAATCTTCCCTCTGTCTCAAGACTCTCTGAACAGCAAGA  
CCCCAATGGCACTTTAGACTTACCCCTGGGATCCTGGACCCCAGTGAGGGCCTAAGG  
CTCCTAATGACTTTTCAAGGTGAGAACAAAAGGAATTGCTCTCCGCCCCACCCCCACC  
TCCTGCTTTCCGCAGGGAGACATGGAAATTCCCAGTTACTAAAATAGATTGTCAATA  
GAGTTATTTATAGCCCTCATTTCCCTCCGGGGACTTGGAAGCTTCAGACAGGGTTTTT  
CATAAACAAAGTCATAACTGATGTGTTTTACAGCATCCTAGAATCCTGGCAGCCTCT  
GAAGTTCTAATTAAGTGAAGCATTTAAGCAACACGTAAAGTACCCCCACTGTGGTA  
TTTGTCTTACTTTTCTGTTTTTAAAGTGTGAGTCACAAGGTAATTGTTGTAACCTG  
TGATATCACTGTTTCTTGTGTCTCTTCTTTCAACTACATCTTTTAAAACAAAACGGT  
GTGGGGTTTGGTTGTTTTGGTGGTAGTGGTAGTGTCTCAGTGGTATCTCCTTAAG  
AAAAAAAAATCATCATGCCAGTGAATTGTTTCTTCAGCCATTTTCAAGATGGGAAGCTGG  
AATAGCCTGTCCCCCAAGCTAAGCCTTCTTCCCTAGCTTTCTGCGTGATTTTACATT  
GAGCATTCCTGTTGCTTTGTTTCTATAACTGTAATGTGGTGATGTCATTGTTAGGGC  
ACTTGAGGGTGGGCGTTCTGGAAGTCCTTTCAGGTTAGTGTTTGGGGGCAGGGTTGC  
TCAGAATACATAAAGGTGCTAACTTAACTGCAGCCATGGAGCTCAGTGAATTCAT  
AACCTTCGGGCTGTCCAAATGTGCACATTAGCTACTGTGACCCCTGTAGGTTAGGGA  
GCCTGAAGCCAGCTCTTTACCTGGTGTTTAGACTCAGCAGAATTTGGAGTCAATGGG

ACCAATGGTTGTGAAATTAAGATTGGAAGTGTGCATCTTATTTTATCACCATCTGC  
CCAACAAAACCTTCAGAAAATGCCTTTGAAGCACAAAAATGTAATCGTTTATGTGAAA  
TCTCTGAGTTGCATTTAGATGCCCATTGCAGCAAGGTGGCTCTCTCACAGATTCCAC  
ACCTTAGCCTAAGATAACCAGACAGCAGGACAGAGAGAGAAAAGTCCTTCTGCTGTGCA  
AACTTCCTTACACTGGACCTCGCCTCTCAGGTGTGTGATTGGTAGGCCAAATCCCGA  
TAGCCAATCGGTGTTGGGTGCTTTGTCTGCTCTACTGGGAGTCCAGTGGTACAATGG  
ATTCTGGCAAAATGCTGCCATCTTGGCCCTCGCTGGGCTGCTTTCTAGGATATTCAT  
AGAGAAAGGGCCGTCCAGATCCAGTATCCTAAAATCCTGAGAGGAGAATATAAGTTA  
GTGTGTCTCACTATAACTATCTCTATGATCGGTCACATTACTATCTAACAGTTACCA  
AATACTATATGCCTAATACTGGTAAGCATTTTATACACACCATTGGATTGAATCCTC  
TCAAAATCCTCAAAAAGGAAGTTATTAATACCTCCATAGGCAAGGAGCCCAGAACCC  
AGAGAGGTCAGGCAGTCTAGTTATAGATGCCTGCTTTGTTTAGAAGTGAACAAGAGC  
ATCAAATTATTAATGTGCCCTGGTTATTAATGCGCCCTGGTTACCTGCTGGATGGAA  
CATCAAGGTGGACTTTTGGCAGTTGCATACACCCAGAGGTATTTTGGCTATTCACGG  
ATTAATTTACACGAAGTGTTTCAGAGACATGTGTAGGGGAAGTCCGGGTTCAGGGG  
GCCTAAGATTCAAACCTCTAGCTTAGCTACGTCTGACCTCCCTAAGCACTAACTTACT  
ATCAAAAGAATGAGCAGTAAAAGAATGGTGTTTACTGCCTGCCTTTATCAGGCAGTG  
AACGTGCAGCGGGCAACGAATGCTTGATAAGTGTGTGTCAGTGTGAAGTCCCATGTA  
CCAGCCGCTGTCCCCACTGCAAAAGCAGCAGAGCGCTCAGACATCATCAGCTGATTT  
ACCAGCAGCAGATTTCTTCTTCTAGTCCCATCCCTGAAGAAGCTTCCAGCCTAGGTA  
CATTGCATGGGCTTTGTGCTCCAGGAGTTCCTACACAGCCCTCAACTTCAACACAGG  
CAAAGTGCTTACTGATCCTCATGTATCTTACAGGGTCCCCTCTACCCACAATACCTC  
ATTGCTGGAACCTTCAAATCTTCCTGAATAAAAGCTTGCCCGTGGTTTAATTA (SEQ ID  
NO: 7) >NP\_067371.1 programmed cell death 1 ligand 2 precursor [*Mus  
musculus*]

MLLLLPILNLSLQLHPVAALFTVTAPKEVYTVDVVGSSVSLECDFDRRECTELEGIRA  
SLQKVENDTSLQSERATLLEEQLPLGKALFHIPSVQVRDSGQYRCLVICGAAWDYKY  
LTVKVKASYMRIDTRILEVPGTGEVQLTCQARGYPLAEVSWQNVSPANTSHIRTP  
GLYQVTSVLRLKPQPSRNFSCMFWNAMKELTSIIDPLSRMEPKVPRTWPLHVFIP  
ACTIALIFLAIVIIQRKRI (SEQ ID NO: 8) Human >NM\_005214.5 *Homo sapiens*  
cytotoxic T-lymphocyte CTLA-4 associated protein 4 (CTLA4), transcript variant  
1, mRNA (CD152)

GCTTTCTATTCAAGTGCCTTCTGTGTGTGCACATGTGTAATACATATCTGGGATCAA  
AGCTATCTATATAAAGTCCTTGATTCTGTGTGGGTCAAACACATTTCAAAGCTTCA  
GGATCCTGAAAGGTTTTGCTCTACTTCCTGAAGACCTGAACACCGCTCCCATAAAGC  
CATGGCTTGCCTTGGATTTACAGCGGCACAAGGCTCAGCTGAACCTGGCTACCAGGAC  
CTGGCCCTGCACTCTCCTGTTTTTTCTTCTCTTCATCCCTGTCTTCTGCAAAGCAAT  
GCACGTGGCCCAGCCTGCTGTGGTACTGGCCAGCAGCCGAGGCATCGCCAGCTTTGT  
GTGTGAGTATGCATCTCCAGGCAAAGCCACTGAGGTCCGGGTGACAGTGCTTCGGCA  
GGCTGACAGCCAGGTGACTGAAGTCTGTGCGGCAACCTACATGATGGGGAATGAGTT  
GACCTTCCTAGATGATTCCATCTGCACGGGGCACCTCCAGTGGAATCAAGTGAACCT  
CACTATCCAAGGACTGAGGGCCATGGACACGGGACTCTACATCTGCAAGGTGGAGCT  
CATGTACCCACCGCCATACTACCTGGGCATAGGCAACGGAACCCAGATTTATGTAAT  
TGATCCAGAACCGTGCCCAGATTCTGACTTCCTCCTCTGGATCCTTGACAGCAGTTAG  
TTCGGGGTTGTTTTTTTATAGCTTTCTCCTCACAGCTGTTTCTTTGAGCAAAATGCT  
AAAGAAAAGAAGCCCTCTTACAACAGGGGTCTATGTGAAAATGCCCCAACAGAGCC  
AGAATGTGAAAAGCAATTTACGCCTTATTTTATTCCCATCAATTGAGAAACCATTAT  
GAAGAAGAGAGTCCATATTTCAATTTCCAAGAGCTGAGGCAATTCTAACTTTTTTGC  
TATCCAGCTATTTTTATTTGTTTGTGCATTTGGGGGGAATTCATCTCTCTTTAATAT  
AAAGTTGGATGCGGAACCCAAATTACGTGTACTACAATTTAAAGCAAAGGAGTAGAA

MACLGFRHKAQLNLATRTWPCTLLFFLLFIPVFCAMHVAQPAVVLASSRGIASFV  
CEYASPGKATEVRVTVLRQADSQVTEVCAATYMMGNELTFLDDSICTGTSSGNQVNL  
TIQGLRAMDTGLYICKVELMYPPPYLIGINGTQIYVIDPEPCPDSDFLWLAAVS  
SGLFFYSFLLTAVSLSKMLKKRSPLTTGVYVKMPPTPEPECEKQFQPYFIPIN (SEQ ID  
NO: 10) Mouse CTLA- >NM\_009843.4 *Mus musculus* cytotoxic T-lymphocyte- 4  
(CD152) associated protein 4 (Ctla4), transcript variant 1, mRNA  
CTACACATATGTAGCACGTACCTTGGATCAAAGCTGTCTATATAAAGTCCCCGAGTC  
TGTGTGGGTTCAAACACATCTCAAGGCTTCTGGATCCTGTTGGGTTTTACTCTGCTC  
CCTGAGGACCTCAGCACATTTGCCCCCAGCCATGGCTTGTCTTGGACTCCGGAGGT  
ACAAAGCTCAACTGCAGCTGCCTTCTAGGACTTGGCCTTTTGTAGCCCTGCTCACTC  
TTCTTTTCATCCCAGTCTTCTCTGAAGCCATACAGGTGACCCAACCTTCAGTGGTGT  
TGGCTAGCAGCCATGGTGTGCGCCAGCTTTCCATGTGAATATTCACCATCACACAACA  
CTGATGAGGTCCGGGTGACTGTGCTGCGGCAGACAAATGACCAAATGACTGAGGTCT  
GTGCCACGACATTCACAGAGAAGAATACAGTGGGCTTCCTAGATTACCCCTTCTGCA  
GTGGTACCTTTAATGAAAGCAGAGTGAACCTCACCATCCAAGGACTGAGAGCTGTTG  
ACACGGGACTGTACCTCTGCAAGGTGGAACCTCATGTACCCACCGCCATACTTTGTGG  
GCATGGGCAACGGGACGCAGATTTATGTCATTGATCCAGAACCATGCCCGGATTCTG  
ACTTCCTCCTTTGGATCCTTGTCGCAGTTAGCTTGGGGTTGTTTTTTTACAGTTTCC  
TGGTCACTGCTGTTTCTTTGAGCAAGATGCTAAAGAAAAGAAGTCCTCTTACAACAG  
GGGTCTATGTGAAAATGCCCCCAACAGAGCCAGAATGTGAAAAGCAATTCAGCCTT  
ATTTTATTTCCCATCAACTGAAAGGCCGTTTATGAAGAAGAAGGAGCATACTTCAGTC  
TCTAAAAGCTGAGGCAATTTCAACTTTCTCTTCTCTCCAGCTATTTTTTACCTGTTT  
GTATATTTTAAGGAGAGTATGCCTCTCTTTAATAGAAAGCTGGATGCAAAATTCCAA  
TTAAGCATACTACAATTTAAAGCTAAGGAGCATGAACAGAGAGCTGGGATATTTCTG  
TTGTGTCAGAACCATTTTACTAAAAGCATCACTTGAAGCAGCATAAGGATATAGCA  
TTATGGTGTGGGGTCAAGGGAACATTAGGGAATGGCACAGCCCCAAAGAAAGGAAGGG  
GGTGAAGGAAGAGATTATATTGTACACATCTTGTATTTACCTGAGAGATGTTTATGA  
CTTAAATAATTTTTAAATTTTTCATGCTGTTATTTTCTTTAACAATGTATAATTACA  
CGAAGGTTTAAACATTTATTCACAGAGCTATGTGACATAGCCAGTGGTTCCAAAGGT  
TGTAAGTGTTCGAAGATGTATTTTAAAGTAATATTGTACATGGGTGTTTCATGTGCTG  
TTGTGTATTTGCTGGTGGTTTGAATATAAACACTATGTATCAGTGTCGTCACAGT



SAAGAAALATVDPSSAFAFGQGRLLHLHLSAGQRLGLVHLETHARHAWQL  
TQGATVLGLFRVTPEIPAGLPSRSE (SEQ ID NO: 14) Mouse 4-1BBL  
>NM\_009404.3 *Mus musculus* tumor necrosis factor (ligand) (CD137L) superfamily,  
member 9 (Tnfsf9), mRNA  
ATAAAGCACGGGCACTGGCGGGAGACGTGCACTGACCGACCGTGGTAATGGACCAGC  
ACACACTTGATGTGGAGGATACCGCGGATGCCAGACATCCAGCAGGTACTTCGTGCC  
CCTCGGATGCGGGCGCTCCTCAGAGATACCGGGCTCCTCGCGGACGCTGCGCTCCTCT  
CAGATACTGTGCGCCCCACAAATGCCGCGCTCCCCACGGATGCTGCCTACCCTGCGG  
TTAATGTTGCGGATCGCGAGGCCGCGTGGCCGCCTGCACTGAACTTCTGTTCCCGCC  
ACCCAAAGCTCTATGGCCTAGTCGCTTTGTTTGTGCTGCTTCTGATCGCCGCCTGTG  
TTCCTATCTTCACCCGCACCGAGCCTCGGCCAGCGCTCACAATCACCACCTCGCCCA  
ACCTGGGTACCCGAGAGAATAATGCAGACCAGGTCACCCCTGTTTCCCACATTGGCT  
GCCCCAACACTACACAACAGGGCTCTCCTGTGTTTCGCCAAGCTACTGGCTAAAAACC  
AAGCATCGTTGTGCAATACAACTCTGAACTGGCACAGCCAAGATGGAGCTGGGAGCT  
CATACCTATCTCAAGGTCTGAGGTACGAAGAAGACAAAAAGGAGTTGGTGGTAGACA  
GTCCCGGGCTCTACTACGTATTTTGGAACTGAAGCTCAGTCCAACATTCACAAACA  
CAGGCCACAAGGTGCAGGGCTGGGTCTCTCTTGTGTTTGAAGCAAAGCCTCAGGTAG  
ATGACTTTGACAACTTGGCCCTGACAGTGGAAGCTGTTCCCTTGCTCCATGGAGAACA  
AGTTAGTGGACCGTTCCTGGAGTCAACTGTTGCTCCTGAAGGCTGGCCACCGCCTCA  
GTGTGGGTCTGAGGGCTTATCTGCATGGAGCCCAGGATGCATACAGAGACTGGGAGC  
TGTCTTATCCCAACACCACCAGCTTTGGACTCTTTCTTGTGAAACCCGACAACCCAT  
GGGAATGAGAACTATCCTTCTTGTGACTCCTAGTTGCTAAGTCCTCAAGCTGCTATG  
TTTTATGGGGTCTGAGCAGGGGTCCCTTCCATGACTTTCTCTTGTCTTTAACTGGAC  
TTGGTATTTATTCTGAGCATAGCTCAGACAAGACTTTATATAATTCAGTAGATAGCA  
TTAGTAAACTGCTGGGCAGCTGCTAGATAAAAAAAATTTCTAAATCAAAGTTTATA  
TTTATATTAATATATAAAAAATAAATGTGTTTGT (SEQ ID NO: 15) >NP\_033430.1  
tumor necrosis factor ligand superfamily member 9 [*Mus musculus*]  
MDQHTLDVEDTADARHPAGTSCPSDAALLRDTGLLADAALLSDTVRPTNAALPTDAA  
YPAVNVRDREAAWPPALNFCSRHPKLYGLVALVLLLLIAACVPIFTRTEPRPALTIT  
TSPNLGTRENNADQVTPVSHIGCPNTTQQGSPVFAKLLAKNQASLCNTTLNWHQSQDG  
AGSSYLSQGLRYEEDKKELVVDSPGLYVVFLELKLSPFTNTGHKVQGWVSLVLQAK  
PQVDDFDNLALTVELFPCSMENKLVDRSWSQLLLKAGHRLSVGLRAYLHGAQDAYR  
DWELSYPNTTSFGLFLVKPDNPWE (SEQ ID NO: 16) Human >NM\_003820.4  
*Homo sapiens* TNF receptor superfamily member HVEM 14 (TNFRSF14),  
transcript variant 1, DNA (CD270)  
ATACCGGCCCTTCCCCTCGGCTTTGCCTGGACAGCTCCTGCCTCCCGCAGGGCCCCAC  
CTGTGTCCCCCAGCGCCGCTCCACCCAGCAGGCCTGAGCCCCTCTCTGCTGCCAGAC  
ACCCCTGCTGCCCACTCTCCTGCTGCTCGGGTTCTGAGGCACAGCTTGTACACCG  
AGGCGGATTCTCTTTCTCTTTCTCTTTCTCTTCTGGCCCACAGCCGCAGCAATGGCG  
CTGAGTTCCTCTGCTGGAGTTCATCCTGCTAGCTGGGTTCCCGAGCTGCCGGTCTGA  
GCCTGAGGCATGGAGCCTCCTGGAGACTGGGGGCCTCCTCCCTGGAGATCCACCCCC  
AAAACCGACGTCTTGAGGCTGGTGCTGTATCTCACCTTCTGAGCCCCCTGCTAC  
GCCCCAGCTCTGCCGTCCTGCAAGGAGGACGAGTACCCAGTGGGCTCCGAGTGCTGC  
CCAAGTGCAGTCCAGGTTATCGTGTGAAGGAGGCCTGCGGGGAGCTGACGGGCACA  
GTGTGTGAACCCTGCCCTCCAGGCACCTACATTGCCACCTCAATGGCCTAAGCAAG  
TGTCTGCAGTGCCAAATGTGTGACCCAGCCATGGGCCTGCGCGCGAGCCGGAAGTGC  
TCCAGGACAGAGAACGCCGTGTGTGGCTGCAGCCCAGGCCACTTCTGCATCGTCCAG  
GACGGGGACCACTGCGCCGCGTGCCGCGCTTACGCCACCTCCAGCCCCGGGCCAGAGG  
GTGCAGAAGGGAGGCACCGAGAGTCAGGACACCCTGTGTGCAAGTGCACCCCCGGGG  
ACCTTCTCTCCCAATGGGACCCTGGAGGAATGTCAGCACCAAGTGCAGCTGG



CTGGTGACGAAGGCCGGAGCTGGGACCAGCAGCTCCCACTGGGTATGGTGGTTC  
TCAGGGAGCCTCGTCATCGTCATTGTTTGTCTCCACAGTTGGCCTAATCATATGTGTG  
AAAAGAAGAAAGCCAAAGGGGTGATGTAGTCAAGGTGATCGTCTCCGTCCAGCGGAAA  
AGACAGGAGGCAGAAGGTGAGGCCACAGTCATTGAGGCCCTGCAGGCCCTCCGGAC  
GTCACCACGGTGGCCGTGGAGGAGACAATAACCCTCATTCACGGGGAGGAGCCCAAAC  
CACTGACCCACAGACTCTGCACCCCGACGCCAGAGATACTGGAGCGACGGCTGCTG  
AAAGAGGCTGTCCACCTGGCGGAACCACCGGAGCCCGGAGGCTTGGGGGCTCCGCCC  
TGGGCTGGCTTCCGTCTCCTCCAGTGGAGGGAGAGGTGGGGCCCCTGCTGGGGTAGA  
GCTGGGGACGCCACGTGCCATTCCCATGGGCCAGTGAGGGCCTGGGGCCTCTGTTCT  
GCTGTGGCCTGAGCTCCCCAGAGTCCTGAGGAGGAGCGCCAGTTGCCCTCGCTCAC  
AGACCACACACCCAGCCCTCCTGGGCCAGCCCAGAGGGCCCTTCAGACCCAGCTGT  
CTGCGCGTCTGACTCTTGTGGCCTCAGCAGGACAGGCCCCGGGCACTGCCTCACAGC  
CAAGGCTGGACTGGGTTGGCTGCAGTGTGGTGTTTAGTGGATAACCATCGGAAGTG  
ATTTTCTAAATTGGATTTGAATTCGGCTCCTGTTTTCTATTTGTCATGAAACAGTGT  
ATTTGGGGAGATGCTGTGGGAGGATGTAAATATCTTGTCTCCTCAA (SEQ ID  
NO: 17) >NP\_003811.2 tumor necrosis factor receptor superfamily member 14  
isoform 1 precursor [*Homo sapiens*]

MEPPGDWGWPPPWRSTPKTDVLRVLVLYLTFLGAPCYAPALPSCKEDEYPVGSECCPKC  
SPGYRVKEACGELTGTVCPCPPGTYIAHLNGLSKCLQCQMCDPAMGLRASRNCST  
ENAVCGCSPGHFCIVQGDHCAACRAYATSSPGQVRVQKGGTESQDTLCQNCPPGTFS  
PNGTLEECQHQTCKSWLVTKAGAGTSSSHWVWWFLSGSLVIVIVCSTVGLIICVKRR  
KPRGDVVKVIVSVQRKRQEAEGEATVIEALQAPPDVTTVAVEETIPSFTGRSPNH (SEQ  
ID NO: 18) Mouse HVEM >NM\_178931.2 *Mus musculus* tumor necrosis factor  
receptor (CD270) superfamily, member 14 (herpesvirus entry mediator) (Tnfrsf14),  
mRNA

GCTCTTGGCCTGAAGTTTCTTGATCAAGAAAATGGAACCTCTCCCAGGATGGGGGTC  
GGCACCTTGAGCCAGGCCCTACAGACAACACCTTCAGGCTGGTGCCTTGTGTCTT  
CCTTTTGAACCTTGCTGCAGCGCATCTCTGCCCAGCCCTCATGCAGACAGGAGGAGTT  
CCTTGTGGGAGACGAGTGCTGCCCCATGTGCAACCCAGGTTACCATGTGAAGCAGGT  
CTGCAGTGAGCATAACAGGCACAGTGTGTGCCCCCTGTCCCCACAGACATATACCGC  
CCATGCAAATGGCCTGAGCAAGTGTCTGCCCTGCGGAGTCTGTGATCCAGACATGGG  
CCTGCTGACCTGGCAGGAGTGCTCCAGCTGGAAGGACACTGTGTGCAGATGCATCCC  
AGGCTACTTCTGTGAGAACCAGGATGGGAGCCACTGTTCCACATGCTTGCAGCACAC  
CACCTGCCCTCCAGGGCAGAGGGTAGAGAAGAGAGGGACTCACGACCAGGACACTGT  
ATGTGCTGACTGCCTAACAGGGACCTTCTCACTTGGAGGGACTCAGGAGGAATGCCT  
GCCCTGGACCAACTGCAGTGCATTTCAACAGGAAGTAAGACGTGGGACCAACAGCAC  
AGACACCACCTGCTCCTCCCAGGTCGTCTACTACGTTGTGTCCATCCTTTTGCCACT  
TGTGATAGTGGGAGCTGGGATAGCTGGATTCTCATCTGCACGCGAAGACACCTGCA  
CACCAGCTCAGTGGCCAAGGAGCTGGAGCCTTTCCAGGAACAACAGGAGAACACCAT  
CAGGTTTCCAGTCACCGAGGTTGGGTTTGCTGAGACCGAGGAGGAGACAGCCTCCAA  
CTGAACAAATTCTGGGTGACAAGACACCGAGGAGACGT (SEQ ID NO: 19)

>NP\_849262.1 tumor necrosis factor receptor superfamily member 14 precursor  
[*Mus musculus*]

MEPLPGWGSAPWSQAPTDNTFRLVPCVFLNLLQRISAQPSCRQEEFLVGDECCPMC  
NPGYHVKQVCSEHTGTVCAPCPPTQTYTAHANGLSKCLPCGVCDPDMGLLTWQECSSW  
KDTVCRICIPGYFCENQDGS HCSTCLQH TTCPPGQ RVEKRGTHDQD TVCADCLTGTFS  
LGGTQEECLPWTNCSAFQQEVRRGTNSTDTTCSSQVVYYVVSILLPLVIVGAGIAGF  
LICTRRHLHTSSVAKELEPFQEQQENTIRFPVTEVGFAETEEETASN (SEQ ID NO: 20)  
Human FGL1 >NM\_004467.4 *Homo sapiens* fibrinogen like 1 (FGL1), transcript  
variant 1, mRNA

AAATGCTTACGATCTGAGTCTGGAAGAGAGTGTGGAATGGAGTGGTCT  
CTGACTCACTTCTGACTTTAGTTTTTCAAGGGGGAACATGGCAAAGGTGTTTCAGTT  
TCATCCTTGTTACCACCGCTCTGACAATGGGCAGGGAAATTTTCGGCGCTCGAGGACT  
GTGCCCAGGAGCAGATGCGGCTCAGAGCCCAGGTGCGCCTGCTTGAGACCCGGGTCA  
AACAGCAACAGGTCAAGATCAAGCAGCTTTTGCAGGAGAATGAAGTCCAGTTCCTTG  
ATAAAGGAGATGAGAATACTGTCATTGATCTTGGAAGCAAGAGGCAGTATGCAGATT  
GTTTCAGAGATTTTCAATGATGGGTATAAGCTCAGTGGATTTTACAAAATCAAACCTC  
TCCAGAGCCCAGCAGAATTTTCTGTTTATTGTGACATGTCCGATGGAGGAGGATGGA  
CTGTAATTCAGAGACGATCTGATGGCAGTGAAAACCTTTAACAGAGGATGGAAAGACT  
ATGAAAATGGCTTTTGGAATTTTGTCCAAAAACATGGTGAATATTGGCTGGGCAATA  
AAAATCTTCACTTCTTGACCACTCAAGAAGACTACACTTTAAAAATCGACCTTGCAG  
ATTTTGAAAAAATAGCCGTTATGCACAATATAAGAATTTCAAAGTTGGAGATGAAA  
AGAATTTCTACGAGTTGAATATTGGGGAATATTCTGGAACAGCTGGAGATTCCCTTG  
CGGGGAATTTTCATCCTGAGGTGCAGTGGTGGGCTAGTCACCAAAGAATGAAATTCA  
GCACGTGGGACAGAGATCATGACAACTATGAAGGGAAGTGCAGAGAAGAAGATCAGT  
CTGGCTGGTGGTTTAAACAGGTGTCCTCTGCAAACCTGAATGGTGTATACTACAGCG  
GCCCCTACACGGCTAAAACAGACAATGGGATTGTCTGGTACACCTGGCATGGGTGGT  
GGTATTCTCTGAAATCTGTGGTTATGAAAATTAGGCCAAATGATTTTATTCCAAATG  
TAATTTAATTGCTGCTGTTGGGCTTTCGTTTCTGCAATTCAGCTTTGTTTAAAGTGA  
TTTGAAAAATACTCATTCTGAACATATCCATGCGCAATCATGATAACTGTTGTGAGT  
AGTGCTTTTTCATTCTTCTCACTTGCCTTTGTTACTTAATGTGCTTTCAGTACAGCAG  
ATATGCAATATTCACCAAATAAATGTAGACTGTGTTAATA (SEQ ID NO: 21)

>NP\_004458.3 fibrinogen-like protein 1 precursor [*Homo sapiens*]

MAKVFSFILVTTALTMGREISALEDCAQEQMRLRAQVRLLETRVKQQQVKIKQLLQE  
NEVQFLDKGDENTVIDLGSKRQYADCSEIFNDGYKLSGFYKIKPLQSPAEFVYCDM  
SDGGGWTVIQRRSDGSENFNRGWKDYENGFGNFVQKHGEYWLGNKNLHFLTQEDYT  
LKIDLADFEKNSRYAQYKNFKVGDEKNFYELNIGEYSGTAGDSLGNFHFPEVQWWAS  
HQRMKFSTWDRDHDNYEGNCAEEDQSGWWFNRCHSANLNGVYYSGPYTAKTDNGIVW  
YTWHGWWYSLKSVVMKIRPNDIFIPNVI (SEQ ID NO: 22) Mouse FGL1

>NM\_145594.2 *Mus musculus* fibrinogen-like protein 1 (Fgl1), mRNA

GTTAGAAGTTCCTGGGAGGCTCTGTGTGGATGGACTGAGCCTAGCTAAGTCCTGATT  
CATTTTGACTTGAGTTCTCTCAGTGGGAAGAATGGGAAAGATTTACAGCTTCGTCCT  
GGTCGCCATTGCTCTGATGATGGGAAGGGAAGGTTGGGCCCTCGAGAGTGAGAACTG  
CTTGCGGGAGCAGGTGAGGCTCAGGGCTCAGGTGCACCAGCTTGAGACCCGGGTCAA  
ACAACAACAGACCATGATTGCACAGCTCTTGCATGAGAAGGAAGTCCAGTTTCTGGA  
TAAAGGATCGGAGAACAGTTTCATTGACCTTGGAGGCAAGAAGCAGTATGCAGATTG  
TTCAGAGATTTACAATGACGGATTTAAGCAGAGTGGATTTTACAAAATCAAACCTCT  
TCAGAGCCTGGCAGAATTCTCTGTTTATTGTGACATGTCTGATGGAGGGGGATGGAC  
TGTAATTCAGAGACGATCTGATGGCAGTGAGAACTTTAACAGGGGTTGGAATGACTA  
TGAAAATGGCTTTTGAAACTTTGTCCAAAACAATGGCGAATACTGGCTGGGTAACAA  
AAACATTAACCTTGCTAACTATTCAAGGAGACTACACTTTAAAAATCGACCTGACAGA  
TTTTGAGAAAAACAGCAGCTTCGCACAATACCAAAGTTTTTAAAGTTGGTGATAAAAA  
GTCTTTTTATGAACTAAATATTGGAGAATATTCTGGCACAGCTGGAGATTCCCTGTC  
AGGAACTTTTCATCCTGAAGTACAGTGGTGGGCTAGTCACCAAAGGATGAAGTTCAG  
CACGTGGGACAGAGATAACGACAATTACCAAAGGAACTGTGCTGAGGAAGAGCAGTC  
TGGCTGGTGGTTTAAACAGGTGTCCTCTGCAAACCTGAACGGTGTCTTACTACCGTGG  
TTCCTACAGGGCAGAAACGGATAATGGTGTGTGTGGTACACCTGGCATGGGTGGTG  
GTATTCCTTGAAATCTGTGGTTATGAAAATTAGGCCAAGTGATTTTATTCCAAATAT  
TATTTAGTTGCCCTCATTGGGATCTCCTTTCTGTAATTCATCTTGGTTTACTTGAAA  
ATAAATATTTGAAAAAGATATAATTCTGAATAACACA (SEQ ID NO: 23)

>NP\_63569.2 fibrogen-like protein 1 precursor [Mus musculus]  
MGKIYSFVLVAIALMMGREGWALESENCLREQVRLRAQVHQLETRVKQQQTMIQQLL  
HEKEVQFLDKGSENSFIDLGGKKQYADCSEIYNDGFKQSGFYKIKPLQSLAEFSVYC  
DMSDGGGWTVIQRRSDGSENFNRGWNDYENGFGNFVQNNGEYWLGNKNINLLTIQGD  
YTLKIDLTDFEKNSSFAQYQSFKVGDKKSFYELNIGEYSGTAGDSLSTGFHPEVQWW  
ASHQRMKFSTWDRDNDNYQGNCAEEEEQSGWWFNRCHSANLNGVYYRGSYRAETDNGV  
VWYTWHGWWYSLKSVVMKIRPSDFIPNII (SEQ ID NO: 24) Human OX-2  
>NM\_005944.7 Homo sapiens CD200 molecule (CD200), (CD200) transcript variant  
1, mRNA  
AGAGCTCCAGGCGCACATCCGCAGTCAGCCACCTCGCGCGCGCCTCCAGGAGCAAGG  
ATGGAGAGGCTGGTGATCAGGATGCCCTTCTCTCATCTGTCTACCTACAGCCTGGTT  
TGGGTCATGGCAGCAGTGGTGCTGTGCACAGCACAAAGTGCAAGTGGTGACCCAGGAT  
GAAAGAGAGCAGCTGTACACACCTGCTTCCCTTAAAATGCTCTCTGCAAAATGCCCAG  
GAAGCCCTCATTGTGACATGGCAGAAAAAGAAAGCTGTAAGCCCAGAAAACATGGTC  
ACCTTCAGCGAGAACCATGGGGTGGTGATCCAGCCTGCCTATAAGGACAAGATAAAC  
ATTACCCAGCTGGGACTCCAAAACCTCAACCATCACCTTCTGGAATATCACCTTGGAG  
GATGAAGGGTGTTACATGTGTCTCTTCAATACCTTTGGTTTTGGGAAGATCTCAGGA  
ACGGCCTGCCTCACCGTCTATGTACAGCCCATAGTATCCCTTCACTACAAATTCTCT  
GAAGACCACCTAAATATCACTTGCTCTGCCACTGCCCCGCCAGCCCCCATGGTCTTC  
TGGAAGGTCCCTCGGTCAGGGATTGAAAATAGTACAGTGACTCTGTCTCACCCAAAT  
GGGACCACGTCTGTTACCAGCATCCTCCATATCAAAGACCCTAAGAATCAGGTGGGG  
AAGGAGGTGATCTGCCAGGTGCTGCACCTGGGGACTGTGACCGACTTTAAGCAAACC  
GTCAACAAAGGCTATTGGTTTTTCAGTTCCGCTATTGCTAAGCATTGTTTCCCTGGTA  
ATTCTTCTCGTCCTAATCTCAATCTTACTGTACTGGAAACGTCACCGGAATCAGGAC  
CGAGAGCCCTAAATAAGTCACACAGCACCTGAAAGTGATTCCCTGGTCTACTTGAA  
TTTGACACAAGAGAAAAGCAGGAGGAAAAGGGGCCATTCTCCAAAGGACCTGAAAGA  
GCAAAAGAGGTGGGAGCGAAAGCCTTAAGGATCCCACGACTTTTTACTGCCATCTGA  
GCTACTCAGTGTTTGAATCCCAAGAGGAAGTCAGTTTACCTCTCAGGTCTGTTGTAG  
GACTTGATTTTGTAAAGCAATGCCATGTTATGTGGTTGAAAGGGCACTGGACTTAGT  
TAGTATCAGGAGCACTGAGCTCACAGACTGACTTGGGCTCCTACTGGTGGGGACCTC  
TGTTAGTCACTTTACCTCATCCAAAGTATAAAGGAATTGGACCAAATAATTTACCAC  
ATAGCTCTAAAACTTAATTTAAAATGTAATTCCAGAAAAAAAAGGGAATAAGCAAA  
GGGGGAAGAATTGAAAGAGAGAGAGAAGAAAGAAATACAGAGAGCTTACCTTTTGCCT  
TTCTGTTGATGTTACATCTCTTCTTCCCTATGTTCTTAGGTCTATGAGTCTGTTTCCC  
CATCATTTGGTATCTAGTCCAGTTCCTGCTTACTGCTTTGCTAATAGCTGGCCTTGC  
TAGAATCCTTGGTTTTCACTGCTGTTCTTCATGTGCTTCTATGAGATTTACTCCAACA  
CAAATAGGACTGAATTTATTGTGAAGTAACATTGGCAATCTTAACTTATTCATTTAA  
CTTATTTTTATAGCTAGATAAATATTGTTAGTCTTAGACAATAGCTCACATTTTTTG  
AGAAGCATGCCCTCCCTGTCCATTTGTCTTATAACATGACCCAGCCCTATTTTACGT  
CATTCTAAATTCAGCCTCATATAATGAAAATACATTATGAAAACAGATGTTTAGGAG  
ATTTCTGTATAGCAGTCAGCCAATTCATATGCTTTGTCTCTGCTGGCTTCTTTTTTC  
CATGCGTTAACTTTTTCCCAATAGCAGAGGAGGCAAATATGAGCATAACAATCCCTTTG  
TTCTAAAGATATTGTTCCAGCTAGTGGAATGATGTTGAATCTTTAATAACCATAATT  
AGTTGCTTTTTTCAGTATCTTCTGCTTTGTCTGTGTCTATCCAGTGGCCTAGGAATTA  
AAGTGTAAGTTGTTTTTCGCTGTTAAATTGGATATTTATATATATATATAGCAAGATT  
TTCATGTGTTATTTAATTCTGTATTGTTTCTTATATTTGTAGTAAAATATTGAACAA  
TTAAAAGTGTTGACTCCAAA (SEQ ID NO: 25)>NP\_005935.4 OX-2 membrane  
glycoprotein isoform a precursor [Homo sapiens]  
MERLVIRMPFSLSTYSLVWVMAAVVLCTAQVQVVTQDEREQLYTPASLKCSLQNAQ  
EALIVTWQKKKAVSPENMVTFSENHGVVIQPAYKDKINITQLGLQNSTITFWNITLE

DEGCYMLFNGTFCYCLTGYVQPIVSLHYKFSEDLNITCSATARPAPMVF  
WKVPRSGIENSTVTLSHPNGTTSVTSILHIKDPKNQVGKEVICQVLHLGTVTDFKQT  
VNKGYWFSVPLLLSIVSLVILLVLISILLYWKRHRNQDREP (SEQ ID NO: 26)  
Mouse OX-2 >NM\_010818.3 *Mus musculus* CD200 antigen (Cd200), (CD200)  
transcript variant 1, mRNA  
GGGCGTGGTTGGTTGGTCTCTCTCCTCCACACTAGAGGAGCTGTAGAGTCTGCCT  
GTGCAGTGGAGGGGGCTCTCTCTACGGCGAATAGTAGTGTCCCTGCTCACAGGTGTT  
GCGGAGATATCCTCCATCGTGGAAGAGCTCAGACCCCGAGAAGCTGGTGTCTAGCTG  
CGGCCCAGAGCAAGGATGGGCAGTCTGGTATTACAGGAGACCTTTCTGCCATCTCTCC  
ACCTACAGCCTGATTTGGGGCATGGCAGCAGTAGCGCTGAGCACAGCTCAAGTGGA  
GTGGTGACCCAGGATGAAAGAAAGGCGCTGCACACAACCTGCATCCTTACGATGTTCT  
CTAAAAACATCCCAGGAACCCTTGATTGTGACATGGCAGAAAAAGAAAGCCGTGAGC  
CCAGAAAAACATGGTCACCTACAGCAAAACCCATGGGGTTGTAATCCAGCCTGCCTAC  
AAAGACAGGATAAATGTCACAGAGCTGGGACTCTGGAACCTCAAGCATCACCTTCTGG  
AACACAACATTGGAAGATGAGGGCTGCTACATGTGTCTCTTCAACACGTTTGGTTCT  
CAGAAGGTCTCAGGAACAGCTTGCCCTTACCCTCTATGTACAGCCCATAGTACACCTT  
CACTACAACATTTTTGAAGACCACCTAAACATCACTTGCTCTGCGACTGCCCGTCCA  
GCCCCTGCCATCTCCTGGAAGGGTACTGGGACAGGAATTGAGAATAGTACCGAGAGT  
CACTTCCATTCAAATGGGACTACATCTGTCACCAGCATCCTCCGGGTCAAAGACCCC  
AAAACCTCAAGTTGGGAAGGAAGTGATCTGCCAGGTTTTTATACCTGGGGAATGTGATT  
GACTACAAGCAGAGTCTGGACAAAGGATTTTTGGTTTTTCAGTTCCACTGTTGCTAAGC  
ATTGTTTTCTCTGGTAATTCTTCTGATCTTGATCTCCATCTTACTATACTGGAAACGT  
CACCGAAATCAGGAGCGGGGTGAATCATCACAGGGGATGCAAAGAATGAAATAAGAG  
CTCTAAAGAAATTATACAGAACCCTGAACGTGTTTCCCTGGTCTACTTGAATCTGAT  
GTGAAAGAAAAGCAGGAGGGGAAAAGGCCATTCTCCATAGGACCTAAGGAGAGCAAAA  
GACCAGACACGAGCCTGTGAGGGATTTGACTTTTTTGTCTGTTGTCCCAGGTCCTCGGT  
GTTTGCATTCCAAGAGGAAGTCGAGTGCCTCGGGTCTGTTGTAGGACTTGATTTTTT  
TTTTTTTTGTAGAGCAATGCAGTGCCATGCTGTTAGAAAGGCTCCAGACTTAGAACC  
ACCAGTGCCAAGCCAGCTCTCAGACCGACTAGGGCTCCCATCGGAGGAACAAATCGT  
AGTCAACTTACCTCACAGAGCTCTCTGGTCCTTACACAAAGTAGAAAGGAGTGGGAC  
CAGAAAATTGGCCATGTCTGAAATCTGATGGAATTTTTAGGAAGAAAACCTGAAGAAT  
AAGCAAAAGAAGAAAGAACACAGAAGGGTCCAAAGAGCTTCTGAGAGTACCTTTTGC  
CTTTCTGTTGGTGTCCCAGCTCTGGTTTTTGTCTTAGGTCCGCCAGTGTGTTTTCCCT  
GTTGTTTGAGTATCTAGTTGACTACCTGCTACTGTTCTGCTGATGGTTGGCCTTGCT  
AGAATCCCTGACTCCCCTGCCGTTCTCTATGTGCTTCTATGAGGGTTACTATGATGA  
AAATAGAGCAGAAGATAGTGTGAAGTAACATTGGCAACTGTAATGTGTCCATTTAAC  
TTATTTTTATAGCACTTAGGCAATATTGTTAGTCTTAGTGAGTAGTTCACATCTTTA  
CAAAAGCATGCTCTCCCTATCCATTGGGCCCACAATAACACTCTCTTTGAGGCCATT  
CTGAATCCTGTCTCGTGTAATGATAATATATTATGAAAACAGATACTTTAAGAATTT  
CCTGTACAGCAGTCAGTTGTTTATTCTCTCTCTCTCTCTCTCTCTCTCTCTCTCTCCCTCC  
CCCACCCAGCTTCTTTTTCTGTGACTTTGTTTTTCATAAAGAGAAGGCATCTCCTG  
AATACAATCGCTTTGTTCTGAAGACATCGTGAACCTATTAATTCTTAACCCTTTGACA  
AACTAGTGAAAGTTGTTTTCTGTATCTTTTGCTTCATCTGTCTTTATAGAGTGACCT  
AGGAATTCAAGTGTAAGTTGTTTCCATTGTTGAACTGGATATTTATATACTTGGTAT  
GCTTTTCACGTGTTATTTAATTCTGTATAATTTCCCTATATTTGTATTAAAATATTGA  
GCAATTAAAAGTGTCAACTAAATATTTGATGTGGCATTCCCTTGAGAAATATAGAAA  
TAAAGAATAAAAAAAAAAAAAAAAAAAAAA (SEQ ID NO: 27) >NP\_034948.3 OX-2  
membrane glycoprotein isoform 1 precursor [*Mus musculus*]  
MGSLVFRRPFCHLSTYSLIWGMAAVALSTAQVEVVTQDERKALHTTASLRCSLKTSQ  
EPLIVTWQKKKAVSPENMVTYSKTHGVVIQPAYKDRINVTELGLWNSSITFWNTTLE



CACTTCAACCAACCGGTTGAGGAGAGGAGGATGTGGTGGTGGCAACGAAAC  
GGACAGTGGGGTCCTGAGGAGAGAAAGATGCAGATGCCCTTCCAGAAGGGGATGCCC  
TTTGAGCTTTGCTTCCTGGTGCAGAGGTCAGAGTTCAAGGTGATGGTGAACAAGAAA  
TTCTTTGTGCAGTACCAACACCGCGTACCCTACCACCTCGTGGACACCATCGCTGTC  
TCCGGCTGCTTGAAGCTGTCCTTTATCACCTTCCAGAACTCTGCAGCCCCTGTCCAG  
CATGTCTTCTCCACAGTGCAGTTCTCTCAGCCAGTCCAGTTCCCACGGACCCCTAAG  
GGGCGCAAACAGAAACTCAGAACTTTCGTCCTGCCCACCAGGCACCCATGGCTCAA  
ACTACCATCCATATGGTTCACAGCACCCCTGGACAGATGTTCTCTACTCCTGGAATC  
CCTCCTGTGGTGTACCCACCCACCCAGCCTATACCATACCTTTCTACACCCCCATTCCA  
AATGGGCTTTACCCGTCCAAGTCCATCATGATATCAGGCAATGTCTTGCCAGATGCT  
ACGAGGTTCCATATCAACCTTCGCTGTGGAGGTGACATTGCTTTCCACCTGAACCCC  
CGTTTCAATGAGAATGCTGTTGTCCGAAACACTCAGATCAACAACCTCTGGGGGCAG  
GAAGAGCGAAGTCTGCTTGGGAGGATGCCCTTCAGTCGAGGCCAGAGCTTCTCGGTG  
TGGATCATATGTGAAGGTCAGTCTTCAAGGTAGCTGTGAATGGTCAACACATGTGT  
GAATATTACCACCGCCTGAAGAACTTGCAGGATATCAACACTCTAGAAGTGGCGGGT  
GATATCCAGCTGACCCACGTGCAGACATAGGCAAGGTCTCTGGCCTAGGGATAAGGG  
CTGGAGCACTCTGCCTGTGTCTTATCTTTCCCTGTCTCAGCCCTGGCACCATCAGA  
AGAGATCATCACTTATAGGAATTCCAGGAAGGTGAAATTCCCAATTGACTCCCTCCA  
CAAAGGGGGTTTTCTAGGCTGTGTGGCACATGGCTGTCAGCCCATAGTCTGAGCCAT  
TGCCCCCAAGCTAGCTATATACTGAGGGAAGTGACCCTCCTGGGTTTGCTCAGATCT  
CTGATCGTTCCCCCCTCTGTGGCCCTTTTCTTTACCCCTCCAGGAGAGCCACCCTG  
ATATCATCCCCTGACCTCCAACCTGACCCACAATGTCCACAGTAACCTTTCCCCCATT  
CTCACCCAGTATCCATAAAATAAAGAAATAATATTGCTTGTCTACAC (SEQ ID NO:  
31)>NP\_034838.2 galectin-9 isoform 1 [*Mus musculus*]  
MALFSAQSPYINPIIPFTGPIQGGLEGLQVTLQGTTKSFAQRVFNQNSFNGNDI  
AFHFNPRFEEGGYVVCNTKQNGQWGPEERKMQMPPFQKGMPFELCFLVQRSEFKVMVN  
KKFFVQYQHRVPYHLVDTIASGCLKLSFITFQNSAAPVQHVFSSTVQFSQPVPQFPRT  
PKGRKQKTQNFPAHQAPMAQTTHMVHSTPGQMFSTPGIPPVVYPTPAYTIPFYTP  
IPNGLYPSKSIMISGNVLPDATTRFHINLRCGGDIAFHLNPRFNENAVVRNTQINNSW  
GQEERSLLGRMPFSRQSFVSWIICEGHCFKVAVNGQHMCEYYHRLKNLQDINTLEV  
AGDIQLTHVQT (SEQ ID NO: 32) Human PVR>NM\_006505.5 *Homo sapiens*  
PVR cell adhesion molecule (CD155) (PVR), transcript variant 1, mRNA  
AGTCACTTGTCTGGAGCTTGAAGAAGTGGGTATTCCCCTTCCCACCCAGGCACTGG  
AGGAGCGGCCCCCGGGGATTCCAGGACCTGAGCTCCGGGAGCTGGACTCGCAGCGA  
CCGCGGCAGAGCGAGCGGGCGCCGGGAAGCGAGGAGACGCCCGCGGGAGGCCAGCT  
GCTCGGAGCAACTGGCATGGCCCGAGCCATGGCCGCGCGTGGCCGCTGCTGCTGGT  
GGCGCTACTGGTGTCTGCTGGCCACCCCCAGGAACCGGGGACGTCGTCGTGCAGGC  
GCCCACCCAGGTGCCCGGCTTCTTGGGCGACTCCGTGACGCTGCCCTGCTACCTACA  
GGTGCCCAACATGGAGGTGACGCATGTGTACAGCTGACTTGGGCGCGGCATGGTGA  
ATCTGGCAGCATGGCCGTCTTCCACCAAACGCAGGGCCCCAGCTATTCGGAGTCCAA  
ACGGCTGGAATTCGTGGCAGCCAGACTGGGCGCGGAGCTGCGGAATGCCTCGCTGAG  
GATGTTCCGGGTGCGCGTAGAGGATGAAGGCAACTACACCTGCCTGTTTCGTACGTT  
CCCGCAGGGCAGCAGGAGCGTGGATATCTGGCTCCGAGTGCTTGCCAAGCCCCAGAA  
CACAGCTGAGGTTCAAGAAGGTCCAGCTCACTGGAGAGCCAGTGCCCATGGCCCGCTG  
CGTCTCCACAGGGGGTTCGCCCCGCCAGCCCAAATCACCTGGCACTCAGACCTGGGCGG  
GATGCCCAATACGAGCCAGGTGCCAGGGTTCCTGTCTGGCACAGTCACTGTCACCAG  
CCTCTGGATATTGGTGCCCTCAAGCCAGGTGGACGGCAAGAATGTGACCTGCAAGGT  
GGAGCACGAGAGCTTTGAGAAGCCTCAGCTGCTGACTGTGAACCTCACCGTGTACTA  
CCCCCAGAGGTATCCATCTCTGGCTATGATAACAACCTGGTACCTTGGCCAGAATGA  
GGCCACCCTGACCTGCGATGCTCGCAGCAACCCAGAGCCCACAGGCTATAATTGGAG

CACGACCATGGGTCCCTGACCTTTGCTGTGGCCAGGCGCCAGGCGCTCGAT  
CCGTCCTGTGGACAAACCAATCAACACAACCTTTAATCTGCAACGTCACCAATGCCCT  
AGGAGCTCGCCAGGCAGAACTGACCGTCCAGGTCAAAGAGGGACCTCCCAGTGAGCA  
CTCAGGCATGTCCCGTAACGCCATCATCTTCCTGGTTCTGGGAATCCTGGTTTTTCT  
GATCCTGCTGGGGATCGGGATTTATTTCTATTGGTCCAAATGTTCCCGTGAGGTCCT  
TTGGCACTGTCATCTGTGTCCCTCGAGTACAGAGCATGCCAGCGCCTCAGCTAATGG  
GCATGTCTCCTATTCAGCTGTGAGCAGAGAGAACAGCTCTTCCCAGGATCCACAGAC  
AGAGGGCACAAGGTGACAGCGTCGGGACTGAGAGGGGAGAGAGACTGGAGCTGGCAA  
GGACGTGGGCCTCCAGAGTTGGACCCGACCCCAATGGATGAAGACCCCCTCCAAAGA  
GACCAGCCTCCCTCCCTGTGCCAGACCTCAAAACGACGGGGGCAGGTGCAAGTTCAT  
AGGTCTCCAAGACCACCCTCCTTTTCATTTGCTAGAAGGACTCACTAGACTCAGGAAA  
GCTGTTAGGCTCACAGTTACAGTTTATTACAGTAAAAGGACAGAGATTAAGATCAGC  
AAAGGGAGGAGGTGCACAGCACACGTTCCACGACAGATGAGGCGACGGCTTCCATCT  
GCCCTCTCCCAGTGGAGCCATATAGGCAGCACCTGATTCTCACAGCAACATGTGACA  
ACATGCAAGAAGTACTGCCAATACTGCCAACCAGAGCAGCTCACTCGAGATCTTTGT  
GTCCAGAGTTTTTTTGTGTTGTCTTGAGACAGGGTCTGGCTCTGTTGGCAGACTAGAGT  
ACAGTGGTGAGATCACAGTTCATTGCAGCCTTGACTTCTCAACGCCAAGTCATCCTC  
CCACCTCAGCCTCCTGAGTAGCTATGACTACAGGTATGTGCCACCACGTCTGGCTAA  
TCTTTTTATTATTTGTAAAGTCGAGGTTTCCCTGTGTTGCCCAGGCTGGTCTTGAAC  
TCTTGGCTCCAAGTGATACTTCTGCCTTGGCCTCCCAAAGTGCTGAATTAAGCAGCT  
CACCATCCACACGGCTGACCTCATACATCAAGCCAATACCGTGTGGCCCAAGACCCC  
CACCATAAATCACATCATTAGCATGAACCACCCAGAGTGGCCCAAGACTCCAAGATC  
AGCTACCAGGCAGGATATTCCAAGGGCTTAGAGATGAATGCCCAGGAGCTGAGGATA  
AAGGGCCCGATCTTTCTTTGGGCAAGGTTAAGCCTTTACTGCATAGCAGACCACACA  
GAAGGGTGTGGGCCACCAGAGAATTTTGGTAAAAAATTTGGCCTCTGGCCTTGAGCTT  
CTAAATCTCTGTATCCGTCAGATCTCTGTGGTTACAAGAAACAGCCACTGACCCTGG  
TCACCAGAGGCTGCAATTCAGGCCGCAAGCAGCTGCCTGGGGGGTGTCCAAGGAGCA  
GAGAAAATACTAGATGTGAACCTTGAAGAAGGTTGTCAGCTGCAGCCACTTTCTGCC  
AGCATCTGCAGCCACTTTCTGCCAGCATCTGCAGCCAGCAAGCTGGGACTGGCAGGA  
AATAACCCACAAAAGAAGCAAATGCAATTTCCAACACAAGGGGGAAGGGATGCAGGG  
GGAGGCAGCGCTGCAGTTGCTCAGGACACGCTCCTATAGGACCAAGATGGATGCGAC  
CCAAGACCCAGGAGGCCCAGCTGCTCAGTGCAACTGACAAGTTAAAAAAGGTCTATGA  
TCTTGAGGGCAGACAGCAGAATTCCTCTTATAAAGAAAACTGTTTGGGAAAATACGT  
TGAGGGAGAGAAGACCTTGGGCCAAGATGCTAAATGGGAATGCAAAGCTTGAGCTGC  
TCTGCAAGAGAAAATAAGCAGGACAGAGGATTTGCTCTGGACAGAGATGGAAGAGCC  
GGGAACAGAGAAGTGTGGGGAAGAGATAGGAACCAGCAGGATGGCAGGGGGCAAAGGG  
CTCAAGGGTGAGGAGGCCAGTGGGACCCACAGAGTTGGGGAGATAAAGGAACATTG  
GTTGCTTTGGTGGCACGTAAGCTCCTTGTCTGTCTCCAGCACCCAGAATCTCATTA  
AGCTTATTTATTGTACCTCCAGCGGCTGTGTGCAATGGGGTCTTTTGTGGAAATCAA  
GGAGCAGACAGGTTTCATGTGTACTGTCACCACGTGGGATGGAACCAGAGGCATGGA  
AGCAAGACGCTAAATGAAGAGGGCCATAAGGGCTGGGATTCCCAGGCACCTTAGGAA  
CAGCTTGTCTTTTTTTTTTCTCTCCAAAAAAAATGTTTAAAGGGACGGTGTCTCCT  
GTCACCCAGGCTGGAGTGCAATGGCACGATCATAGCTCATTGCAGCCTCTAACTCCG  
GGGCTCAAGCAATCCTCCCACCTCAGCCTACCAAGTAGCTGTGACCACAGCTGCCCC  
TCACCATGCTAAGCTAATTTTTTTAATTAGATAGTACATAAACGTCCCAAAATTAGA  
AGATAAAAAGACATGAGGGATCCATTCTAATTTGTGTTTGGAGTGTAATGGTCCAGC  
TCCATTCTTCTGCACATGGATATCCAGTTTTTACACAACACTGTGAATGTAATGAATG  
CCACTGAATCATACACTCAAAAATAGCTAAAATGGCAAATTGTCTGTTATCTCTTTT  
TAACCACCATTTTTTGAAAATTAATTATACCAAAAAACCATTGAATAGTGCACCTTAT  
TTATTTATTTATTTGTTTATTTATTTATTTATTTTAGAAATAAGAGTCTCACTTTGT

>NP\_006496.4 poliovirus receptor isoform alpha precursor [*Homo sapiens*]  
MARAMAAAWPLLLVALLVLSWPPPGTGDVVVQAPTQVPGFLGDSVTLPCYLQVPNME  
VTHVSQLTWARHGESGSMVVFHQQTQGPSYSESKRLEFVAARLGAELRNASLRMFGLR  
VEDEGNYTCLFVTFTPQGSRSVDIWLRLVLAKPQNTAEVQKVQLTGEPVPMARCVSTGG  
RPPAQITWHSDLGGMPNTSQVPGFLSGTVTVTSLWILVPSSQVDGKNVTCKVEHESF  
EKPQLLTVNLTVYYPPPEVSISGYDNNWYLGQNEATLTCDARSNPEPTGYNWSTTMGP  
LPPFAVAQGAQLLIRPVDKPINTTLCINVTNALGARQAELTVQVKEGPPSEHSGMSR  
NAIIFLVLGILVFLILLGIGIYFYWSKCSREVLWHCHLCPSTEHASASANGHVSYS  
AVSRENSSSQDPQTEGTR (SEQ ID NO: 34) Mouse PVR >NM\_027514.2 *Mus*  
*musculus* poliovirus receptor (Pvr), mRNA (CD155)



GGTTGTCTTCCACCCCAAGGAGAGGGGACCAACATCAAAGAGCCAGAGAGGGTGA AATT  
CTTGGCTGCCCCAACAGGATCTGAGGAACGCATCTCTGGCCATCTCGAACTTAAGTGT  
AGAAGACGAAGGCATCTATGAATGTCAGATTGCCACATTCCCCAGAGGCAGTAGAAG  
CACCAATGCCTGGCTGAAGGTGCAAGCCCGACCTAAGAACACTGCAGAGGCCCTGGA  
GCCCTCTCCCACCTTGATACTGCAGGATGTGGCTAAATGCATCTCTGCCAATGGTCA  
CCCTCCTGGACGAATCTCTTGGCCCTCGAATGTGAATGGAAGTCACCGTGAAATGAA  
GGAACCAGGGTCCCAGCCGGGCACCACCACAGTTACCAGCTACCTCTCCATGGTACC  
TTCTCGCCAGGCAGACGGCAAGAACATCACCTGCACGGTGGAGCATGAAAGCTTACA  
GGAGCTGGACCAGCTGCTGGTGACCCTTTCCCAACCCTATCCACCTGAAAACGTGTC  
CATCTCTGGCTATGACGGCAACTGGTATGTTGGCCTCACTAACTTGACCCTGACCCTG  
TGAAGCTCACAGCAAACCAGCGCCTGACATGGCTGGATATAACTGGAGCACGAACAC  
GGGTGACTTTCCCAACTCTGTTAAGCGCCAGGGCAATATGCTTCTAATCTCCACCGT  
AGAGGATGGTCTCAATAACACGGTCATTGTGTGCGAAGTCACCAATGCCCTAGGGTC  
TGGGCAGGGCCAAGTGCACATCATTGTTAAAGAGAAACCTGAGAATATGCAGCAAAA  
TACAAGATTACACCTAGGCTACATCTTTCTTATCGTCTTTGTCTCTCGCTGTAGTCAT  
CATCATCGCAGCACTATACACTATACGAAGATGCAGGCATGGTCGTGCTCTGCAGTC  
CAATCCCTCAGAGAGGGGAGAACGTCCAGTATTCATCTGTGAACGGCGACTGTAGACT  
GAACATGGAGCCAAACAGCACAAAGGTGACGGTGCTGGGTAGACAGAACTAAGGAACT  
TGAAGGCATAGCAACTGGAACCCTACTCTCATAAATGAAGAAGCCTCCAGAGAGACT  
GGCTGCTCAGTGTGATGAGCATAGCAAGTTTGGGGGGTCTCCCAGGATGCTGCCGAA  
TTCCACGTTGTCAAAAGGACCCATGGAGGCCAGTGTGTTGGCTCACTCTTGACATCT  
CAGCAAGCTGGGGGGGGGGGGGGGAGCATAAAGCAAGGTTGAGTCTAGCTTTGGGCTA  
TAGAGCAAAGCCCTGTCCATACACAAACAAGCTAAGGGGGCTTTGAGACGGTCAGAAA  
CTGAAGTCTTGCTTTGGGTAAAGTAAATCCTCTACCGCATGTATGTGCTAGACTTGA  
AAGACTTCCACACAGACCTCTTTATAAGTTGACTCCATTGGGGCTATCCCCTCCTCT  
CTGGACAAGGTCTCTGTATGTAGCCAAGGCTAGGCTCAAACCTCACAGAGATATGTCT  
GCTTCTACCTCCCCAGTGCTAGAGTTGAAAGTATTTGTGCCACTGCACTTTTCTAGG  
TCTTCTTTTAATGAAGTAAAGTATATATTTATAAAAAGCTATTTAGTTATATATATA  
TATATTTTGTAGACTATTTTCATAGAGCCCAAGCTAACCTCAAACCTTACTATGTAGCC  
AAGAGTGATGGTAAACTAATTTATTTTAATTTATTTGTCTTCAATTTTAACCATCAC  
CCAACCCCTGCTCCCTTCCATATCTTCTTTCAATCCATTTCATTGTCTTTTTTCTTCC  
CAGACACTATTCTGACTTACGTCTCCATTACAAACATTTTATTGAACTACATAAAAA  
TGTGTGAACCACAAAAAAAATGTATTTGTCAAAATTGTAGTTGTCTTTCTGAGGC  
TGACCTGAGTTCTCTGATACCATTCTCTCCAGTTGTATCCAGTTTCCTGTAAACAAT  
GTGACTTTGTTTTTCTCAGTAGCTAAACATCCCAATTATGTGAGTGTACACTTTCT  
TACTCATTCCTCTGTGGGCCACCAGCTGGGTTGGTTCCATATCTGAGCTATTGTGC  
ATGGAATTGTCTCTGTGGTGGGTTTAGTAAACTCCCAGGAATGCCTGTACATGTTTG  
TAGAGGCCAGAAGAAGGCACAAAATCTTGAGCCAGGCTTACATGCACTTGTGAGTAG  
CCCCACATAGGTGCTAAGAACCCAGTTCAGGTCCTCTGCTGTGGGATGGTGGGCTGT  
GCACAGAAAGCCTGGTCCCGGTCTAGCAAAGGTCTGGAACCTCCGGAGCCGGTGGGCT  
GTGATTTACACCAGCATGGGATGGAAGGAGTTGGACCTCGCCTCCTGGGCACCTGGC  
TCCTGTCACATAGCTACAGCCTCCCACAGCCCCCCTATAGGGAGGTATGCAGCATCA  
ATCACATAGTAGCTGCACTAAGCCCTCCCACATGCAAATAAGGTTTCCCCAAACTCT  
CAGTCCAAGCCAATGAAAAGTACCTGCTGTCAAACCCTAAATCATCCCCAAACTCT  
GTAAGTCCTATCAGGGAATAAAATGTGTGTGAAAACCTAAAAAAAAAAAAAAAAA (SEQ  
ID NO: 35)>NP\_081790.1 poliovirus receptor precursor [*Mus musculus*]  
MAQLARATRSPLSWLLLLFCYALRKAGGDIRVLVPYNSTGVLGGSTTLHCSLTSNEN  
VTITQITWMKKDSGGSHALVAVFHPKKGPNIKEPERVKFLAAQQDLRNASLAISNLS  
VEDEGIYECQIATFPRGSRSTNAWLKVQARPKNTAEALEPSPTLILQDVAKCISANG  
HPPGRISWPSNVNGSHREMKEPGSQPGTTTTVTSYLSMVPSPROADGKNITCTVEHESL

QELDQLVLTLSQPYPPENVSIGYDGNWYVGLTNLTLTCEAHSKPAPDMAGYNWSTN  
TGDFPNSVKRQGNMLLISTVEDGLNNTVIVCEVTNALGSGQGQVHIIVKEKPENMQQ  
NTRLHLGYIFLIVFVLAVVIII AALYTIRRCRHGRALQSNP SERENVQYSSVNGDCR  
LNMEPNSTR (SEQ ID NO: 36) Human >NM\_002856.3 *Homo sapiens* nectin  
cell adhesion molecule 2 Nectin-2 (NECTIN2), transcript variant alpha, mRNA  
(CD112)

GTGACGTCAGCGGGTTCGAACCGCCGGAGCTGAGCGAGAGGCCGGGGGTGCCGAGCC  
isoform alpha  
GGGCGGGGAGAGCTGGGCCGGGAGAGCAGAACAGGGAGGCTAGAGCGCAGCGGGAAC  
CGGCCCGGAGCCGGAGCCGGAGCCCCACAGGCACCTACTAAACCGCCCAGCCGATCG  
GCCCCACAGAGTGGCCCGCGGGCCTCCGGCCGGGGCCAGTCCCCTCCCGGGCCCTC  
CATGGCCCGGGCCGCTGCCCTCCTGCCGTCGAGATCGCCGCCGACGCCGCTGCTGTG  
GCCGCTGCTGCTGCTGCTCCTGGAAACCGGAGCCCAGGATGTGCGAGTTCAAGT  
GCTACCCGAGGTGCGAGGCCAGCTCGGGGGCACCGTGGAGCTGCCGTGCCACCTGCT  
GCCACCTGTTCTTGACTGTACATCTCCCTGGTGACCTGGCAGCGCCCAGATGCACC  
TGCGAACCACCAGAATGTGGCCGCCTTCCACCCTAAGATGGGTCCCAGCTTCCCCAG  
CCCGAAGCCTGGCAGCGAGCGGCTGTCCTTCGTCTCTGCCAAGCAGAGCACTGGGCA  
AGACACAGAGGCAGAGCTCCAGGACGCCACGCTGGCCCTCCACGGGCTCACGGTGGA  
GGACGAGGGCAACTACACTTGCGAGTTTGCCACCTTCCCCAAGGGGTCCGTCCGAGG  
GATGACCTGGCTCAGAGTCATAGCCAAGCCCAAGAACCAAGCTGAGGCCCAGAAGGT  
CACGTTTCAGCCAGGACCCTACGACAGTGGCCCTCTGCATCTCCAAAGAGGGCCGCCC  
ACCTGCCCCGGATCTCCTGGCTCTCATCCCTGGACTGGGAAGCCAAAGAGACTCAGGT  
GTCAGGGACCCTGGCCGGAAGTGTCACTGTCACCAGCCGCTTCACCTTGGTGCCCTC  
GGGCCGAGCAGATGGTGTCACGGTCACCTGCAAAGTGGAGCATGAGAGCTTCGAGGA  
ACCAGCCCTGATACCTGTGACCCTCTCTGTACGCTACCCTCCTGAAGTGTCCATCTC  
CGGCTATGATGACAACTGGTACCTCGGGCGTACTGATGCCACCCTGAGCTGTGACGT  
CCGCAGCAACCCAGAGCCCACGGGGCTATGACTGGAGCACGACCTCAGGCACCTTCCC  
GACCTCCGCAGTGGCCCCAGGGCTCCCAGCTGGTCATCCACGCAGTGGACAGTCTGTT  
CAATACCACCTTCGTCTGCACAGTCACCAATGCCGTGGGCATGGGCCGCGCTGAGCA  
GGTCATCTTTGTCCGAGAAACCCCCAGGGCCTCGCCCCGAGATGTGGGGCCCGCTGGT  
GTGGGGGGCCGTGGGGGGGACACTGCTGGTGCTGCTGCTTCTGGCTGGGGGGTCCTT  
GGCCTTCATCCTGCTGAGGGTGAGGAGGAGGAGGAAGAGCCCTGGAGGAGCAGGAGG  
AGGAGCCAGTGGCGACGGGGGATTCTACGATCCGAAAGCTCAGGTGTTGGGAAATGG  
GGACCCCGTCTTCTGGACACCAGTAGTCCCTGGTCCCATGGAACCAGATGGCAAGGA  
TGAGGAGGAGGAGGAGGAGGAAGAGAAGGCAGAGAAAGGCCTCATGTTGCCTCCACC  
CCCAGCACTCGAGGATGACATGGAGTCCCAGCTGGACGGCTCCCTCATCTCACGGCG  
GGCAGTTTATGTGTGACCTGGACACAGACAGAGACAGAGCCAGGCCCCGGCCCTCCCG  
CCCCGACCTGACCACGCCGGCCTAGGGTTCCAGACTGGTTGGACTTGTTTCGTCTGG  
ACGACACTGGAGTGGAACACTGCCTCCCCTTTCTTGGGACTTGGAGGGAGGTGGAA  
CAGCACACTGGACTTCTCCCGTCTCTAGGGCTGCATGGGGAGCCCCGGGGAGCTGAGT  
AGTGGGGATCCAGAGAGGACCCCCGCCCCCAGAGACTTGGTTTTGGCTCCAGCCTTC  
CCCTGGCCCCGTGACACTCAGGAGTTAATAAATGCCTTGGAGGAAAACA (SEQ ID  
NO: 37) >NP\_002847.1 nectin-2 isoform alpha precursor [*Homo sapiens*]  
MARAAALLPSRSPPTPLLWPLLLLLLLETGAQDVRVQVLPEVRGQLGGTVELPCHLL  
PPVPGLYISLVTWQRPDAPANHQNVAAAFHPKMGPSFPSPKPGSERLSFVSAKQSTGQ  
DTEAELQDATLALHGLTVEDEGNYTCEFATFPKGSVRGMTWLRVIAKPKNQAEAKV  
TFSQDPTTVALCISKEGRPPARISWLSSLDWEAKETQVSGTLAGTVTVTSRFTLVPS  
GRADGVTVTCKVEHESFEEPALIPVTLSVRYPPPEVSISGYDDNWYLGRTDATALSCDV  
RSNPEPTGYDWSTTSGTFPTSAVAQGSQVLVIHAVDSLNTTFVCTVTNAVGMGRAEQ  
VIFVRETPRASPRDVGPLVWGAVGGTLLVLLLLAGGSLAFILLRVRRRRKSPGGAGG

GASGDYFNGDPVFWFVTPVPGPMEPDGKDEEEEEEEKAEKGLMLPPP  
PALEDDMESQLDGSLISRRVYV (SEQ ID NO: 38) Mouse Nectin-  
>NM\_001159724.1 *Mus musculus* nectin cell adhesion 2 (CD112) molecule 2  
(Nectin2), transcript variant 2, mRNA isoform alpha  
GAGCCCTAGGATCGGCTTGGCGAAGAGGGGCGGGGCCTGTGACGTCATGAGTCCGGC  
CCGCTGGAGCTAAGCGAGGGGCGGGGGGCGCGGATCCTGAGAGCCAGGCGAGGGAA  
AGCTGGGCCGAACGAACCTGATCCGGGGAGCCGTGAGCGGCGGAAGCCGGCCTGGAGC  
CGGACACTTCAGACCCCTGACTGCCCTCCCAGCCGATCGGTACACGAAGAGTGGTCC  
CTAGGCACCCCTGCCCCGGGCCCAGTCCCTCCCCGGGCCCCCATGGCCCCGGGCCG  
AGTCCTCCCGCCGTCCAGATTGTCACCGACGCTGCCGTTGTTGCCGCTGCTACTGCT  
CCTGCTTCAGGAAACAGGAGCCCAAGATGTGCGGGTACGAGTGCTTCCCGAGGTCCG  
GGGCCGCTTGGGAGGCACCGTGGAGTTACCGTGCCACCTGCTCCCACCCACGACGGA  
GCGCGTCTCTCAGGTGACCTGGCAGCGCCTGGATGGCACAGTTGTGGCTGCTTTCCA  
CCCATCCTTCGGAGTGGATTTCCCCAACTCTCAGTTCAGCAAGGACCGTCTGTCCTT  
TGTCAGAGCGAGACCAGAAACAAACGCAGACCTGCGGGATGCCACACTGGCCTTCCG  
GGGACTGAGGGTAGAGGACGAGGGCAATTACACCTGCGAGTTTGCCACGTTTCCCAA  
CGGTACCCGCGAGGGGGGTGACCTGGCTCAGAGTCATAGCCCAGCCTGAGAACCACGC  
TGAAGCCCAGGAGGTCACAATTGGCCCCCAGTCGGTGGCTGTAGCCCGCTGTGTCTC  
CACTGGGGGGCCGCCCCCTGCCGAATCACCTGGATCTCATCTCTGGGTGGAGAGGC  
CAAAGATACTCAGGAGCCAGGGATACAGGCTGGCACCGTCACTATCATCAGCCGATA  
CTCCTTGGTGCCCCGTGGGCCGAGCGGATGGCGTCAAGGTCACGTGTAGAGTGGAACA  
CGAGAGCTTCGAAGAGCCGATCCTGCTGCCAGTGACCCTCTCTGTGCGCTACCCTCC  
AGAAGTATCCATCTCCGGCTATGATGACAACTGGTACCTTGCCCGCAGTGAGGCCAT  
ACTGACCTGTGATGTACGAAGCAACCCAGAGCCCACAGACTATGACTGGAGCACGAC  
CTCGGGCGTCTTCCCAGCCTCTGCAGTGGCCCAGGGCTCTCAGCTGCTTGTCCACTC  
TGTGGATCGAATGGTCAACACTACCTTCATCTGTACAGCCACCAACGCTGTGGGGAC  
AGGCCGTGCTGAGCAGGTCATCCTGGTGCGAGACACCCCCCAGGCCTCCCGAGATGT  
GGGTCCGCTGGTGTGGGGGGCCGTGGGGGGGAACATTGCTGGTGCTACTCCTGGCTGG  
GGGGTTCTTGGCCTTGATCCTGCTGAGGGGGAGGAGGAGGCGGAAGAGCCCTGGAGG  
AGGAGGAAATGATGGCGACAGAGGATCCTACGATCCAAAGACTCAGGTGTTTGGGAA  
CGGGGGTCTGTCTTCTGGAGGTCAGCATCCCCTGAGCCCATGAGGCCAGATGGCAG  
GGAGGAAGATGAGGAGGAGGAGGAAGAAATGAAGGCAGAGGAAGGTCTCATGCTACC  
TCCACACGAGTCACCTAAGGACGACATGGAGTCCCATCTGGATGGCTCCCTCATCTC  
TCGGCGGGCAGTTTACGTGTGACCCTACGATATAGACACTGGACACATGGAAACACC  
AAGTTCCACCCTCACTGCCAACCACACCAATGCCAGCCAGCAACGATGGCTAGGGAC  
CGGTTGGACTGGTTCTTCTGGGGCACACTGGAGTTGGAAGGGCACCGCCCCTGCTTT  
CAGGATAGAGGACAAGTGGAACCACACAGACTCCTATCTTTAGGGCCTCATGGAGTA  
GGGGACCCAGGAGCGCCATGGTGCACACTCAGGACTCCTCAGAGCTTGCTTTCGGC  
CCCAGCCTAGCCCTGGCCCCGAAACACTCAGGAGCTAATAAATGCCTTGTTCGAAAA  
AAAAAAAAAAAAAAAA (SEQ ID NO: 39)>NP\_001153196.1 nectin-2 isoform 2  
precursor [*Mus musculus*]  
MARAAVLPPSRLSPTLPLLPLLLLLLQETGAQDVRVRVLPEVRGRLGGTVELPCHLL  
PPTTERVSQVTWQRLDGTVVAAFHPSFGVDFPNSQFSKDRLSFVRARPETNADLRDA  
TLAFRGLRVEDEGNYTCEFATFPNGTRRGVTWLRVIAQPENHAEAEQVTIGPQSVAV  
ARCVSTGGRPPARITWISSLGGEAKDTQEPGIQAGTVTIISRYSLVPVGRADGVKVT  
CRVEHESFEPIILLPVTLSVRYPPEVSISGYDDNWYLGRSEAILTCDVRSNPEPTDY  
DWSTTSGVFPASAVAQGSQLLVHSVDRMVNTTFICTATNAVGTGRAEQVILVRDTPQ  
ASRDVGPLVWGAVGGTLLVLLLAGGFLALILLRGRRRRKSPGGGGNDGDRGSYDPKT  
QVFGNGGPVFWRSASPEPMRPDGREDEEEEEEMKAEGLMLPPHESPKDDMESHLD  
GSLISRRVYV (SEQ ID NO: 40) Human >NM\_001042724.2 *Homo sapiens*

nectin cell adhesion Nectin-2 molecule 2 (NECTIN2), transcript variant delta, mRNA (CD112)

GTGACGTCAGCGGGTTCGAACCGCCGGAGCTGAGCGAGAGGCCGGGGGTGCCGAGCC

isoform delta

GGGCGGGGAGAGCTGGGCCGGGAGAGCAGAACAGGGAGGCTAGAGCGCAGCGGGAAC  
CGGCCCCGGAGCCGGAGCCGGAGCCCCACAGGCACCTACTAAACCGCCCAGCCGATCG  
GCCCCCACAGAGTGGCCCCGCGGGCCTCCGGCCGGGCCCAGTCCCCTCCCGGGGCCCTC  
CATGGCCCCGGGCCGCTGCCCTCCTGCCGTCGAGATCGCCGCCGACGCCGCTGCTGTG  
GCCGCTGCTGCTGCTGCTGCTCCTGGAAACCGGAGCCCAGGATGTGCGAGTTCAAGT  
GCTACCCGAGGTGCGAGGCCAGCTCGGGGGCACCGTGGAGCTGCCGTGCCACCTGCT  
GCCACCTGTTCTCTGGACTGTACATCTCCCTGGTGACCTGGCAGCGCCCAGATGCACC  
TGCGAACCACCAGAATGTGGCCGCCTTCCACCCTAAGATGGGTCCCAGCTTCCCCAG  
CCCGAAGCCTGGCAGCGAGCGGCTGTCTTTCGTCTCTGCCAAGCAGAGCACTGGGCA  
AGACACAGAGGCAGAGCTCCAGGACGCCACGCTGGCCCTCCACGGGCTCACGGTGGA  
GGACGAGGGCAACTACACTTGCAGAGTTTGCCACCTTCCCCAAGGGGTCCGTCCGAGG  
GATGACCTGGCTCAGAGTCATAGCCAAGCCCAAGAACCAAGCTGAGGCCCAGAAGGT  
CACGTTTCAGCCAGGACCCTACGACAGTGGCCCTCTGCATCTCCAAGAGGGGCCGCCC  
ACCTGCCCCGGATCTCCTGGCTCTCATCCCTGGACTGGGAAGCCAAAGAGACTCAGGT  
GTCAGGGACCCTGGCCGGAAGTGTACTGTCAACAGCCGCTTCACCTTGGTGCCCTC  
GGGCCGAGCAGATGGTGTACCGTCACTGCAAAGTGGAGCATGAGAGCTTCGAGGA  
ACCAGCCCTGATACCTGTGACCCTCTCTGTACGCTACCCTCCTGAAGTGTCCATCTC  
CGGCTATGATGACAACTGGTACCTCGGCCGTACTGATGCCACCCTGAGCTGTGACGT  
CCGCAGCAACCCAGAGCCCACGGGCTATGACTGGAGCACGACCTCAGGCACCTTCCC  
GACCTCCGCAGTGGCCCAGGGCTCCCAGCTGGTCATCCACGCAGTGGACAGTCTGTT  
CAATACCACCTTCGTCTGCACAGTCACCAATGCCGTGGGCATGGGCCGCGCTGAGCA  
GGTCATCTTTGTCCGAGAGACCCCCAACACAGCAGGCGCAGGGGGCCACAGGCGGCAT  
CATCGGGGGGCATCATCGCCGCCATCATTGCTACTGCTGTGGCTGCCACGGGCATCCT  
TATCTGCCGGCAGCAGCGGAAGGAGCAGACGCTGCAGGGGGCAGAGGAGGACGAAGA  
CCTGGAGGGACCTCCCTCCTACAAGCCACCGACCCCCAAAAGCGAAGCTGGAGGCACA  
GGAGATGCCCTCCCAGCTCTTCACTCTGGGGGGCCTCGGAGCACAGCCCCTCAAGAC  
CCCCTACTTTGATGCTGGCGCCTCATGCACTGAGCAGGAAATGCCTCGATACCATGA  
GCTGCCCACCTTGGAAGAACGGTCAGGACCCTTGACCCCTGGAGCCACAAGCCTGGG  
GTCCCCCATCCCGGTGCCTCCAGGGCCACCTGCTGTGGAAGACGTTTCCCTGGATCT  
AGAGGATGAGGAGGGGGAGGAGGAGGAAGAGTATCTGGACAAGATCAACCCCATCTA  
TGATGCTCTGTCTATAGCAGCCCCTCTGATTCCTACCAGGGCAAAGGCTTTGTCAT  
GTCCCGGGGCCATGTATGTGTGAGCTGCCATGCGCCTGGCGTCTCACATCTCACCTGT  
TGATCCCTTAGCTTTTCTTGCCAAGGATCTAGTGCCCCCTGACCTCTGGCCAGGCCAC  
TGTCAGTTAACACATATGCATTCCATTTGTGATGTCTACCTTGGTGGCTCCACTATG  
ACCCCTAACCCATGAGCCCAGAGAAATTCACCGTGATAATGGAATCCTGGCAACCTT  
ATCTCATGAGGCAGGAGGTGGGGAAGGTGCTTCTGCACAACCTCTGATCCCAAGGAC  
TCCTCTCCCAGACTGTGACCTTAGACCATACTCTACCCCCCAATGCCTCGACTCC  
CCCAAAATCACAAAGAAGACCCTAGACCTATAATTTGTCTTCAGGTAGTAAATTCCC  
AATAGGTCTGCTGGAGTGGGCGCTGAGGGCTCCCTGCTGCTCAGACCTGAGCCCTCC  
AGGCAGCAGGGTCCCCTTACCCCCCTCCCCACCCTGTTCCCCAAAGGTGGGAAAGAG  
GGGATTCCCCAGCCCAAGGCAGGGTTTTCCCAGCACCCCTCCTGTAAGCAGAAGTCTC  
AGGGTCCAGACCCTTCCCTGAGCCCCCACCCTTCCCTGCTACCAAGCA  
AGCAGCCCCAGCCTAGGGTCAGACAGGGTGAGCCTCATACTGTGCCTTGATGG  
CCCCAGCCTTGGGAGAAGAATTTACTGTTAACCTGGAAGACTACTGAATCATTTTAC  
CCTTGCCCAGTGGAATAGGACCTAAACATCCCCCTTCCGGGGAAAGTGGGTCATCTG  
AATTGGGGGTAGCAATTGATACTGTTTTGTAAACTACATTTCTACAAAATATGAAT

TTATACTTTGA (SEQ ID NO: 41) >NP\_001036189.1 nectin-2 isoform delta precursor [*Homo sapiens*]

MARAAALLPSRSPPTPLWPLLLLLLLETGAQDVRVQVLPEVRGQLGGTVELPCHLL  
PPVPGLYISLVTWQRPDAPANHQNVAAAFHPKMGPSFSPKPGSERLSFVSAKQSTGQ  
DTEAELQDATLALHGLTVEDEGNYTCEFATFPKGSVRGMTWLRVIAKPKNQAEAQKV  
TFSQDPTTVALCISKEGRPPARISWLSSLDWEAKETQVSGTLAGTVTVTSRFTLVPS  
GRADGVTVTCKVEHESFEEPALIPVTL SVRYPPEVSISGYDDNWYLGRTDRTLSCDV  
RSNPEPTGYDWSTTSGTFPTSAVAQGSQ LVIHAVDSL FNTTFVCTVTNAVGMGRAEQ  
VIFVRETPNTAGAGATGGIIGGIIAIIATAVAATGILICRQQRKEQTLQGAEEDED  
LEGPPSYKPPTPKAKLEAQEMPSQLFTLGASEHSPLKTPYFDAGASCTEQEMPRYHE  
LPTLEERSGPLHPGATSLGSPIPVPPGPPAVEDVSLDLEDEEGEEEEEEYLDKINPIY

DALSYSSPSDSYQGKGFVMSRAMYV (SEQ ID NO: 42) Mouse Nectin-  
>NM\_008990.3 *Mus musculus* nectin cell adhesion molecule 2 2 (CD112)  
(Nectin2), transcript variant 1, mRNA isoform beta

GAGCCCTAGGATCGGCTTGGCGAAGAGGGGCGGGGCCTGTGACGTCATGAGTCCGGC  
CCGCTGGAGCTAAGCGAGGGGCGGGGGGCGCGGATCCTGAGAGCCAGGCGAGGGAA  
AGCTGGGCCGAACGAACCTGATCCGGGGAGCCGTGAGCGGCGGAAGCCGGCCTGGAGC  
CGGACACTTCAGACCCCTGACTGCCCTCCCAGCCGATCGGTACACGAAGAGTGGTCC  
CTAGGCACCCCTGCCCCGGGCCCAGTCCCTCCCCGGGCCCCCATGGCCCCGGGCCG  
AGTCCTCCCGCCGTCCAGATTGTCACCGACGCTGCCGTTGTTGCCGCTGCTACTGCT  
CCTGCTTCAGGAAACAGGAGCCCAAGATGTGCGGGTACGAGTGCTTCCCGAGGTCCG  
GGGCCGCTTGGGAGGCACCGTGAGTTACCGTGCCACCTGCTCCCACCCACGACGGA  
GCGCGTCTCTCAGGTGACCTGGCAGCGCCTGGATGGCACAGTTGTGGCTGCTTTCCA  
CCCATCCTTCGGAGTGGATTTCCCCAACTCTCAGTTCAGCAAGGACCGTCTGTCCTT  
TGTCAGAGCGAGACCAGAAACAAACGCAGACCTGCGGGATGCCACACTGGCCTTCCG  
GGGACTGAGGGTAGAGGACGAGGGCAATTACACCTGCGAGTTTGCCACGTTTCCCAA  
CGGTACCCGCAGGGGGGTGACCTGGCTCAGAGTCATAGCCCAGCCTGAGAACCACGC  
TGAAGCCCAGGAGGTCACAATTGGCCCCCAGTCGGTGGCTGTAGCCCGCTGTGTCTC  
CACTGGGGGGCCGCCCCCTGCCGAATCACCTGGATCTCATCTCTGGGTGGAGAGGC  
CAAAGATACTCAGGAGCCAGGGATACAGGCTGGCACCGTCACTATCATCAGCCGATA  
CTCCTTGGTGCCCGTGGGCCGAGCGGATGGCGTCAAGGTCACGTGTAGAGTGGAACA  
CGAGAGCTTCGAAGAGCCGATCCTGCTGCCAGTGACCCTCTCTGTGCGCTACCTTCC  
AGAAGTATCCATCTCCGGCTATGATGACAACTGGTACCTTGGCCGCAGTGAGGCCAT  
ACTGACCTGTGATGTACGAAGCAACCCAGAGCCCACAGACTATGACTGGAGCACGAC  
CTCGGGCGTCTTCCCAGCCTCTGCAGTGGCCCAGGGCTCTCAGCTGCTTGTCCACTC  
TGTGGATCGAATGGTCAACACTACCTTCATCTGTACAGCCACCAACGCTGTGGGGAC  
AGGCCGTGCTGAGCAGGTCATCCTGGTGCGAGAGTCACCCAGCACAGCAGGAGCAGG  
GGCCACTGGTGGCATCATTGGAGGTATTATCGCTGCCATCATCGCCACCGCAGTGGC  
TGGCACAGGCATCCTCATCTGCCGACAACAGCGGAAGGAGCAGAGGCTTCAAGCTGC  
GGATGAGGAAGAAGAAGTGGAAAGGACCTCCCTCCTATAAACCACCCACCCCGAAGGC  
CAAGCTGGAGGAACCAGAGATGCCCTCTCAACTCTTACCTTGGGGGGCCTCAGAGCA  
CAGCCCAGTGAAGACGCCATACTTTGATGCTGGTGTCTCTTGTGCTGATCAGGAGAT  
GCCTCGGTATCACGAGCTGCCCACTCTGGAAGAGCGGTCAGGGCCCCTGCTGTTGGG  
GGCTACAGGCCTGGGACCTTCTCTTCTGGTGCTCCAGGACCCAATGTTGTGGAGGG  
GGTTTCCCTGAGTCTCGAAGATGAGGAGGAAGATGATGAGGAGGAAGACTTCCCTGGA  
TAAATCAACCCTATTTATGATGCCCTGTCCTACCCAGCCCCTCTGACTCCTACCA  
GAGCAAAGACTTTTTTGTGTCACGGGCCATGTATGTGTGAGGGAGGCACAGGGGCTC  
TGACGTCTCACCTTTCACCTTGACCCATGAGCTTTCACACAGTAATCTAGGACACT  
CTGACTTCCAGGCAGACCAGGGACAACCTATCACCCATTGCAATCCACCTGTGACTTC  
TTAGTGACTCCACCATGACGTCCAATCTATGATGTCTGAGGCAGGCAAACCTGCACA

ACTTGGAAATGAGGATTTTATCTTACCTTGGAGGAGCTTACCATATGCCTTATGC  
ACCACCTGTGACCCCCCCCCCCCCCAAGGACTCCTAAGACTACGACCCTTTGACC  
ATGCCACTCAGTATCTCAAGAACCCTTAAAGTCCCAAAGGAATCGGACCTTGCACTT  
GTCCTCAGGCAATAGAGTCCAACAGATATGCAAGAACGGGATCAGGGGCTCCCTGTT  
GCTCAGACCTGAGCCCTCCAGGCAGCAGAAGCTCACCTGATCCCTCCCCACCCTGCT  
CCCCAAAGGTGAAAAGGAGAGGATTCCCCAATGTAAGGTAGGACCTCCCCATCTCCA  
CCTACTCCTGCAGGCAGGAATCTCAGGTTTCTCACACCCTCTCCTCAGCACCCAGGT  
TCCTGTCTCCAGAGCATGAATTCCAGGTCCAATGCTAGAGGGGAGAACCTAATGCAA  
GTGTGCCTTGCCACCCCAAGTTTGGGAGACTCTGCTCTTATCCTGAGGACTACTGAA  
TTCTTTTAAACCCCTACCCAGTGAGATGAGAACTACATATCCCTCTTTAGGGGATGGT  
GTGTGTATGTGTGTGTGATGGAGAATCTGGGCATCTGGGTTGGGAATTTTATTTTGT  
AAGCATTTCTACATAATATGAGTTTCTACTTTGATAAAGTCTTGTGTTTTCTGTG (SEQ  
ID NO: 43) >NP\_033016.3 nectin-2 isoform 1 precursor [*Mus musculus*]  
MARAAVLPPSRLSPTLPLLPLLLLLLQETGAQDVRVRVLPEVRGRLGGTVELPCHLL  
PPTTERVSQVTWQRLDGTVVAAFHPSFGVDFPNSQFSKDRLSFVRARPETNADLRDA  
TLAFRGLRVEDEGNYTCEFATFPNGTRRGVTWLRVIAQPENHAEAEQVTIGPQSVAV  
ARCVSTGGRPPARITWISSLGGEAKDTQEPGIQAGTVTIISRYSLVPVGRADGVKVT  
CRVEHESFEEPILLPVTLSVRYPPEVSISGYDDNWYLGRSEAILTCDVRSNPEPTDY  
DWSTTSGVFPASAVAQGSQLLVHSVDRMVNTTFICTATNAVGTGRAEQVILVRES  
TAGAGATGGIIGGIIAAIIATAVAGTGILICRQQRKEQRLQAADEEEEEELEGPPSYKP  
PTPKAKLEEPEMPSQLFTLGASEHSPVKTPYFDAGVSCADQEMPRYHELPTLEERSG  
PLLLGATGLGPSLLVPPGPNVVEGVSLSLEDEEEEDDEEEDFLDKINPIYDALSYSP  
SDSYQSKDFFVSRAMYV (SEQ ID NO: 44) Human IL-10 >NM\_000572.3 *Homo  
sapiens* interleukin 10 (IL10), transcript variant 1, mRNA  
ACACATCAGGGGCTTGCTCTTGCAAAACCAAACCACAAGACAGACTTGCAAAAGAAG  
GCATGCACAGCTCAGCACTGCTCTGTTGCCTGGTCCTCCTGACTGGGGTGAGGGCCA  
GCCCAGGCCAGGGCACCCAGTCTGAGAACAGCTGCACCCACTTCCCAGGCAACCTGC  
CTAACATGCTTCGAGATCTCCGAGATGCCTTCAGCAGAGTGAAGACTTTCTTTCAA  
TGAAGGATCAGCTGGACAACCTTGTTGTTAAAGGAGTCCTTGCTGGAGGACTTTAAGG  
GTTACCTGGGTTGCCAAGCCTTGCTCTGAGATGATCCAGTTTTACCTGGAGGAGGTGA  
TGCCCCAAGCTGAGAACCAAGACCCAGACATCAAGGCGCATGTGAACTCCCTGGGGG  
AGAACCTGAAGACCCTCAGGCTGAGGCTACGGCGCTGTCATCGATTCTTCCCTGTG  
AAAACAAGAGCAAGGCCGTGGAGCAGGTGAAGAATGCCTTTAATAAGCTCCAAGAGA  
AAGGCATCTACAAAGCCATGAGTGAGTTTGACATCTTCATCAACTACATAGAAGCCT  
ACATGACAATGAAGATACGAAACTGAGACATCAGGGTGCGGACTCTATAGACTCTAG  
GACATAAATTAGAGGTCTCCAAAATCGGATCTGGGGCTCTGGGATAGCTGACCCAGC  
CCCTTGAGAAACCTTATTGTACCTCTCTTATAGAATATTTATTACCTCTGATACCTC  
AACCCCCATTTCTATTTATTTACTGAGCTTCTCTGTGAACGATTTAGAAAGAAGCCC  
AATATTATAATTTTTTTTCAATATTTATTATTTTACCTGTTTTTTAAGCTGTTTCCAT  
AGGGTGACACACTATGGTATTTGAGTGTTTTAAGATAAATTATAAGTTACATAAGGG  
AGGAAAAAAAATGTTCTTTGGGGAGCCAACAGAAGCTTCCATTCCAAGCCTGACCAC  
GCTTTCTAGCTGTTGAGCTGTTTTCCCTGACCTCCCTCTAATTTATCTTGTCTCTGG  
GCTTGGGGCTTCCTAACTGCTACAAATACTCTTAGGAAGAGAAACCAGGGAGCCCCCT  
TTGATGATTAATTCACCTTCCAGTGTCTCGGAGGGATTCCCCTAACCTCATTCCCCA  
ACCACTTCATTCTTGAAAGCTGTGGCCAGCTTGTTATTTATAACAACCTAAATTTGG  
TTCTAGGCCGGGCGCGGTGGCTCACGCCTGTAATCCCAGCACTTTGGGAGGCTGAGG  
CGGGTGATCACTTGAGGTCAGGAGTTCCTAACCAAGCCTGGTCAACATGGTGAAACC  
CCGTCTCTACTAAAAATACAAAAATTAGCCGGGCATGGTGGCGCGCACCTGTAATCC  
CAGCTACTTGGGAGGCTGAGGCAAGAGAATTGCTTGAACCCAGGAGATGGAAGTTGC  
AGTGAGCTGATATCATGCCCCTGTACTCCAGCCTGGGTGACAGAGCAAGACTCTGTC

TCAAAGAAATAAATAAATAAATTGTTCTTCAAGAACTGTTTATTAAGAA  
TTTATTCAATTCCTCTGGGAATGTTACATTGTTTGTCTGTCTTCATAGCAGATTTA  
ATTTTGAATAAATAAATGTATCTTATTCACATCA (SEQ ID NO: 45) >NP\_000563.1  
interleukin-10 isoform 1 precursor [*Homo sapiens*]

MHSSALLCCLVLLTGVRASPGQGTQSENSCTHFPGNLPNMLRDLRDAFSRVKTFFQM  
KDQLDNLLLKESLLEDFKGYLGCCQALSEMIQFYLEEVMPPQAENQDPDIKAHVNSLGE  
NLKTLRLRLRRCHRFLPCENKSKAVEQVKNAFNKLQEKGIYKAMSEFDIFINYIEAY  
MTMKIRN (SEQ ID NO: 46) Mouse IL-10 >NM\_010548.2 *Mus musculus*  
interleukin 10 (Il10), mRNA

ACATTTAGAGACTTGCTCTTGCCTACTACCAAAGCCACAAGGCAGCCTTGCAGAAAAGA  
GAGCTCCATCATGCCTGGCTCAGCACTGCTATGCTGCCTGCTCTTACTGACTGGCAT  
GAGGATCAGCAGGGGCCAGTACAGCCGGGAAGACAATACTGCACCCACTTCCCAGT  
CGGCCAGAGCCACATGCTCCTAGAGCTGCGGACTGCCTTCAGCCAGGTGAAGACTTT  
CTTTCAAACAAAGGACCAGCTGGACAACATACTGCTAACCGACTCCTTAATGCAGGA  
CTTTAAGGGTTACTTGGGTTGCCAAGCCTTATCGGAAATGATCCAGTTTTACCTGGT  
AGAAGTGATGCCCCAGGCAGAGAAGCATGGCCCAGAAATCAAGGAGCATTTGAATTC  
CCTGGGTGAGAAGCTGAAGACCCTCAGGATGCGGCTGAGGCGCTGTCATCGATTTCT  
CCCCTGTGAAAATAAGAGCAAGGCAGTGGAGCAGGTGAAGAGTGATTTTAATAAGCT  
CCAAGACCAAGGTGTCTACAAGGCCATGAATGAATTTGACATCTTCATCAACTGCAT  
AGAAGCATACATGATGATCAAAATGAAAAGCTAAAACACCTGCAGTGTGTATTGAGT  
CTGCTGGACTCCAGGACCTAGACAGAGCTCTCTAAATCTGATCCAGGGATCTTAGCT  
AACGGAAACAACCTCCTTGAAAACCTCGTTTGTACCTCTCTCCGAAATATTTATTAC  
CTCTGATACCTCAGTTCCCATTCTATTTATTCACTGAGCTTCTCTGTGAACTATTTA  
GAAAGAAGCCCAATATTATAATTTTACAGTATTTATTATTTTAACTGTGTTTAAG  
CTGTTTCCATTGGGGACACTTTATAGTATTTAAAGGGAGATTATATTATATGATGGG  
AGGGGTTCTTCCTTGGAAGCAATTGAAGCTTCTATTCTAAGGCTGGCCACACTTGA  
GAGCTGCAGGGCCCTTTGCTATGGTGTCTTTCAATTGCTCTCATCCCTGAGTTCAG  
AGCTCCTAAGAGAGTTGTGAAGAACTCATGGGTCTTGGAAGAGAAACCAGGGAGA  
TCCTTTGATGATCATTCTGCAGCAGCTCAGAGGGTTCCTACTGTCATCCCCCAG  
CCGCTTCATCCCTGAAAACCTGTGGCCAGTTTGTATTATTAACCACCTAAAATTAGT  
TCTAATAGAACTCATTTTTTAAGTAGAAGTAATGCAATTCCTCTGGGAATGGTGTATT  
GTTTGTCTGCCTTTGTAGCAGACTCTAATTTTGAATAAATGGATCTTATTCG (SEQ ID  
NO: 47) >NP\_034678.1 interleukin-10 precursor [*Mus musculus*]

MPGSALLCCLLLLTGMRISRGQYSREDNNCTHFPVQGSHMLLELRTAFSQQVKTFFQT  
KDQLDNILLTDSLMQDFKGYLGCCQALSEMIQFYLVEVMPPQAEKHGPEIKEHLNSLGE  
KLKTLRMRLRRCHRFLPCENKSKAVEQVKSDFNKLQDQGVYKAMNEFDIFINCIEAY  
MMIKMKS (SEQ ID NO: 48) Human TSG-6 >NM\_007115.3 *Homo sapiens*  
TNF alpha induced protein 6 (TNFAIP6), mRNA

AGTCACATTTTCAGCCACTGCTCTGAGAATTTGTGAGCAGCCCCTAACAGGCTGTTAC  
TTCCTACTACAAGTACGATATGATCATCTTAATTTACTTATTTCTCTTGCTATGGGAA  
GACACTCAAGGATGGGGATTCAAGGATGGAATTTTTCATAACTCCATATGGCTTGAA  
CGAGCAGCCGGTGTGTACCACAGAGAAGCACGGTCTGGCAAATACAAGCTCACCTAC  
GCAGAAGCTAAGGCGGTGTGTGAATTTGAAGGCGGCCATCTCGCAACTTACAAGCAG  
CTAGAGGCAGCCAGAAAAATTGGATTTTCATGTCTGTGCTGCTGGATGGATGGCTAAG  
GGCAGAGTTGGATACCCCATTTGTGAAGCCAGGGCCCAACTGTGGATTTGGAAAAACT  
GGCATTATTGATTATGGAATCCGTCTCAATAGGAGTGAAAGATGGGATGCCTATTGC  
TACAACCCACACGCAAAGGAGTGTGGTGGCGTCTTTACAGATCCAAAGCAAATTTTT  
AAATCTCCAGGCTTCCCAAATGAGTACGAAGATAACCAAATCTGCTACTGGCACATT  
AGACTCAAGTATGGTCAGCGTATTCACCTGAGTTTTTTTAGATTTTGACCTTGAAGAT  
GACCCAGGTTGCTTGGCTGATTATGTTGAAATATATGACAGTTACGATGATGTCCAT

GGCTTTGGTGGAGATACGTGGAGATGCTGATGCATCATCAGTACAGGA  
AATGTCATGACCTTGAAGTTTCTAAGTGATGCTTCAGTGACAGCTGGAGGTTTCCAA  
ATCAAATATGTTGCAATGGATCCTGTATCCAAATCCAGTCAAGGAAAAAATACAAGT  
ACTACTTCTACTGGAAATAAAAACTTTTTAGCTGGAAGATTTAGCCACTTATAAAAA  
AAAAAAAAAAGGATGATCAAAACACACAGTGTTTATGTTGGAATCTTTTGGAACCTCCT  
TTGATCTCACTGTTATTATTAACATTTATTTATTATTTTTCTAAATGTGAAAGCAAT  
ACATAATTTAGGGAAAATTGGAAAATATAGGAACTTTAAACGAGAAAATGAAACCT  
CTCATAATCCCACTGCATAGAAATAACAAGCGTTAACATTTTCATATTTTTTTCTTT  
CAGTCATTTTTCTATTTGTGGTATATGTATATATGTACCTATATGTATTTGCATTTG  
AAATTTTGGAATCCTGCTCTATGTACAGTTTTGTATTATACTTTTTTAAATCTTGAAC  
TTTATAAACATTTTCTGAAATCATTGATTATTCTACAAAAACATGATTTTAAACAGC  
TGTAATAATTTCTATGATATGAATGTTTTATGCATTATTTAAGCCTGTCTCTATTGT  
TGGAATTTTCAGGTCATTTTCATAAATATTGTTGCAATAAATATCCTTGAACACACAA  
AAAAAAAAAAAAAAAA (SEQ ID NO: 49)>NP\_009046.2 tumor necrosis factor-

inducible gene 6 protein precursor [*Homo sapiens*]

MIILIYFLLLWEDTQGWGFKDGFHNSIWLERAAGVYHREARSGKYKLTYAEAKAV  
CEFEGGHLATYKQLEAARKIGFHVCAAGWMAKGRVGYPIVKPGPNCGFGKTGIIDYG  
IRLNRSERWDAYCYNPHAKECGGVFTDPKQIFKSPGFPNEYEDNQICYWHIRLKYGQ  
RIHLSFLDFDLEDDPGCLADYVEIYDSYDDVHGFVGRYCGDELPPDDIISTGNVMTLK  
FLSDASVTAGGFQIKYVAMDPVSKSSQGKNTSTTSTGNKNFLAGRFSHL (SEQ ID  
NO: 50) Mouse TSG-6 >NM\_009398.2 *Mus musculus* tumor necrosis factor alpha  
induced protein 6 (Tnfaip6), mRNA

CCGCTGCTCTGAGAATTTCTGTGTGGGCAGCCCCGACATTGTAACCGGCTCTGCAACC  
GAAGAGATGGTCGTCCTCCTTTGCTTATGCGTCTTGCTGTGGGAAGAGGCTCACGGA  
TGGGGATTCAAGAACGGGATCTTTCATAACTCCATATGGCTTGAACAAGCAGCGGGC  
GTATACCACAGAGAAGCTCGGGCTGGCAGATACAAGCTCACCTACGCCGAAGCCAAG  
GCCGTATGTGAATTTGAAGGTGGTCGTCTCGCAACCTACAAGCAGCTAGAGGCAGCC  
AGAAAAATTGGATTCCATGTCTGTGCTGCTGGATGGATGGCCAAGGGTAGAGTCGGA  
TACCCCATTTGTGAAACCTGGGCCCCAACTGTGGATTTGGGAAAACGGGTATCATCGAT  
TATGGAATCCGGCTCAACAGGAGTGAGCGATGGGATGCCTATTGCTACAACCCACAT  
GCAAAGGAGTGTGGTGGTGTCTTCACAGATCCGAAGCGAATTTTTTAAATCCCCGGGC  
TTCCCAAATGAGTACGATGACAACCAGGTCTGCTACTGGCACATTCGGCTCAAGTAC  
GGTCAGCGAATTCACCTGAGCTTTTTGGACTTTGACCTTGAACATGATCCAGGCTGC  
TTGGCTGACTATGTAGAAATCTATGACAGTTATGATGACGTCCACGGCTTTGTAGGA  
AGATACTGTGGTGATGAACTTCCAGAAGACATCATTAGCACAGGAAATGTCATGACC  
TTGAAGTTTCTGAGTGATGCATCCGTCACGGCTGGAGGCTTCCAGATTAAATACGTC  
ACAGTGGATCCTGCATCTAAATCCAGTCAAGCCAAAAATACAAGTACTACTGGAAAT  
AAGAAGTTCTTACCTGGAAGGTTTAGCCATCTATAAAAAATTTTTTTTAAATGTT  
CAAAACATCCAGTACAATATTTATATTTGTTTTTGTGTTGTTGTTGTTGGTTTTTTTT  
TTTTATTTTGTTTTTGTTTTGTTTTTTTTGAGACGGGGTTTCTCTGTATAGCCTTGGCT  
GTCCTGGAACCTCACTTTGAAGACCAGGCTGGCCTCGAACTCAGAAATCCACCTGCCT  
CCGCCTACCAAGTGCTGGGATTAAAGGCGTCCACCACCACCGCCCGGCTTCAATATT  
TATATTTGTAGCTCTTGGACCTCGTTTGTCTCTTTTGTATTTTTTATTATTAACATG  
TATTTATTATTTTTCCAAATGTGAAAGCCATATGTAATTATGTGGAATTTGACAAA  
TAAATACAGAGAACTTCAAATGAGTTTTTTTTTTTTTAAATCTCATAATTGTACTACACA  
GAAATAACTAATGTTAAAGTTTTTAAATGTTTGTCTTTCATTCATTTTTCTACTTGT  
AGTATATGTACATATGTAACCTCTATGATTTGCGTTTTGAATTTTGGCATTCTGCCTTT  
TGTAACCTGATATTTTTAACCTTGACATTGTATAGCTCAAGCACTTCCCAAGATCTC  
TGAGTTTTCTACAAAATGGGACTTTGTAAATATGATTGTTCCCTGCTTTATTTAAGC  
TGAATTTATATTAGGATTTAAGGTTGTTTTTCATAAATATTGCTGTAATAAATACTTT



TGGAT (SEQ ID NO: 51) >NP\_033424.1 tumor necrosis factor-inducible gene 6  
protein precursor [*Mus musculus*]  
MVVLLCLCVLLWEEAHGWGFKNGIFHNSIWLEQAAGVYHREARAGRYKLTYAEAKAV  
CEFEGGRLATYKQLEAARKIGFHVCAAGWMAKGRVGYPIVKPGPNCGFGKTGIIDYG  
IRLNRSERWDAYCYNPHAKECGGVFTDPKRIFKSPGFPNEYDDNQVCYWHIRLKYGQ  
RIHLSFLDFDLEHDPGCLADYVEIYDSYDDVHGFVGRYCGDELPEDIISTGNVMTLK  
FLSDASVTAGGFQIKYVTVDPAKSSQAKNTSTTGNKKFLPGRFSHL (SEQ ID NO: 52)  
Human B7-H3 >NM\_001024736.2 *Homo sapiens* CD276 molecule (CD276),  
(CD276) transcript variant 1, mRNA

ATTCGGGGCCGGGCTCGCTGCGGGCGGCGACTGAGCCAGGCTGGGGCCGCGTCCCTGAG  
TCCCAGAGTCGGCGCGGGCGCGGCAGGGGCAGCCTTCCACCACGGGGAGCCCAGCTGT  
CAGCCGCCTCACAGGAAGATGCTGCGTCGGCGGGGCAGCCCTGGCATGGGTGTGCAT  
GTGGGTGCAGCCCTGGGAGCACTGTGGTTCTGCCTCACAGGAGCCCTGGAGGTCCAG  
GTCCCTGAAGACCCAGTGGTGGCACTGGTGGGCACCGATGCCACCCTGTGCTGCTCC  
TTCTCCCCTGAGCCTGGCTTCAGCCTGGCACAGCTCAACCTCATCTGGCAGCTGACA  
GATACCAAACAGCTGGTGCACAGCTTTGCTGAGGGCCAGGACCAGGGCAGCGCCTAT  
GCCAACCGCACGGCCCTCTTCCCGGACCTGCTGGCACAGGGCAACGCATCCCTGAGG  
CTGCAGCGCGTGCGTGTGGCGGACGAGGGCAGCTTCACCTGCTTCGTGAGCATCCGG  
GATTCGGCAGCGCTGCCGTCAGCCTGCAGGTGGCCGCTCCCTACTCGAAGCCCAGC  
ATGACCCTGGAGCCCAACAAGGACCTGCGGGCCAGGGGACACGGTGACCATCACGTGC  
TCCAGCTACCAGGGCTACCCTGAGGCTGAGGTGTTCTGGCAGGATGGGCAGGGTGTG  
CCCCTGACTGGCAACGTGACCACGTCGCAGATGGCCAACGAGCAGGGCTTGTTTGAT  
GTGCACAGCATCCTGCGGGTGGTGTGCTGGGTGCAAATGGCACCTACAGCTGCCTGGTG  
CGCAACCCCGTGCTGCAGCAGGATGCGCACAGCTCTGTCACCATCACACCCCAGAGA  
AGCCCCACAGGAGCCGTGGAGGTCCAGGTCCCTGAGGACCCGGTGGTGGCCCTAGTG  
GGCACCGATGCCACCCTGCGCTGCTCCTTCTCCCCCGAGCCTGGCTTCAGCCTGGCA  
CAGCTCAACCTCATCTGGCAGCTGACAGACACCAAACAGCTGGTGCACAGTTTCACC  
GAAGGCCGGGACCAGGGCAGCGCCTATGCCAACCGCACGGCCCTCTTCCCGGACCTG  
CTGGCACAAAGGCAATGCATCCCTGAGGCTGCAGCGCGTGCGTGTGGCGGACGAGGGC  
AGCTTCACCTGCTTCGTGAGCATCCGGGATTTCGGCAGCGCTGCCGTCAGCCTGCAG  
GTGGCCGCTCCCTACTCGAAGCCCAGCATGACCCTGGAGCCCAACAAGGACCTGCGG  
CCAGGGGACACGGTGACCATCACGTGCTCCAGCTACCGGGGCTACCCTGAGGCTGAG  
GTGTTCTGGCAGGATGGGCAGGGTGTGCCCTGACTGGCAACGTGACCACGTGCGAG  
ATGGCCAACGAGCAGGGCTTGTTTGATGTGCACAGCGTCCTGCGGGTGGTGTGGGT  
GCGAATGGCACCTACAGCTGCCTGGTGCGCAACCCCGTGCTGCAGCAGGATGCGCAC  
GGCTCTGTACCATCACAGGGCAGCCTATGACATTCCCCCCAGAGGCCCTGTGGGTG  
ACCGTGGGGCTGTCTGTCTGTCTCATTGCACTGCTGGTGGCCCTGGCTTTCGTGTGC  
TGGAGAAAGATCAAACAGAGCTGTGAGGAGGAGAATGCAGGAGCTGAGGACCAGGAT  
GGGGAGGGAGAAGGCTCCAAGACAGCCCTGCAGCCTCTGAAACACTCTGACAGCAAA  
GAAGATGATGGACAAGAAATAGCCTGACCATGAGGACCAGGGAGCTGCTACCCCTCC  
CTACAGCTCCTACCCTCTGGCTGCAATGGGGCTGCACTGTGAGCCCTGCCCCCAACA  
GATGCATCCTGCTCTGACAGGTGGGCTCCTTCTCCAAAGGATGCGATACACAGACCA  
CTGTGCAGCCTTATTTCTCCAATGGACATGATTCCCAAGTCATCCTGCTGCCTTTTT  
TCTTATAGACACAATGAACAGACCACCCACAACCTTAGTTCTCTAAGTCATCCTGCC  
TGCTGCCTTATTTACAGTACATACATTTCTTAGGGACACAGTACACTGACCACATC  
ACCACCCTCTTCTTCCAGTGCTGCGTGGACCATCTGGCTGCCTTTTTTCTCCAAAAG  
ATGCAATATTCAGACTGACTGACCCCTGCCTTATTTACCAAAGACACGATGCATA  
GTCACCCCGGCCTTGTTTCTCCAATGGCCGTGATACTAGTGATCATGTTCAGCCC  
TGCTTCCACCTGCATAGAATCTTTTCTTCTCAGACAGGGACAGTGCGGCCTCAACAT  
CTCCTGGAGTCTAGAAGCTGTTTCCTTTCCCTCCTTCCTCCTCTTGCTCTAGCCTT

ATAACTGGCCTTTTCCCTTCCCTTCCCTGAGGACCTGCGGCACTCTGCGCCCA  
CATGCACAGCTGTGCATGGAGACCTGCAGGTGCACGTGCTGGAACACGTGTGGTTCC  
CCCCTGGCCCAGCCTCCTCTGCAGTGCCCTCTCCCCTGCCCATCCTCCCCACGGAA  
GCATGTGCTGGTCACACTGGTTCTCCAGGGGTCTGTGATGGGGCCCCTGGGGGTGAG  
CTTCTGTCCCTCTGCCTTCTCACCTCTTTGTTCTTTCTTTTCATGTATCCATTGAG  
TTGATGTTTATTGAGCAACTACAGATGTCAGCACTGTGTTAGGTGCTGGGGGGCCCTG  
CGTGGGAAGATAAAGTTCCTCCCTCAAGGACTCCCCATCCAGCTGGGAGACAGACAA  
CTAACTACACTGCACCCTGCGGTTTGCAGGGGGGCTCCTGCCTGGCTCCCTGCTCCAC  
ACCTCCTCTGTGGCTCAAGGCTTCCTGGATACCTCACCCCCATCCCACCCATAATC  
TTACCCAGAGCATGGGGTTGGGGCGGAAACCTGGAGAGAGGGACATAGCCCCTCGCC  
ACGGCTAGAGAATCTGGTGGTGTCCAAAATGTCTGTCCAGGTGTGGGCAGGTGGGCA  
GGCACCAAGGCCCTCTGGACCTTTCATAGCAGCAGAAAAGGCAGAGCCTGGGGCAGG  
GCAGGGCCAGGAATGCTTTGGGGACACCGAGGGGACTGCCCCCACCACCACCATGG  
TGCTATTCTGGGGCTGGGGCAGTCTTTTCTGGCTTGCCCTCTGGCCAGCTCCTGGCC  
TCTGGTAGAGTGAGACTTCAGACGTTCTGATGCCTTCCGGATGTCATCTCTCCCTGC  
CCCAGGAATGGAAGATGTGAGGACTTCTAATTTAAATGTGGGACTCGGAGGGATTTT  
GTAAACTGGGGGTATATTTTGGGGAAAATAAATGTCTTTGTAAAAA (SEQ ID NO:  
53) >NP\_001019907.1 CD276 antigen isoform a precursor [*Homo sapiens*]  
MLRRRGSPGMGVHVGAAALGALWFCLTGALEVQVPEDPVVALVGTDATLCCSFSPGPG  
FSLAQLNLIWQLTDTKQLVHSFAEGQDQGSAYANRTALFPDLLAQGNASLRLQVRV  
ADEGSFTCFVSIRDFGSAAVSLQVAAPYSKPSMTLEPNKDLRPGDVTITCSSYQGY  
PEAEVFWQDQGQGVPLTGNVTTSQMANEQGLFDVHSILRVVLGANGTYSLVRNPVLQ  
QDAHSSVTITPQRSPTGAVEVQVPEDPVVALVGTDATLRCSFSPEPGFSLAQLNLIW  
QLTDTKQLVHSFTEGRDQGSAYANRTALFPDLLAQGNASLRLQVRVVADEGSFTCFV  
SIRDFGSAAVSLQVAAPYSKPSMTLEPNKDLRPGDVTITCSSYRGYPEAEVFWQDG  
QGVPLTGNVTTSQMANEQGLFDVHSVLRVVLGANGTYSLVRNPVLQQDAHGSVTIT  
GQPMTFPPEALWVTVGLSVCLIALLLVALAFVCWRKIKQSCEEENAGAEDQDGEGE  
KTALQPLKHSDSKEDDGQEIA (SEQ ID NO: 54) Mouse B7-H3 >NM\_133983.4  
*Mus musculus* CD276 antigen (Cd276), mRNA (CD276)  
CGGCGCGGCGCGCCAAAGTGACCTGGTACAGCCTGGACCCCAAGCTCATCGGCTTTG  
TCTGGCTGGCCGCTGGCCTCTTCCCCTTGGATTGGATGATCCTGAGGCCTTTGG  
AGGAACTTCGAGACAAAGGCCCTCTTCTCTTCCACGGGCAGGAGCAGCCATTGCG  
CACGGAGAGCCCAGCTGTCAGCTGTCTCACAGGAAGATGCTTCGAGGATGGGGTGGC  
CCCAGTGTGGGTGTGTGTGTGCGCACAGCACTGGGGGTGCTGTGCCTCTGCCTCACA  
GGAGCTGTGGAAGTCCAGGTCTCTGAAGACCCCGTGGTGGCCCTGGTGGACACGGAT  
GCCACCCTACGCTGCTCCTTTTCCCCAGAGCCTGGCTTCAGTCTGGCACAGCTCAAC  
CTCATCTGGCAGCTGACAGACACCAAACAGCTGGTGCACAGCTTCACGGAGGGCCGG  
GACCAAGGCAGTGCCTACTCCAACCGCACAGCGCTCTTCCCTGACCTGTTGGTGCAA  
GGCAATGCGTCCTTGAGGCTGCAGCGCGTCCGAGTAACCGACGAGGGCAGCTACACC  
TGCTTTGTGAGCATCCAGGACTTTGACAGCGCTGCTGTTAGCCTGCAGGTGGCCGCC  
CCCTACTCGAAGCCCAGCATGACCCTGGAGCCCAACAAGGACCTACGTCCAGGGAAC  
ATGGTGACCATCACGTGCTCTAGCTACCAGGGCTATCCGGAGGCCGAGGTGTTCTGG  
AAGGATGGACAGGGAGTGCCCTTGACTGGCAATGTGACCACATCCCAGATGGCCAAC  
GAGCGGGGCTTGTTTCGATGTTTACAGCGTGCTGAGGGTGGTGGTGGTGGTAAACGGC  
ACCTACAGCTGCCTGGTACGCAACCCGGTGTTGCAGCAAGATGCTCACGGCTCAGTC  
ACCATCACAGGGCAGCCCCTGACATTCCCCCCTGAGGCTCTGTGGGTAAACCGTGGGG  
CTCTCTGTCTGTCTTGTGGTACTACTGGTGGCCCTGGCTTTTCGTGTGCTGGAGAAAG  
ATCAAGCAGAGCTGCGAGGAGGAGAATGCAGGTGCCGAGGACCAGGATGGAGATGGA  
GAAGGATCCAAGACAGCTCTACGGCCTCTGAAACCCTCTGAAAACAAAGAAGATGAC  
GGACAAGAAATTGCTTGATTGGGAGCTGCTGCCCTTCCCAGGTGGGGGGCCACCCT

CTGGCAGTGTGAGCTTCAATGCCAGCCCTTCCCCCAACGAATGGGTTTGTCCCACA  
GATCTACCCGTTTCGTCAAAGGACGTGGTCCATAGACCACCCACAGCCTTACTTTTCC  
AATGGACTTAATTCCCATCATCCTGCAGCCTCATTTTCTCCAGTGACACGATACACGA  
ACCATCCTGCGGCCTTATTTCCACGGACACGACACAAAGATGTCCCTCCTCGGTGT  
TCCTCCAGAGTCGTCTGGTGGCCTTGTGATACGGCGTGAACCTTCTTCCTTCTGCCT  
TACGTCTAATGGACACACACGCACCACCCCCACACCCTTGCTCCTCCAAAGCCATGC  
AGACTGTGTAAGTGTCTATTATTCTCCAAGGGGCATCCTGTGCAGATGAAACCCTGCT  
TTATTTCCCTGAAGACAGCTGCACAGTGACCTCTTAGTTCTTGCTCCCATGGCCCTG  
ATGTATCCTAGTTACCAGCCCTCAACCTCAGTTCTGAGGGTGGGATCCCATCGCTCA  
GCAAGGCTTCATCCTGACCTCCCTGCCCTGATCTGATCTGGCCCTGGCTTTTGTTGT  
CTCGCTCCCTGACTAAGTGAGATGGGGCACTCTCCCGCCCCCGCCCCCCCCCAGGTCA  
CAGATACCTACCTGCAGCTGTGCGTGCTGGATCACGCACATACTTGCCTTGCATGGT  
CTCCTGGCTGCCCTGGGCTGTGCCTGTTCTTCCATAGGAAGCAAGTTCTTGTCTCCC  
TGGTTCTCAGGGCCCCCTCAGGGGCTCAGCCTTCAGCCCTGTGCTTCCCCATGTTGGG  
AATCTTTGTTACCTTTTTTCTTCTTTGTAAATTAACATCTGATAACAACCACAGGGTC  
CAATGGGACTTTCACAGACCTGCCAGCTAGATAAATAATGACAACAGAAGTTTATTA  
ATATTTTAAGACTTAGGCCTTTTGCTGGGCAGCCTCCCAACTATTCTATCCTGACTA  
ATCCTGGCACTATGTCCCACCACATGGCCAGGTCTACCTCTCTGCTCCACTCTCCAT  
CCACCTCCATGTCTGCCAGCAAATCTCCCGTGATTCAAGTTCTTCTCCCAGAGTCCCT  
ATCTCTGCCCAGAAGTACCATCTTCGACTTCCTGCCCAACTATTGGCCGTCAGCTCT  
TCATTAAAGCCGATCAGATGTAATTCTAGATTGCCTTAGGCAGGTGAGGAAGAAACA  
AGTATTTGTAAAATATGAGACCAGCAATGGGCCATAGAAATAACAGCACCAGATCCT  
GCCAGCATTTAGCCCTCTGTTGGTACAAAATTAACAATTGAATATACAGAGACCTAC  
TTCCAGAGTGTACCCCAACAACAGGCGTGAGCATGGTGCTGGGTACTAGGGTCCTGC  
TGGAAAATCAGAGACCTTACCTACAGCTGGGACATGACCTTGCTTCCGACTTACCCA  
CCACTTCTGGATACCTCACCTCAGCCCACACTATCCCTGGCCTAGGGGCCAGGGTA  
GAGCCAGAAACATGGAGAAAGCATGGCCCCTTGCCGTACCTGGAGAACTGGGTATTT  
TCCAGAGTCTTTATAGATGTGGACTGGAAGGCAGGTGGCCACAGCCGTGCAGACCTG  
GGTCAGGTCAGAAACCTATGCCATGCTGGGACCTACTCAACAGCAGAAGCATGAAGA  
GGGCCTGAGGACAAGAAAGGCCTTCTTACCATGGTGCTATTCTGGAGCTGGGATATA  
TACCTGGCTTGTCTCTGACTGCCCTGGCTTCTGGCAGAACTTCTGATGTCCTCCTGA  
AGGCCTCTCTCCCACCCCAGTACCTGAGAACCTGAGGATAATTTAAACATGGGACTC  
TGGCCAGCACCTGGGAGAGACAGGTAGATCTCTGATTTTTTGACTCAGCCTGGTCTAT  
CGAGTGAGTTCCAGGACATCTGGGGCTACACAGAGAAACCATCTTAAAGACTAAAAA  
TAATAAACATGAGACTGTAAACTGGGTGTATTTTGGGAGAAATAAATGTCTTTTTTCT  
TTCAA (SEQ ID NO: 55)>NP\_598744.1 CD276 antigen precursor [*Mus  
musculus*]

MLRGWGGPSVGVCVRTALGVLCLCLTGAVEVQVSEDPVVALVDTDATLRCSFSPEPG  
FSLAQLNLIWQLTDTKQLVHSFTEGRDQGSAYSNRTALFPDLLVQGNASLRLQRVRV  
TDEGSYTCFVSIQDFDSA AVSLQVAAPYSKPSMTLEPNKDLRPGNMV TITCSSYQGY  
PEAEVFWKDGQGVPLTGNVTTSQMANERGLFDVHSVLRVVLGANGTYSLVRNPVLQ  
QDAHGSVTITGQPLTFPPEALWVTVGLSVCLVLLVALAFVCWRKIKQSCEEENAGA  
EDQDGDGEGSKTALRPLKPSENKEDDGQEIA (SEQ ID NO: 56) Human B7-H4  
>NM\_024626.4 *Homo sapiens* V-set domain containing T cell (VTCN1) activation  
inhibitor 1 (VTCN1), transcript variant 1, mRNA  
GTGAGTCACCAAGGAAGGCAGCGGCAGCTCCACTCAGCCAGTACCCAGATACGCTGG  
GAACCTTCCCCAGCCATGGCTTCCCTGGGGCAGATCCTCTTCTGGAGCATAATTAGC  
ATCATCATTATTCTGGCTGGAGCAATTGCACTCATCATTGGCTTTGGTATTTTCAGGG  
AGACACTCCATCACAGTCACTACTGTCGCCTCAGCTGGGAACATTGGGGAGGATGGA  
ATCCTGAGCTGCACTTTTGAACCTGACATCAAACCTTTCTGATATCGTGATACAATGG

CTGAAGGCTGTTAGTTGTTTGGCTTGGTGGTGAAGCAAGGAACTGAAGTTCG  
TCGGAGCAGGATGAAATGTTTCAGAGGCCGGACAGCAGTGTTTGCTGATCAAGTGATA  
GTTGGCAATGCCTCTTTGCGGCTGAAAAACGTGCAACTCACAGATGCTGGCACCTAC  
AAATGTTATATCATCACTTCTAAAGGCAAGGGGAATGCTAACCTTGAGTATAAACT  
GGAGCCTTCAGCATGCCGGAAGTGAATGTGGACTATAATGCCAGCTCAGAGACCTTG  
CGGTGTGAGGCTCCCCGATGGTTCCCCCAGCCCACAGTGGTCTGGGCATCCCAAGTT  
GACCAGGGAGCCAACTTCTCGGAAGTCTCCAATACCAGCTTTGAGCTGAACTCTGAG  
AATGTGACCATGAAGGTTGTGTCTGTGCTCTACAATGTTACGATCAACAACACATAC  
TCCTGTATGATTGAAAATGACATTGCCAAAGCAACAGGGGATATCAAAGTGACAGAA  
TCGGAGATCAAAGGCGGAGTCACCTACAGCTGCTAAACTCAAAGGCTTCTCTGTGT  
GTCTCTTCTTTCTTTGCCATCAGCTGGGCACTTCTGCCTCTCAGCCCTTACCTGATG  
CTAAAATAATGTGCCTCGGCCACAAAAAAGCATGCAAAGTCATTGTTACAACAGGGA  
TCTACAGAACTATTTCAACCACCAGATATGACCTAGTTTTATATTTCTGGGAGGAAAT  
GAATTCATATCTAGAAGTCTGGAGTGAGCAAACAAGAGCAAGAAACAAAAAGAAGCC  
AAAAGCAGAAGGCTCCAATATGAACAAGATAAATCTATCTTCAAAGACATATTAGAA  
GTTGGGAAAATAATTCATGTGAACTAGACAAGTGTGTTAAGAGTGATAAGTAAAATG  
CACGTGGAGACAAGTGCATCCCCAGATCTCAGGGACCTCCCCCTGCCTGTCACCTGG  
GGAGTGAGAGGACAGGATAGTGCATGTTCTTTGTCTCTGAATTTTTAGTTATATGTG  
CTGTAATGTTGCTCTGAGGAAGCCCCCTGGAAAGTCTATCCCAACATATCCACATCTT  
ATATTCCACAAATTAAGCTGTAGTATGTACCCTAAGACGCTGCTAATTGACTGCCAC  
TTCGCAACTCAGGGGCGGCTGCATTTTAGTAATGGGTCAAATGATTCACCTTTTTATG  
ATGCTTCCAAAGGTGCCTTGGCTTCTCTTCCCAACTGACAAATGCCAAAGTTGAGAA  
AAATGATCATAATTTTAGCATAAACAGAGCAGTCGGCGACACCGATTTTATAAATAA  
ACTGAGCACCTTCTTTTTAAACAAACAAATGCGGGTTTATTTCTCAGATGATGTTCA  
TCCGTGAATGGTCCAGGGAAGGACCTTTCACCTTGTCTATATGGCATTATGTCATCA  
CAAGCTCTGAGGCTTCTCCTTTCCATCCTGCGTGGACAGCTAAGACCTCAGTTTTCA  
ATAGCATCTAGAGCAGTGGGACTCAGCTGGGGTGATTTGCCCCCATCTCCGGGGG  
AATGTCTGAAGACAATTTTGGTTACCTCAATGAGGGAGTGGAGGAGGATACAGTGCT  
ACTACCAACTAGTGGATAGAGGCCAGGGATGCTGCTCAACCTCCTACCATGTACAGG  
ACGTCTCCCCATTACAACCTACCCAATCCGAAGTGTCAACTGTGTCAGGGGCTAAGAAA  
CCCTGGTTTTTGAGTAGAAAAGGGCCTGGAAAGAGGGGAGCCAACAAATCTGTCTGCT  
TCCTCACATTAGTCATTGGCAAATAAGCATTCTGTCTCTTTGGCTGCTGCCTCAGCA  
CAGAGAGCCAGAACTCTATCGGGCACCAGGATAACATCTCTCAGTGAACAGAGTTGA  
CAAGGCCTATGGGAAATGCCTGATGGGATTATCTTCAGCTTGTTGAGCTTCTAAGTT  
TCTTTCCCTTCATTCTACCCTGCAAGCCAAGTTCTGTAAGAGAAATGCCTGAGTTCT  
AGCTCAGGTTTTTCTTACTCTGAATTTAGATCTCCAGACCCTGCCTGGCCACAATTCA  
AATTAAGGCAACAAACATATACCTTCCATGAAGCACACACAGACTTTTGAAAGCAAG  
GACAATGACTGCTTGAATTGAGGCCTTGAGGAATGAAGCTTTGAAGGAAAAGAATAC  
TTTGTTCAGCCCCCTTCCCACACTCTTCATGTGTTAACCCTGCCTTCCTGGACC  
TTGGAGCCACGGTGACTGTATTACATGTTGTTATAGAAAACCTGATTTTAGAGTTCTG  
ATCGTTCAAGAGAATGATTAAATATACATTTCCCTACACCA (SEQ ID NO: 57)

>NP\_078902.2 V-set domain-containing T-cell activation inhibitor 1 isoform 1 precursor [*Homo sapiens*]

MASLGQILFWSIISIIILAGAIALIIGFGISGRHSITVTTVASAGNIGEDGILSCT  
FEPDIKLSDIVIQWLKEGV LGLVHEFKEGKDELSEQDEMFRGRTAVFADQVIVGNAS  
LRLKNVQLTDAGTYKCYIITSKGKGNANLEYKTGAFSMPEVNVDYNASSETLRCEAP  
RWFPQPTVVWASQVDQGANFSEVSNTSFELNSENVTMKVVS VLYNVTINNTYSCMIE  
NDIAKATGDIKVTESEIKRRSHLQLLSKASLCVSSFFAISWALLPLSPYLM LK (SEQ ID  
NO: 58) Mouse B7-H4 >NM\_178594.3 *Mus musculus* V-set domain containing T  
cell (VTCN1) activation inhibitor 1 (Vtcn1), mRNA

GTGAGTCACAAACCCAGGAGGAGCAGGCAGGCAGGCAGGCTCCATCACCAAATC  
TGGCCCCACACACAGCAGGACTGTGGGAAGGAACTCCCTCTCCATGGCTTCCTTGGG  
GCAGATCATCTTTTGGAGTATTATTAACATCATCATCCTGGCTGGGGCCATCGC  
ACTCATCATTGGCTTTGGCATTTCAGGCAAGCACTTCATCACGGTCACGACCTTCAC  
CTCAGCTGGAAACATTGGAGAGGACGGGACCCTGAGCTGCACTTTTGAACCTGACAT  
CAAACCTCAACGGCATCGTCATCCAGTGGCTGAAAGAAGGCATCAAAGGTTTGGTCCA  
CGAGTTCAAAGAAGGCCAAAGACGACCTCTCACAGCAGCATGAGATGTTTCAAGGGCCG  
CACAGCAGTGTTTGCTGATCAGGTGGTAGTTGGCAATGCTTCCCTGAGACTGAAAAA  
CGTGCAGCTCACGGATGCTGGCACCTACACATGTTACATCCGCACCTCAAAAGGCCAA  
AGGGAATGCAAACCTTGAGTATAAGACCGGAGCCTTCAGTATGCCAGAGATAAATGT  
GGACTATAATGCCAGTTCAGAGAGTTTACGCTGCGAGGCTCCTCGGTGGTTCCCCCA  
GCCCCACAGTGGCCTGGGCATCTCAAGTCGACCAAGGAGCCAATTTCTCAGAAGTCTC  
CAACACCAGCTTTGAGTTGAACTCTGAGAATGTGACCATGAAGGTCGTATCTGTGCT  
CTACAATGTCACAATCAACAACACATACTCCTGTATGATTGAAAACGACATTGCCAA  
AGCCACCGGGGACATCAAAGTGACAGATTTCAGAGGTCAAAGGCCGAAGTCAGCTGCA  
GTTGCTGAACTCTGGGCCTTCCCCGTGTGTTTTTTCTTCTGCCTTTGTGGCTGGCTG  
GGCACTCCTATCTCTCTCCTGTTGCCTGATGCTAAGATGAGGGGGCCCTGGCTACACA  
AAAGCATGCAACGTTGCTGGTCCAACAGAATCCCGGAGAACTACAGAAATATTTTCC  
TCAAGACATGACCTAGTTTTATATTTCTAGAAGAAGATGAAATCATGTCTAGAAGTC  
TGGAGAGAGCAGACAGGAACAAGATGTGGAAGGAAAACAAAAGTAACCCACAGACAC  
CCCCGATCGGAACAAGATGGACCTAGAAAATAATTCAACCAAACCTAGAGTATACTAA  
GTGTGCTGTTACAATGTGTGTAGGGTAGGTGTCCTCCACATCTCAGGGGGCCTCCCC  
TGGTCCACCAGCTCCTGAGTTAGGATGGGCTGTTATGATGTCACTCTGAAGGTTCTCT  
GGATGGTTCCTACTGCCATATACTCATTTTATATTCAGCACATTAAACCATAGTGAA  
TGCTATGAAAAGCTGCTAATCAGCTGCCACTCCGAGATTCCGAGGTGGCAACGTCTG  
AGTGACAGGTCCAGTGATTTCGCTTCTCCTTAGGATGCTTTTACAAGCTCTTTGGCGT  
CTCCTCCCACCTGGCAAATGCCAAATGCATAGGGGAGGGTGATCATCATTCTAGGGC  
AAACAAAATAGTTGAGGGATGCTGATTTCCCAAATCATCCGAATCACTTCTCCCTTG  
AGCAAACAAGCGCCCTGTTATTTCTCAAATGCTGCTTTGTGAATCAGTCCAGGGCAA  
GGCGCTCTCCTCATCCCGCTATGTGGCCTTAAGTCATCGTAAGGTTTGAAGTTTCTA  
CTTTTCGATCCTGCATGGAGAGCTATAATCTCAGCTCCCCCGCCCCCCCCACACACAC  
CTCTGCACACACACCCCCCCCCCAACACTGGGAGTAAACCAGGATGATGTCCGTCTTC  
TCATTCCCCATGTGACCGTTGGCAGTGTAGAGAGACTGATTGTACAGCTAAAGGAA  
GAGGGACAACAGGGTCACTGGTGTCTACAGAGATTATATTCTACGTGTCTCACTGAA  
TTTACACAACCTCCAAGTGCCAACCACATCAAGGTCAGGAAATCCTGAACTGGAATAA  
GAAAGACCCAGAAGATGAATGTGAACAGATCCATTTGCTTCCCGACAGTGGGCACAG  
ACTTCAGTCTCTGGCTACTGTTCCAAGACCCAGGGCTCTGCAATTGTGTGACATCCT  
TCAGTGAACCCACATGGGAAATTCTCCATGGAATTATCTTCAGCCCACTGTACTTCT  
GAATCCCTCTTCCTTCTGTGCCACACAGCAAGTCTGGCTTAAATGCTGCCTGA  
TCTCCATTTCAAGTTTTCTGCCTCTGGATTTTTAGATCTCAAGACCATGGACGAAAC  
ATCAGTTACAGCAACAAAAGTGAATTTCCGTGCAGAGACTTCTAGGGGTTCTGTTT  
GTTTTTCAGGGTGCTAGAGATCACACTCAGATGCTCATATATGTTAGGTAAATGTTCT  
CCCCTGAGTTACAGCCCAGCTCACACAGAGACTTCTAAAAGAAAATACGGCCATGC  
TCTTTGAAATGGAGCATTGAGGGATGAAGTTTGGATGGCGAAGAAAACCTTCTACCA  
GCTCTCTCCCCACATTCGTGCCAAGCACTGCCTCCCTAGACTTCGGGGTCACCATATC  
TGTACTACGTTTTTGATACAGAAGGCTCGAGACCATTCAAGAGAATTATTTAGTACAC

(SEQ ID NO: 59)>NP\_848709.2 V-set domain containing T-cell activation inhibitor 1 precursor [*Mus musculus*]

MASLGQIIFWSIINIILLAGAIALIIGFGISGKHFITVTTFTSAGNIGEDGTLST

FEPDIKLNIGIVIQWLKEGIKGLVHEFKEGKDDLSQLHEMFRGRITAVFADQVVVGNAS

LRKYNQVLTDTYRTQYCTYRTSKGKNANLEYKKTGAFSMPEINVDYNASSESLRCEAP  
RWFPQPTVAWASQVDQGANFSEVSNTSFELNSENVTMKVVSVLNVNTINNTYSCMIE  
NDIAKATGDIKVTDSEVKRRSQLQLLNSGPSPCVFSSAFVAGWALLSLSCCLMLR (SEQ  
ID NO: 60) Human B7-H5 >NM\_022153.2 *Homo sapiens* V-set immunoregulatory  
receptor (VISTA) (VSIR), mRNA  
AGTCGCGGGAGGCTTCCCCGCGCCGGCCGCGTCCCCGCCCGCTCCCCGGCACCAGAAG  
TTCCTCTGCGCGTCCGACGGCGACATGGGCGTCCCCACGGCCCTGGAGGCCGCGCAGC  
TGGCGCTGGGGATCCCTGCTCTTCGCTCTCTTCCTGGCTGCGTCCCTAGGTCCGGTG  
GCAGCCTTCAAGGTCGCCACGCCGTATTCCTGTATGTCTGTCCCGAGGGGCGAGAAC  
GTCACCCTCACCTGCAGGCTCTTGGGCCCTGTGGACAAAGGGCACGATGTGACCTTC  
TACAAGACGTGGTACCGCAGCTCGAGGGGCGAGGTGCAGACCTGCTCAGAGCGCCGG  
CCCATCCGCAACCTCACGTTCCAGGACCTTCACCTGCACCATGGAGGCCACCAGGCT  
GCCAACACCAGCCACGACCTGGCTCAGCGCCACGGGCTGGAGTCGGCCTCCGACCAC  
CATGGCAACTTCTCCATACCATGCGCAACCTGACCCTGCTGGATAGCGGCCTCTAC  
TGCTGCCTGGTGGTGGAGATCAGGCACCACCACTCGGAGCACAGGGTCCATGGTGCC  
ATGGAGCTGCAGGTGCAGACAGGCAAAGATGCACCATCCAACCTGTGTGGTGTACCCA  
TCCTCCTCCCAGGATAGTGAAAACATCAGGCTGCAGCCCTGGCTACGGGTGCCTGC  
ATCGTAGGAATCCTCTGCCTCCCCCTCATCCTGCTCCTGGTCTACAAGCAAAGGCAG  
GCAGCCTCCAACCGCCGTGCCCAGGAGCTGGTGCGGATGGACAGCAACATTCAAGGG  
ATTGAAAACCCCGGCTTTGAAGCCTCACCACTGCCCAGGGGATACCCGAGGCCAAA  
GTCAGGCACCCCCTGTCCTATGTGGCCCAGCGGCAGCCTTCTGAGTCTGGGCGGCAT  
CTGCTTTCGGAGCCCAGCACCCCCCTGTCTCCTCCAGGCCCCGGAGACGTCTTCTTC  
CCATCCCTGGACCCTGTCCCTGACTCTCCAAACTTTGAGGTCATCTAGCCCAGCTGG  
GGGACAGTGGGCTGTTGTGGCTGGGTCTGGGGCAGGTGCATTTGAGCCAGGGCTGGC  
TCTGTGAGTGGCCTCCTTGGCCTCGGCCCTGGTTCCCTCCCTCCTGCTCTGGGCTCA  
GATACTGTGACATCCCAGAAGCCCAGCCCCTCAACCCCTCTGGATGCTACATGGGGA  
TGCTGGACGGCTCAGCCCCTGTTCCAAGGATTTTGGGGTGCTGAGATTCTCCCCTAG  
AGACCTGAAATTCACCAGCTACAGATGCCAAATGACTTACATCTTAAGAAGTCTCAG  
AACGTCCAGCCCTTCAGCAGCTCTCGTTCTGAGACATGAGCCTTGGGATGTGGCAGC  
ATCAGTGGGACAAGATGGACACTGGGGCCACCCTCCCAGGCACCAGACACAGGGCACG  
GTGGAGAGACTTCTCCCCCGTGGCCGCCTTGGCTCCCCCGTTTTGCCCCAGGGCTGCT  
CTTCTGTCAGACTTCCTCTTTGTACCACAGTGGCTCTGGGGCCAGGCCTGCCTGCCC  
ACTGGCCATCGCCACCTTCCCCAGCTGCCTCCTACCAGCAGTTTCTCTGAAGATCTG  
TCAACAGGTAAAGTCAATCTGGGGCTTCCACTGCCTGCATTCCAGTCCCCAGAGCTT  
GGTGGTCCCGAAACGGGAAGTACATATTGGGGCATGGTGGCCTCCGTGAGCAAATGG  
TGTCTTGGGCAATCTGAGGCCAGGACAGATGTTGCCCCACCCACTGGAGATGGTGCT  
GAGGGAGGTGGGTGGGGCCTTCTGGGAAGGTGAGTGGAGAGGGGGCACCTGCCCCCCG  
CCCTCCCCATCCCCTACTCCCCTGCTCAGCGCGGGCCATTGCAAGGGTGCCACACA  
ATGTCTTGTCCACCCTGGGACACTTCTGAGTATGAAGCGGGATGCTATTAAAACTA  
CATGGGGAAACAGGTGCAAACCCTGGAGATGGATTGTAAGAGCCAGTTTAAATCTGC  
ACTCTGCTGCTCCTCCCCCACCCCCACCTTCCACTCCATACAATCTGGGCCTGGTGG  
AGTCTTCGCTTCAGAGCCATTTCGGCCAGGTGCGGGTGATGTTCCCATCTCCTGCTTG  
TGGGCATGCCCTGGCTTTTGTTTTTTATACACATAGGCAAGGTGAGTCCTCTGTGGAAT  
TGTGATTGAAGGATTTTAAAGCAGGGGAGGAGAGTAGGGGGCATCTCTGTACACTCT  
GGGGGTAAAACAGGGAAGGCAGTGCCTGAGCATGGGGACAGGTGAGGTGGGGCTGGG  
CAGACCCCCTGTAGCGTTTAGCAGGATGGGGGGCCCCAGGTACTGTGGAGAGCATAGT  
CCAGCCTGGGCATTTGTCTCCTAGCAGCCTACACTGGCTCTGCTGAGCTGGGCCTGG  
GTGCTGAAAGCCAGGATTTGGGGCTAGGCGGGAAGATGTTTCGCCCAATTGCTTGGGG  
GGTTGGGGGGGATGGAAAAGGGGAGCACCTCTAGGCTGCCTGGCAGCAGTGAGCCCTG  
GGCCTGTGGCTACAGCCAGGGAACCCACCTGGACACATGGCCCTGCTTCTAAGCCC

CCAGTTCAGTCCAAAGTGGTCCACTGAGGGCCTCCTGCTCTGCCTGGGCTGGG  
CCAGGGGCTTTGAGGAGAGGGTAAACATAGGCCCGGAGATGGGGCTGACACCTCGAG  
TGGCCAGAATATGCCCAAACCCCGGCTTCTCCCTTGTCCCTAGGCAGAGGGGGGTCC  
CTTCTTTTGTTCCTCTGGTCAACCACAATGCTTGATGCCAGCTGCCATAGGAAGAGG  
GTGCTGGCTGGCCATGGTGGCACACACCTGTCCTCCCAGCACTTTGCAGGGCTGAGG  
TGGAAGGACCGCTTAAGCCCAGGTGTTCAAGGCTGCTGTGAGCTGTGTTTCGAGCCAC  
TACTCTCCAGCCTGGGGACGGAGCAAACTTTGCCTCAAAACAAATTTTAAAAAGAA  
AGAAAGAAGGAAAGAGGGTATGTTTTTCACAATTCATGGGGGCCTGCATGGCAGGAG  
TGGGGACAGGACACCTGCTGTTCCCTGGAGTCGAAGGACAAGCCCACAGCCCAGATTC  
CGGTTCTCCCAACTCAGGAAGAGCATGCCCTGCCCTCTGGGGAGGCTGGCCTGGCCC  
CAGCCCTCAGCTGCTGACCTTGAGGCAGAGACAACCTTCTAAGAATTTGGCTGCCAGA  
CCCCAGGCCTGGCTGCTGCTGTGTGGAGAGGGAGGCGGCCCGCAGCAGAACAGCCAC  
CGCACTTCCTCCTCAGCTTCCTCTGGTGCGGGCCCTGCCCTCTCTTCTCTGGACCCTT  
TTACAACCTGAACGCATCTGGGCTTCGTGGTTTCCTGTTTTTCAGCGAAATTTACTCTG  
AGCTCCCAGTTCCATCTTCATCCATGGCCACAGGCCCTGCCTACAACGCACTAGGGA  
CGTCCCTCCCTGCTGCTGCTGGGGAGGGGCAGGCTGCTGGAGCCGCCCTCTGAGTTG  
CCCGGGATGGTAGTGCCTCTGATGCCAGCCCTGGTGGCTGTGGGCTGGGGTGCATGG  
GAGAGCTGGGTGCGAGAACATGGCGCCTCCAGGGGGCGGGAGGAGCACTAGGGGGCTG  
GGGCAGGAGGCTCCTGGAGCGCTGGATTCTGTGGCACAGTCTGAGGCCCTGAGAGGGA  
AATCCATGCTTTTAAGAACTAATTCATTGTTAGGAGATCAATCAGGAATTAGGGGCC  
ATCTTACCTATCTCCTGACATTCACAGTTTAATAGAGACTTCCTGCCTTTATTCCCT  
CCCAGGGAGAGGCTGAAGGAATGGAATTGAAAGCACCATTTGGAGGGTTTTGCTGAC  
ACAGCGGGGACTGCTCAGCACTCCCTAAAAACACACCATGGAGGCCACTGGTGACTG  
CTGGTGGGCAGGCTGGCCCTGCCTGGGGGAGTCCGTGGCGATGGGCGCTGGGGTGGA  
GGTGCAGGAGCCCCAGGACCTGCTTTTCAAAAGACTTCTGCCTGACCAGAGCTCCCA  
CTACATGCAGTGGCCCAGGGCAGAGGGGCTGATACATGGCCTTTTTTCAGGGGGTGCT  
CCTCGCGGGGTGGACTTGGGAGTGTGCAGTGGGACAGGGGGGCTGCAGGGGGTCCTGCC  
ACCACCGAGCACCAACTTGGCCCCCTGGGGTCCTGCCTCATGAATGAGGCCTTCCCCA  
GGGCTGGCCTGACTGTGCTGGGGGCTGGGTTAACGTTTTCTCAGGGAACCAACAATGC  
ACGAAAGAGGAACCTGGGGTTGCTAACCAGGATGCTGGGAACAAAGGCCTCTTGAAGC  
CCAGCCACAGCCCAGCTGAGCATGAGGCCCAGCCCATAGACGGCACAGGCCACCTGG  
CCCATTCCCTGGGCATTCCCTGCTTTGCATTGCTGCTTCTCTTCACCCCATGGAGGC  
TATGTCACCCTAACTATCCTGGAATGTGTTGAGAGGGATTCTGAATGATCAATATAG  
CTTGGTGAGACAGTGCCGAGATAGATAGCCATGTCTGCCTTGGGCACGGGAGAGGGA  
AGTGGCAGCATGCATGCTGTTTCTTGGCCTTTTCTGTTAGAATACTTGGTGCTTTCC  
AACACACTTTTACATGTGTTGTAACCTTGTTTGATCCACCCCTTCCCTGAAAATCCT  
GGGAGGTTTTATTGCTGCCATTTAACACAGAGGGCAATAGAGGTTCTGAAAGGTCTG  
TGTCTTGTCAAAACAAGTAAACGGTGGAACTACGACTAAA (SEQ ID NO: 61)

>NP\_071436.1 V-type immunoglobulin domain-containing suppressor of T-cell activation precursor [*Homo sapiens*]

MGVPTALEAGSWRWGSLFLAASLGPVAAFKVATPYSLYVCPEGQNVTLTCRLL  
GPVDKGHDTVTFYKTWYRSSRGEVQTCSERRPIRNLTFQDLHLHHGGHQAANTSHDLA  
QRHGLEASDHHGNFSITMRNLTLDSGLYCCLVVEIRHHHSEHRVHGAMELQVQTG  
KDAPSNCVVYPSSSQDSENITAAALATGACIVGILCLPLILLVYKQRQAASNRRAQ  
ELVRMDSNIQGIENPGFEASPPAQGIPEAKVRHPLSYVAQRQPSESGRHLLSEPSTP  
LSPPGPGDVFFPSLDPVPDSPNFEVI (SEQ ID NO: 62) Mouse B7-H5

>NM\_028732.4 *Mus musculus* V-set immunoregulatory receptor (VISTA) (Vsir), transcript variant 1, mRNA

GGGGGCGCTGCTGGGCGGGGAGCTTGCTCGGCCGCCTGCCTCGCCTTGGGCTCAGCA  
TTCCTCTAGCGAGCGAGCGGCGTGTACAGCCGGCTCCCTGGGCTCCTGGAGTCCCG

CTTGCTTCCAAGCGCACTCCAGCAGTCTCTTTCTGCTCTTTGCCCGGCTCGACAGGCGAC  
ATGGGTGTCCCCGCGGTCCCAGAGGCCAGCAGCCCCGCGCTGGGGAACCCCTGCTCCTT  
GCTATTTTCCTGGCTGCATCCAGAGGTCTGGTAGCAGCCTTCAAGGTCACCACTCCA  
TATTCTCTCTATGTGTGTCCCGAGGGACAGAATGCCACCCTCACCTGCAGGATTCTG  
GGCCCCGTGTCCAAAGGGCACGATGTGACCATCTACAAGACGTGGTACCTCAGCTCA  
CGAGGCGAGGTCCAGATGTGCAAAGAACACCGGCCCATACGCAACTTCACATTGCAG  
CACCTTCAGCACCACGGAAGCCACCTGAAAGCCAACGCCAGCCATGACCAGCCCCAG  
AAGCATGGGCTAGAGCTAGCTTCTGACCACCACGGTAACTTCTCTATCACCCCTGCGC  
AATGTGACCCCAAGGGACAGCGGCCTCTACTGCTGTCTAGTGATAGAATTA AAAAAC  
CACCACCCAGAACAACGGTTCTACGGGTCCATGGAGCTACAGGTACAGGCAGGCAAA  
GGCTCGGGGTCCACATGCATGGCGTCTAATGAGCAGGACAGTGACAGCATCACGGCT  
GCGGCCCTGGCCACCGGCGCCTGCATCGTGGGAATCCTCTGCCTCCCCCTTATCCTG  
CTGCTGGTCTATAAGCAGAGACAGGTGGCCTCTCACCGCCGTGCCCAGGAGTTGGTG  
AGGATGGACAGCAGCAACACCCAAGGAATCGAAAACCCAGGCTTCGAGACCACTCCA  
CCCTTCCAGGGGATGCCTGAGGCCAAGACCAGGCCGCCACTGTCCTATGTGGCCCAG  
CGGCAACCTTCGGAGTCAGGACGGTACCTGCTCTCTGACCCCAGCACACCTCTGTCTG  
CCTCCAGGCCCTGGGGACGTCTTTTTTCCCATCCCTAGATCCAGTCCCTGACTCCCCT  
AACTCTGAAGCCATCTAAACCAGCTGGGGAACCATGAACCATGGTACCTGGGTCAGG  
GATATGTGCACTTGATCTATGGCTGGCCCTTGGACAGTCTTTTAGGCACTGACTCCA  
GCTTCCTTGCTCCTGCTCTGAGCCTAGACTCTGCTTTTACAAGATGCACAGACCCTC  
CCCTATCTCTTTCAGACGCTACTTGGGGGGCAGGGAGAAGATGTTGGATTGCTCATT  
GCTGTTCTCAAGATCTTGGGATGCTGAGTTCTCCCTAGAGACTTGACTTCGACAGCC  
ACAGATGTCAGATGACCTGCATCCTATGAACGTCCGGCTTGGCAAGAGCCTTTCTTC  
ATGGAAACCAGTAGCCCGGAGGGGATGAGGTAGGCACCTTGCCACCCTCCCGGGAGA  
GAGACACAAGATGTGAGAGACTCCTGCTCACTGTGGGGGTGTGGCTGGCCTGCTTGT  
TTGCCTGAGGATGCTCCTCTGTTGGA CTGACTCTATCCCCCTGGATTCTGGAGCTTG  
GCTGGCCTATGTCCCACCAGAGGAGCATCTCAGCAGCCTTCCACCAGCAACCTGAGG  
GCCTGCCAGCTTCGTGGCTCTGGGCTCTCATTACCTGTATGGCCGTCCACAGAGCTC  
AGTGGCCAGAGGCTTTGAAACAGGAAGTACATGTCAGG TTCAGGAACCACTGTGAGC  
TCATTAGTGTCTTGAGCAATGTGAGGCCTGGACCAGTGGACACGGAGGGAGGGTGGC  
GAGAGGATGATGGGGATGATGAGGGGAACACGCTCCCTTCTGTCTTGT CATCCAC  
CACTACCACTATT CAGTGTGGAGCAGTGGCAAAGGTGACCGACCTCCACAATGTCT  
AGTGATGCTGGACCA TTTCTAAGTGTGAAAGAGATGCTATTA AAAACAGTATGTGGC  
AATGGCTGCCAACAGCTGAGTGGACTGGAGGCACTGGCTTTAAGGCCCTGGAGGTGC  
AGGGCCCCGTATGGGGATAGGGATGGGAGTTTCAGTGAGGGCCTAGGGATCACTCCG  
CTTCTGACCACTCTTCTTCTGAGCCTCACCTCAGGGTGACCTTCAGGCACACAGAAG  
AGCTTGCCCCCTGGTCCGATACTACTCTTGGCTCTCATCTCCAGGGTTTGGCATGACC  
TGGGCACACAGGGGGAGTCTTCAGAAAGGATTTTAAAGCATGAAAAGAAAGGGTAGT  
TCTTGTGAGGTAGGGATGGGCAGCTGATGTTTGAGAGTGAGGAGGGATACGGCTGGG  
CAGATCACTCTCCAGTCTCTAGAGGGAAAGTAGCTCTAAGTCTGGGAGAGCAGCAGC  
CCAGTGGTACCATATGTCTTCTTGCAGCTTCCACTGGCTGGGCTGAACTGGGCATGG  
GTAGGAAAGCTCCTGTTOTGGGCCTGCAGCCAGGGAGAACCCCATTCATTCCCTGAG  
GACAGATGGGTGGGGAGAGAAGAGAGAGTTTCAGGCCGGGAAGCAGCAATAAGCTAT  
CTGCTGGGGACCCAGACAAGTTGTCTGATGAGGTCCAAGATGTGGGATGCCAGTTAT  
ACCTGGGGCTTGGGGATCCTTAGAGGC TTTGTATCATCATCATAGGAGTGTCTGGGGT  
GGCCAGGGCATCAAAGCCATGACCCCTGTTTTATCCTCAGGGTCCACTCTTCTGCAC  
CATCCATTGCTCTAGATCTATGCAGTTACTATAGACAGAATGTGTTGTTCTGTTTGG  
CTTTGGGGATAATGGCCTGGCGAACTGCCAGCTGTT CAGTGGCAGGGCTGTGAGGCC  
AGTCAAAGACTAGAACCCACAGACCAGCTGAACGATGAGTATAGCCTGTCCCCTGGG  
GGAGCCTGACCTGTCTCCAGCCCTAAGCTTCAGACCTCACC ACTCAGATGACTTCTA



(SEQ ID NO: 63) >NP\_083008.1 V-type immunoglobulin domain-containing suppressor of T-cell activation isoform 1 precursor [*Mus musculus*]  
MGVPAVPEASSPRWGTL~~LL~~AIFLAASRGLVAAFKVTTPYSLYVCPEGQNATLTCRIL  
GPVSKGHDVTIYK~~TW~~YLSSRGEVQMCKEHRPIR~~NFT~~LQHLQH~~HG~~SHLKANASHDQPQ  
KHGLELASDHHGNFSITLRNVT~~PR~~DSGLYCCLVIELKNHHPEQRFYGS~~MEL~~QVQAGK  
GSGSTCMASNEQDS~~DS~~SITAAALATGACIVGILCLPLILLLVYKQRQVASHRRAQELV  
RMDSSNTQGIENPGFETTPPFQGMPEAKTRPPLSYVAQRQPSESGRYLLSDPSTPLS  
PPGPGDVFFPSLDPVPDSPNSEAI (SEQ ID NO: 64) Human B7-H7 >NM\_007072.4  
*Homo sapiens* HERV-H LTR-associating 2 (HHLA2) (HHLA2), transcript variant  
1, mRNA

TGTTCTGCACAAAGATCAAGAGACAGACACTGCTCTTTCTCTGCTCATATA  
ACATCTCTGAGTGGATCTCAAGGCATATTCCCTTTGGCTTTCTTCATTTATGTTTCCT  
ATGAATGAACAAATCGTCATTGGAAGACTTGATGAAGATATAATTCTCCCTTCTTCA  
TTTGAGAGGGGATCCGAAGTCGTAATACACTGGAAGTATCAAGATAGCTATAAGGTT  
CACAGTTACTACAAAGGCAGTGACCATTTGGAAAGCCAAGATCCCAGATATGCAAAC  
AGGACATCCCTTTTCTATAATGAGATTCAAAATGGGAATGCGTCGCTATTTTTTCAGA  
AGAGTAAGCCTTCTGGACGAAGGAATTTACACCTGCTATGTAGGAACAGCAATTCAA  
GTGATTACAAACAAAGTGGTGCTAAAGGTGGGAGTTTTTTCTCACACCCGTGATGAAG  
TATGAAAAGAGGAACACAAACAGCTTCTTAATATGCAGCGTGTTAAGTGTTTATCCT  
CGTCCAATTATCACGTGGAAAATGGACAACACACCTATCTCTGAAAACAACATGGAA  
GAAACAGGGTCTTTGGATTCTTTTTTCTATTAAACAGCCCACTGAATATTACAGGATCA  
AATTCATCTTATGAATGTACAATTGAAAATTCCTGCTGAAGCAAACATGGACAGGG  
CGCTGGACGATGAAAGATGGCCTTCATAAAATGCAAAGTGAACACGTTTCACTCTCA  
TGTC AACCTGTAAATGATTATTTTTCACCAAACCAAGACTTCAAAGTTACTTGGTCC  
AGAATGAAAAGTGGGACTTTCTCTGTCTGGCTTACTATCTGAGCTCCTCACAAAAT  
ACAATTATCAATGAATCCCGATTCTCATGGAACAAAGAGCTGATAAACCAGAGTGAC  
TTCTCTATGAATTTGATGGATCTTAATCTTTCAGACAGTGGGGAATATTTATGCAAT  
ATTTCTTCGGATGAATATACTTTACTTACCATCCACACAGTGCATGTAGAACCGAGC  
CAAGAAACAGCTTCCCATAACAAAGGCTTATGGATTTTGGTGCCCTCTGCGATTTTG  
GCAGCTTTTCTGCTGATTTGGAGCGTAAAATGTTGCAGAGCCCAGCTAGAAGCCAGG  
AGGAGCAGACACCCTGCTGATGGAGCCCAACAAGAAAGATGTTGTGTCCCTCCTGGT  
GAGCGCTGTCCCAGTGCACCCGATAATGGCGAAGAAAATGTGCCTCTTTCAGGAAAA  
GTATAGGAAATGAGAGAAGACTGTGACAACCTCATGACCTGCATCCTTAATATCCAGT  
GACTTCATCTCCCCTTTCTTCACCACAATTCCAGGCAATGGCCTGTCTGGAGCAGACA  
ATTCTACCACTGCAAAGAGTTGTAACCATTTTCTGGTATCACATTTATTTTTCAAGA  
CATACTTTTCAAGACATCATTCACTGACCCACTACCTGCATTGAGTATAAATGCCTG  
GATGTTAAGGATTCCAATTTAACTTTGAAAAGAACTGTCTCATTCAATTACATTTCT  
GTTACAGTCAGCCCAGGAGGTTACAGTGAGCTCTCCACTAAGAATCTGGAAGAAATG  
CATCACTAGGGGTTGATTCCCAATCTGATCAACTGATAATGGGTGAGAGAGCAGGTA  
AGAGCCAAAGTCACCTTAGTGGAAGGTTAAAAACCAGAGCCTGGAAACCAAGATGA  
TTGATTTGACAAGGTATTTTAGTCTAGTTTTATATGAACGGTTGTATCAGGGTAACC  
AACTCGATTTGGGATGAATCTTAGGGCACCAAAGACTAAGACAGTATCTTTAAGATT  
GCTAGGGGAAAAGGGCCCTATGTGTCAGGCCTCTGAGCCCAAGCCAAGCATCGCATCC  
CCTGTGATTTGCACGTATACATCCAGATGGCCTAAAGTAACTGAAGATCCACAAAAG  
AAGTAAAAATAGCCTTAACTGATGACATTCCACCATTGTGATTTGTTCCCTGCCCCAC  
CCTAACTGATCAATGTACTTTGTAATCTCCCCCACCCTTAAGAAGGTACTTTGTAAT  
CTTCCCCACCCTTAAGAAGGTTCTTTGTAATTCTCCCCACCCTTGAGAATGTACTTT  
GTGAGATCCACCCTGCCCACAAAACATTGCTCTTAACTTCACCGCCTAACCCAAAAC  
CTATAAGAACTAATGATAATCCATCACCTTCGCTGACTCTCTTTTCGGACTCAGCC  
CACCTGCACCCAGGTGAAATAAACAGCTTTATTGCTCACACAAA (SEQ ID NO: 65)

>NP\_009003.1 HERV-H LTR-associating protein 2 isoform a precursor [*Homo sapiens*] MKAQTALSFLLILITSLSGSQGIFPLAFFIYVPMNEQIVIGRLDEDIILPSSFERGS  
EVVIHWKYQDSYKVHSYYKGS DHLESQDP RYANRTSLFYNEIQNGNASLFFRRVSL  
DEGIYTCYVGTAIQVITNKVVLKVG VFLTPVMKYEK RNTNSFLICSVLSVYPRPIIT  
WKMDNTPISENNMEETGSLDSFSINSPLNITGSNSSYECTIENSLLKQ TWTGRWTMK  
DGLHKMQSEHVSLSQPVNDYFSPNQDFKVTWSRMKSGTFSVLAYYLSSSQNTIINE  
SRFSWNKELINQSDFSMNLMDLNLSDSGEYLCNISSDEYTL LTIHTVHV EPSQETAS  
HNKGLWILVPSAILAAFLLIWSVKCCRAQLEARRSRHPADGAQ QERCCVPPGERCPS  
APDNGEENVPLSGKV (SEQ ID NO: 66 Mouse BTNL1 >NM\_001111094.1 *Mus musculus* butyrophilin-like 1 (Btnl1), mRNA

ACCTTAAATAAGAGCTGAGAGCTGAGCCTTTCTCCTAGACTCCTCCAGGAGAA  
ACTCTAAAGCCAGAGCCTGGGGGCGAGCATTGTGTGTCCACCTTGCCACTGAGAACAT  
CTACGGAAATTGGACACTCTGGCCCCAGCATCCACACGCTTGACTGTTGGCCACAGT  
AACACAGGTGTGGATGGTCCCCAGAGCCAGGGTCCAGGAGTGCAGTGGAGGATCCCTG  
GGGCTTCAAGGAACCCACAGCTCTGTCCAGACGGGAATTTTTTTCCTGAGAACTTTC  
ACCTGTTGCCCTCCTATGGTGAACCTGGACTTGACCTTCCACTCTGATGATGAAGGG  
CTCCCCCTCCGTCCCTCCAGCTGGTTGTCTCCTCCCTCTGCTCCTCCTGCTGTTTAC  
CGGAGTCTCTGGAGAAGTGTCTTGGTTTTCTGTGAAGGGACCAGCTGAGCCCATCAC  
TGTCCTGCTGGGGACTGAAGCCACCCTGCCCTGCCAGCTGTCTCCTGAACAGAGTGC  
AGCTCGCATGCACATCCGATGGTACCGTGCCCAGCCCACCCCTGCTGTGCTGGTGT  
CCACAACGGACAGGAGCAGGGAGAGGTGCAGATGCCGGAATACAGGGGCGAGGACCCA  
GATGGTGAGACAAGCCATTGACATGGGAAGTGTGGCTCTGCAGATACAGCAGGTCCA  
GGCCTCTGATGATGGCCTGTACCACTGTCAGTTTACAGATGGCTTCACCTCCCAAGA  
GGTCTCCATGGAGCTTCGAGTCATAGGTTTAGGCTCTGCCCCCTCTTGTTACATGAC  
AGGACCTGAGAATGATGGGATCCGAGTGTTGTGCTCCTCAAGTGGCTGGTTCCCAA  
ACCCAAAGTGCAATGGAGAGACACCTCCGGGAACATGCTACTGTCCTCCTCTGAGTT  
GCAGACCCAAGACAGAGAAGGGCTCTTCCAGGTGGAAGTGTCTCTTTTGGTCACAGA  
TAGAGCTATTGGCAATGTGATCTGCTCCATCCAAAATCCCATGTATGACCAGGAGAA  
ATCGAAGGCCATCCTCCTCCCAGAGCCCTTCTTCCCCAAGACGTGTCCATGGAAAGT  
AGCCCTGGTTTGTCTGTCTCATACTATTGGTCCTGCTCGGTGGGATCAGCCTTG  
AATCTGGAAAGAACATCAAGTCAAAAGGAGAGAAATTAATAAATGGTCAAAGGAACA  
TGAAGAAATGCTTCTGTTGAAGAAGGGGACAAAATCTGTACTGAAGATCAGAGATGA  
CCTCCAGGCCGACCTAGATCGGAGGAAGGCGCTGTACAAAGAAGACTGGAAGAAGGC  
CTTGCTGTACCCTGACTGGAGGAAGGAGCTGTTCCAGGAGGCTCCTGTGAGGATAAA  
TTATGAAATGCCTGACCAGGACAAGACAGACTCAAGGACAGAAGAGAACAGAGGTGA  
GGAGACTGTCAGCAGCTCACAAGTAGACCACAACCTCATCACACTCTCCCAGGAAGG  
CTTCATGTTGGGAAGATACTACTGGGAGGTGGATGTCAAGGACACAGAGGAGTGGAC  
ACTAGGAGTTTATGAGCTGTGCACTCAGGATGCATCACTTACAGACCCCTTGAGGAA  
ATTCAGAGTCCTGGAAAAGAATGGAGATGGATACAGGGCTCTTGACTTCTGTTCCCA  
AAACATTAATTCGGAAGAACCTCTGCAACTGAAGACACGTCCGCTGAAGATCGCCAT  
CTTCTTGATCAGGAAGACAATGACCTCTCTTTCTACAACATGACCGATGAGACACA  
CATCTTTTCCTTTGCCCAGGTCCCTTTCTTGGGATCACCTATCCTTACTTCACACG  
TAATTCCATGGGGCTCTCTGCAACAGCACAGCCCTAAGTGATGTGCACAGGGAATTC  
AATGGGTGGGTGCTGCAGCGTGCTACCCGTAAGGCCCTCTTAGGCAGGCACAGGGGG  
CCTCTGACCAAGAGGCCTCTTAACCTGAGACTCCATGAGCCTCGGGGATCAGATCCT  
GGACAAGATTCTCGGACCATCTGTGTCGTGCATGGTGTATAGTTATTAATAGCCTT  
CCTTCTTTTGACAAAAATGTGTTTAATCATTCTAAGATAAATGAATCCATGGCTTT  
CTGA (SEQ ID NO: 67)>NP\_001104564.1 butyrophilin-like protein 1 precursor  
[*Mus musculus*]

MMKGSPSVPPAGCLLPLLLLFTGVSGEVSWFSVKGPAEPITVLLGTEATLPCQLSP  
EQSAARMHIRWYRAQPTPAVLVFHNGQEQQGEVQMPEYRGRQTQMVQRQAIDMGVALQI  
QQVQASDDGLYHCQFTDGFTSQEVSMELRVIGLGSAPLVHMTGPENDGIRVLCSSSG  
WFPKPKVQWRDTSNMLLSSELQTQDREGLFQVEVSLLVTDRAIGNVICSIQNP  
DQEKSKAILLPEPFFPKTCPWKVALVCSVLILLVLLGGISLGIWKEHQVKRREIKKW  
SKEHEEMLLLKKGTKSVLKIRDDLQADLDRRKALYKEDWKKALLYPDWRKELFQEAP  
VRINYEMPQDKTDSRTEENRGEETVSSSQVDHNLITLSQEGFMLGRYYWEVDVKDT  
EEWTLGVYELCTQDASLTDPLRKFRVLEKNGDGYRALDFCSQNINSEEPLQLKTRPL  
KIAIFLDQEDNDLSFYNMTEDETHIFSFAQVPFLGSPYPYFTRNSMGLSATAQP (SEQ ID  
NO: 68) Human VSIG8>NM\_001013661.1 *Homo sapiens* V-set and immunoglobulin  
domain containing 8 (VSIG8), mRNA

MRVGGAFHLLLVCLSPALLSAVRINGDGQEVLYLAEGDNVRLGCPYVLDPEDYGPNG  
LDIEWMQVNSDPAHHRENVFLSYQDKRINHGSLPHLQQRVRFAASDPSQYDASINLM  
NLQVSDTATYECRVKKTTMATRKVIVTVQARPAVPMCWTEGHMTYGNDVVLKCYASG  
GSQPLSYKWAKISGHHYPYRAGSYTSQHSYHSELSYQESFHSSINQGLNNGDLVLKD  
ISRADDGLYQCTVANNVGYSVCVVEVKVSDSRRIGVIIIGIVLGSLLALGCLAVGIWG  
LVCCCCGGSGAGGARGAFGYGNGGGVGGGACGDLASEIREDAVAPGCKASGRGSRVT  
HLLGYPTQNVSRSLRRKYAPPPCGGPEDVALAPCTAAACEAGPSPVYVKVKS AEPA  
DCAEGPVQCKNGLLV (SEQ ID NO: 70) Mouse VSIG8 >NM\_177723.4 *Mus  
musculus* V-set and immunoglobulin domain containing 8 (Vsig8), transcript  
variant 1, mRNA



TTAGTTCCTCCCTTCTGCTCCCAAAATCCAGGATCCGACCAAGGATCCGACCAAGG  
GATGTCATCCTGCTCTGTAGCTCAGAGGAAGGCATTCCTCGACCAACTTACCTTTGG  
GAGAAGTTAGACAATACCCTCAAACACTACCTCCAACAGCTACTCAGGACCAGGTCCAG  
GGAACAGTCACCATCCGGAACATCAGTGCCCTGTCTTCAGGTTTGTACCAGTGCGTG  
GCTTCTAATGCTATTGGAACCAGCACCTGTCTTCTGGATCTCCAGGTTATTTACCCC  
CAGCCCAGGAACATTGGACTAATAGCTGGAGCCATTGGCACTGGTGCAGTTATTATC  
ATTTTTTGCATTGCACTAATTTTAGGGGCATTCTTTTACTGGAGAAGCAAAAATAAA  
GAGGAGGAAGAAGAAGAAATTCCTAATGAAATAAGAGAGGATGATCTTCCACCCAAG  
TGTTCTTCTGCCAAAGCATTTCACACTGAGATTTCTCCTCGGACAACAACACACTA  
ACCTCTTCCAATGCCTACAACAGTCGATACTGGAGCAACAATCCAAAAGTTCATAGA  
AACACAGAGTCAGTCAGCCACTTCAGTGACTTGGGCCAATCTTTCTCTTTCCACTCA  
GGCAATGCCAACATAACCATCCATTTATGCTAATGGGACCCATCTGGTCCCGGGTCAA  
CATAAGACTCTGGTAGTGACAGCCAACAGAGGGTCATCACCACAGGTGATGTCCAGG  
AGCAATGGCTCAGTCAGTAGGAAGCCTCGGCCTCCACACACTCATTCCTACACCATC  
AGCCACGCAACACTGGAACGAATTGGTGCAGTACCTGTCATGGTACCAGCCCAGAGT  
CGGGCCCGGGTCCCTTGGTATAGGACATGAGGAAATGTTGTGTTTCAGAAATGAATAAAT  
GGAATGCCCTCATAACAAGGGGGAGGGTGGGGTGGGGAGTGCTGGGAAAGAAACACTT  
CCTTATAATTATATTAGTAAAATGCACAAAGAAGAAGGCAGTGCTGTTACTTGGCCA  
CTAAGATGTGTAAAATGGACTGAAATGCTCCATCATGAAGACTTGCTTCCCCACCAA  
AGATGTCCTGGGATTCTGCTGGATCTCAAAGATGTGCCAAGCCAAGGAAAAAGATAC  
AAGAGCAGAATAGTACTTAAAATCCAAACTGCCGCCCAGATGGGCTTGTTCTTCATG  
CCTAACTTAATAATTTTTAAGAGATTAAAGTGCCAGATGGAGTTTAAATATTGAAAT  
TATTTTAAAGGTAGGTGTCTTTAAGAAAATAACAAGCAACCCTGTGATATGTTCCG  
TCTCTCCCAATTCCCTCGTTATATAGAGGGCTTAATGGTATAAATGGTTAATATTGG  
TCCCAACAGGGCTGACTCTTCTATCATATAATCAAACTTTTTTACATGAGCAAAATT  
CAGTAAGAAATGGGGGAAGACAAAGGAAACGTCTTTGAGAAGCCCCTTCATATTTAT  
TTATTTATCTCTTCCTGAACCATGAATTTTCATATGTGGAATATTGCTATATTGACAG  
ATTCTTGCCTGTCTGTGTTATTCTAGGATCTGTTACAGGTCCATGGCAATTACTGTT  
TATTTTTTCTGGA AAAAATATTTTTTTTATAAAAGGCTTTTTTTTTTTTTTAAATACA  
TGAGAGGCATTGGGCTAAGAAAGAAAAGACTGTTGTATAATACCTTGTTCAATGGTT  
GTATTTAGTGAGCTCATAGAGGTCCATCATATCATGACCGAGCTAGGTTGTGTGGGC  
AGGAAGGTAGGGCTAAGGGGTTGTAGCCTTGCTGGGCAGCCTCTCAGAGCAAGGTTG  
TTCAGATCTCCCTTGCTATTACAGTAGGTTACTATTAATGAGGGCAGCACCTGATGC  
CTTTTGTACTGAGGTATGTAAC TTTCTCCTTATTTGACAAGTAGAAGTTAACTTACT  
TGTCAGGGAGGGCAGACGTTTTTTTGTTCTGTTTCGTTTTTCAAATAATGCTTTTT  
GCAAAAGAGGTAAGACTGAGACTAAAGGTGTTATCTTCTGGTGTGCTCCTGGAAGTG  
TCTACCCTACATTTGTGTCAGCTCAGGGTTGCAGTGTTGCCCAGATGCATTTTACAT  
CACTGTAAAGAGATTACTTTTGTGGTTACTACCTGGCTTGGCTGGCCTTGCGGTTCA  
CCAGATTAATTTACAAACTCCCCCACTTTATTTTGTGCTATGTAGATCTGGCCATAC  
TTGCATTAGTGACTGTCTTGCCTTAACCACACTTAAGCAACCCACAAATTTCTTCTC  
AGATTTGTTTCCTAGATTACTTATGATACTCATCCCATGTCTCAATAAGAGTGTCTT  
TTCTTTCTGGATGTGTTCTCTTACTCCCTCTTACCACCATACTTTTTTGCTCTCTTCT  
CCTGCAAGCGTAGTCTTCACAGGGAGTGGCTTCCTGACATTTTTTTTCAGTTATGTGA  
ATGAATGGAACCAACAGCTGCTGCAAACACTGTTTTTCCAAGAAGGCTACACTCAG  
AACCTAACCATTGCCAACCATTTCAGTATTGATAAAAAGCTGAATTTACTTTAGCAT  
TACTTATTTTTTTTTCCATTTGATGGTTCTTACTTTGTAAAAATTTAAATAAATGAA  
TGTCTATACTTTTTATAAAGAAAAGTGAAAATACCATGACACTGAAAAGATGATGCT  
ATCAGATGCTGTTTAGAAAGCATTATCTTGCATTTCTTTATTCTTTCTAATTATCT  
AAAATTCAATAAAATTTTATTCATATAAAATAAGTTGTCATTAATTATCAATACTAA  
CGAGTATGTCATTTTAAAACTTAGTATTCTCTTTAATGTTACAAGA (SEQ ID NO: 73)

CGGCTGGTGGTGGCCGCGGCGGCCGGCGAGCCCCGGGACGCCCCGAGCCTGCCCCGAGC  
CTCGGCGGAGCGGAGTGCCCTCGGCGCTCCCGTGTCCTCGCTTGGTCCCACGCTGCAC  
CCCGCCGCCAGGAGCCCCGGCGGACGGCGGCTCCCCCGGCGGCTCCGGCATGACTCG  
GCGGCGCTCCGCTCCGGCGTCTTGGCTGCTCGTGTCGCTGCTCGGTGTCGCAACATC  
CCTGGAAGTGTCCGAGAGCCCAGGCAGTGTCCAGGTGGCCCCGGGGCCAGACAGCAGT  
CCTGCCCTGCGCCTTCTCCACCAGTGCTGCCCTCCTGAACCTCAATGTCATTTGGAT  
GGTCATTCCCCTCTCCAATGCAAACCAGCCCCGAACAGGTCATTCTTTATCAGGGTGG  
ACAAATGTTTGACGGCGCCCTCCGGTTCCACGGGAGGGTAGGATTTACCGGCACCAT  
GCCTGCTACCAATGTCTCGATCTTCATCAATAACACACAGCTGTCAGATACGGGCAC  
GTACCAGTGCTTGGTGAATAACCTTCCAGACAGAGGGGGCAGAAACATCGGGGGTCAC  
TGGCCTCACAGTGTTAGTCCCCCCTTCTGCTCCACAATGCCAAATCCAAGGATCCCA  
GGACCTCGGCAGTGACGTCATCCTTCTGTGTAGTTCAGAGGAAGGCATCCCTCGGCC  
CACGTACCTTTGGGAGAAGTTAGATAATACGCTCAAGCTACCTCCAACAGCCACTCA  
GGACCAGGTCCAGGGAACAGTCACCATCCGGAATATCAGTGCCCTCTCTTCCGGTCT  
GTACCAGTGTGTGGCTTCTAATGCCATCGGGACCAGCACCTGTCTGCTGGACCTCCA  
GGTTATCTCACCCAGCCCCGGAGCGTTGGAGTAATAGCCGGAGCGGTTGGCACCGG  
TGCTGTTCTTATCGTCATCTGCCTTGCCTAATTTAGGGGGCGTTCTTTTACTGGAG  
AAGCAAAAACAAAGAGGAGGAGGAGGAAGAAATTCCTAATGAAATCAGAGAGGATGA  
TCTTCCCCCTAAATGCTCTTCTGCCAAAGCCTTCCACACGGAGATATCCTCCTCAGA  
AAATAACACGCTGACCTCTTCCAATACCTACAACAGTCGATACTGGAACAACAATCC  
AAAACCCCATAGAAACACAGAGTCTTTCAACCACTTCAGTGACTTACGCCAGTCTTT  
CTCTGGCAATGCAGTTATCCCATCAATCTATGCAAATGGGAACCATCTGGTTTTGGG  
TCCACATAAGACTCTGGTAGTTACAGCCAACAGAGGGTCATCACCTCAGGTCTTGCC  
CAGGAACAATGGTTCAGTCAGCAGGAAGCCTTGGCCTCAACACACTCATTCCTACAC  
AGTAAGCCAAATGACCCTGGAGCGCATCGGTGCAGTGCCTGTCATGGTGCCTGCCCA  
GAGTCGAGCAGGGTCCCTGGTATAGGATGACTGAGGAAACCATGTTCAGAAGAGAAT  
AAATGGACCGCCTTCAGGCAAGGGGGGAGCACTGCCTTCAGGCAAGGGGGGAGCACT  
GCCTTCAGGCAAGAGGGAGAGTGGGATGGGTGAGTGCTGAAAAATAAACTTTTTGTTA  
CGATTCCATTAGCAAAAAGCACAAAGAGGAGGCGTGTGTGAAGTGGCCTGGGGTTGT  
TCCATAATGAAGACTCAAGAAGACTGTTTCCCCACCACAGATGTCCTGAGATTCAGT  
TAAAACGAAACATGCTGCATCTCCAGAGATGTGCCAAGCCAAGGAGAATGCTAGAAG  
CAGAGTAAAGCTTACCCCCCAAACCTGTGGTCCAGCTGGACCCCTTCTTTAATTCTTG  
CCTAACTTAATTATTTTCAGGACCCTTCAAGTGCCAGGTGGAATTTACATAATGAAA  
TTATTTTTTTAAAAATAGGTGTCCTTAGGGAGAGAAAACAGGAGCAAGCTCATGGTCT  
GGCCTAGTCTCCCTCTCCCACTCCTTCTGATGACACTAGCAATGCATTCCATCTGAC  
CTGACTTTATCATAGAGGCAAAATTGTTCAGAACACTGGCTGGAGATGGGGAGAAAT  
AAGGAAACTTCTTGTGAACACCCTACACACACACACACACACACACACACACACA  
CACACACACACACACACACACACACACACACACATTTATTTACCTCCTCCTGAACCATGAA

TCGTATTGGTGTGATTCTGTATTGACAGATTCTCATCTGTTACACTCTAGGATCTC  
TCACAGGTCTGTGGCAATTACTGTTTCATGATTTTCCTGAAAAAATATTTTTTTTAAAAG  
AAAACATATTTTTTTTTTAAATACTAGAGAGACAGTGGACTAGGAAAGCGAGAACTTGCC  
GCCTTGTCTAGTGACTGTATTCAATGACTGAACAGAGGCCCCCCCCCCACCATAACAAGA  
GTTTTAGGTGATTGAGTGGGTGGAACCAGCTGGAGCCAGGTGGGAGGGGGCCTTTACA  
TTGCCAGCAGGGCCCCCAAAGAATTGAGATTGTGTATGGCAACCGTTAATGAGGACAG  
CGCCTGATGCCTTTTTGTACCGAGGAAGATAATTGCCTCTTGTTTGACAAGTAGAGTT  
TAGTAGGTTATTACAAAAAGGGCAAGAGTTGTTTTGGTTTTGTTTCTTTCAAATAA  
TTTTTTTTTCAAAGAATAACAAGGGTTAGGCCAAATGGGGGACCTTCCTGTGTGCTCT  
TGGGGGTCTGCTCAGCATCTGGAATTTGGGTGTGCGATTTTCCCTGAACACATTGC  
ATACCAGTGTA AAAAGACTCTGCCTCCCCCTTTTTGGCTTTTTTACTGGGCTTGGC  
TGGCCTTGCAGTTTACCAGATTCATTTACAGACTCTCTGCTCTGTATGGCGCCGCCT  
GCCATGTCTGTCTTGGTGACTATCCTGCCTTAATCACTTTGCTTTAGGGCAACTCAT  
GGTGATCTCTTCCAAGATCTGTTTTTAAATTGTTTGGACTACTTGAGCCACA ACTCT  
CAGAGGACATTCCTTTTTTTTTTTTTTTTTTTTTCTCCTTTCTTCCATTGCTTTGTC  
CCTCTTCCCCTGTGCTTCCTGCCTTCTTTCCCTGTCCCATGGGCACAGTCCTCACAG  
GGAGTGGCCTCCTCTCTCCAGTGATGTAAGTGAATGGAAGCCATCACTGGCTGCACA  
TACCTTTTTTCAAAGGGGACACTCGGGAAGTCACTGCTGTGACCGTTTCGATGTTGAT  
AAGAAGGTGAATTTACTGTAGTGTTACCACCTTCTCCCCACTTGATGGTTCTTGACT  
TTGTAAAAAATTTAAATAAATGAATGTCTATACTTTTTTAAGGAAAAGAGAAAATACCA  
TGTCACAGAAAAGGTGAAACTATTAGATGCTGTTTAGAAAGCATTATCTTGCATTT  
CTTTATTCTTTCTAATTACCTAAAATTCAATAAAAGTTTATTCATATAAAAAAAAAA  
AAAAAAAAAAAA (SEQ ID NO: 75) >NP\_733548.2 immunoglobulin superfamily  
member 11 precursor [*Mus musculus*]

MTRRRSAPASWLLVSLLGVATSLEVSESPGSVQVARGQTAVLPCAFSTSAALLNLNV  
IWMVIPLSNANQPEQVILYQGGQMF DGALRFHGRVGF TGTMPATNVSIFINNTQLSD  
TGTYQCLVNNLPDRGGRNIGVTGLTVLVPPSAPQCQIQGSQDLGSDVILLCSSEGI  
PRPTYLWEKLDNTLKL PPTATQDQVQGTVTIRNISALSSGLYQCVASNAIGTSTCLL  
DLQVISPPQRSVGVIAGAVGTGAVLIVICLALISGAFFYWR SKNKEEEEEIEIPNEIR  
EDDLPPKCSSAKAFHTEISSSENNTLTSSNTYNSRYWNNNPKPHRNTESFNHFSDLR  
QSFSGNAVIPSIIYANGNHLVLGPHKTLVVTANRGSSPQVLPRNNGSVSRKPWPQH TH  
SYTVSQMTLERIGAVPVMVPAQSRAGSLV (SEQ ID NO: 76) Human VSIG4  
>NM\_007268.3 *Homo sapiens* V-set and immunoglobulin domain containing 4  
(VSIG4), transcript variant 1, mRNA

ACAGACGCTGGCGGCCACCAGAAGTTTGAGCCTCTTTGGTAGCAGGAGGCTGGAAGA  
AAGGACAGAAGTAGCTCTGGCTGTGATGGGGATCTTACTGGGCCTGCTACTCCTGGG  
GCACCTAACAGTGGACACTTATGGCCGTCCCATCCTGGAAGTGCCAGAGAGTGTAAC  
AGGACCTTGGAAGGGGATGTGAATCTTCCCTGCACCTATGACCCCTGCAAGGCTA  
CACCCAAGTCTTGGTGAAGTGGCTGGTACAACGTGGCTCAGACCCTGTCACCATCTT  
TCTACGTGACTCTTCTGGAGACCATATCCAGCAGGCAAAGTACCAGGGCCGCCTGCA  
TGTGAGCCACAAGGTTCCAGGAGATGTATCCCTCCAATTGAGCACCTGGAGATGGA  
TGACCGGAGCCACTACACGTGTGAAGTCACCTGGCAGACTCCTGATGGCAACCAAGT  
CGTGAGAGATAAGATTACTGAGCTCCGTGTCCAGAACTCTCTGTCTCCAAGCCCAC  
AGTGACAACTGGCAGCGGTTATGGCTTCACGGTGCCCCAGGGAATGAGGATTAGCCT  
TCAATGCCAGGCTCGGGGTTCTCCTCCCATCAGTTATATTTGGTATAAGCAACAGAC  
TAATAACCAGGAACCCATCAAAGTAGCAACCCTAAGTACCTTACTCTTCAAGCCTGC  
GGTGATAGCCGACTCAGGCTCCTATTTCTGCACTGCCAAGGGCCAGGTTGGCTCTGA  
GCAGCACAGCGACATTGTGAAGTTTGTGGTCAAAGACTCCTCAAAGCTACTCAAGAC  
CAAGACTGAGGCACCTACAACCATGACATACCCCTTGAAAGCAACATCTACAGTGAA  
GCAGTCCTGGGACTGGACCACTGACATGGATGGCTACCTTGGAGAGACCAGTGCTGG



GGCAGGAAAGAGCGTGGCTGTCTTGTCTCATCTTCTTGTCTGTAT  
GGTGGTTTTTACCATGGCCTATATCATGCTCTGTCTGGAAGACATCCCAACAAGAGCA  
TGTCTACGAAGCAGCCAGGGCACATGCCAGAGAGGCCAACGACTCTGGAGAAACCAT  
GAGGGTGGCCATCTTCGCAAGTGGCTGCTCCAGTGATGAGCCAACTTCCCAGAATCT  
GGGCAACAACACTACTCTGATGAGCCCTGCATAGGACAGGAGTACCAGATCATCGCCCA  
GATCAATGGCAACTACGCCCCGCTGCTGGACACAGTTCCTCTGGATTATGAGTTTCT  
GGCCACTGAGGGCAAAAGTGTCTGTAAAAAATGCCCCATTAGGCCAGGATCTGCTGA  
CATAATTGCCTAGTCAGTCCTTGCCTTCTGCATGGCCTTCTTCCCTGCTACCTCTCT  
TCCTGGATAGCCCAAAGTGTCCGCCTACCAACACTGGAGCCGCTGGGAGTCACTGGC  
TTTGCCCTGGAATTTGCCAGATGCATCTCAAGTAAGCCAGCTGCTGGATTGCTCT  
GGGCCCTTCTAGTATCTCTGCCGGGGGCTTCTGGTACTCCTCTCTAAATACCAGAGG  
GAAGATGCCCATAGCACTAGGACTTGGTCATCATGCCTACAGACACTATTCAACTTT  
GGCATCTTGCCACCAGAAGACCCGAGGGAGGCTCAGCTCTGCCAGCTCAGAGGACCA  
GCTATATCCAGGATCATTCTCTTTCTTCAGGGCCAGACAGCTTTTAATTGAAATTG  
TTATTTACAGGCCAGGGTTCAGTTCTGCTCCTCCACTATAAGTCTAATGTTCTGAC  
TCTCTCCTGGTGCTCAATAAATATCTAATCATAACAGCAA (SEQ ID NO: 77)

>NP\_009199.1 V-set and immunoglobulin domain-containing protein 4 isoform 1 precursor [*Homo sapiens*]

MGILLGLLLLGHLLTVDTYGRPILEVPESVTGPWKGDVNLPCITYDPLQGYTQVLVKWL  
VQRGSDPVTIFLRDSSGDHIQQAQYQGRHLVSHKVPDVSLSLSTLEMDDRSHYTCE  
VTWQTPDGNQVVRDKITELRVQKLSVSKPTVTTGSGYGFTVPQGMRISLQCQARGSP  
PISYIWYKQQTNNEPIKVATLSTLLFKPAVIADSGSYFCTAKGQVVGSEQHSDIVKF  
VVKDSSKLLKTKTEAPTMTYPLKATSTVKQSWDWTDMDGYLGETSAGPGKSLPVF  
AIIILSLCCMVVFTMAYIMLCRKTSQQEHVYEAARAHAREANDSGETMRVAIFASG  
CSSDEPTSQNLGNNYSDEPCIGQEYQIIAQINGNYARLLDTPLDYEFATEGKSVC (SEQ  
ID NO: 78) Mouse VSIG4 >NM\_177789.5 *Mus musculus* V-set and  
immunoglobulin domain containing 4 (Vsig4), mRNA

AGCTACCAGCACTTCCAGGTTCTTCAGCAGCAAGAGGATGGAAGGATGAATAGAAGT  
AGCTTCAAATAGGATGGAGATCTCATCAGGCTTGCTGTTTCTGGGCCACCTAATAGT  
GCTCACCTATGGCCACCCCACCCTAAAAACACCTGAGAGTGTGACAGGGACCTGGAA  
AGGAGATGTGAAGATTCAGTGCATCTATGATCCCCTGAGAGGCTACAGGCAAGTTTT  
GGTGAAATGGCTGGTAAGACACGGCTCTGACTCCGTCACCATCTTCTCTACGTGACTC  
CACTGGAGACCATATCCAGCAGGCAAAGTACAGAGGCCGCTGAAAGTGAGCCACAA  
AGTTCCAGGAGATGTGTCCCTCCAAATAAATACCCTGCAGATGGATGACAGGAATCA  
CTATACATGTGAGGTCACCTGGCAGACTCCTGATGGAAACCAAGTAATAAGAGATAA  
GATCATTGAGCTCCGTGTTTCGGAAATATAATCCACCTAGAATCAATACTGAAGCACC  
TACAACCCTGCACTCCTCTTTGGAAGCAACAACCTATAATGAGTTCAACCTCTGACTT  
GACCACTAATGGGACTGGAAAACCTTGAGGAGACCATTGCTGGTTCAGGGAGGAACCT  
GCCAATCTTTGCCATAATCTTCATCATCTCCCTTTGCTGCATAGTAGCTGTCACCAT  
ACCTTATATCTTGTTCCGCTGCAGGACATTCCAACAAGAGTATGTCTATGGAGTGAG  
CAGGGTGTTTGCCAGGAAGACAAGCAACTCTGAAGAAACCACAAGGGTGACTACCAT  
CGCAACTGATGAACCAGATTCCCAGGCTCTGATTAGTGACTACTCTGATGATCCTTG  
CCTCAGCCAGGAGTACCAAATAACCATCAGATCAACAATGTCTATTCTGCTGCTG  
AACACAGTTTCCAGAACTAAGAAGTTCTTGCTACTGAAGAAAATAACATCTGCTAA  
AATGCCCCTACTAAGTCAAGGTCTACTGGCGTAATTACCTGTTACTTATTTACTACT  
TGCCTTCAACATAGCTTTCTCCCTGGCTTCCTTTCTTCTTAGACAACCTAAAGTATC  
TATCTAGTCTGCCAATTCTGGGGCCATTGAGAAATCCTGGGTTTGGCTAAGAATATA  
CTACATGCACCTCAAGAAATCTAGCTTCTGGGCTTCACCCAGAACAAATTTTCTTCT  
AGGGCCTTCACAACCTCTTCTCCAAACAGCAGAGAAATTCCATAGCAGTAGAGGTTCT  
TTATCATGCCTCCAGACAGCGTGAGTCTCAGTCCTACAAACTCAGACAAGCACATGG

GCTCTAGGATTACTCTCTTTCTCTCTAGGGCCAGATGACTTTTATGATATTACTATT  
GCTACATTATGAATCTAATGCACATGTATTCTTTTGTGTTAATAAATGTTTAATCA  
TGACATCAA(SEQ ID NO: 79)>NP\_808457.1 V-set and immunoglobulin domain-  
containing protein 4 precursor [*Mus musculus*]  
MEISSGLLFLGHLIVLTYGHPTLKTPESVTGTWKGDVKIQCIYDPLRGYRQVLVKWL  
VRHGSDSVTIFLRDSTGDHIQQAKYRGRLKVSHKVPGDVSLQINTLQMDDRNHYTCE  
VTWQTPDGNQVIRDKIIELRVRKYNPPRINTEAPTTLHSSLEATTIMSSSTSDLTTNG  
TGKLEETIAGSGRNLPIFAIIFIISLCCIVAVTIPYILFRCRTFQQEYVYGVSRVFA  
RKTSNSEETTRVTTIATDEPDSQALISDYSDDPCLSQEYQITIRSTMSIPAC (SEQ ID  
NO: 80) Human Tim-3>NM\_032782.5 *Homo sapiens* hepatitis A virus cellular  
(HAVCR2) receptor 2 (HAVCR2), mRNA  
ATTTGGAGAGTTAAACTGTGCCTAACAGAGGTGTCCTCTGACTTTTCTTCTGCAAG  
CTCCATGTTTTTCACATCTTCCCTTTGACTGTGTCTGCTGCTGCTGCTGCTACTACT  
TACAAGGTCCTCAGAAGTGGAATACAGAGCGGAGGTTCGGTCAGAATGCCTATCTGCC  
CTGCTTCTACACCCCAGCCGCCCCAGGGAACCTCGTGCCCGTCTGCTGGGGCAAAGG  
AGCCTGTCCTGTGTTTGAATGTGGCAACGTGGTGCTCAGGACTGATGAAAGGGATGT  
GAATTATTGGACATCCAGATACTGGCTAAATGGGGATTTCCGCAAAGGAGATGTGTC  
CCTGACCATAGAGAATGTGACTCTAGCAGACAGTGGGATCTACTGCTGCCGGATCCA  
AATCCCAGGCATAATGAATGATGAAAAATTTAACCTGAAGTTGGTCATCAAACCAGC  
CAAGGTCACCCCTGCACCGACTCGGCAGAGAGACTTCACTGCAGCCTTTCCAAGGAT  
GCTTACCACCAGGGGACATGGCCCAGCAGAGACACAGACACTGGGGAGCCTCCCTGA  
TATAAATCTAACACAAATATCCACATTGGCCAATGAGTTACGGGACTCTAGATTGGC  
CAATGACTTACGGGACTCTGGAGCAACCATCAGAATAGGCATCTACATCGGAGCAGG  
GATCTGTGCTGGGCTGGCTCTGGCTCTTATCTTCGGCGCTTTAATTTTCAAATGGTA  
TTCTCATAGCAAAGAGAAGATACAGAATTTAAGCCTCATCTCTTTGGCCAACCTCCC  
TCCCTCAGGATTGGCAAATGCAGTAGCAGAGGGAATTCGCTCAGAAGAAAACATCTA  
TACCATTGAAGAGAACGTATATGAAGTGGAGGAGCCCAATGAGTATTATTGCTATGT  
CAGCAGCAGGCAGCAACCCTCACAACCTTTGGGTTGTGCTTTTGCAATGCCATAGAT  
CCAACCACCTTATTTTTGAGCTTGGTGTTTTGTCTTTTTTCAGAACTATGAGCTGTG  
TCACCTGACTGGTTTTTGAGGTTCTGTCCACTGCTATGGAGCAGAGTTTTTCCCATTT  
TCAGAAGATAATGACTCACATGGGAATTGAACTGGGACCTGCACTGAACTTAAACAG  
GCATGTCATTGCCTCTGTATTTAAGCCAACAGAGTTACCCAACCCAGAGACTGTAA  
TCATGGATGTTAGAGCTCAAACGGGCTTTTATATACACTAGGAATTCTTGACGTGGG  
GTCTCTGGAGCTCCAGGAAATTCGGGCACATCATATGTCCATGAACTTCAGATAAA  
CTAGGGAAACTGGGTGCTGAGGTGAAAGCATAACTTTTTTGGCACAGAAAGTCTAA  
AGGGGCCACTGATTTTCAAAGAGATCTGTGATCCCTTTTTTGTTTTTTGTTTTTGAGA  
TGGAGTCTTGCTCTGTTGCCCAGGCTGGAGTGCAATGGCACAATCTCGGCTCACTGC  
AAGCTCCGCCTCCTGGGTTCAAGCGATTCTCCTGCCTCAGCCTCCTGAGTGGCTGGG  
ATTACAGGCATGCACCACCATGCCCAGCTAATTTGTTGTATTTTTAGTAGAGACAGG  
GTTTCACCATGTTGGCCAGTGTGGTCTCAAACCTCCTGACCTCATGATTTGCCTGCCT  
CGGCCTCCCAAAGCACTGGGATTACAGGCGTGAGCCACCACATCCAGCCAGTGATCC  
TTAAAAGATTAAAGAGATGACTGGACCAGGTCTACCTTGATCTTGAAGATTCCCTTG  
AATGTTGAGATTTAGGCTTATTTGAGCACTGCCTGCCCAACTGTCAGTGCCAGTGCA  
TAGCCCTTCTTTTGTCTCCCTTATGAAGACTGCCCTGCAGGGCTGAGATGTGGCAGG  
AGCTCCCAGGGAAAAACGAAGTGCAATTTGATTGGTGTGTATTGGCCAAGTTTTGCTT  
GTTGTGTGCTTGAAAGAAAATATCTCTGACCAACTTCTGTATTCGTGGACCAAACCTG  
AAGCTATATTTTTACAGAGAAGAAGAAGCAGTGACGGGGACACAAATTCTGTTGCCTG  
GTGGAAAGAAGGCAAAGGCCTTCAGCAATCTATATTACCAGCGCTGGATCCTTTGAC  
AGAGAGTGGTCCCTAAACTTAAATTTCAAGACGGTATAGGCTTGATCTGTCTTGCTT  
ATTGTTGCCCCCTGCGCCTAGCACAATTCTGACACACAATTGGAACCTACTAAAAAT

TTTTTTTACTGTT (SEQ ID NO: 81) >NP\_116171.3 hepatitis A virus cellular  
receptor 2 precursor [*Homo sapiens*]

MFSHLPFDCVLLLLLLLLLRSSEVEYRAEVGQNAYLPCFYTPAAPGNLVPVCWGKGA  
CPVFECGNVVLRTDERDVNYWTSRYWLNNGDFRKGDVSLTIENVTLADSGIYCCRIQI  
PGIMNDEKFNLKLVIPAKVTPAPTRQRDFTAAFPRLTTRGHGPAETQTLGSLPDI  
NLTQISTLANELRDSRLANDLRDSGATIRIGIYIGAGICAGLALALIFGALIFKWYS  
HSKEKIQNLSLISLANLPPSGLANAVAEGIRSEENIYTIENVYEEVEEPNEYCYVS  
SRQQPSQPLGCRFAMP (SEQ ID NO: 82) Mouse Tim-3 >NM\_134250.2 *Mus  
musculus* hepatitis A virus cellular (HAVCR2) receptor 2 (Havcr2), mRNA  
ACCATTTTAAACCGAGGAGCTAAAGCTATCCCTACACAGAGCTGTCCTTGGATTTCCTC  
CTGCCAAGTACTCATGTTTTTCAGGTCTTACCCTCAACTGTGTCCTGCTGCTGCTGCA  
ACTACTACTTGCAAGGTCATTGGAAAATGCTTATGTGTTTGAGGTTGGTAAGAATGC  
CTATCTGCCCTGCAGTTACACTCTATCTACACCTGGGGCAGTTGTGCCTATGTGCTG  
GGGCAAGGGATTCTGTCCTTGGTCACAGTGTACCAACGAGTTGCTCAGAACTGATGA  
AAGAAATGTGACATATCAGAAATCCAGCAGATACCAGCTAAAGGGCGATCTCAACAA  
AGGAGACGTGTCTCTGATCATAAAGAATGTGACTCTGGATGACCATGGGACCTACTG  
CTGCAGGATACAGTTCCCTGGTCTTATGAATGATAAAAAATTAGAACTGAAATTAGA  
CATCAAAGCAGCCAAGGTCCTCCAGCTCAGACTGCCCATGGGGACTCTACTACAGC  
TTCTCCAAGAACCCTAACCACGGAGAGAAATGGTTCAGAGACACAGACACTGGTGAC  
CCTCCATAATAACAATGGAACAAAAATTTCCACATGGGCTGATGAAATTAAGGACTC  
TGGAGAAACGATCAGAACTGCTATCCACATTGGAGTGGGAGTCTCTGCTGGGTTGAC  
CCTGGCACTTATCATTGGTGTCTTAATCCTTAAATGGTATTCCTGTAAGAAAAAGAA  
GTTATCGAGTTTGAGCCTTATTACACTGGCCAACCTGCCTCCAGGAGGGTTGGCAAA  
TGCAGGAGCAGTCAGGATTCGCTCTGAGGAAAATATCTACACCATCGAGGAGAACGT  
ATATGAAGTGGAGAATTCAAATGAGTACTACTGCTACGTCAACAGCCAGCAGCCATC  
CTGACCGCCTCTGGACTGCCACTTTTAAAGGCTCGCCTTCATTTCTGACTTTGGTAT  
TTCCCTTTTTTGAAAACCTATGTGATATGTCACTTGGCAACCTCATTGGAGGTTCTGAC  
CACAGCCACTGAGAAAAGAGTTCCAGTTTTCTGGGGATAATTAACCTCACAAGGGGAT  
TCGACTGTAACCTCATGCTACATTGAAATGCTCCATTTTATCCCTGAGTTTCAGGGAT  
CGGATCTCCCACTCCAGAGACTTCAATCATGCGTGTTGAAGCTCACTCGTGCTTTCA  
TACATTAGGAATGGTTAGTGTGATGTCTTTGAGACATAGAGGTTTGTGGTATATCTG  
CAAAGCTCCTGAACAGGTAGGGGGAATAAAGGGCTAAGATAGGAAGGTGAGGTTCTT  
TGTTGATGTTGAAAATCTAAAGAAGTTGGTAGCTTTTCTAGAGATTTCTGACCTTGA  
AAGATTAAGAAAAAGCCAGGTGGCATATGCTTAACACTATATAACTTGGGAACCTTA  
GGCAGGAGGGTGATAAGTTCAAGGTCAGCCAGGGCTATGCTGGTAAGACTGTCTCAA  
AATCCAAAGACGAAAATAAACATAGAGACAGCAGGAGGCTGGAGATGAGGCTCGGAC  
AGTGAGGTGCATTTTGTACAAGCACGAGGAATCTATATTTGATCGTAGACCCACAT  
GAAAAAGCTAGGCCTGGTAGAGCATGCTTGTAGACTCAAGAGATGGAGAGGTAAGG  
CACAACAGATCCCCGGGGCTTGCCTGCAGTCAGCTTAGCCTAGGTGCTGAGTTCCAA  
GTCCACAAGAGTCCCTGTCTCAAAGTAAGATGGACTGAGTATCTGGCGAATGTCCAT  
GGGGGTTGTCCTCTGCTCTCAGAAGAGACATGCACATGAACCTGCACACACACACAC  
ACACACACACACACACACACACACACACACACACACATGAAATGAAGGTTT  
TCTCTGTGCCTGCTACCTCTCTATAACATGTATCTCTACAGGACTCTCCTCTGCCTC  
TGTTAAGACATGAGTGGGAGCATGGCAGAGCAGTCCAGTAATTAATTCCAGCACTCA  
GAAGGCTGGAGCAGAAGCGTGGAGAGTTCAGGAGCACTGTGCCCAACACTGCCAGAC  
TCTTCTTACAGAAGAAAAAGGTTACCCGCAAGCAGCCTGCTGTCTGTAAAAGGAAAC  
CCTGCGAAAGGCAAACCTTTGACTGTTGTGTGCTCAAGGGGAACTGACTCAGACAACT  
TCTCCATTCTGGAGGAACTGGAGCTGTTTCTGACAGAAGAACAACCGGTGACTGG  
GACATACGAAGGCAGAGCTCTTGCAGCAATCTATATAGTCAGCAAAATATTCTTTGG  
GAGGACAGTCGTCACCAAATTGATTTCCAAGCCGGTGGACCTCAGTTTCATCTGGCT

TACAGCTGCCTGCCAGTGGCCTTGATCTGTGCTGGCTCCCATCTATAACAGAAATCA  
AATTAAATAGACCCCGAGTGAAAATATTAAGTGAGCAGAAAGGTAGCTTTTGTTCAAA  
GATTTTTTTTGCATTGGGGAGCAACTGTGTACATCAGAGGACATCTGTTAGTGAGGAC  
ACCAAAACCTGTGGTACCGTTTTTTCATGTATGAATTTTGTGTGTTAGGTTGCTTCT  
AGCTAGCTGTGGAGGTCCTGGCTTTCTTAGGTGGGTATGGAAGGGAGACCATCTAAC  
AAAATCCATTAGAGATAACAGCTCTCATGCAGAAGGGGAAAATAATCTCAAATGTTT  
TAAAGTAATAAACTGTACTGGCAAAGTACTTTGAGCATATTTAAA (SEQ ID NO:  
83)>NP\_599011.2 hepatitis A virus cellular receptor 2 homolog precursor [*Mus  
musculus*]

MFSGLTLNVCVLLLLQLLLARSLENAYVFEVGKNAYLPCSYTLSTPGALVPMCWGKGF  
CPWSQCTNELLRTDERNVTYQKSSRYQLKGDNLKGDVSLIKNVTLDDHGTYYCCRIQ  
FPGLMNDKKLELKLDIKAAKVTPAQTAHGDSTTASPRTLTERNGSETQTLVTLHNN  
NGTKISTWADEIKDSGETIRTAIHIGVGVSAGLTLALIIGVLILKWYSCKKKKLSSL  
SLITLANLPPGGLANAGAVRIRSEENIYTIENNVYEVENSNYYCYVNSQQPS (SEQ ID  
NO: 84) Human Tim-4 >NM\_138379.3 *Homo sapiens* T cell immunoglobulin  
and mucin (TIMD4) domain containing 4 (TIMD4), transcript variant 1, mRNA  
AGACTCCTGGGTCCGGTCAACCGTCAAAATGTCCAAAGAACCTCTCATTCTCTGGCT  
GATGATTGAGTTTTTGGTGGCTTTACCTGACACCAGTCACTTCAGAGACTGTTGTGAC  
GGAGGTTTTTGGGTCACCGGGTGACTTTGCCCTGTCTGTACTCATCCTGGTCTCACAA  
CAGCAACAGCATGTGCTGGGGGAAAGACCAGTGCCCTACTCCGGTTGCAAGGAGGC  
GCTCATCCGCACTGATGGAATGAGGGTGACCTCAAGAAAGTCAGCAAAATATAGACT  
TCAGGGGACTATCCCGAGAGGTGATGTCTCCTTGACCATCTTAAACCCCAGTGAAAG  
TGACAGCGGTGTGTACTGCTGCCGCATAGAAGTGCCTGGCTGGTTCAACGATGTAAA  
GATAAACGTGCGCCTGAATCTACAGAGAGCCTCAACAACCACGCACAGAACAGCAAC  
CACCACCACACGCAGAACAACAACAAGCCCCACCACCACCCGACAAATGACAAC  
AACCCCAGCTGCACTTCCAACAACAGTCGTGACCACACCCGATCTCACAACCGGAAC  
ACCACTCCAGATGACAACCATTTGCCGTCTTCACAACAGCAAACACGTGCCTTTCACT  
AACCCCAAGCACCTTCCGGAGGAAGCCACAGGTCTTCTGACTCCCGAGCCTTCTAA  
GGAAGGGGCCCATCCTCACTGCAGAATCAGAACTGTCCTCCCCAGTGATTCTTGAG  
TAGTGTTGAGTCTACTTCTGCTGACACTGTCCTGCTGACATCCAAAGAGTCCAAAGT  
TTGGGATCTCCCATCAACATCCCACGTGTCAATGTGGAAAACGAGTGATTCTGTGTC  
TTCTCCTCAGCCTGGAGCATCTGATACAGCAGTTCCTGAGCAGAACAAAACAACAAA  
AACAGGACAGATGGATGGAATACCCATGTCAATGAAGAATGAAATGCCCATCTCCCA  
ACTACTGATGATCATCGCCCCCTCCTTGGGATTTGTGCTCTTCGCATTGTTTGTGGC  
GTTTCTCCTGAGAGGGAACTCATGGAAACCTATTGTTTCGCAGAAACACACAAGGCT  
AGACTACATTGGAGATAGTAAAAATGTCCTCAATGACGTGCAGCATGGAAGGGGAAGA  
CGAAGACGGCCTTTTTACCCTCTAACAACGCAGTAGCATGTTAGATTGAGGATGGGG  
GCATGACACTCCAGTGTCAAATAAGTCTTAGTAGATTTCCTTGTTTCATAAAAAAG  
ACTCACTTATTCCATGGATGTCATTGATCCAGGCTTGCTTTAGTTTCATGAATGAAG  
GGTACTTTAGAGACCACAA (SEQ ID NO: 85)>NP\_612388.2 T-cell  
immunoglobulin and mucin domain-containing protein 4 isoform 1 precursor  
[*Homo sapiens*]

MSKEPLILWLMIEFWWLYLTPVTSETVVTEVLGHRVTLPCLYSSWSHNSNSMCWGKD  
QCPYSGCKEALIRTDGMRVTSRKSAYRLQGTIPRGDVSILTILNPSESDSGVYCCRI  
EVPGWFNVDVKINVRNLQRASTTTHRTATTTTRRTTTTSPTTTTRQMTTTPAALPTTV  
VTTPDLTTGTPLQMTTIAVFTTANTCLSLTPSTLPPEATGLLTPEPSKEGPILTAES  
ETVLPSDSWSSVESTSADTVLLTSKESKVWDL PSTSHVSMWKTSDSVSSPQPGASDT  
AVPEQNKTTKTGQMDGIPMSMKNEPISQLLMIIAPSLGFLVLFALFVAFLLRGKLME  
TYCSQKHTRLDYIGDSKNVLNDVQHGREDGLFTL (SEQ ID NO: 86) Mouse  
Tim-4 >NM\_178759.4 *Mus musculus* T cell immunoglobulin and mucin (TIMD4)

domain containing 4 (Timd4), mRNA

AGATCCTATCAAAATGTCCAAGGGGCTTCTCCTCCTCTGGCTGGTGACGGAGCTCTG  
GTGGCTTTATCTGACACCAGCTGCCTCAGAGGATACAATAATAGGGTTTTTGGGCCA  
GCCGGTGACTTTGCCTTGTCTTACCTCTCGTGGTCCCAGAGCCGCAACAGTATGTG  
CTGGGGCAAAGGTTTCATGTCCCAATTCCAAGTGCAATGCAGAGCTTCTCCGTACAGA  
TGGAACAAGAATCATCTCCAGGAAGTCAACAAAATATACACTTTTGGGGGAAGGTCCA  
GTTTGGTGAAGTGTCTTGACCATCTCAAACACCAATCGAGGTGACAGTGGGGTGTA  
CTGCTGCCGTATAGAGGTGCCTGGCTGGTTCAATGATGTCAAGAAGAATGTGCGCTT  
GGAGCTGAGGAGAGCCACAACAACCAAAAAACCAACAACAACCACCCGGCCAACCAC  
CACCCCTTATGTGACCACCACCACCCACAGAGCTGCTTCCAACAACAGTCATGACCAC  
ATCTGTTCTCCCAACCACCACACCACCCACAGACACTAGCCACCACTGCCTTCAGTAC  
AGCAGTGACCACGTGCCCCCTCAACAACACCTGGCTCCTTCTCACAAGAAACCACAAA  
AGGGTCCGCCTTCACTACAGAATCAGAACTCTGCCTGCATCCAATCACTCTCAAAG  
AAGCATGATGACCATATCTACAGACATAGCCGTACTCAGGCCACAGGCTCTAACCC  
TGGGATTCTCCCATCCACTTCACAGCTGACGACACAGAAAACAACATTAACAACAAG  
TGAGTCTTTGCAGAAGACAATAATCACATCAGATCAACAGCAGACAGACCATCTT  
GATCATTGCCTGCTGTGTGGGATTTGTGCTAATGGTGTATTGTTTCTGGCGTTTCT  
CCTTCGAGGGAAAGTCACAGGAGCCAACCTGTTTGCAGAGACACAAGAGGCCAGACAA  
CACTGAAGATAGTGACAGCGTCCTCAATGACATGTCACACGGGAGGGATGATGAAGA  
CGGGATCTTCACTCTCTGACTCACCATCTTTATTTAGGATTAAGGATAGGGAATGGC  
ACTTGAATTGTCAAATAAGTTTGGGGACATTGTAATTTCCGTTTTAAAGTCTCACTC  
TGTTTACTGATGCTGTGGGTCTGTCTGGTTGTATCTTCCCACATGAAGGTGCTTIA  
GAGACACATTTTCCCTGCCTCGTGCCTTAGTCCTCTTTGTTGTTTTGTGGCTAGGTG  
ACTTTTCACACTGGGCTTGAACACTGTCAGTGATGGTGAAATCCTTGCCACAGCTTT  
GGGAGTCTCTTGCAGTCTCCCAGCAGTAGAGGGAGTTAGAAATATCCAGAGGGGAAA  
AAAAAATCTCTCTTTTTCAGACAGTATCTGCTTTATTGGTGGTAGCTGAACTTCATTT  
ATACAGAGCTCCTTTAACCTGTCTGTCTTCTTCTTGGTATCTAAGCTGCCTTTTGT  
TTTGTTTTTGTTTTTGTTTTTATGATATTAACCTCTTTTTCACATTCAAGTTTCTTTA  
AAGTTGACTATAGTGCCTTCTGAACTCTTGCAGAGAGTTTGGATTTTGGAAAGCTGCC  
AGGTACCCATCACAGCAGGGGTGCCAGTGACAAGGATGGTGTACAAATGAAACACTG  
AAGCTATCCAAATAAATTCCTCTAAGTGTAATTCATTTTACTGCAGCACAGGAAGAA  
CAAATTTGTCTTACAACTTTAATAATTAGTACCATTATGAACCCTAGGAGAGAAATA  
AGAGCAAATACCTGTTGAATAAATGAATGTAAGAAAATGTGTGTCTGAGCAAGAATA  
CTCTGTCTGGCTACTATGGGAAGCTAGCTAGATCTGAAAGACATTCTCAGACTATCC  
TCATGTTCAAGGCATTAAAGGAATAAGCCTCCAGCCCCTAACCTTAGGAGAATTCTG  
CAGTCAAGTGAGGAGTTTTTAAAACAGGAATCTCTAGGTTCCAGTCCTCTAGCTATT  
CTTTTATGCTTAGTCCAGGTAATGAGTTGAACATCCAAGTATTTTTTAAGGACCCAA  
AGAAATGCAACCAGAGCTATTACCAGAATTTTGGAGTGGTCCTCCTAGAGTTGCCGC  
ATGTTGCTGGGAAAATTGGGGTCTTAGAGTTCTTAGTCTACTTAATAAAAGAATTTT  
AAAAAATGG (SEQ ID NO: 87) >NP\_848874.3 T-cell immunoglobulin and

mucin domain-containing protein 4 precursor [*Mus musculus*]

MSKGLLLLWLVTTELWWLYLTPAASEDTIIGFLGQPVTLPCHYLSWSQSRNSMC  
WGKGSCPNKCNALLRTDGTRIISRKSTKYTLLGKVQFGEVSLTISNTNRGD  
SGVYCCRIEVPWFNDVKKNVRLRRATTTKKPTTTTRPTTTPYVTTTTPEL  
LPTTVMTTSVLPPTTTPPQTLATTAFASTAVTTCPTSTPGSFSQETTKGSFTTE  
SETLPASNHSQRSMMTISTDIAVLRPTGSNPGILPSTSQTTLTQKTTLTSES  
LQKTTKSHQINSRQTILIIACCVGFVLMVLLFLAFLLRGKVTGANCLQRHKRPD  
NTEDSDSVLNDMSHGRDDEGIFTL (SEQ ID NO: 88) Human >NM\_001712.5

*Homo sapiens* CEA cell adhesion molecule CEACAM1 1 (CEACAM1), transcript  
variant 1, mRNA

ACGACAGAGTAGAGTAGTGGAAAGAACAGCAGAGGAGGTGACAGAGCAGGCGCTGCTCGAAGCGT  
TCCTGGAGCCCAAGCTCTCCTCCACAGGTGAAGACAGGGGCCAGCAGGAGACAC  
CATGGGGCACCTCTCAGCCCCACTTCACAGAGTGCCTGTACCCTGGCAGGGGC  
TTCTGCTCACAGCCTCACTTCTAACCTTCTGGAACCCGCCACCCTGCCCAG  
CTCACTACTGAATCCATGCCATTCAATGTTGCAGAGGGGAAGGAGGTTCTTCT  
CCTTGTCCACAATCTGCCCCAGCAACTTTTTTGGCTACAGCTGGTACAAAGGGG  
AAAGAGTGATGGCAACCGTCAAATTGTAGGATATGCAATAGGAACTCAACAA  
GCTACCCCAAGGGCCCCGCAAACAGCGGTGCGAGAGACAATATACCCCAATGCATC  
CCTGCTGATCCAGAACGTACCCAGAATGACACAGGATTCTACACCCTACAAG  
TCATAAAGTCAGATCTTGTGAATGAAGAAGCAACTGGACAGTTCCATGTATAC  
CCGGAGCTGCCCCAAGCCCTCCATCTCCAGCAACAACCTCCAACCCTGTGGAGGA  
CAAGGATGCTGTGGCCTTCACCTGTGAACCTGAGACTCAGGACACAACCTACC  
TGTGGTGGATAAACAATCAGAGCCTCCCGGTGAGTCCCAGGCTGCAGCTGTCC  
AATGGCAACAGGACCCTCACTCTACTCAGTGTCAACAAGGAATGACACAGGACC  
CTATGAGTGTGAAATACAGAACCCAGTGAGTGCGAACCGCAGTGACCCAGTCA  
CCTTGAATGTCACCTATGGCCCCGGACACCCCCACCATTTCCTTTCAGACACC  
TATTACCGTCCAGGGGGCAAACCTCAGCCTCTCCTGCTATGCAGCCTCTAACCC  
ACCTGCACAGTACTCCTGGCTTATCAATGGAACATTCCAGCAAAGCACACAAG  
AGCTCTTTATCCCTAACATCACTGTGAATAATAGTGGATCCTATACCTGCCAC  
GCCAATAACTCAGTCACTGGCTGCAACAGGACCACAGTCAAGACGATCATAGT  
CACTGAGCTAAGTCCAGTAGTAGCAAAGCCCCAAATCAAAGCCAGCAAGACCA  
CAGTCACAGGAGATAAGGACTCTGTGAACCTGACCTGCTCCACAAATGACACT  
GGAATCTCCATCCGTTGGTTCTTCAAAAACCAGAGTCTCCCGTCTCTCGGAGAG  
GATGAAGCTGTCCCAGGGGCAACACCACCCTCAGCATAAACCTGTCAAGAGGG  
AGGATGCTGGGACGTATTGGTGTGAGGTCTTCAACCCAATCAGTAAGAACCAA  
AGCGACCCCATCATGCTGAACGTAAACTATAATGCTCTACCACAAGAAAATGG  
CCTCTCACCTGGGGCCATTGCTGGCATTGTGATTGGAGTAGTGGCCCTGGTTG  
CTCTGATAGCAGTAGCCCTGGCATGTTTTCTGCATTTCTGGGAAGACCGGCAGG  
GCAAGCGACCAGCGTGATCTCACAGAGCACAAACCCTCAGTCTCCAACCACAC  
TCAGGACCACTCCAATGACCCACCTAACAAGATGAATGAAGTTACTTATTCTA  
CCCTGAACTTTGAAGCCCAGCAACCCACACAACCAACTTCAGCCTCCCCATCC  
CTAACAGCCACAGAAATAATTTATTCAGAAGTAAAAAAGCAGTAATGAAACCT  
GTCCTGCTCACTGCAGTGCTGATGTATTTCAAGTCTCTCACCCTCATCACTAG  
GAGATTCTTTTCCCTGTAGGGGTAGAGGGGTGGGGACAGAAACAACCTTTCTC  
CTACTCTTCCTTCCTAATAGGCATCTCCAGGCTGCCTGGTCACTGCCCCTCTC  
TCAGTGTCAATAGATGAAAGTACATTGGGAGTCTGTAGGAAACCCAACCTTCT  
TGTCATTGAAATTTGGCAAAGCTGACTTTGGGAAAGAGGGGACCAGAACTTCCC  
CTCCCTTCCCCTTTTCCCAACCTGGACTTGTTTTAACTTGCCTGTTTCAGAGC  
ACTCATTCTTCCCACCCCCAGTCCTGTCCTATCACTCTAATTCGGATTGTC  
ATAGCCTTGAGGTTATGTCCTTTTCCATTAAGTACATGTGCCAGGAAACAAGA  
GAGAGAGAAAGTAAAGGCAGTAATGCCTTCTCCTATTTCTCCAAAGCCTTGTG  
TGAATCACCAAACACAAGAAAATCAAATATATAACCAATAGTGAAATGCCAC  
ACCTTTGTCCACTGTCAGGGTTGTCTACCTGTAGGATCAGGGTCTAAGCACCT  
TGGTGTCTTAGCTAGAATACCACCTAATCCTTCTGGCAAGCCTGTCTTCAGAGA  
ACCCACTAGAAGCAACTAGGAAAATCACTTGCCAAAATCCAAGGCAATTCCTG  
ATGGAAAATGCAAAAGCACATATATGTTTTAATATCTTTATGGGCTCTGTTCA  
AGGCAGTGCTGAGAGGGAGGGGTATAGCTTCAGGAGGGAACCAGCTTCTGAT  
AAACACAATCTGCTAGGAACTTGGGAAAGGAATCAGAGAGCTGCCCTTCAGCG  
ATTATTTAAATTATTGTTAAAGAATACACAATTTGGGGTATTGGGATTTTCT  
CCTTTTCTCTGAGACATTCCACCATTTTAATTTTTGTAAGTCTTATTTATGT

GGTAAAGGTTATTATTAGTCTAGCTCAATCCGTAATCCGCTTA  
GGTGAAAGAAACCACCGAAATCCCTCAGGTCCCTTGGTCAGGAGCCTCTCAAG  
ATTTTTTTTGTGTCAGAGGCTCCAAATAGAAAATAAGAAAAGGTTTTCTTCATTC  
ATGGCTAGAGCTAGATTTAACTCAGTTTCTAGGCACCTCAGACCAATCATCAA  
CTACCATTCTATTCCATGTTTGCACCTGTGCATTTTCTGTTTGCCCCCATTCA  
CTTTGTGTCAGGAAACCTTGGCCTCTGCTAAGGTGTATTTGGTCCTTGAGAAGTG  
GGAGCACCTACAGGGACACTATCACTCATGCTGGTGGCATTGTTTACAGCTA  
GAAAGCTGCACTGGTGCTAATGCCCCTTGGGGAAATGGGGCTGTGAGGAGGAG  
GATTATAACTTAGGCCTAGCCTCTTTTAAACAGCCTCTGAAATTTATCTTTTCT  
TCTATGGGGTCTATAAATGTATCTTATAATAAAAAGGAAGGACAGGAGGAAGA  
CAGGCAAATGTACTTCTCACCCAGTCTTCTACACAGATGGAATCTCTTTGGGG  
CTAAGAGAAAGGTTTTATTCTATATTGCTTACCTGATCTCATGTTAGGCCTAA  
GAGGCTTTCTCCAGGAGGATTAGCTTGGAGTTCTCTATACTCAGGTACCTCTT  
TCAGGGTTTTCTAACCTGACACGGACTGTGCATACTTTCCCTCATCCATGCT  
GTGCTGTGTTATTTAATTTTTCTGGCTAAGATCATGTCTGAATTATGTATGA  
AAATTATTCTATGTTTTTATAATAAAAATAATATATCAGACATCGA (SEQ ID NO: 89)  
>NP\_001703.2 carcinoembryonic antigen-related cell adhesion molecule 1 isoform  
1 precursor [*Homo sapiens*]

MGHLSAPLHRVRVPWQGLLLTASLLTFWNPPTTAQLTTESMPFNVAEGKEVLL  
LVHNLPPQLFGYSWKGERVDGNRQIVGYAIGTQQATPGPANSGRETIYPNAS  
LLIQNVTQNDTGfYTLQVIKSDLVNEEATGQFHVYPELPKPSISSNNSNPVED  
KDAVAFTCEPETQDTTYLWWINNQSLPVSRLQLSNGNRTLTLSSVTRNDTGP  
YECEIQNPVSANRSDPVTNLNVTYGPDTPTISPSDTYYRPGANLSLSCYAASNP  
PAQYSWLINGTFQQSTQELFIPNITVNNSGSYTCHANNSVTGCNRTTVKTIIV  
TELSPVVAKPQIKASKTTVTGDKDSVNLTCSTNDTGISIRWFFKNQSLPSSER  
MKLSQGNTTLSINPVKREDAGTYWCEVFNPISKNQSDPIMLNVNYNALPQENG  
LSPGAIAGIVIGVVALVALIAVALACFLHFGKTGRASDQRDLTEHKPSVSNHT  
QDHSNDPPNKMNEVTYSTLNFEAQPTQPTSASPSLTATEIHYSEVKKQ (SEQ ID NO:  
90) Mouse >NM\_001039185.1 *Mus musculus* carcinoembryonic CEACAM1 antigen-  
related cell adhesion molecule 1 (Ceacam1), transcript variant 1, mRNA  
AAAGCTCCTTTAAGAAAAGCAGGGCAGATATCAGGGCAGCCTGGCTTAGCAGT  
AGTGTTGGAGAAGAAGCTAGCAGGCAGGCAGCAGAGACATGGAGCTGGCCTCA  
GCACATCTCCACAAAGGGCAGGTTCCCTGGGGAGGACTACTGCTCACAGCCTC  
ACTTTTAGCCTCCTGGAGCCCTGCCACCACTGCTGAAGTCACCATTGAGGCTG  
TGCCGCCCCAGGTTGCTGAAGACAACAATGTTCTTCTACTTGTTCAACAATCTG  
CCCCTGGCGCTTGGAGCCTTTGCCTGGTACAAGGGAAACACTACGGCTATAGA  
CAAAGAAATTGCACGATTTGTACCAAATAGTAATATGAATTCACGGGGCAAG  
CATAACGGGCAGAGAGATAATATACAGCAATGGATCCCTGCTCTTCCAAATG  
ATCACCATGAAGGATATGGGAGTCTACACACTAGATATGACAGATGAAACTA  
TCGTCGTACTCAGGCGACTGTGCGATTTCATGTACACCCCATATTATTAAAGC  
CCAACATCACAAGCAACAACCTCCAATCCCGTGGAGGGTGACGACTCCGTATCA  
TTAACCTGTGACTCTTACACTGACCCTGATAATATAAACTACCTGTGGAGCAG  
AAATGGTGAAAGCCTTTCAGAAGGTGACAGGCTGAAGCTGTCTGAGGGCAACA  
GGACTCTCACTTTACTCAATGTCACGAGGAATGACACAGGACCCTATGTGTGT  
GAAACCCGGAATCCAGTGAGTGCAACCGAAGTGACCCATTCAGCCTGAACAT  
TATCTATGGTCCGGACACCCCGATTATATCCCCCTCAGATATTTATTTGCATC  
CAGGGTCAAACCTCAACCTCTCCTGCCATGCAGCCTCTAACCCACCTGCACAG  
TACTTTTGGCTTATCAATGAGAAGCCCCATGCATCCTCCCAAGAGCTCTTTAT  
CCCCAACATCACTACTAATAATAGCGGAACCTATACCTGCTTCGTCAATAACT  
CTGTCACTGGCCTCAGTAGGACCACAGTCAAGAACATTACAGTCCTTGAGCCA

[illegible]



AAATGTATTATATGACTGATAGCA (SEQ ID NO: 91) >NP\_001034274.1  
carcinoembryonic antigen-related cell adhesion molecule 1 isoform 1 precursor  
[*Mus musculus*]  
MELASAHHLHKGQVPWGGLLLTASLLASWSPATTAEVTTIEAVPPQVAEDNNVLL  
LVHNLPLALGAFAWKGNNTTAIDKEIARFVPNSNMNFTGQAYSGREIYSNGS  
LLFQMITMKDMGVYTLDMTDENYRRTQATVRFHVHPILLKPNITSNNSNPVEG  
DDSVSLTCDSTDPDNINYLWSRNGESLSEGDRLKLSEGNRTLTLNLTNRNDT  
GPYVCETRNPVSVNRSDPFSLNIIYGPDTPIISPSDIYLHPGSNLNLSCHAAS  
NPPAQYFWLINEKPHASSQELFIPNITTNNSGTYTFCFVNSVTGLSRTTVKNI  
TVLEPVTQPFLQVTNTTVKELDSVTTLTCLSDIGANIQWLENSQSLQLTERMT  
LSQNNILRIDPIKREDAGEYQCEISNPVSVRRSNSIKLDIIFDPTQGGLSDG  
AIAGIVIGVVAGVALIAGLAYFLYSRKSGGGSDQRDLTEHKPSASNHNLAPSD  
NSPNKVDDVAYTVLNFNSQQPNRPTSAPSSPRATETVYSEVKKK (SEQ ID NO: 92)

Human >NM\_007048.6 *Homo sapiens* butyrophilin subfamily 3 BTN3A1 member  
A1 (BTN3A1), transcript variant 1, mRNA  
ATTCCTCACGATGACCCGACAGTCTCTGCTTTCTTTTTCCTTTCTTCCAGAAG  
GAGATTTAACCATAGTAGAAAGAATGGAGAACTATTAAGTGCCTTTCTTCTGT  
GGGCTGTGATTTTCAGAGGGGAATGCTAAGAGGTGATTTTCAATGTTGGGACT  
CAAAGGTGAAGACACTGAAGGACAGAATTTTGGCAGAGGAAAGATCTTCTTC  
GGTCACCATACTTGAGTTAGCTCTAGGGAAGTGGAGGTTTCCATTTGGAATTC  
TATAGCTTCTTCCAGGTCATAGTGTCTGCCCCCACCTTCCAGTATCTCCTGA  
TATGCAGCATGAATGAAAATGGCAAGTTTCTGCTGGCCTTCTTCTGCTCAACTT  
TCGTGTCTGCCTCCTTTTGCTTCAGCTGCTCATGCCTCACTCAGCTCAGTTTT  
CTGTGCTTGGACCCTCTGGGCCCATCCTGGCCATGGTGGGTGAAGACGCTGAT  
CTGCCCTGTCACCTGTTCCCGACCATGAGTGCAGAGACCATGGAGCTGAAGTG  
GGTGAGTTCAGCCTAAGGCAGGTGGTGAACGTGTATGCAGATGGAAAGGAAG  
TGGAAGACAGGCAGAGTGCACCGTATCGAGGGAGAACTTCGATTCTGCGGGAT  
GGCATCACTGCAGGGAAGGCTGCTCTCCGAATACACAACGTCACAGCCTCTGA  
CAGTGGAAAGTACTTGTGTTATTTCCAAGATGGTGACTTCTATGAAAAAGCCC  
TGGTGGAGCTGAAGGTTGCAGCACTGGGTTCTGATCTTCACGTTGATGTGAAG  
GGTTACAAGGATGGAGGGATCCATCTGGAGTGCAGGTCCACTGGCTGGTACCC  
CCAACCCCAAATACAGTGGAGCAACAACAAGGGAGAGAACATCCCGACTGTGG  
AAGCACCTGTGGTTGCAGACGGAGTGGGCCTGTATGCAGTAGCAGCATCTGTG  
ATCATGAGAGGCAGCTCTGGGGAGGGTGTATCCTGTACCATCAGAAGTTCCCT  
CCTCGGCCTGGAAAAGACAGCCAGCATTTCATCGCAGACCCCTTCTTCAGGA  
GCGCCCAGAGGTGGATCGCCGCCCTGGCAGGGACCCTGCCTGTCTTGCTGCTG  
CTTCTTGGGGGAGCCGGTACTTCCTGTGGCAACAGCAGGAGGAAAAAAGAC  
TCAGTTCAGAAAGAAAAAGAGAGAGCAAGAGTTGAGAGAAATGGCATGGAGCA  
CAATGAAGCAAGAACAAGCACAAAGAGTGAAGCTCCTGGAGGAACTCAGATGG  
AGAAGTATCCAGTATGCATCTCGGGGAGAGAGACATTCAGCCTATAATGAATG  
GAAAAAGGCCCTCTTCAAGCCTGCGGATGTGATTCTGGATCCAAAAACAGCAA  
ACCCCATCCTCCTTGTCTTCTGAGGACCAGAGGAGTGTGCAGCGTGCCAAAGGAG  
CCCCAGGATCTGCCAGACAACCCTGAGAGATTTAATTGGCATTATTGTGTTCT  
CGGCTGTGAGAGCTTCATATCAGGGAGACATTACTGGGAGGTGGAGGTAGGGG  
ACAGGAAAGAGTGGCATATAGGGGTGTGCAGTAAGAATGTGCAGAGAAAAGGC  
TGGGTCAAATGACACCTGAGAATGGATTCTGGACTATGGGGCTGACTGATGG  
GAATAAGTATCGGACTCTAACTGAGCCCAGAACCAACCTGAAACTTCCTAAGC  
CCCCTAAGAAAGTGGGGGTCTTCTGACTATGAGACTGGAGATATCTCATTC  
TACAATGCTGTGGATGGATCGCATATTCATACTTTCTGACGTCTCCTTCTC  
TGAGGCTCTATATCCTGTTTTTCAGAATTTTGACCTTGGAGCCCACGGCCCTGA

CTATTGCTCCAGCGTGAAGAAAGAGTAAGTTCCTGCAATTCTGACCGCTG  
CTGATCATTCCCTAGAGACACCAGTAACCCCGGGCTTAGCTAACGAAAGTGGG  
GAGCCTCAGGCTGAAGTAACCTTTTCTCTGCTTCTCCCTGCCCAGCTCAGAGCT  
GAGGGCCTCCCCCTCCACAGCAACCAATCACAACCATAAAGCTACAAGCACGC  
ACTGAAGCACTTTACTGATACTCATTCAATTATTCATATGACAGTTGTTTGAG  
TTTGGTACCATCTTATTTTCCCCTTATACAGATAAGGAAACTGGGGTGCAGAA  
AAGTGAATTGACTACAAAGTAGACATGACTAGTTAACAACACAGCTGGGATCT  
AAACAGCAATAACTAACATTAATGGAGAACTTAAAATGCTCTGAGTGCTGTGT  
TATGAGCTTTGGTGGATGTCACCTTTAATCCTCGCAACACCCTGTCGGGTA  
GTCTCATTTAGCAAGTATGGAAGTTGAGGCAGGGCAACATTAAGCAACTTACA  
TAACTCATGCAGTAATTTCTGCAGTTGGGAGATGTTTCACTTCAGTCCCCGGC  
CCTATGGCCGTTCTTTTCCACCCTGTTTCTTCCCCCATAGGAAGAACCCACCT  
GTAGCCCTGAGGTTCTTTTCCCAGGATGGCTCCAGGATAAGGATCACTGTAGG  
TGGTTGTGGAGTTGACACCCCTGTTGACTCCTTCCCAGCTGATTGTCAGAGCC  
TTAGACCCAGCACGCCTTGGATTAGCTCTGCAGAGTGTCTTGGTTGAGAGAAT  
AACCTCACCGTACCCACATGACACGTGATTTGGAAAGAGACTAGAGGCCACAC  
TTGATAAATCATGGGGAACAGATGTGTTCCACCCAACAAATGTGATAAGTGAT  
CATGCAGCCAGAGCCAGCCTTCCTTCAATCAAGGTTTCCAGGCAGAGCAAATA  
CCCTAGAGATTCTCTGTGATATAGGAAATTTGGATGAAGGGAGCTAGAAGAAA  
TACAGGGATTTTTTTTTTTTTTTTAAAGATGGAGTCTTACTCTGTTGCTAGGCTG  
GAGTGCAGTGGTGCATCTCAGCTCCCTGCAACCTCCACCTCCTGGGTTCAAA  
CAATTCTCCTGCCTCAGCCTCCCGAGTACTGGGAATATAGGTGCACGCCACCA  
CACCCAACAAATTTTTGTACTTTTAGTACAGATGAGGGTTCCTATGTTGGCC  
AGGATGGTCTCGATCTCTTGACCTCATGATCCACCCACCTCGGTCTCCCAAAG  
TGCTGGGATTACAGGCTTGAGCCACCGGGTGACCGGCTTACAGGGATATTTT  
AATCCCGTTATGGACTCTGTCTCCAGGAGAGGGGTCTATCCACCCCTGCTCAT  
TGGTGGATGTTAAACCAATATTCCTTTCAACTGCTGCCTGCTAGGGAAAAACT  
ACTCCTCATTATCATCATTATTATTGCTCTCCACTGTATCCCTCTACCTGGC  
ATGTGCTTGTCAAGTTCTAGTTGTTCAATAAATTTGTTAATAATGCTGA (SEQ ID  
NO: 93) >NP\_008979.3 butyrophilin subfamily 3 member A isoform a precursor  
[*Homo sapiens*]

MKMASFLAFLLLNFRVCLLLLQLLMPHSAQFSVLGPSGPILAMVGEDADLPCH  
LFPTMSAETMELKWVSSSLRQVVNVYADGKEVEDRQSAPYRGRTSILRDGITA  
GKAALRIHNVTASDSGKYLCYFQDGDIFYEKALVELKVAALGSDLHVDVKGYKD  
GGIHLECRSTGWYPQPQIQWSNNKGENIPTVEAPVVADGVGLYAVAASVIMRG  
SSGEGVSC TIRSSLLGLEKTASISIA DPF FRS AQRWIAALAGTLPVLLLLLLGG  
AGYFLWQQQEEKKTQFRKKKREQELREMAWSTMKQEQSTRVKLLEELRWRSIQ  
YASRGERHSAYNEWKKALFKPADVILDPKTANPILLVSEDQRSVQRAKEPQDL  
PDNPERFNWHYCVLGCE SFISGRHYWEVEVGDRKEWHIGVCSKNVQRKGWVKM  
TPENGFWTMGLTDGNKYRTLTEPRTNLKLPKPPKKVG VFLDYETGDISFYNAV  
DGSIH TFLDVSFSEALYPVFRILTLEPTALTICPA (SEQ ID NO: 94) Human  
>NM\_007047.5 *Homo sapiens* butyrophilin subfamily 3 BTN3A2 member A2  
(BTN3A2), transcript variant 1, mRNA

GACTCTTACTGTTTCTCATGGTGAGAAGACAATATTTGCTTTTCTTTTTCT  
TTCTTCCGGATGAGAGGCTAAGCCATAATAGAAAGAATGGAGAATTATTGATT  
GACCGTCTTTATTCTGTGGGCTCTGATTCTCCAATGGGAATACCAAGGGATGG  
TTTTCCATACTGGAACCCAAAGGTAAAGACACTCAAGGACAGACATTTTTGGC  
AGAGCATAGATGAAAATGGCAAGTTCCTTGCTTTTCTTCTGCTCAACTTTCA  
TGTCTCCCTCCTCTTGGTCCAGCTGCTCACTCCTTGCTCAGCTCAGTTTTCTG  
TGCTTGGACCCTCTGGGCCCATCCTGGCCATGGTGGGTGAAGACGCTGATCTG

CCCTGTACACCTGTTTCCCGACCATGAGTGACAGACACCATGGAGCTGAAGTGGGT  
AAGTTCCAGCCTAAGGCAGGTGGTGAACGTGTATGCAGATGGAAAGGAAGTGG  
AAGACAGGCAGAGTGCACCGTATCGAGGGAGAACTTCGATTCTGCGGGATGGC  
ATCACTGCAGGGAAGGCTGCTCTCCGAATACACAACGTCACAGCCTCTGACAG  
TGGAAAGTACTTTGTGTTATTTCCAAGATGGTGAAGTTCTATGAAAAAGCCCTGG  
TGGAGCTGAAGGTTGCAGCACTGGGTTCTAATCTTCACGTCGAAGTGAAGGGT  
TATGAGGATGGAGGGATCCATCTGGAGTGCAGGTCCACCGGCTGGTACCCCCA  
ACCCCCAAATACAGTGGAGCAACGCCAAGGGAGAGAACATCCCAGCTGTGGAAG  
CACCTGTGGTTGCAGATGGAGTGGGCCTATATGAAGTAGCAGCATCTGTGATC  
ATGAGAGGGCGGCTCCGGGGGAGGGTGTATCCTGCATCATCAGAAATTCCTCTCT  
CGGCCTGGAAGAGACAGCCAGCATTTCATCGCAGACCCCTTCTTCAGGAGCG  
CCCAGCCCTGGATCGCAGCCCTGGCAGGGACCCTGCCTATCTTGCTGCTGCTT  
CTCGCCGGAGCCAGTTACTTCTTGTGGAGACAACAGAAGGAAATAACTGCTCT  
GTCCAGTGAGATAGAAAGTGAGCAAGAGATGAAAGAAATGGGATATGCTGCAA  
CAGAGCGGGAAATAAGCCTAAGAGAGAGCCTCCAGGAGGAACTCAAGAGGAAA  
AAAATCCAGTACTTGAAGTCTGTGGAGAGGAGTCTTCGTCCGATACCAATAAGTC  
AGCCTGATGCTCTAATGGAAAAATGGCCCTCTTCAAGCCTGGTGAGGAAATGC  
TTCAGATGAGGCTCCACCTTGTTAAATAAATTGGATGTATGGAAAAATAGACT  
GCAGAAAAGGGGAACTCATTAGCTCACGAGTGGTTCGAGTGAAGATTGAAAAT  
TAACCTCTGAGGGCCAGCACAGCAGCTCATGCCTGTAATCCTAGCACTTTGGA  
AGGCTGAGGAGGGCGGATCACAAAGGTCAGGAGATCAAGACCATCCTGGCTAAC  
ACGGTGAAACCCCGTCTCTACTAAAAATACAAAAAATAAAAAATTAGCCGGGC  
ATGGTGACGGGCACCTGTAGTCCCAGCTACTCGGGAGGCTGAGGCAGGAGAAT  
GGCATGAACCCGGAAGGCAGAGCTTGCAGTGAGCCGAGATCACGCCACTGCAC  
TCCAGCCTGGGAGACAGAGCGAGACTCTGTCTCAAGAAAAAAAAAAAAAAAAAAAA  
AAAAGAAAAGAAAATTAACCTCTGAGTATAAAGCATCAGTGGGCAGAATCAAT  
GTGGGGAGGGGAAACAACAAAAATGTAGAAAGAGGATCCTTGTTGCTTCTTGGG  
GCCGCATCAGGGTATTGGGTAGGCAGATACTGACCTTACTTTCAATTTCCCCT  
CTGGTCACTAGACCCCTGGGGCTTTCACCAATGACATTGATGAGAGAATCACA  
TTCAGGGCAGGCTAGGGACACGGGGTTCTGGAAGGACCTCCTCAGCATGGCCC  
AAGCCTTGTCATGCTGTGGCTCTTAAATCCAGGAAAAATGGCTGACCCCATGGA  
CACCTCCTCAAACCTCTCTGCAGCAGATGTAATTCTGTATCCAGACATGGCAAA  
TGCCATCCTCCTTGTCTTCTGAGGACCAGAGGAGTGTACAGCGTGCTGAGGAGC  
CCCATGACCTACCAGACAACCCTGAGAGATTTGAATGGCGTTACTGTGTGCTT  
GGCTGTGAAAGCTTCATGTCAGAGAGACACTACTGGGAGGTGGAAGTGGGGGA  
CAGAAAAGAGTGGCATATTGGGGTATGTAGTAAGAACGTGGAGAGGAAAAAAG  
TTTGGGTCAAAATGACACCGGAGAACGGATACTGGACTATGGGCCTGACTGAT  
GGGAATAAGTATCGGGCTCTCACTGAGCCCAGAACCAACCTGAAACTTCCTGA  
GCCTCCTAGGAAAGTGGGGGTCATCCTGGACTATGAGACTGGACATATCTCGT  
TCTACAATGCCACGGATGGATCTCATATCTACACATTTCTGCACGCCTCTTCC  
TCTGAGCCTCTGTATCCTGTATTCAGAATTTTGACCTTGGAGCCCACTGCCCT  
GACCGTTTGCCCAATAACCAAAAGTAGAGAGTTCCCCCGATCCCGACCTAGTGC  
CTGATCATTCCCTGGAGATAACCACTGACCCCAAGGCTTAGCTAATGAAAGTGGG  
GAGCCTCAGGCTGAAGTAACATCTCTGCTTCTCCCTGCCCAGCCTGGAGCTAA  
GGGTCTCACCCCTCCACAACAGCCAGTCAGAACCATAAAGCTACAGGCACACAC  
TGAAGCACTTTACTGATATTCATTCAATTATTCATAGGACAGTTGTTTGAGT  
TTGGTGCCACCTTATTGGCCCCCTTTATACAGATAAGGAAACTGGGGTGTAGAA  
AAGTGTATTGACTTTACAAAGCAGACAGGAATAGTGAACAACAGAGCTGGGAT  
CTGAACAACAATGACTAACATTAATGGAGAATTTAAAACGTTCTGAGTGCTGT  
GTTATGAGCTTTGGTGGGTGTCACTCCTTTAATCCTCACAACACCCTGTCAGG

TAGTCTCATTGTTGGCAAGTATGGAAGGACAGGAGGCAAGGAGTCAAGTACCTTA  
CATAACTCACACGGTAATTTGTGCAGTTGGGAGATGTTTCAGCTTCAGTCCCTG  
GCCAATTGCCCCGTTCTTTTCCAGCCTGATTTTTCCTGCATGGGAAGAGCCCAC  
ATGTAGCCCTGAGGTTCCCTTCCCAGGACAGCTCCAGGATCGAGATCACTGTG  
AGTGGTTGTGGAGTTAAGACCCCTATGGACTCCTTCCCAGCTGATTATCAGAG  
CCTTAGACCCAGCACTCCTTGGATTGGCTCTGCAGAGTGTCTTGGTTGAGAGA  
ATAACGTTGCAGTTCCCACAGGGGCATGTGACTTTGAAAGAGACTAGAGGCCAC  
ACTCAGTTAATAATGGGGCACAGATGTGTTCCCACCCAACAAATGTGATAAGT  
GATCGTGCAGCCAGAGCCAGCCTTCCTTCAGTCAAGGTTTCCAGGCAGAGCAA  
ATACCCTAGAGATTCTCTGTAATATTGGTAATTTGGATGAAGGAAGCTAGAAG  
AATTACAGGGATGTTTTTAATCCCAGTATGGACTCAGTCTCCTGGAAAAGGAT  
CTGTCCACTCCTGGTCATTGGTGGATGTTAAACCCATATTCCTTTCAACTGCT  
GCCTGCTAGGGAAAAGTCTCCTCATTATCATCACTATTATTGCTCACCCTG  
TATCCCCTCTACTGGGCAAGTGCTTGTCAAGTTCTAGTTGTTCAATAAATTTG  
TTAATAATGCTGA (SEQ ID NO: 95)>NP\_008978.2 butyrophilin subfamily 3  
member A2 isoform a precursor [*Homo sapiens*]  
MKMASSLAFLLLNFHVSLLVQLLTPCSAQFSVLGPSGPILAMVGEDADLPCH  
LFPTMSAETMELKWVSSSLRQVVNVYADGKEVEDRQSAPYRGRTSILRDGITA  
GKAALRIHNVTASDSGKYLCYFQDGDIFYEKALVELKVAALGSNLHVEVKGYED  
GGIHLECRSTGWYPQPQIQWSNAKGENIPAVEAPVVADGVGLYEVAASVIMRG  
GSGEGVSCIIRNSLLGLEKTASISIADPFFRSAQPWIAALAGTLPILLLLLAG  
ASYFLWRQQKEITALSSEIESEQEMKEMGYAATEREISLRESLQEELKRKKIQ  
YLTRGESSSDTNKSA (SEQ ID NO: 96) Human>NM\_007049.5 *Homo sapiens*  
butyrophilin subfamily 2 BTN2A1 member A1 (BTN2A1), transcript variant 1,  
mRNA AGATTTTCGTTTCCTGCATCTCCAAACATGGCGACCTAGGAGAAGGGGAAGAAC  
AATTTTTTCTCCTCTTTTGGGAAGGTTTGTGTCTAGTAGTGCCTGTGCCCTG  
GGCAGATTGGAGAGAAGAGGGGACGACTGGAGAATCGTCGAGAACCAGCGGAGA  
AAAGAAAAAGCAACGTTTAATTCTAGAAGGCCTCCTGTCCCTGCCTGCTCTGG  
GTGCTCATGGAATCAGCTGCTGCCCTGCACTTCTCCCGGCCAGCCTCCCTCCT  
CCTCCTCCTCCTCAGCCTGTGTGCACTGGTCTCAGCCCAGTTTATTGTCGTGG  
GGCCCACTGATCCCATCTTGGCCACGGTTGGAGAAAACACTACGTTACGCTGC  
CATCTGTCACCCGAGAAAAATGCTGAGGACATGGAGGTGCGGTGGTTCCGGTC  
TCAGTTCTCCCCCGCAGTGTTTGTGTATAAAGGTGGCAGAGAGAGAACAGAGG  
AGCAGATGGAGGAGTACCGAGGAAGAACCACCTTTGTGAGCAAAGACATCAGC  
AGGGGCAGCGTGGCCCTGGTCATACACAACATCACAGCCCAGGAAAACGGGCAC  
CTACCGCTGTTACTTCCAAGAAGGCAGGTCCTACGATGAGGCCATCCTGCACC  
TCGTAGTGGCAGGACTAGGCTCTAAGCCCCTCATTTCAATGAGGGGGCCATGAA  
GACGGGGGGCATCCGGCTGGAGTGCATATCTAGAGGGTGGTACCCAAAGCCCCT  
CACAGTGTGGAGGGACCCCTACGGTGGGGTTGCGCCTGCCCTGAAAGAGGTCT  
CCATGCCTGATGCAGACGGCCTCTTCATGGTCACCACGGCTGTGATCATCAGA  
GACAAGTCTGTGAGGAACATGTCCTGCTCTATCAACAACACCCTGCTCGGCCA  
GAAGAAAGAAAGTGTCAATTTTATTCCAGAATCCTTTATGCCCAGTGTGTCTC  
CCTGTGCAGTGGCCCTGCCTATCATTGTGGTTATTCTGATGATACCCATTGCC  
GTATGCATCTATTGGATCAACAACTCCAAAAGGAAAAAAAGATTCTGTCAGG  
GGAAAAGGAGTTTGAACGGGAAACAAGAGAAATTGCTCTAAAGGAACTGGAGA  
AAGAACGTGTGCAAAAAGAGGAAGAAGTCAAGTAAAAGAGAAACTTCAAGAA  
GAATTGCGATGGAGAAGAACATTCTTACATGCTGTTGATGTGGTCCTGGATCC  
AGACACCGCTCATCCCGATCTCTTCCTGTCAGAGGACCGGAGAAGTGTGAGAA  
GGTGCCCTTCAGGCACCTAGGGGAGAGCGTGCCTGACAACCCAGAGAGATTC  
GACAGTCAGCCTTGTGTCTAGGCCGGGAGAGCTTCGCTTCAGGGAAACATTA

CTGGAGGAGGTGGAGAGGTGGAAGGAGGAGGTCCTGCTGATTCCTCAGAATGGCTTCTGG  
ACAGTGTTGAGAGGAAAGGGGAGGTCCTGCTGATTCCTCAGAATGGCTTCTGG  
ACCTTGGAGATGCATAAAGGGCAATACCGGGCCGTGTCCTCCCCTGATAGGAT  
TCTCCCTTTGAAGGAGTCCCTTTGCCGGGTGGGCGTCTTCCTGGACTATGAAG  
CTGGAGATGTCTCCTTCTACAACATGAGGGACAGATCGCACATCTACACATGT  
CCCCGTTTCAGCCTTTTCCGTGCCTGTGAGGCCCTTCTTCAGGTTGGGGTGTGA  
GGACAGCCCCATCTTCATCTGCCCTGCACTCACAGGAGCCAATGGGGTTCACGG  
TGCCTGAAGAGGGCCTGACACTTCACAGAGTGGGGACCCACCAGAGCCTATAG  
AATCAATTCCTTGGTCTCACAGCCATGTAGACAAGCCCTGGTTCATCTCAGCAG  
CCACCGCACAAACACCCCTGGTGGAAAGACACGCCCTCCTCCCCTCTGGTCACAC  
AAGAGAACATCTTCCAGCTGCCTCTTTTCACACCCACTACAGACCTCAGCCCCA  
GTTTTCTCCTCCTCACTAGGCTGTGTTTTTAGTAGTTCCTTTGCTTGTAATA  
TGGGATGGGATCCAGGCATAGGGAAGTAGTTGTTACACAGCTCCCAGCCAAGA  
AGAAAGTGTGAGAAGTTGATGGGCAGCAAACCTGCTGTTTAACATCAGGGTGA  
CCACATTAAGCCCAGTATTCCAGTTGGCACCAGAAGATATGGACTTGGAATGA  
GGCCTACAGGGTTCACCAGGATGTAAGAGGAGAGAGGAATCCACAGGACCACC  
AGAGAGGAGAGGGAACCAGATATGCAGATCAGAGATAGAGGAAGTGGAAACCAG  
AGAGCTGGGAGGGACCAAGGTTGTAAGGGTGGCTAAGTCCCACCATAACAGCT  
AAGGGGACCTGGGAGATGATGGCTCATTTCCACCCAGCCCCAGGATTTCCAGA  
GCGCACATCCACAGGCCTGGACCTGGGATGAAGATGAATGAAGAACATGGATG  
CACGTGGATGTAGTTTGGCTCAGGTGTCCCTGCAGTTGGCAAGGAGTCAGTAC  
TCAGTCCCTGAGTGTGGCTGAAATTTGAGGTCCTGGCTGAGCCAAGGAGTAAT  
GGACCAGATCTACCTCAGTATTCAAGTTCAGTGGGGACACCAGTGGCTTCAAA  
CTTCCTGGTTTCATGATATCTTGAGACGCCTTACAAATGATGGAGGATTCCAA  
AGAGTTTTTGTATTATTGGGTAAATATTTGTTGGTATTTATGGCATTGAGAT  
TGAAACTAAGAAATGTTTTAATTTATTACCTTTACAACATTTATTACATTAC  
ATACATACATTTACAACATTTATTAATTTATATTAATAAATAGCATGAATAAGCC  
AATTATAGGTAAATATAAGTAGAATGTTTGTGAAAAATAAGTATGGTATCCAA  
AGCAAAATAAATTTTATTGTGAAGTGTG (SEQ ID NO: 97) >NP\_008980.1  
butyrophilin subfamily 2 member A1 isoform 1 precursor [*Homo sapiens*]  
MESAAALHFSRPASLLLLLLSLCALVSAQFIVVGPTDPILATVGENTTLRCHL  
SPEKNAEDMEVRWFRSQFSPAVFVYKGGREERTEEQMEEYRGRTTFVSKDISRG  
SVALVIHNITAQENGTYRCYFQEGRSYDEAILHLVVAGLGSKPLISMGRGHEDG  
GIRLECISRGWYPKPLTVWRDPYGGVAPALKEVSMPDADGLFMVTTAVIIRDK  
SVRNMSCSINNTLLGQKKESVIFIPESFMPSVSPCAVALPIIVVILMIPIAVC  
IYWINKLQKEKKILSGEKEFERETREIALKELEKERVQKEEELQVKEKLQEEL  
RWRRTFLHAVDVVLDPDTAHPDLFLSEDRRSVRRCPFRHLGESVPDNPFRFDS  
QPCVLGRESFASGKHYYEVEVENVIEWTVGVCRDSVERKGEVLLIPQNGFWTL  
EMHKGQYRAVSSPDRLPLKESLCRVGVFLDYEAGDVSFYNMRDRSHIYTCPR  
SAFSVPVRPFFRLGCEDSPIFICPALTGANGVTVP EEGTLHRVGTHQSL (SEQ ID NO:  
98) Human >NM\_001040462.3 *Homo sapiens* butyrophilin like 8 BTNL8 (BTNL8),  
transcript variant 2, mRNA  
AGAACAGCGCAGTTTGGCCTCCGCTCACGCAGAGCCTCTCCGTGGCTTCCGCA  
CCTTGAGCATTAGGCCAGTTCTCCTCTTCTCTCTAATCCATCCGTCACCTCTC  
CTGTCATCCGTTTCCATGCCGTGAGGTCCATTACAGAACACATCCATGGCTC  
TCATGCTCAGTTTGGTTCTGAGTCTCCTCAAGCTGGGATCAGGGCAGTGGCAG  
GTGTTTGGGCCAGACAAGCCTGTCCAGGCCTTGGTGGGGGAGGACGCAGCATT  
CTCCTGTTTCCTGTCTCCTAAGACCAATGCAGAGGCCATGGAAGTGCGGTTCT  
TCAGGGGCCAGTTCTCTAGCGTGGTCCACCTCTACAGGGACGGGAAGGACCAG  
CCATTTATGCAGATGCCACAGTATCAAGGCAGGACAAAACCTGGTGAAGGATTC

MALMLSLVLSLLKLGSGQWQVFGPDKPVQALVGEDAAFSCFLSPKTNAEAMEV  
RFFRGQFSSVVHLYRDGKDQPFMQMPQYQGR TKLVKDSIAEGRISLRLENITV  
LDAGLYGCRISQSYYQKAIWELQVSALGSVPLISITGYVDRDIQLLCQSSGW  
FPRPTAKWKGPQGQDLSTDSRTNRDMHGLFDVEISLTVQENAGSISCSMRHAH  
LSREVESRVQIGDTFFEPISWHLATKVLGILCCGLFFGIVGLKIFFSKFQWKI  
QAELDWRRKHGQAELRDARKHAVEVTLPETAHPKLCVSDLKTVTHRKAPEV  
PHSEKRFTRKSVVASQSFQAGKHYWEVDGGHNRWRVGVCRDDVDRRKEYVTL  
SPDHGYWVLRLNGEHL YFTLNPRFISVFPRTPTTKIGVFLDYECGTISFFNIN  
DQSLIYTLTCRFEGLLRPYIEYPSYNEQNGTPIVICPVTQESEKEASWQRASA  
IPETSNSESSSQATTPFLPRGEM (SEQ ID NO: 100) Human >NM\_006995.5 *Homo  
sapiens* butyrophilin subfamily 2 BTN2A2 member A2 (BTN2A2), transcript  
variant 1, mRNA

GGTCTGTTCTCCCCCGTGTGTAAGAGGTTGGGAGAGAGAGAACA  
GAGGAGCAGATGGAGGAGTACCGGGGAAGAATCACCTTTGTGAGCAAAGACAT  
CAACAGGGGGCAGCGTGGCCCTGGTCATACATAACGTCACAGCCCAGGAGAATG  
GGATCTACCGCTGTTACTTCCAAGAAGGCAGGTCCTACGATGAGGCCATCCTA  
CGCCTCGTGGTGGCAGGCCTTGGGTCTAAGCCCCTCATTGAAATCAAGGCCCA  
AGAGGATGGGAGCATCTGGCTGGAGTGCATATCTGGAGGGTGGTACCCAGAGC  
CCCTCACAGTGTGGAGGGACCCCTACGGTGAGGTTGTGCCCCGCCCTGAAGGAG  
GTTTCCATCGCTGATGCTGACGGCCTCTTCATGGTCACCACAGCTGTGATCAT  
CAGAGACAAGTATGTGAGGAATGTGTCCTGCTCTGTCAACAACACCCTGCTCG  
GCCAGGAGAAGGAAACTGTCATTTTTATTCCAGAATCCTTTATGCCCAGCGCA  
TCTCCCTGGATGGTGGCCCTAGCTGTCATCCTGACCGCATCTCCCTGGATGGT  
GTCCATGACTGTCATCCTGGCTGTTTTTCATCATCTTCATGGCTGTCAGCATCT  
GTTGCATCAAGAACTTCAAAGGGGAAAAAAGATTCTGTCAGGGGAAAAAGAAA  
GTTGAACAAGAGGAAAAAGAAATTGCACAGCAACTTCAAGAAGAATTGCGATG  
GAGAAGAACATTCTTACATGCTGCTGATGTGGTCCTGGATCCAGACACCGCTC  
ATCCCGAGCTCTTCCTGTCAGAGGACCGGAGAAGTGTGAGGCGGGGCCCTAC  
AGGCAGAGAGTGCCTGACAACCCAGAGAGATTTCGACAGTCAGCCTTGTGTCCT  
GGGATGGGAGAGCTTCGCCTCAGGGAAACATTACTGGGAGGTGGAGGTGGAAA  
ACGTGATGGTGTGGACTGTGGGGGTCTGCAGACACAGTGTTGAGAGGAAAGGG  
GAGGTCTGCTGATTCCTCAGAATGGCTTCTGGACCCTGGAGATGTTTGGAAA  
CCAATACCGGGGCCCTGTCCTCCCCTGAGAGGATTCTCCCTTTGAAGGAGTCCC  
TTTGCCGGGTGGGCGTCTTCCTGGACTATGAAGCTGGAGATGTCTCCTTCTAC  
AACATGAGGGACAGATCGCACATCTACACATGTCCCCGTTACGCCTTTACTGT  
GCCTGTGAGGCCCTTCTTCAGGTTAGGGTCTGATGACAGCCCCATCTTCATCT  
GCCCTGCACTCACAGGAGCCAGTGGGGTTCATGGTGCCTGAAGAGGGCCTGAAA  
CTTCACAGAGTGGGGACCCACCAGAGCCTATAGAATCAATTCCTTGGACTCAC  
AGCCATGCAGATAAGCCCTGGCCATCTCAGCAGCCACCGCACAACCCCCCTAA  
TGAAAGACACGCCCTCCTCCCCTCTGGTCACGTAAGAGAACATCTTCCAGCTG  
CCTTTTTTCACACCCACTCCAGCCCTCTGCCCCAGTTTTTCTCCTCCTACTAGT  
CTGTGGCTTTAGTAGTTCCCTTTGCTTGTAATTATGGGATGGGATCCAGGCATA  
GGGAAGTAGTTGTTTCATAGCTCCCAGTCAAAAAGAAAGTGAGAGAAGCTGTT  
GGGCAGCGAACCTACTGTTTAAAATCAGGATAACACATTAAGCCCAATATGC  
CAGTTGGCACCAGATGCTGTGGACTTGGAATGAGGCCAACAGGGTTCACCAGG  
ATGAGAGAGGAGAGAGGAATCCACAGGACCACCAGAAGGGAGAGGGAACCAGA  
TATGCAGATCAGAGATAGAGGAAGTGGAACCAGAGAGCTGGGAGGGACCAAGG  
TTGTAAGGATGGCTAAGTCCCACCATAAGAGCTAAAGGGTCCTGGGAGATGAT  
GGCTCATTTCCACCCAACCCCAGGATTTCCACAGCACACACCCACAGGCCTGG  
ACCTGGGATGAAGATGAATGAAGAACATGGACTCATGTGGATGTGGTTTGGCT  
CAGATGTCCCTGCAATAAACAAGGGGTGAGTACTTAGTCCCTGAGTGTGGTTG  
AGGTTTGAGGTCCTGGTCGAGCAGGGCAGTACTGGACCAGGTCTACGTCAGCA  
TTCAGGTTCAATGGGGACACCAGTGGCTTCAAACCTTCCTGATCTAATTATGTT  
TTTAGACACTTAGAAGTTATTGAGGACTTTAAAGAGCTTTTGTTTATTTGGGT  
TAATATTTATGACATTTGACATTGAAACAAAAATTTAAAATGTTATCTTTTAA  
TTTATGTTAAAATAGCATTAAATAAATCAGTTATAGGTAAATGTAGATAGGATG  
TTTTGTGAAAAAGCAATCTATTGTGTCCAAATAAAAAAACAAGTGTGACA  
CTGGTTAACTTTTTCCAGATCTCATGTCTGGCTTAATAAGAGATATTTGTATT  
ATCATATCTGCCTTTGTATTAAACCTATTGGTATATCATAGGTCATGTTAGCT  
CAAAAAAATTTACTGCACACTACTGAGAGAATGAGATGAAAAACGATTAATG  
TTTCATTATTATTATTGTGAAAATATTATTAACACTGGGGACTCCTTAAGAGT  
ACATCAGAGTTCTCTCTAGGAATCCCAAACACATTTTGAAACTAGAAATAGT

GGATCTCTGGAAGTGAATCCATCATGTGCTGGTTAATTTTAGATGATGCAACCTGACTG  
GATTAAGGAATACCTAGACAGCTGGTACAACATTATTTCTGGGTGTGTCTGTG  
AGTGTGTTTCCAGAAGAGATTGGCAAGTGAGTCAGTGGGAAATTCTCTCCTTC  
TGTTGGCTGGGTGCCCAATAACAACAAAAAGGCAGAGGAAAGGCCAAATTCTTCT  
CTCCTCTGGAGCTGAGACACTCTTCTTCTTCTGCCCCTTGGACATCAGAACTCC  
TGGCTCTCCGGCCTTTGAACTTCAGGACTTGTACCAGGAGGCCCTGGGTTCTC  
AGGCCTTTGGCTTTGGACTGAGAGTTACACAATCAGCTTCCCTGGTTCTGAGG  
CTTTCAGACTTAAACTGAGCCATGCTACCAGCATCCCAGGGTCTCCAGCCTAC  
AGATGAGCTGTTGTGCGATTTCTTAGCCTCCATAATCACATGAGCCAATCTCC  
TTAATAAATGCCTGCTCATAGATCTGTATCTACATCTATATCTGTATGTGCAT  
CTATATCTATGCCTATATCTATATCTATATCATATTGATTTTGTCTCTCTGGA  
GAACCCTGACTAATAAAAATGAGGCATCTAAAA (SEQ ID NO: 101)>NP\_008926.2  
butyrophilin subfamily 2 member A2 isoform a precursor [*Homo sapiens*]  
MEPAAALHFSLPASLLLLLLLLLLLLSLCALVSAQFTVVGPANPILAMVGENTTL  
RCHLSPEKNAEDMEVRWFRSQFSPAVFVYKGGRETEEQMEEYRGRITFVSKD  
INRGSVLVIHNVTAQENGIYRCYFQEGRSYDEAILRLVVAGLGSKPLIEIKA  
QEDGSIWLECISGGWYPEPLTVWRDPYGEVVPALKEVSIADADGLFMVTTAVI  
IRDKYVRNVSCSVNNTLLGQEKETVIFIPESFMPSASPWMVALAVILTASPWM  
VSMTVILAVFIIFMAVSICCIKKLQREKKILSGEKKVEQEEKEIAQQLQEELR  
WRRTFLHAADVVLDPDTAHPFLFSEDRRSVRRGPYRQRPDPNPERFDSQPCV  
LGWESFASGKHYWEVEVENVMVWTVGVCRHSVERKGEVLLIPQNGFWTLEMFG  
NQYRALSSPERILPLKESLCRVGVFLDYEAGDVSFYNMRDRSHIYTCPRSAFT  
VPVRPFFRLGSDDSPIFCALTGASGVMVPEEGLKLHRVGTHQSL (SEQ ID NO: 102)  
Mouse >NM\_175938.3 *Mus musculus* butyrophilin, subfamily 2, BTN2A2 member  
A2 (Btn2a2), transcript variant 1, mRNA



MEPTTSLRSCPIASLLFFLVLSLFLVLSAQFTVIGPAEPILAMVGENTTLHCH  
LSPERNAEEMEVRWFRWRFFPAVLVYRGHQERPEEQMVAYRGRTTFMRTDISK  
GRVALIIHNVTAYDNGIYCCYFQEGRSYDQATMKLMVASLGSEPLIKMKTLED  
GSILLECTSEGWYPEPRAVWRDPYDEVVPALEEEYTADREGLFTVTMTIIIRD  
CSVNRNMTCSVNNTLLSQEVESVILIPESFVPSLPLWMVAVAVTLPVVMLILLT  
SGSICLVKKHRRKKSILSAEKEAEYEEKEAARQLQEELRWRRTLLHAADVVD  
PDTAHPFLSDDQRSVIRGSSRQSVPDNPERFDCRPCVLGRESFSSGKHYYWE  
VEVENVMVWAIGVCRDSVERKGEALLVPQNGFWTLEMFGSQYRALSSPEKIIP  
LKERLHRIAVFLDCEGGDISFYNMRRDRSHIYTCPPVTFTGPLRPFRLGSSDS  
PLFICPAFTGAQGVTIPEGGLFLYKTRPISQSLVRKP(SEQ ID NO: 104) Human  
>NM\_001732.3 *Homo sapiens* butyrophilin subfamily 1 BTN1A1 member A1  
(BTN1A1), mRNA

CTGTCTCTCCGGAACGCGAGACGCGCCGAGACCACTGGAGAGCTACGCTGGTTCCGAAAGA  
AGGTTTCGCCGGCCGTGCTGGTGCATAGGGACGGGCGCGAGCAGGAAGCCGAG  
CAGATGCCCCGAGTACCGCGGGCGGGCGACGCTGGTCCAGGACGGCATCGCCAA  
GGGGCGCGTGCCCTTGAGGATCCGTGGCGTCAGAGTCTCTGACGACGGGGAGT  
ACACGTGCTTTTTTCAGGGAGGATGGAAGCTACGAAGAAGCCCTGGTGCATCTG  
AAGGTGGCTGCTCTGGGCTCTGACCCTCACATCAGTATGCAAGTTCAAGAGAA  
TGGAGAAATCTGTCTGGAGTGCACCTCAGTGGGATGGTACCCAGAGCCCCAGG  
TGCAGTGGAGAACTTCCAAGGGAGAGAAGTTTCCATCTACATCAGAGTCCAGG  
AATCCTGATGAAGAAGGTTTGTTCACCTGTGGCTGCTTCAGTGATCATCAGAGA  
CACTTCTGCGAAAAATGTGTCCTGCTACATCCAGAATCTCCTTCTTGGCCAGG  
AGAAGAAAGTAGAAATATCCATAACCAGCTTCCTCCCTCCCAAGGCTGACTCCC  
TGGATAGTGGCTGTGGCTGTCATCCTGATGGTTCTAGGACTTCTCACCATTGG  
GTCCATATTTTTCACTTGGAGACTATAACAACGAAAGACCCAGAGAGAGGAGGA  
ATGAATTCAGCTCTAAAGAGAGACTCCTGGAAGAACTCAAATGGAAAAAGGCT  
ACCTTGCATGCAGTTGATGTGACTCTGGACCCAGACACAGCTCATCCCCACCT  
CTTTCTTTATGAGGATTCAAAATCTGTTTCGACTGGAAGATTCACGTCAGAAAC  
TGCCTGAGAAAACAGAGAGATTTGACTCCTGGCCCTGTGTGTTGGGCCGTGAG  
ACCTTCACCTCAGGAAGGCATTACTGGGAGGTGGAGGTGGGAGACAGGACTGA  
CTGGGCAATCGGCGTGTGTAGGGAGAATGTGATGAAGAAAGGATTTGACCCCA  
TGACTCCTGAGAATGGGTTCTGGGCTGTAGAGTTGTATGGAAATGGGTACTGG  
GCCCTCACTCCTCTCCGGACCCCTCTCCCATTGGCAGGGCCCCCACGCCGGGT  
TGGGATTTTCTAGACTATGAATCAGGAGACATCTCCTTCTACAACATGAATG  
ATGGATCTGATATCTATACTTTCTCCAATGTCACCTTCTCTGGCCCCCTCCGG  
CCCTTCTTTTGCCTATGGTCTAGCGGTAAAAAGCCCCTGACCATCTGCCCAAT  
TGCTGATGGGCCTGAGAGGGTACAGTCATTGCTAATGCCCAGGACCTTTCTA  
AGGAGATCCCATTTGTCCCCCATGGGGGAGGACTCTGCCCTAGGGATGCAGAC  
ACTCTCCATTCTAAGCTAATCCCTACCCAACCCAGCCAAGGGGCACCTTAAGG  
AATATCTCAGCTCATCTGTTTTCTTTCTCTAACCCCTCTCCTCCATAGCCT  
TCTGAGGCTTCACCTGCTAGCTTTACCCAGTCTGTTTCTTCTGTTGGGTGGC  
AATTAATTAATCCTGTGAAGGTTACATTGCTGCTGCTAGAGAGGGTGGGGATT  
GCACCTTCCAAATCTGTTTCTGTACCAATATTTGGGGGATGGAGGGGTGACTC  
AAACTGCTTCTAGTGTTCTCCTAATCCCTTAAGACTAGAACCTATAGGAACT  
ACTTGGAGCAAACCTCAAAGGACAGATTAGGGATCGAGATTGGGTCAGGTTAGC  
ATGGGGTTGTGGTTGAAATATCTTGGTATCCAGGATAAGGGTATGTGGAAAAA  
CAGGCTTTAGGCAAGTGGAATAATTCAAATGTGCTGTGAAAGGACAATCTCAG  
GCTGAAATCCCATAAAGGAACTTGGAGGGAATATTATGATGGAGGGAAGTGAG  
GTGAATCCAGGCACATGATGAACACCTGGCTCATCCATAGAGTTTTTCACAGCC  
TATATCGCAAATTTTCTAAGCCACGTCCTATAGGACAGAGGAGACTGGCCCCA  
CTTCTATGGGTCTGAGCTGTGGAAAAGGGAGAGCAGAGAGGAACTGAGATGAG  
CAGGGATGAAGGGTCAGGCAGAAAGCGTGATAGAGGAGAGAATTTTTTGACAAA  
ACTCAAAGTTGTTTGCACAGCTGTTCTTTGTACCCTGTTCTTTCTCTGCGC  
CCTCCTGTTTCTCCCTTGCCTGGAAGTCATTCCACCCTCAATTTGTTGATCCA  
CAAGTTTCCAGTTGTCCTCTTCTTTTTTGTATAGCATCTCTCTATTTCAAAGA  
CATTCCTAGAAGTCATCCTTCAGTGATATCACCCTTGCTCAGTCACCATCTC  
AACCTTATGTCACCTCAGCCCTCATCTCAATGCCCAAACCCCTTACACACACC  
TTCAGTTAGCTTCAACTGCCTCCGTTTCCACACTGTGCACCTTTCACCTTTCC  
TACCCAGCTTTCTACATGCTGCCTCTCCTCAGGGTCCCCTGAATGCTGCATC  
ATTGTGTTTCAGTGCAGCTGGACTGATTGCACCTGTGTATTTGCCCTGAGCAC  
TTTCCTTTACACATGTGGCTTGTCTTGCCAATAGACTCCAGGCTTATACCTTC  
CATTTCCATCGTATTCTCCAGTTTCCAGGATAGACGTTGCTCATCGTCTTTAC

CTAATAATAAGTTTGTCTGATTGCTGAAA (SEQ ID NO: 105) >NP\_001723.2  
butyrophilin subfamily 1 member A1 precursor [*Homo sapiens*]  
MAVFPSSGLPRCLLTLLQLPKLDSAPFDVIGPPEPILAVVGEDAELPCRLS  
PNASAEHLELRWFRKKVSPAVLVHRDGREQEAEQMPEYRGRATLVQDGIKGR  
VALRIRGVRVSDDGEYTCFFREDGSYEEALVHLKVAALGSDPHISMQVQENGE  
ICLECTSVGWYPEPQVQWRTSKGEKFPSTSESARNPDEEGLFTVAASVIIRDTS  
AKNVSCYIQNLLLQGEKKVEISIPASSLPRLTPWIVAVAVILMVLGLLTIGSI  
FFTWRLYNERPRERRNEFSSKERLLEELKWKKATLHAVDVTLDPDTAHPHLFL  
YEDSKSVRLED SRQKLPEKTERFDSWPCVLGRETFTSGRHYWEVEVGDRTDWA  
IGVCRENVMKKGFDPMTPENGFWAVELYGNGYWALTPLRTPLPLAGPPRRVGI  
FLDYESGDISFYNMNDGSDIYTF SNVTFSGPLRPFCLWSSGKKPLTICPIAD  
GPERVTVIANAQDLSKEIPLSPMGEDSAPRDADTLH SKLIPTQPSQGAP (SEQ ID NO:  
106) Mouse >NM\_013483.3 *Mus musculus* butyrophilin, subfamily 1, BTN1A1  
member A1 (Btn1a1), mRNA

AACAGCACACAGCCTTCTTCCTTCTGAAGAGCTCTCTCTTTGGCCCCGGGGTG  
ACAAGCAGCCCTTTTCACTTGATCACTGTGGCTCTGGCTCCCTTTTCTCTGTG  
GTCTGTTCGAAATCGCCTGAAGCTCTTGGCGGGCTTCATTGCCCCAGTTAGCTC  
AGAGATGGCAGTTCCCACCAACTCCTGCCTCCTGGTCTGTCTGCTCACCCTCA  
CTGTCCTACAGCTGCCCACGCTGGATTCGGCAGCTCCCTTCGATGTGACCGCA  
CCTCAGGAGCCAGTGTTGGCCCTAGTGGGCTCAGATGCCGAGCTGACCTGTGG  
CTTTTCCCCAAACGCGAGCTCAGAATACATGGAGCTGCTGTGGTTTCGACAGA  
CGAGGTTCGACAGCGGTACTTCTATACCGGGATGGCCAGGAGCAGGAGGGCCAG  
CAGATGACGGAGTACCGCGGGAGGGCGACGCTGGCGACAGCCGGGCTTCTAGA  
CGGCCGCGCTACTCTGCTGATCCGAGATGTCAGGGTCTCAGACCAGGGGGAGT  
ACCGGTGCCTTTTCAAAGACAACGACGACTTCGAGGAGGCCGCCGTATACCTC  
AAAGTGGCTGCTGTGGGTTTCAAGATCCTCAAATCAGTATGACGGTTCAAGAGAA  
TGGAGAAATGGAGCTGGAGTGCACCTCCTCTGGATGGTACCCAGAGCCTCAGG  
TGCAGTGGAGAACAGGCAACAGAGAGATGCTACCATCCACGTCAGAGTCCAAG  
AAGCATAATGAGGAAGGCCTGTTCACTGTGGCAGTTTCAATGATGATCAGAGA  
CAGCTCCATAAAGAACATGTCCTGCTGCATCCAGAATATCCTCCTTGGCCAGG  
GGAAGGAAGTAGAGATCTCCTTACCAGCTCCCTTCGTGCCAAGGCTGACTCCC  
TGGATAGTAGCTGTGGCTATCATCTTACTGGCCTTAGGATTTCTCACCATTGG  
GTCCATATTTTCACTTGGAACTATAACAAGGAAAGATCCAGTCTGCGGAAGA  
AGGAATTTGGCTCTAAAGAGAGACTTCTGGAAGAACTCAGATGCAAAAAGACT  
GTACTGCATGAAGTTGACGTGACTCTGGATCCAGACACAGCCCACCCCCACCT  
CTTCTGTATGAAGATTCAAAGTCAGTTCGATTGGAAGATTCACGTCAGATCC  
TGCCTGATAGACCAGAGAGATTTGACTCCTGGCCCTGTGTGTTGGGCCGTGAG  
ACCTTTACTTCAGGGAGACATTACTGGGAGGTGGAGGTGGGAGATAGAACTGA  
CTGGGCCATTGGTGTGTGTAGGGAGAATGTGGTGAAGAAAGGGTTTGACCCCA  
TGACTCCTGATAATGGGTTCTGGGCTGTGGAGTTGTATGGAAATGGGTACTGG  
GCCCTCACCCCACTCAGGACCTCTCTCCGATTAGCAGGGCCCCCTCGCAGAGT  
TGGGGTTTTTCTGGACTATGACGCAGGAGACATTTCTTCTACAACATGAGTA  
ACGGATCTCTTATCTATACTTTCCCTAGCATCTCTTTCTCTGGCCCCCTCCGT  
CCCTTCTTTTGTCTGTGGTCCTGTGGTAAAAAGCCCCTGACCATCTGTTCAAC  
TGCCAATGGGCCTGAGAAAGTCACAGTCATTGCTAATGTCCAGGACGACATTC  
CCTTGTCCCCGCTGGGGGAAGGCTGTACTTCTGGAGACAAAGACACTCTCCAT  
TCTAAACTGATCCCGTTCTCACCTAGCCAAGCGGCACCATAACAAATATTCCA  
GCTTCACGACTTTGCCTTCCTTTGACTAATCCCTCATGCCCCGAAGCTTCAGC  
TGTTGGCTTCTTGCAGCCCTGCTTCTTCTTCTGGTGGATGGAGATTAATTCACAT  
TGGGAAGGTTAGGTATGTTGCTGCCAGACAAGGCAGGAAGAAAGGCCATCCTA

ACATCTGCTTCCTGTAAGGCCCTCTGGGCAGAAAGCATGCGCTGGTGTCTCCTCC  
TGATCTGGGCCAGGGGCTGAGGCAGGCTCCCCTCGCCTCAGGAATGATGACA  
GGCACAATAGAAACAACGGGGAACATTTCTGCGAGAGAAAGGTGGCTCTATCAT  
CTTACAATGTCACCTCTCCTCCACCACGGCACAAGTGACCCAGGTCAACTGGG  
AGCAGCAGGACCAGCTTCTGGCCATTTGTAATGCTGACTTGGGGTGGCACATC  
TCCCCATCCTTCAAGGATCGAGTGGCCCCAGGTCCCGGCTGGGCCTCACCT  
CCAGTCGCTGACCGTGAACGATACAGGGGAGTACTTCTGCATCTATCACACCT

[illegible]

MRWCLLIWQGLRQAPLASGMMTGTIETTGNISAEKGGSIILQCHLSSTTAQ  
VTQVNWEQQDQLLAICNADLGWHISPSFKDRVAPGPGLGLTLQSLTVNDTGEY  
FCIYHTYPDGTYTGRIFLEVLESSVAEHGARFQIPLLGAMAATLVVICTAVIV  
VVALTRKKKALRIHSVEGDLRRKSAGQEEWSPSAPSPPGSCVQAEAAPAGLCG  
EQRGEDCAELHDYFNVLSYRSLGNCSFFTETG (SEQ ID NO: 110) Mouse TIGIT  
>NM\_001146325.1:98-823 *Mus musculus* T cell immunoreceptor with Ig and  
ITIM domains (Tigit), mRNA

ATGCATGGCTGGCTGCTCCTGGTCTGGGTCCAGGGGCTGATACAGGCTGCCTT  
CCTCGCTACAGGAGCCACAGCAGGCACGATAGATACAAAGAGGAACATCTCTG  
CAGAGGAAGGTGGCTCTGTTCATCTTACAGTGTCACTTCTCCTCTGACACAGCT  
GAAGTGACCCAAGTCGACTGGAAGCAGCAGGACCAGCTTCTGGCCATTATAG  
TGTTGACCTGGGGTGGCATGTCGCTTCAGTCTTCAGTGATCGGGTGGTCCCAG  
GCCCCAGCCTAGGCCTCACCTTCCAGTCTCTGACAATGAATGACACGGGAGAG  
TACTTCTGTACCTATCATACTGATCCTGGTGGGATTACAAAGGGGAGAATATT  
CCTGAAGGTCCAAGAAAGCTCAGTGGCTCAGTTCCAGACTGCCCCGCTTGGAG  
GAACCATGGCTGCTGTGCTGGGACTCATTGCTTAATGGTCACAGGAGTGACT  
GTACTGGCTAGAAAGAAGTCTATTAGAATGCATTCTATAGAAAGTGGCCTTGG  
GAGAACAGAAGCGGAGCCACAGGAATGGAACCTGAGGAGTCTCTCATCCCCTG  
GAAGCCCTGTCCAGACACAAACTGCCCCCTGCTGGTCCCTGTGGAGAGCAGGCA  
GAAGATGACTATGCTGACCCACAGGAATACTTTAATGTCCTGAGCTACAGAAG  
CCTAGAGAGCTTCATTGCTGTATCGAAGACTGGCTAA (SEQ ID NO: 111)

>NP\_001139797.1 T-cell immunoreceptor with Ig and ITIM domains precursor  
[*Mus musculus*]

MHGWLLLVWVQGLIQAFLATGATAGTIDTKRNISAEEGGSVILQCHFSSDTA  
EVTQVDWKQQDQLLAISVDLGWHVASVFSDRVVPGPSLGLTFQSLTMNDTGE  
YFCTYHTYPGGIYKGRIFLKVQESSVAQFQTAPLGGTMAAVLGLICLMVTGVT  
VLARKKSIRMHSIESGLGRTEAEPQEWNLRLSLSSPGSPVQTQTAPAGPCGEQA  
EDDYADPQEYFNVLSYRSLESFIAVSKTG (SEQ ID NO: 112) Human

>NM\_001252.5 *Homo sapiens* CD70 molecule (CD70), CD27L (CD70) transcript  
variant 1, mRNA

AGAGAGGGGGCAGGCTGGTCCCCTGACAGGTTGAAGCAAGTAGACGCCCAGGAG  
CCCCGGGAGGGGGGCTGCAGTTTCCTTCCTTCCTTCCTTCGCGCAGCGCTCCGCGCC  
CCCATCGCCCCTCCTGCGCTAGCGGAGGTGATCGCCGCGGCGATGCCGGAGGA  
GGGTTCGGGCTGCTCGGTGCGGCGCAGGCCCTATGGGTGCGTCCTGCGGGCTG  
CTTTGGTCCCATTGGTTCGCGGGCTTGGTGATCTGCCTCGTGGTGTGCATCCAG  
CGCTTCGCACAGGCTCAGCAGCAGCTGCCGCTCGAGTCACTTGGGTGGGACGT  
AGCTGAGCTGCAGCTGAATCACACAGGACCTCAGCAGGACCCCAGGCTATACT  
GGCAGGGGGGGCCCAGCACTGGGCGGCTCCTTCCTGCATGGACCAGAGCTGGAC  
AAGGGGCAGCTACGTATCCATCGTGATGGCATCTACATGGTACACATCCAGGT  
GACGCTGGCCATCTGCTCCTCCACGACGGCCTCCAGGCACCACCCACCACCC  
TGGCCGTGGGAATCTGCTCTCCCGCCTCCCGTAGCATCAGCCTGCTGCGTCTC  
AGCTTCCACCAAGGTTGTACCATTGCCTCCCAGCGCCTGACGCCCCTGGCCCCG  
AGGGGACACACTCTGCACCAACCTCACTGGGACACTTTTGCCTTCCCGAAACA  
CTGATGAGACCTTCTTTGGAGTGCAAGTGGGTGCGCCCCTGACCACTGCTGCTG  
ATTAGGGTTTTTTTAAATTTTATTTTATTTTATTTAAGTTCAAGAGAAAAAGTG  
TACACACAGGGGGCCACCCGGGGTGGGGTGGGAGTGTGGTGGGGGGTAGTGGT  
GGCAGGACAAGAGAAGGCATTGAGCTTTTTCTTTTCAATTTTCTATTAAAAAAT  
ACAAAAATCA (SEQ ID NO: 113)>NP\_001243.1 CD70 antigen isoform 1

[*Homo sapiens*]

MPEEGSGCSVRRRPYGCVLRAALVPLVAGLVICLVVCIQRFAQAQQQLPLESL

GWDVAELQLNPLRQYQDRLYQWQGSFLHGPDLKGLHQPDLGIYMV  
HIQVTLAICSSTTASRHHPTTLAVGICSPASRSISLLRLSFHQGCTIASQRLT  
PLARGDTLCTNLTGTLPSRNTDETFFGVQWVRP (SEQ ID NO: 114) Mouse CD27L  
>NM\_011617.2 *Mus musculus* CD70 antigen (Cd70), mRNA (CD70)  
GAAGGTGCCAAAAGCTCCAGGGGATTTCCCTGCCCTCCGAGAAGAGGCCCAGT  
TCTTCCCCTGCATCGGACATCCCCGAGGTTCTAAGGGCAGGTCAAGGCAGGCA  
GAAGCTTCAAAGCTCGGCTGAGGAGGCTACAGCTTCCCGCTGCCTTCAGGCC  
GCTGCTTCCGTGCAGGGATGCCGGAGGAAGGTCGCCCTTGCCCCTGGGTTCGC  
TGGAGCGGGACCGCGTTCCAGCGCCAATGGCCATGGCTGCTGCTGGTGGTGT  
TATTACTGTGTTTTGCTGTTGGTTTCATTGTAGCGGACTACTCAGTAAGCAGC  
AACAGAGGCTGCTGGAGCACCTTGAGCCGCACACAGCTGAGTTACAGCTGAAT  
CTCACAGTTCCTCGGAAGGACCCACACTGCGCTGGGGAGCAGGCCCAGCCTT  
GGGAAGGTCCTTCACACACGGACCAGAGCTGGAGGAGGGCCATCTGCGTATCC  
ATCAAGATGGCCTCTACAGGCTGCATATCCAGGTGACACTGGCCAACCTGCTCT  
TCCCCAGGCAGCACCTTGACAGCAGGGGCCACCCTGGCTGTGGGCATCTGCTC  
CCCCGCTGCGCACGGCATCAGCTTGCTGCGTGGGCGCTTTGGACAGGACTGTA  
CAGTGGCATTACAGCGCCTGACATACCTGGTCCACGGAGATGTCCTCTGTACC  
AACCTCACCTTGCTGCTGCCGTCCCGCAACGCTGATGAGACCTTCTTTGG  
AGTTCAGTGGATATGCCCTTGACCACAACCTCCAGGATGACTTGTGAATATTTT  
TTTTCTTTTCAAGTTCTACGTATTTATAAATGTATATAGTACACATA (SEQ ID NO: 115)  
>NP\_035747.1 CD70 antigen [*Mus musculus*]  
MPEEGRPCPWVRWSGTAFQRQWPWLLLVVFITVFCWFHCSGLLSKQQQRLLLE  
HPEPHTAELQLNLTVPKDPTRLRWGAGPALGRSFTHGPELEEGHLRIHQDGLY  
RLHIQVTLANCSSPGSTLQHRATLAVGICSPAAGHGISLLRGRFGQDCTVALQR  
LTYLVHGDVLCTNLTPLPSRNADETFFGVQWICP (SEQ ID NO: 116) Human  
>NM\_001244.4 *Homo sapiens* TNF superfamily member 8 CD30L (TNFSF8),  
transcript variant 1, mRNA (CD153)  
GTCATTTTCCTACGCGCCCTCTGACATCAGCCACCTTCTCTGTAGCTAGTTTC  
TCTGCACACAACCTTAATCCCTGGCAATGAAAAATGAACCTCTCCCCCACCCTT  
GCTGCCGCTCTCGCCTCACGCCCCCAGAGAAGAGTTTCTCCACCAGGCAGCA  
GGTGAAGGTTTTTTTCCAAGTCACATGATTCAGGATTCAGGGGGAGAATCCTT  
CTTGAACAGAGATGGGCCCAGAACTGATCAGATGAAGAGAGATAAGGTGTG  
ATGTGGGGAAGACTATATAAAGAATGGACCCAGGGCTGCAGCAAGCACTCAAC  
GGAATGGCCCCTCCTGGAGACACAGCCATGCATGTGCCGGCGGGCTCCGTGGC  
CAGCCACCTGGGGACCACGAGCCGCAGCTATTTCTATTTGACCACAGCCACTC  
TGGCTCTGTGCCTTGTCTTCACGGTGGCCACTATTATGGTGTGTTGGTCGTTT  
AGGACGGACTCCATTCCCAACTCACCTGACAACGTCCCCCTCAAAGGAGGAAA  
TTGCTCAGAAGACCTCTTATGTATCCTGAAAAGGGCTCCATTCAAGAAGTCAT  
GGGCCTACCTCCAAGTGGCAAAGCATCTAAACAAAACCAAGTTGTCTTGGAAC  
AAAGATGGCATTCTCCATGGAGTCAGATATCAGGATGGGAATCTGGTGATCCA  
ATTCCCTGGTTTGTACTTCATCATTTGCCAACTGCAGTTTCTTGTACAATGCC  
CAAATAATTCTGTGATCTGAAGTTGGAGCTTCTCATCAACAAGCATATCAAA  
AAACAGGCCCTGGTGACAGTGTGTGAGTCTGGAATGCAAACGAAACACGTATA  
CCAGAATCTCTCTCAATTCTTGCTGGATTACCTGCAGGTCAACACCACCATAT  
CAGTCAATGTGGATACATTCCAGTACATAGATACAAGCACCTTTCCTCTTGAG  
AATGTGTTGTCCATCTTCTTATACAGTAATTCAGACTGAACAGTTTCTCTTGG  
CCTTCAGGAAGAAAGCGCCTCTCTACCATACAGTATTTTCATCCCTCCAAACAC  
TTGGGCAAAAAGAAAACCTTTAGACCAAGACAACTACACAGGGTATTAAATAG  
TATACTTCTCCTTCTGTCTCTTGGAAGATACAGCTCCAGGGTTAAAAAGAGA  
GTTTTTAGTGAAGTATCTTTCAGATAGCAGGCAGGGAAGCAATGTAGTGTGGT

GGGCAGACAGACCCACACACAGCAATCAGAAGGAGGATGGAATGGAATGCGATGTCCCAAGCCAAACC  
TCTAATTCACCTGTATGGTCTTGATCTATTTCTTCTGTTTTGAGAGCCTCCAGT  
TAAAATGGGGCTCCAGTACCAGAGCAGCTAGCAACTCTGCCCTAATGGGAAAT  
GAAGGGGAGCTGGGTGTGAGTGTTTACACTGTGCCCTTCACGGGATACTTCTT  
TTATCTGCAGATGGCCTAATACTTAGTTGTCCAAGTCGCGATCAAGGACTCTC  
TCACACAGGAACTTCCCTATACTGGCAGATACACTTGTGACTGAACCATGCC  
CAGTTTATGCCTGTCTGACTGTCACCTCTGGCACTAGGAGGCTGATCTTGTACT  
CCATATGACCCCAACCCCTAGGAACCCCAAGGGAAAACCAGGCTGGGACAGCCC  
CCTGTTCCCTGAGATGGAAAGCACAAATTTAATACACCACCACAATGGAAAACA  
AGTTCAAAGACTTTTACTTACAGATCCTGGACAGAAAGGGGCATAATGAGTCTG  
AAGGGCAGTCCTCCTTCTCTAGGTTACATGAGGCAGGAATAAGAAGTCAGACA  
GAGACAGCAAGACAGTTAACAATGTAGGTAAAGAAATAGGGTGTGGTCACTCT  
CAATTCACCTGGCAAATGCCTGAATGGTCTGTCTGAAGGAAGCAACAGAGAAGT  
GGGGAATCCAGTCTGCTAGGCAGGAAAGATGCCTCTAAGTTCTTGTCTCTGGC  
CAGAGGTGTGGTATAGAACCAGAAACCCATATCAAGGGTGACTAAGCCCGGCT  
TCTGGTATGAGAAATTAACTTGTATACAAAATGGTTGCCAAGGCAACATAAA  
ATTATAAGAATTCACCTATACCTTCCCCTCCCTGGAACCTCAGGATCCAAGTCTA  
GAAAATGAAAGGACTGGGTTTGAATTGCTTCAAAACCTCTTCCATCTCAGAAG  
ACCAGACCCTGGGAACTGAGATTCCAGACACAATTTTGGAAGCTCTCCAACCA  
AAATAAGGCCCCCCTACCCCAAGTATATAATTGAAGACACTAGTAACACCTGAC  
TGCATCTCATCTCAGCAGAGCCAGAATATGGGGACAAGGTTTCAGGGTGCCCTG  
CTGAATGGTGTGAACAGCAGGATCTCAAGGATGTAATGGAAAGAACTACCACA  
CTGACCATCCAGAATCTAAGAGACCATCTGGGTGTTTGGGAAACCATCTGACG  
AGGCCTGACTCTATTCCAGTTAGATTGACAATAATTGAGCAGCAGGCATTTTT  
CATTTCTGGTCAGGAAAGCATTGTGCCTTTAGCAAACAATCAGTGTGCAACAG  
TGATGTGGTCATCTAGCCAGGGAATGGCTGCTCCATCCCCTGCATAATATATT  
CCTGCTTCAAACACCTCTCAGAAAACCAGTTCCGCGAGGGTTTTTATATCCCC  
ACAAAGTTGTTGAGAGACAATGATGACCCTGGAAGTGGGGAGGAGGACTTCTG  
AGAAACAGCAACCTCTCTCCTGATTGGGGTAGCCATGAGATTTCTCTAGCTAT  
ATCCAACCTTGGCATCTGTACATCATCTTTGGAGGAACATCTTATTTGTGGAAG  
GACCTTGACAAGCCGTTTGAGATGGAATGTAGGCCCTGATGTTATGCTTCAGT  
AAAAAAAGATGGAAGCTTCCCTGCTATACCAAAACATGGAGCAAAATTTGCAT  
TTTTCTCAAGAAGGAGAGAAAAGGAGTAGGACTCCAGCAAAGTTTGTGAGAAG  
GAAAGCTAGAAAAGATTTAAAAGAAAAAAAGAAAGAACAATCAGCAGTGGTG  
GTATGGATGAAAGGGACTTGAGAGAACAAAAATGGCTAAGGGAAAATTTTAAG  
TCATCTGCTGAGCAGTGTGCTGTGTCAACCTCCTCCTAGGTCTCCTCTATGAA  
ATATTTAGTAAAGTCTACATTTCTCTTTAACTCTTTCTGTGAGTAGATTCTTT  
GGGAGAAGCAGGCATTGGAAGAGGTGTTGAATTCAGCAAGCCAAATGGTCTGT  
GGTAAAAAACAAAACAGACTTTGAGACTCAAGGCTAAAAAACAGGGAAATGG  
CTGGCATTGAGTCACACACTAACTGCATAGGACAAATGAATCTTGCTTAAAC  
CAACTCATGCATTCTTGAAAAGGTATATGCAACCCAACTGTGTGTAACTAAG  
CAATTTTTTTGCCATCTCACATTCTAACTCGAGAAAGATTCCATTTTCATTTT  
TCACCAACTGTTCTCTGAGCAGAGGTACCTGACTTTTGCCTGTGAGTGGTTT  
CTAATCTCAGTCTCTGTCAAGCAATGCTAAGAAAGCCAACACCTAAAGACACA  
AGGGGTACATCATTTAAATGAATAATGTAACCAAACAAACAAAAAAGAGAAT  
AATCATTAATAACTCAACTGATAGATATGTAGGGAGTAGGCAACCCAGGAAGT  
TTAAACTAAATTCTGTACTCTTGAGGGTTAACCAGCCCCTGGGAATGTTAT  
GAGCAAATGATACTCCATGAGTAAAATGATATCTATGCAAGTAAAATAAATAA  
TTTATCTAACTGGGAA (SEQ ID NO: 117) >NP\_001235.1 tumor necrosis factor  
ligand superfamily member 8 isoform 1 [*Homo sapiens*]



MDPGLQALMGDLMQAPSTGVSFSLYTLTATLSRGLTATLFT  
VATIMVLVVQRTDSIPNSPDNVPLKGGNCSEDLCLILKRAPFKKSWAYLQVAK  
HLNKTCLSWNKDGLHGVRYQDGNLVIQFPGLYFIICQLQFLVQCPNNSVDLK  
LELLINKHIKKQALVTVCESGMQTKHVYQNLSQFLLDYLQVNTTISVNVDTFQ  
YIDTSTFPLENVLSIFLYSNSD (SEQ ID NO: 118) Mouse CD30L >NM\_009403.3  
*Mus musculus* tumor necrosis factor (CD153) (ligand) superfamily, member 8  
(Tnfsf8), mRNA

AGATTAATCCCAGGCGATGAAAAATGAACCTCTCCCCCACCCTTGCAGCCACC  
CTTCGCCTCACGCCCCCAGAGAAGAGTTTCTCCATCCGGCAACTGGTGAAGGC  
TTTTTTCCAAGTCACATGATCCAGGATGCAGGGGAAAATCCTTCTTGGAACAG  
AGCTGGGTACAGAACCGAATCAGATGAGGAGAGATAAGGTGTGATGTGGGACA  
GACTATATAAAGCATGGAGCCAGGGCTGCAACAAGCAGGCAGCTGTGGGGCTC  
CTTCCCCTGACCCAGCCATGCAGGTGCAGCCCGGCTCGGTAGCCAGCCCCTGG  
AGAAGCACGAGGGCCCTGGAGAAGCACAAGTCGCAGCTACTTCTACCTCAGCAC  
CACCGCACTGGTGTGCCTTGTTGTGGCAGTGGCGATCATTCTGGTACTGGTAG  
TCCAGAAAAAGGACTCCACTCCAAATACAACCTGAGAAGGCCCCCCTTAAAGGA  
GGAAATTGCTCAGAGGATCTCTTCTGTACCCTGAAAAGTACTCCATCCAAGAA  
GTCATGGGCCTACCTCCAAGTGTCAAAGCATCTCAACAATACCAAACCTGTCAT  
GGAACGAAGATGGCACCATCCACGGACTCATATACCAGGACGGGAACCTGATA  
GTCCAATTCCCTGGCTTGACTTCATCGTTTGCCAACTGCAGTTCCTCGTGCA  
GTGCTCAAATCATTCTGTGGACCTGACATTGCAGCTCCTCATCAATTCCAAGA  
TCAAAAAGCAGACGTTGGTAACAGTGTGTGAGTCTGGAGTTCAGAGTAAGAAC  
ATCTACCAGAATCTCTCTCAGTTTTTGTCTGCATTACTTACAGGTCAACTCTAC  
CATATCAGTCAGGGTGGATAATTTCCAGTATGTGGATACAAACACTTTCCCTC  
TTGATAATGTGCTATCCGTCTTCTTATATAGTAGCTCAGACTGAATAGTTGTT  
CTTAACCTTTATGAAAATGCTGTCTACCATACAGTACTTCATCTGTCCAAACA  
TGGGCCAAAGAAAATATTAGGACAACCTCAAACCTAAGCATGTGAGTTAGTGCAC  
TTCTCTTTCTGTCTTTGGAAAAATACAAACCCAGGATTTAGAAAGTGGAGTC  
TCCTTCAGATGCACAAACAGGAAAGAATGTGATATGTGCACAGAGACCTACTT  
GGGCACTAGAAGGGGTTGAGTTGTCCCAGTATAACCACTAATTCAGTACCTT  
GAGCCATTTTTCTTCCCCTGGAACCTGGGGTCTGAATCTGGAAAAGTAGGAG  
ATGAGATTTACATTTCCCAATATTTTCTTCAACTCAGAAGACGAGACTGTGG  
AGCTGAGCTCCCTACACAGATGAAGGCCTCCCATGGCATGAGGAAAATGATGG  
TACCAGTAATGTCTGTCTGACTGTCATCTCAGCAAGTCCTAAGGACTTCCATG  
CTGCCTTGTTGAAAGATACTCTAACCTCTTGTAATGGGCAAAGTGATCCTGTC  
TCTCACTGAGGGGAGTAGCTGCTGCCATCTCCTGAGACATACATGGAGACATT  
TTCTGCCCAAATTCCATTCTGTGTGCAGTTTTTAAGTATTCCCCCAAAGTTC  
TTGACAATGAGAACTTTGAATGTGGGAAGAGCTTCTGGACAGCAAACATTAAAC  
AGCTTCTCCTGACCAGAGAGACCATGCAAGCTTGGTCTTAGACCCATCAAGCT  
TGAGGTTTCTACATTGTGGGAGACAGACTTTTGACAAACCATTTGAGTTGATG  
TCTGGGCCCCTGGGAGTTCTCCTTCAGTAAGGAGAGCAAGCCGTTCTAGTGCT  
GTGTCAGAGGATGGAGTAAAATAGACACTTTTCTGAAGGAAAGGAGAACAAAG  
TTCCAGAAAAAGGCTAGAAAATGTTTAAAAGGAAAAGAAAAAACTCAGCTTTT  
CTCATATGAGAGGAACCCAGAAAAACAACACTGAAAAAGAAGAGTGGCTCTGT  
CAACCTCCTCTTAGGTCTCCTCCTCTCTAGTTATTGGGAAAGGAGTTGCATGG  
TACAGGACAAGTTCTGGTGTGTGGTCAAATAGAATCAGATGTGGAGAACACCA  
TGCAGAGAATAAGGAGACCTGTCATATTTGTGTTGTACTCAAATGAGGGGCAA  
ATGAATCTTAGGCTAAATCAAATAACAGTCTCTGTCAAGCTGTGCTCAGAAAG  
TCAACCACTGAAGATGGAGGGTGAGGCACGTCATTTAAAAAAAGTGAAATGTA GC  
(SEQ ID NO: 119) >NP\_033429.1 tumor necrosis factor ligand superfamily

member 8 [Mus musculus]  
MEPGLQQAGSCGAPSPDPAMQVQPGSVASPWSTRPWRSTSRSYFYLSTTALV  
CLVVAVAILVLVVQKKDSTPNTTEKAPLKGGNCSEDLFCTLKSTPSKKSWAY  
LQVSKHLNNTKLSWNEDGTIHLIYQDGNLIVQFPGLYFIVCQLQFLVQCSNH  
SVDLTLQLLINSKIKKQTLVTVCESGVQSKNIYQNLSSQFLLHYLQVNSTISVR  
VDNFQYVDTNTFPLDNVLSVFLYSSSD (SEQ ID NO: 120) Human >NM\_005092.4  
*Homo sapiens* TNF superfamily member 18 GITRL (TNFSF18), mRNA  
ATCACTTGTGAATTTTTGTTTTCCACAGCTCTCATTTCTCCAAAAATGTGTTT  
GAGCCACTTGGAAAATATGCCTTTAAGCCATTCAAGAACTCAAGGAGCTCAGA  
GATCATCCTGGAAGCTGTGGCTCTTTTGCTCAATAGTTATGTTGCTATTTCTT  
TGCTCCTTCAGTTGGCTAATCTTTATTTTTCTCCAATTAGAGACTGCTAAGGA  
GCCCTGTATGGCTAAGTTTGGACCATTACCCTCAAAATGGCAAATGGCATCTT  
CTGAACCTCCTTGCGTGAATAAGGTGTCTGACTGGAAGCTGGAGATACTTCAG  
AATGGCTTATATTTAATTTATGGCCAAGTGGCTCCCAATGCAAACACTACAATGA  
TGTAGCTCCTTTTGAGGTGCGGCTGTATAAAACAAAGACATGATACAAACTC  
TAACAAACAAATCTAAAATCCAAAATGTAGGAGGGACTTATGAATTGCATGTT  
GGGGACACCATAGACTTGATATTCAACTCTGAGCATCAGGTTCTAAAAAATAA  
TACATACTGGGGTATCATTTTACTAGCAAATCCCCAATTCATCTCCTAGAGAC  
TTGATTTGATCTCCTCATTCCCTTCAGCACATGTAGAGGTGCCAGTGGGTGGA  
TTGGAGGGAGAAGATATTCAATTTCTAGAGTTTGTCTGTCTACAAAAATCAAC  
ACAAACAGAACTCCTCTGCACGTGAATTTTCATCTATCATGCCTATCTGAAAG  
AGACTCAGGGGAAGAGCCAAAGACTTTTGGTTGGATCTGCAGAGATACTTCAT  
TAATCCATGATAAAACAAATATGGATGACAGAGGACATGTGCTTTTCAAAGAA  
TCTTTATCTAATTCTTGAATTCATGAGTGGAAAAATGGAGTTCTATTCCCATG  
GAAGATTTACCTGGTATGCAAAAAGGATCTGGGGCAGTAGCCTGGCTTTGTTT  
TCATATTCTTGGGCTGCTGTAATTCATTCTTCTCATACTCCCATCTTCTGAGA  
CCCTCCCAATAAAAAGTAGACTGATAGGATGGCCACAGATATGCCTACCATAC  
CCTACTTTAGATATGGTGGTGTTAGAAGATAAAGAACAATCTGAGAACTATTG  
GAATAGAGGTACAAGTGGCATAAAATGGAATGTACGCTATCTGGAAATTTCTC  
TTGGTTTTATCTTCCTCAGGATGCAGGGTGCTTTAAAAAGCCTTATCAAAGGA  
GTCATTCCGAACCCTCACGTAGAGCTTTGTGAGACCTTACTGTTGGTGTGTGT  
GTCTAAACATTGCTAATTGTAAAGAAAGAGTAACCATTAGTAATCATTAGGTT  
TAACCCCAAGAAATGGTATTATCATTACTGGATTATGTCATGTAATGATTTAGTA  
TTTTTAGCTAGCTTTCCACAGTTTGCAAAGTGCTTTTCGTAAAACAGTTAGCAA  
TTCTATGAAGTTAATTGGGCAGGCATTTGGGGGAAAATTTTAGTGATGAGAAT  
GTGATAGCATAGCATAGCCAACCTTCCTCAACTCATAGGACAAGTGACTACAA  
GAGGCAATGGGTAGTCCCCTGCATTGCACTGTCTCAGCTTTAGAATTGTTATT  
TCTGCTATCGTGTTATAAGACTCTAAAACCTTAGCGAATTCATTTTCAGGAAG  
CATATTTCCCTTTAGCCCAAGGTGAGCAGAGTGAAGCTACAACAGATCTTTCC  
TTTACCAGCACACTTTTTTTTTTTTTTCTGCCTGAATCAGGGAGATCCAGGAT  
GCTGTTCAGGCCTTATCCCAACCAAATTTCCCTCTTCACTTTGCAGGGCCCAT  
CTTAGTCAAATGTGCTAACTTCTAAAATAATAAATAGCACTAATTCAAATTT  
TTGGACTCTTAAATTAGCTACTTGCAGGTTCTTGTTGAAAGGTATATAATATT  
ACATTGTAAACAAATTTAAAATATTTATGGATATTTGTGAAAAGCTGCATTAT  
GTAAATAATATTACATGTAAAGCTATTTAAAAGAGGTTTTTTTTTGTATTTTG  
TTTAACAAAAATTGCTCAGGAGCATGCTAAGCCTGAGGCCAAGTTGTTTCTTA  
GTATGACTTTTTTAAAAAACATCTGCTGAGTAGCTACAGGGCCAAAGACTTGG  
AGAGCTTGTTTCTGTTGCATTTGCATATCTTCTCAGGAAATTAAAGTGTGTCA  
TACATATGTGTGTGTGTGTGTGTGTGTGTGTGTATATGTGTGTGTGTATATAT  
ATGTATACTTATAAAATCTTGGTGTTCTTGATCTTTGTTGTGTTATAAGCAAT

GTGTGCTGGAGTGTGGGCTGGTGCTAGCTTATAAGCACATATTATTATTTTCA  
GGAATGTTGCACTTTAGTTATTAACATATAGGCATTCTTGAAATTGGCTATGGT  
GGGAGTATTTATACCATGTAAATTGGCAAACACTACACATTTTTCCTTTTGGAC  
AGCTAGTTCACCAGCACACCACTGTGAAACTCTCCTTAATGACTCCTCTCTGC  
CCCCGCTTCATTCCTGGGATAATCATAGCAGACTAAGGGAGAAAATGAAATTG  
TAAAAATTTGGCATACTGGTGATTTCTCAGGGCAAGCAGAGGTTACTACAGCT  
GCAGCTAGAGGGATGACTACCAACAGGTGACCTTTACATTTTTCCTGATGTTAT  
AATTTTAGCTTTTGTGTTTCAATGTATACTGTTTTCTCTCCACATAGTA  
GTCTGCATTTTAAATCTATAATAAAACATGCTGATAACTGG (SEQ ID NO: 121)  
>NP\_005083.3 tumor necrosis factor ligand superfamily member 18 [*Homo sapiens*] MCLSHLENMPLSHSRTQGAQRSSWKLWLFCSIVMLLFLCSFSWLIFLQLET  
AKEPCMAKFGPLPSKWQMASSEPPCVNKVSDWKLEILQNGLYLIYGQVAPNAN  
YNDVAPFEVRLYKNKMDMIQTLTNKSKIQNVGGTYELHVGDTIDLIFNSEHQVL  
KNNTYWGIILLANPQFIS (SEQ ID NO: 122) Mouse GITRL >NM\_183391.3 *Mus musculus* tumor necrosis factor (ligand) superfamily, member 18 (Tnfsf18), mRNA  
TTGTGGGTATCTGCTTTCCCCAGTTCTCATTCCATCAGAGAACGAGTTCTAGC  
CTCATGGAGGAAATGCCTTTGAGAGAATCAAGTCCTCAAAGGGCAGAGAGGTG  
CAAGAAGTCATGGCTCTTGTGCATAGTGGCTCTGTTACTGATGTTGCTCTGTT  
CTTTGGGTACACTGATCTATACTTCACTCAAGCCAACTGCCATCGAGTCCTGC  
ATGGTTAAGTTTGAAGTATCATCCTCAAATGGCACATGACATCTCCCAAACC  
TCACTGTGTGAATACGACATCTGATGGGAAGCTGAAGATACTGCAGAGTGGCA  
CATATTTAATCTACGGCCAAGTGATTCTGTGGATAAGAAATACATAAAAGAC  
AATGCCCCCTTCGTAGTACAGATATATAAAAAGAATGATGTCCTACAAACTCT  
AATGAATGATTTTCAAATCTTGCCTATAGGAGGGGTTTATGAAGTGCATGCTG  
GAGATAACATATATCTGAAGTTCAACTCTAAAGACCATATTCAGAAAATAAC  
ACATACTGGGGGATCATCTTAATGCCTGATCTACCATTTCATCTCTTAGAGATT  
GGGTTTGGTCTCCTCATCTTCTTCTTTGTATCCCGAGATGCTGGTGGGTGGGT  
TGGAGGGGGATGATTGATGGCAATGCACACAGTTTGTGAGGGCTTACAAATTG  
ACACAATCAGAGCCTCTTGGCATATAAAATTTTAGCCCTCATATCTGTCTGAA  
GAGGACTCAGCAAATGGGCCAATCCCTAATGTTGGGTCTGCAAATGGACTTGT  
ACAATCCATGATAAAAAGGAGTATGGGCCACAGAAGACAGAAACTCTTCCAAA  
GAATGTCTTTCTAACCTTGATCCCTGGGTAGAATGAGATCCTGTTTCCATGGG  
AGTCTTACTTGGCTTGCAAAAAAGGGTGTAGGGCAGTAGCTTGGCCTTTTTTC  
CATCATAATTTCTTGAGCTGTTTTACCTTAATCCCTCCAAACTCTCACCTTC  
TGAGAGCCTCCTAATGAAACATTGTTAGACTGGTGGGGTGGCCAAGACATGCC  
AACAACACCCTTCTTTAGAGGTGGTGTTTTTAGAGGACAGAGAACATTATGAA  
GCCTAGAGCAGCAGAGGTCAAGATGCCACGAAATGGAATTGATCTGGGAATTT  
TTTTTTTTTTTTCATTCTCAGGATGCAGGTTCAATTCTGAACTTTCCCCTAGGCC  
TTCATTGCTTTTGTGTGTATGTGTGCATAAATTCTGCAAATAGAAAAATGAGA  
GTTTGCACCAGTACTCACTAGATTTAACACCAGAAAGTGGTACTTTTCTGGCT  
GTATTATGCCATGATAGCACATTTTCTGTTGGTGTTCCTAACTGACAAGTAT  
AACAGTTTTCTTAAACCACACAACAATGCTATGATGTTAATGGGGTAGATATT  
TTTGGAAAAAAATTGCACAGTGAGAACATGGGTAGATGAACCCTAAGACTCTT  
ACCTCAATTCAGAACTCGCAAGGAGTTAAGTGAGTGGGGTCTTCATTAGACCA  
TTCACATGGTCTCTGCTTTGAAACTGGCGTTGCTACTGTCTCATTATACATCA  
CTAAATGGAATTAACCTCAACTTTGAAATGGATGCATCGACTTTACCCCAAGG  
TGTCCAGAATGAAGCTACAAGACTTTTACCAGCAGTCATTTTCTTTTGCCTG  
GAGCAAGAAGATCCAGGATACTGTTGGAAGAGTTCATCTCACTCAACCATGCT  
GACTTTCCAAAGTAATAATGAACATTTGTGTTCAAATTTTGGATTCTGTTAAA  
TTTAGCCAGCTTGTGAGTTCTTGTGCGAAAAGTATTTTAAACCAATTTACACTA

TTTATGGTATTGTGAAACGCTAAGTATTTTATATATACTAATTT  
AAAATATTTTTATTTTATGTAACAAAAATACTATAGGCTAAGCTATTTCTTCT  
TATTTTTTTATGAATACTTGCTGAATTGCCATAGGGGCACAAAGACTCTTCTGT  
TTGCATATCTTCTCAGGAAATTAATAATTGTATCACATGTATTTATAAGAA (SEQ ID  
NO: 123) >NP\_899247.3 tumor necrosis factor ligand superfamily member 18  
[*Mus musculus*]

MEEMPLRESSPQRAERCKKSWLLCIVALLMLLCSLGTLIYTSLKPTAIESCM  
VKFELSSSKWHMTSPKPHCVNTTSDGKLKILQSGTYLIYGQVIPVDKKYIKDN  
APFVVQIYKKNDVLQTLMNDFQILPIGGVYELHAGDNIYLKFNSKDHQKNT  
YWGIILMPDLPFIS (SEQ ID NO: 124) Human >NM\_000074.3 *Homo sapiens*  
CD40 ligand (CD40LG), mRNA CD40L

AATCCTGAGTAAGGTGGCCACTTTGACAGTCTTCTCATGCTGCCTCTGCCACC (CD154)  
TTCTCTGCCAGAAGATAACCATTTCAACTTTAACACAGCATGATCGAAACATAC  
AACCAAACCTTCTCCCCGATCTGCGGCCACTGGACTGCCCATCAGCATGAAAAT  
TTTTATGTATTTACTTACTGTTTTTCTTATCACCCAGATGATTGGGTCAGCAC  
TTTTTGCTGTGTATCTTCATAGAAGGTTGGACAAGATAGAAGATGAAAGGAAT  
CTTCATGAAGATTTTGTATTCATGAAAACGATACAGAGATGCAACACAGGAGA  
AAGATCCTTATCCTTACTGAACTGTGAGGAGATTAAAAGCCAGTTTGAAGGCT  
TTGTGAAGGATATAATGTTAAACAAAGAGGAGACGAAGAAAGAAAACAGCTTT  
GAAATGCAAAAAGGTGATCAGAATCCTCAAATTGCGGCACATGTCATAAGTGA  
GGCCAGCAGTAAAACAACATCTGTGTTACAGTGGGCTGAAAAAGGATACTACA  
CCATGAGCAACAACCTTGTAACCCTGGAAAATGGGAAACAGCTGACCGTTAAA  
AGACAAGGACTCTATTATATCTATGCCCAAGTCACCTTCTGTTCCAATCGGGA  
AGCTTCGAGTCAAGCTCCATTTATAGCCAGCCTCTGCCTAAAGTCCCCCGGTA  
GATTCGAGAGAATCTTACTCAGAGCTGCAAATACCCACAGTTCCGCCAAACCT  
TGCGGGCAACAATCCATTCCTTGGGAGGAGTATTTGAATTGCAACCAGGTGC  
TTCGGTGTTTGTCAATGTGACTGATCCAAGCCAAGTGAGCCATGGCACTGGCT  
TCACGTCCTTTGGCTTACTCAAACCTCTGAACAGTGTCACCTTGCAAGGCTGTGG  
TGGAGCTGACGCTGGGAGTCTTCATAATACAGCACAGCGGTAAAGCCCACCCC  
CTGTAACTGCCTATTTATAACCCTAGGATCCTCCTTATGGAGAACTATTTAT  
TATACTCCAAGGCATGTAGAAGTGTAAATAGTGAATTACAGGTCACATGAA  
ACCAAACCGGGCCCTGCTCCATAAGAGCTTATATATCTGAAGCAGCAACCCCA  
CTGATGCAGACATCCAGAGAGTCCTATGAAAAGACAAGGCCATTATGCACAGG  
TTGAATTCTGAGTAAACAGCAGATAACTTGCCAAGTTCAGTTTTGTTTCTTTG  
CGTGCAGTGTCTTTCCATGGATAATGCATTTGATTTATCAGTGAAGATGCAGA  
AGGGAAATGGGGAGCCTCAGCTCACATTCAGTTATGGTTGACTCTGGGTTCCT  
ATGGCCTTGTTGGAGGGGGCCAGGCTCTAGAACGTCTAACACAGTGGAGAACC  
GAAACCCCCCCCCCCCCCCCCCGCCACCCTCTCGGACAGTTATTCATTCTCTTTC  
AATCTCTCTCTCTCCATCTCTCTCTTTTCACTCTCTCTCTCAACCTCTTTCT  
TCCAATCTCTCTTTCTCAATCTCTCTGTTTCCCTTTGTCAGTCTCTTCCCTCC  
CCCAGTCTCTCTTCTCAATCCCCCTTTCTAACACACACACACACACACACACA  
CACACACACACACACACACACACACACACAGAGTCAGGCCGTTGCTAGTCAGT  
TCTCTTCTTTCCACCCTGTCCCTATCTCTACCACTATAGATGAGGGTGAGGAG  
TAGGGAGTGCAGCCCTGAGCCTGCCCCTCCTCATTACGAAATGACTGTATTT  
AAAGGAAATCTATTGTATCTACCTGCAGTCTCCATTGTTTCCAGAGTGAACCT  
GTAATTATCTTGTTATTTATTTTTTTGAATAATAAAGACCTCTTAACATTA (SEQ ID  
NO: 125) >NP\_000065.1 CD40 ligand [*Homo sapiens*]

MIETYNQTSRSAATGLPISMKIFMYLLTVFLITQMIGSALFAVYLRRLDKI  
EDERNLHEDFVFMKTIQRCNTGERSLSLLNCEEIKSQFEGFVKDIMLNKEETK  
KENSFEMQKGDQNPQIAAHVISEASSKTTSVLQWAEKGYTMSNNLVTLENGK

QLTVKRLGLYYIYAQVTFCSNREPSSQAPFIKLSPLGRFILLRAANTH  
SSAKPCGQQSIHLGGVFELQPGASVFNVTDP SQVSHGTGFTSFGLLLKL (SEQ ID NO:  
126) Mouse CD40L >NM\_011616.2 *Mus musculus* CD40 ligand (Cd40lg), mRNA  
CTTTCAGTCAGCATGATAGAAACATACAGCCAACCTTCCCCCAGATCCGTGGC  
AACTGGACTTCCAGCGAGCATGAAGATTTTTATGTATTTACTTACTGTTTTCC  
TTATCACCCAAATGATTGGATCTGTGCTTTTTTGCTGTGTATCTTCATAGAAGA  
TTGGATAAGGTCGAAGAGGAAGTAAACCTTCATGAAGATTTTGTATTTCATAAA  
AAAGCTAAAGAGATGCAACAAAGGAGAAGGATCTTTATCCTTGCTGAACTGTG  
AGGAGATGAGAAGGCAATTTGAAGACCTTGTCAAGGATATAACGTTAAACAAA  
GAAGAGAAAAAAGAAAACAGCTTTGAAATGCAAAGAGGTGATGAGGATCCTCA  
AATTGCAGCACACGTTGTAAGCGAAGCCAACAGTAATGCAGCATCCGTTCTAC  
AGTGGGCCAAGAAAGGATATTATACCATGAAAAGCAACTTGGTAATGCTTGAA  
AATGGGAAACAGCTGACGGTTAAAAGAGAAGGACTCTATTATGTCTACACTCA  
AGTCACCTTCTGCTCTAATCGGGAGCCTTCGAGTCAACGCCCATTCATCGTCG  
GCCTCTGGCTGAAGCCCAGCAGTGGATCTGAGAGAATCTTACTCAAGGCGGCA  
AATACCCACAGTTCCTCCCAGCTTTGCGAGCAGCAGTCTGTTCACTTGGGCGG  
AGTGTTTGAATTACAAGCTGGTGCTTCTGTGTTTGTCAACGTGACTGAAGCAA  
GCCAAGTGATCCACAGAGTTGGCTTCTCATCTTTTGGCTTACTCAAACCTCTGA  
ACAGTGCGCTGTCCTAGGCTGCAGCAGGGCTGATGCTGGCAGTCTTCCCTATA  
CAGCAAGTCAGTTAGGACCTGCCCTGTGTTGAACTGCCTATTTATAACCCTAG  
GATCCTCCTCATGGAGAACTATTTATTATGTACCCCCAAGGCACATAGAGCTG  
GAATAAGAGAATTACAGGGCAGGCCAAAATCCCAAGGGACCCTGCTCCCTAAG  
AACTTACAATCTGAAACAGCAACCCCACTGATTCAGACAACCAGAAAAGACAA  
AGCCATAATACACAGATGACAGAGCTCTGATGAAACAACAGATAACTAATGAG  
CACAGTTTTGTTGTTTTATGGGTGTGTCGTTCAATGGACAGTGTACTTGACTT  
ACCAGGGAAGATGCAGAAGGGCAACTGTGAGCCTCAGCTCACAATCTGTTATG  
GTTGACCTGGGCTCCCTGCGGCCCTAGTAGG (SEQ ID NO: 127) >NP\_035746.2  
CD40 ligand [*Mus musculus*]

MIETYSQPSRSVATGLPASMKIFMYLLTVFLITQMIGSVLFAVYLHRRLDKV  
EEEVNLHEDFVFIKKLKRCNKGEGLSLLNCEEMRRQFEDLVKDITLNKEEKK  
ENSFEMQRGDEDPQIAAHVVSEANSNAASVLQWAKKGYTMMKSNLVMLENGKQ  
LTVKREGLYYVYTQVTFCSNREPSSQRPFIIVGLWLKPSSGSERILLKAANTHS  
SSQLCEQQSVHLGGVFELQAGASVFNVTESQVIHRVGFSSFGLLKL (SEQ ID NO:  
128) Human >NM\_003807.5 *Homo sapiens* TNF superfamily member 14 LIGHT  
(TNFSF14), transcript variant 1, mRNA (CD258)  
CGAGACTCCATCTCAAAAACAAAACAAATAAACGAACAAAAAAACCCACAACG  
TATTATTTTCTTGTTTACGAGGTTTCTTGTCTCTCTGGCTCCACCAGAAGAGG  
AGCAGGGACCCTTCTTGCTGTTGTTTCATTGCTGCATCCCCCACACCGAGAGCA  
GAGCCTGGCATGGGCAGAAAGTCCTCAGTCGATATTTGGTGGCCCCAAGCGAA  
TGAAGCATCCAAGAAGGGAAAGCTGGGGGCTCCCCACTGCACTTGCCACCTGA  
GTCACATTTTCAGAAGCCTCTGGAAAGTCGTGCACAGCCCAGGAGTGTTGAGC  
AATTCGGTTTTCTCTGAGGTTGAAGGACCCAGGCGTGTGAGCCCTGCTCCAG  
ACACCTTGGGCATGGAGGAGAGTGTCGTACGGCCCTCAGTGTTTGTGGTGAT  
GGACAGACCGACATCCCATTCACGAGGCTGGGACGAAGCCACCGGAGACAGTC  
GTGCAGTGTGGCCCCGGGTGGGTCTGGGTCTCTTGCTGTTGCTGATGGGGGCGG  
GGCTGGCCGTCCAAGGCTGGTTCCTCCTGCAGCTGCACTGGCGTCTAGGAGAG  
ATGGTCACCCGCCTGCCTGACGGACCTGCAGGCTCCTGGGAGCAGCTGATACA  
AGAGCGAAGGTCTCACGAGGTCAACCCAGCAGCGCATCTCACAGGGGCCAACT  
CCAGCTTGACCGGCAGCGGGGGGCCGCTGTTATGGGAGACTCAGCTGGGCCTG  
GCCTTCCTGAGGGGCCTCAGCTACCACGATGGGGCCCTTGTGGTCACCAAAGC

[illegible]



GTATTTTGTGGAGGAGGAGGACCTCTCCAGGACTCAGGACTCAAAACCCAGCAATAGAGGTT  
TGAAGTCCTCCCTTTAAGGAGCCCTGAACTCTGCAGTGCTCGGGGCGGTGTAG  
ACTGOTGACCTGCTTTGGGCAATCTTCAAATCAGAGACCTGGAGACTTGGGGC  
GTGGAGCCCAGGAGCGAGGGGTCAGCTCATTTCCTGATATTCAGGAAGAAAG  
AATCAAGCTGGGGTATTTATGCTTCTGATGCAAACACTGAGATTTTCGGCTTTC  
TGGGTTTTTGAGCTGGAGGCAAGAAACCTTCCCAGAGTGTCATCAGGACCATGT  
TGGCAGGACTTGGGGCTCCAGACTTGCCACCACACTCTGGCCTCTCCCATCCA  
TCCGCTGCATTGGTTTCCAGCCACCAAAACAGCACTGGCCCCCTGGCTGCAAC  
TGGCCAGGTACGAGCTTCTGAGCACCTACATTCCTCAGGGACATCTTGATGAG  
ATCTCAGTACTCAGTCCAATGCGCAGCAGCGACAGACATGCCAGGAATGGTTG  
GTCAGAAGGGAAGGGAGGAAAGGGAGGAAAGAAGGGAATGCAGAAGAGAAGGG  
GGGAAAACAAGACCAAAACAAAACAGCAACAACAAAGCGGCAGGGAGGAGGTG  
ACACCCTTGGGGATACTTTAGTCAACACACTTAGAACAGATTGTGCCAGGCCT  
GTTGGATTCCTGGAGTTGATGGGATCGTGGGAAGGCACAATGGGGAGCAAGTG  
GGCTTGGGTTATGGCTCAGTGGGTAAAGTGCAATTATGGGGATCTGAGTTTGA  
ATCCCTGGTACCCATATAAAGACACAGATGCGGTGATGGGCACTTGTGACAAT  
GAGATCATCAATAGGGAATGGAGACAGGAGGGACCTCTGGGGTTCACTGGCCA  
GGCAGTCTAGCTGAATCAAAGAGCTCCAAGTTCAGTCGATAGCTCCTGAAGAT  
GACAACTGAGGCTATTCTCCAAACCCACACGCAGGACACATGCGTAATAAAT  
AAAATTTTAAAAAT (SEQ ID NO: 131) >NP\_062291.1 tumor necrosis factor  
ligand superfamily member 14 [*Mus musculus*]  
MESVVQPSV FVVDGQTDIPFRRLEQNHRRRRRCGT VQVSLALVLLLGAGLATQG  
WFLRLHQLRGDIVAHL PDGGKGSWEKLIQDQRSHQANPAHLTGANASLIGI  
GGPLLWETRLGLAFLRGLTYHDGALVTMEPGY YYYVYSKVQLSGVGC PQGLANG  
LPITHGLYKRTSRYPKLELLVSRRSPCGRANSSRVWWDSSFLGGVVHLEAGE  
EVVVRVPGNRLVRPRDGTRSYFGAFMV (SEQ ID NO: 132) Human TL1  
>NM\_005118.4 *Homo sapiens* TNF superfamily member 15 (TNFSF15), transcript  
variant 1, mRNA  
AGAGGTGCCTCCAGGAGCAGCAGGAGCATGGCCGAGGATCTGGGACTGAGCTT  
TGGGGAAACAGCCAGTGTGGAAATGCTGCCAGAGCACGGCAGCTGCAGGCCCA  
AGGCCAGGAGCAGCAGCGCACGCTGGGCTCTCACCTGCTGCCTGGTGTGCTC  
CCCTTCCTTG CAGGACTCACCACATACCTGCTTGTCAGCCAGCTCCGGGCCCCA  
GGGAGAGGCCTGTGTGCAGTTCCAGGCTCTAAAAGGACAGGAGTTTGCACCTT  
CACATCAGCAAGTTTATGCACCTCTTAGAGCAGACGGAGATAAGCCAAGGGCA  
CACCTGACAGTTGTGAGACAACTCCCACACAGCACTTTAAAAATCAGTTCCC  
AGCTCTGCACTGGGAACATGAACTAGGCCTGGCCTTCACCAAGAACC GAATGA  
ACTATACCAACAAATTCCTGCTGATCCCAGAGTCGGGAGACTACTTCATTAC  
TCCCAGGTCACATTCCGTGGGATGACCTCTGAGTGCAGTGAAATCAGACAAGC  
AGGCCGACCAACAAGCCAGACTCCATCACTGTGGTCATCACCAAGGTAACAG  
ACAGCTACCCTGAGCCAACCCAGCTCCTCATGGGGACCAAGTCTGTATGCGAA  
GTAGGTAGCAACTGGTTCCAGCCCATCTACCTCGGAGCCATGTTCTCCTTGCA  
AGAAGGGGACAAGCTAATGGTGAACGTCAGTGACATCTCTTTGGTGGATTACA  
CAAAAGAAGATAAAACCTTCTTTGGAGCCTTCTTACTATAGGAGGAGAGCAAA  
TATCATTATATGAAAGTCCTCTGCCACCGAGTTCCTAATTTTCTTTGTTCAA  
TGTAATTATAACCAGGGGTTTTCTTGGGGCCGGGAGTAGGGGGGCATTCCACAG  
GGACAACGGTTTAGCTATGAAATTTGGGGCCCCAAAATTTACACTTCATGTGC  
CTTACTGATGAGAGTACTAACTGGAAAAAGGCTGAAGAGAGCAAATATATTAT  
TAAGATGGGTTGGAGGATTGGCGAGTTTCTAAATATTAAGACACTGATCACTA  
AATGAATGGATGATCTACTCGGGTCAGGATTGAAAGAGAAATATTTCAACACC  
TTCCTGCTATACAATGGTCAACAGTGGTCCAGTTATTGTTCAATTTGATCATA



AAATTGCTTCAATTCAGAGCTTTGAGTCCCTAAGGAAAGCTTCAAAAA  
CAGTATAAACTTTTCAGAGGCAAAATCCTTCACCAATTTTTTCCACATACTTTCA  
TGCCTTGCCTAAAAAAAATGAAAAGAGAGTTGGTATGTCTCATGAATGTTTCAC  
ACAGAAGGAGTTGGTTTTTCATGTCTACAGCATATGAGAAAAGCTACCTTT  
CTTTTGATTATGTACACAGATATCTAAATAAGGAAGTATGAGTTTCACATGTA  
TATCAAAAATACAACAGTTGCTTGTATTTCAGTAGAGTTTTCTTGCCACCTAT  
TTTGTGCTGGGTTCTACCTTAACCCAGAAGACACTATGAAAAACAAGACAGAC  
TCCACTCAAATTTATATGAACACCACTAGATACTTCCTGATCAAACATCAGT  
CAACATACTCTAAAGAATAACTCCAAGTCTTGGCCAGGCGCAGTGGCTCACAC  
CTGTAATCCCAACACTTTGGGAGGCCAAGGTGGGTGGATCATCTAAGGCCGGG  
AGTTCAAGACCAGCCTGACCAACGTGGAGAAACCCCATCTCTACTAAAAATAC  
AAAATTAGCCGGGCGTGGTAGCGCATGGCTGTAATCCTGGCTACTCAGGAGGC  
CGAGGCAGAAGAATTGCTTGAACCTGGGGAGGCAGAGGTTGCGGTGAGCCCAGA  
TCGCGCCATTGCACTCCAGCCTGGGTAACAAGAGCAAACTCTGTCCAAAAAA  
AAAAAAATAAAATAATAACTCCAAGCCTTTAAAAAATATCATCTGAACTGTT  
ACATCAGATTTCTGGCACTCTACTGACTGTGGAAGATAGCCAGCTGACTGGAA  
GATAGCCAGCTGATTAGTTCCCTGAAGAAACCTGAAGACAGATACCTGGTTAA  
CTAGATCAACTACACTGCCAACTTGTTTGATGCTGAGAGACAATGGACTTATT  
CCATGGGGGAAGGGAAAAAAGAAGTCAATCACCAAATCTGAAGAAGTTAACCT  
AGATCTTTGAGGTTTGATTTGCAACTTTATATGCAGAGTATTATGTGGGTATT  
TTCCCTTAAATATTCAAAGGGATTACATATGGGATTAGCTAATGAGCCTAG  
CCAAGACCTTCCCTGGAGGACAGGCTGGTCATTGCGGAGGTCCCTTCTGTGCT  
TCAGTGGGTTCATATCCTCTAGTCCGTATGATTTTCCTACGCTAATATGTCAA  
GGGCAGGAGAGGCAGCTCTGTTCTCCTAGCCTTTGTTGACTTGTCTGCAAAGC  
AGGAATCTGCCCATTTGTTTCCAAGGAGCAAATGAGCTCATGAGAATGAAAGA  
TGTTAACTTCATGCATTCTGTGCCATCTGAGCATTTCGGTATTATATGACTGG  
TGACCCTTGGCCCGTATTATAAATGCTTCCTATCCTGGGAGACCTCATGGATG  
AGTCTGAGAGGAAATTTGGCACCAAAATCACTCTCACTCTGGTTTCCAGTAGA  
CTATAGAGGCAGAGAGGCATTTGAGAGGCTCCTGAGCAAAGTGTCCAGTGTA  
CAGGAGCACTTCATTAATATTTATTGAGTTATAATTAAATAAAAATTAATTT  
TGATTTCTCAGTTTGGAGGTTAAGGCTCTAAATATATTTTCTAACCTCTGCTA  
GGCTAACTTAAGCCAGGCCTTTTTCTTGCTTCCCTTTCTCAAAACAGTCAGC  
ACAGACTCAGTGGGAGCACAGAGGAGTGTGGTCACCTCCACCTGGCTCACCAG  
AGTCTTCATAGAGGAAGTGAAGCCTGGAAGAACTGGGCGGGCCCCAGATGAC  
CACAGGGAAAGGGCATCTCAGATGGAGGAATTACCCTTGACTTAAAGCAGAAA  
AGAAAGATTTCTCAGTAACTCCAAAACCTTGCTTGATAGGAGAATATTCCCTCA  
ACCAATTCCTAGGACAATATTTATTGGTAGATCAAGAATGTTTCCTCAATAAC  
TCTAGTCTAGCTCCATGATCAGAACTAACACCCATTAAAAACATAAAATGTT  
TTTCTGAACCGGTCTTCATGGTGCGTGAGAGCACCAAGCAGCTTTGGTATGCA  
GGAGGAGTTTTGCACAGAAGAGTGGCCTGCTCAAACCTGCCCCTGTTCTGTA  
GGTGATCTGGTGGATCTGGAAATTTATCCCAAGACAGGAATTTCTAATATTC  
GAAGACATTTGAGGCTTTGGGAAATTTCTCTGCTGTGCATTTATTTGGCTCCTG  
TCATAAGCTTGTTTTTTTAAAGAATGTATCATAGCTCAAGTTTTTACTGCTGAT  
TTTGTTAAATTCTGTATAGTATATTTTTTACGGAAAGGCACAGTCAGACATTC  
CTAATAGGGCTCATGTCAGAACTTCTGTTCCCAAGGCATTATCTCCATAGCAA  
AAATTAGTGCCTGTTTTTCAAAGTGAGGTGGGAAAATGCTTTTTAAGATCATG  
TGATGTTCCCTAAAAGGGGTAAATGGGGTGTATTCAGGGTTTGGGAGGGAGG  
AAGAAGCATGCTTTAGAAAACAGTAAATTTAGGGAGAAAATGCTTTGTTGGTT  
AAATGTCCTCAAAGGCTGAATTCAAATCAATTCACAAACATTTACTGAGT  
ACCTACTGCCCCTGGGGACACAGAGATAAATTATTTAGTCTCAGACACACTCA

TTCTAAGTTCCTCCAGCAGCACTCTCTAATCTTCTTAACTTTGTTT  
GTATTAGCTAATTAATTCGTAAACTTTTAGGCACATGGATCTATTCTCATTATG  
AAAATGGATGCCATTTGATTAAGGCTGATGACTAACAAAATGATTTGTGTTA  
CTCGAAGTGTTTTTTTTTAAAAATAGCTACTCAAGGATAGTTTTCCATAAATCAA  
GAAGGTAAAAAAGTTCCCATTTTTTTATTGTAGAATCCATTATTTAAACTACAT  
GTAGAGACAGGTTATTATTTGCTATATTCAAGTTTGGTCATCAATACCCTTAA  
AAATATTAGAATTTTATGGATGACCCAGAAATGCTTTGAAAATCTGTGTTCT  
CAGCAAATACAGAGACCATGATCAAAATGCACAGAATCACTAACATTTTGATG  
CTAGCATGGTTTCAGTCTATTTGGCAGAACAGAATTGATTATGCTACTAAAAT  
TTCTTTTTCTTTTTTTTTTTTTTTTTTTTTTTTGGAGACAGAGTCTTGCTTTGTCAC  
CCAGGCTGAAGTGCAGTGGCAGGATCTCAGTTCCTGCAACCTCTGCCTCCCA  
GGTTCACGCCATTCTCCTGCTTCAGCCTCCCGAGTAGCTGGGACTACAGGCTC  
CCACCACCATGCCCGGCTAATTTTTTGCATTTTTTAGTAGAGACGGGGTTTCAC  
CGTGTTAGCCAGGATGGTCTCGATCTCCTGACCTCGTGATCCGCCCCGCCTCAG  
CCTTCCAAAGTGCTGGGATTACAGGCGTGAGCCACTGTGCCCGGACTCTGATT  
TTTTTTTTTACTAAGGTACAGTAAGAAAAGGGAAAAGTGACGTTTTCACTTCC  
TGAAATATGTCAGGTTGAATCAATAATAGAGCACACCAGAACTCTTGGCTCCA  
TTTCAACCTAAACTATTCAGTTCTCATCACCCCAGAGGAAATTCCGCCTCTGT  
GCTGGTCAGTAATCCCCCTGGATTATAAAAGTTTAACTAACTCACTGTGCACA  
AGGCACGGCCATTGCCAACATTCTCTTGCAAGGTATTTTCCCAAGCCCTTACC  
CAATTCTGTTTCCATGATTGTGACATTGGGGATTAATTCTGCAAGACAGAACT  
GTTTATATTCTGTACCTTAAAAACACATGCAAACATCTCTTGCTTAAGATTT  
CTGGCTTTTCTATGGCCCAGAGTCCTAGAAGTGTTTTGATATTTGTAGCAGAA  
TTTTCAAGTGTACATCCTTATCCTGGATATTAACATTTTTTGCATCATATTGGC  
AGCTGGACCTACAGAGAATTTAGTAGACTGTAAACCTAATAAGCCTTGAATCC  
TTTTGCACCAGTGGTGAGAGAATGTGGATCAGAGCCATCACCTCCATGCCCCG  
TCACCCTCTAACAACCACATTTACAACCTTCCCCAGCTCTGAGACACACTTGCC  
TCCACCCCTTCCATCACCCCATTTTAAAGATGAAAATACCACACCAGCCTGGAA  
GGAAGAAGTTACTTGCCCAGGGCCACATAGTGAGTTAAGGGCTGATCTAGAGC  
TAGGAAGCTGTCTTCCTGAACCATAATCCTGGACTCTTCTAACCTCTCTACTC  
ATCGCAAATAGAGTTCATTTTAGTGATTTGAAGGAAGATGGGACAAGTATTTT  
CAAACACCTGTAGGACAACATGGAAGTGGGAGGAGACTTCTACTGTAGCTCCC  
CAGAGAAGAGAGCTAGGGCTACAGAGTTGCAGTTACAAGGTTGCCCTCTCTGG  
CTTGATCCCCAAAGGAATTTTCTACTCCAAAATAGAATTTTTTCTAGGATGCTA  
TTTCTCAGTCCCTGGAGATACTCAAACAAAGGGCTTGTCACAAGGGTTTTTGT  
AGAAGCTATTCTTCACAGAGGTTGGGGGAGAGATTAAAGCCAAAGGATCTCTGA  
GGTCTTTTTCAAATCTATAATTATGTGGCCTTTTGTTTCATTGACTTCCATGTG  
TTCTAGTTGATCATTACAAACCTGGCAGGCCTTCTCAAGGGTTCAGTAATTAG  
CTGTCAATTTCCCATTTGTCCAGAGAGTGTTCCAACACAAAATAACCCCTAAGATC  
TTGGCCAATAGAGAAATGTCATGGAATTTTAGAAATGACAGTATCTGCGGAGT  
TTATTCCAAGTTATATCATTTCAAAGATGAAGAAACCCAGGCTCAGAGGGAGC  
CATCACATCCACACCCTGTCACCCTTCGTGGCCAGTGCCAGACAGTAGCTAGT  
TGGATGCTAAAAGTAGAATTTAGATATCTTAACAATAAGCCCAGCAGTCTTTC  
AACTTCATTCGTAAATCATTTTTGTTTTGAGCATCTGTCACGTGGCAGCACTT  
GCCTGGATACTGGAGAGCTGAGAAGGAATGCGACAGGCAAGTCCTACTCTCAC  
AGTGTATACATTCAGGAGGAACAAGACACACAGTGCCAAGTAAATAAAGTAGC  
TGAACCTCATCAAATGATTTTATTCTTAAAGTCATTAAAGCATGTAATGTTCC  
CCTTTTTTTGTTTCAGGGGTGTACAGATTGAAGAAGTGATAGGTGTTTATGTGG  
TTTTAGTGACAAACCCCATGTGCTTTCATTGATTTTATGTTTTATGTTAAAAC  
ATCAACCGCAAGGTAAAATGCATATTGTATGTTGTTGGATACGTACTTAACTG

GTATGCATCCCCATCCATCTTCTTGGGTTAGTGTATGTAATCTTCTAATCTCTGTGTA  
 GAAATGTTGTATGTGTTAATATATTTAATAGATGTAACCTTAATAAACTGGCAT  
 TGAAGACTGAA (SEQ ID NO: 133) >NP\_005109.2 tumor necrosis factor  
 ligand superfamily member 15 isoform VEGI-251 precursor [*Homo sapiens*]  
 MAEDLGLSFGETASVEMLPEHGSCRPKARSSSARWALTCCLVLLPFLAGLTTY  
 LLVSQLRAQGEACVQFQALKGQEFAPSHQQVYAPLRADGDKPRAHLTVVRQTP  
 TQHFKNQFPALHWEHELGLAFTKNRMNYTNKFLIPESGDYFIYSQVTFRGMT  
 SECSEIRQAGRPNKPDSITVVITKVTDSTYPEPTQLLMGTKSVCEVGSNWFQPI  
 YLGAMFSLQEGDKLMVNVSDISLVDYTKEDKTFFGAFLL (SEQ ID NO: 134) Mouse  
 TL1 >NM\_177371.4 *Mus musculus* tumor necrosis factor (ligand) superfamily,  
 member 15 (Tnfsf15), mRNA  
 ATCAGAAGTCTCTCCAAGACAGCAGAAGGATGGCAGAGGAGCTGGGGTTGGGC  
 TTCGGAGAAGGAGTCCCAGTGGAAGTGCTGCCGGAAGGCTGTAGACACAGGCC  
 AGAGGCCAGGGCCGGGCTAGCTGCCAGGAGCAAAGCCTGCCTGGCTCTCACCT  
 GCTGCCTGTTGTCATTTCCCATCCTCGCAGGACTTAGCACCCCTCCTAATGGCT  
 GGCCAGCTCCGGGTCCCCGGAAAAGACTGTATGCTTCGGGGCCATAACAGAAGA  
 GAGATCTGAGCCTTCACCACAGCAAGTTTACTCACCTCCCAGAGGCAAGCCGA  
 GAGCACACCTGACAATTAAGAAACAAACCCCAGCACCATCTGAAAAATCAG  
 CTCTCTGCTCTACACTGGGAACATGACCTAGGGATGGCCTTCACCAAGAACGG  
 GATGAAGTACATCAACAAATCCCTGGTGATCCCAGAGTCAGGAGACTATTTCA  
 TCTACTCCCAGATCACATTCCGAGGGACCACATCTGTGTGTGGTGACATCAGT  
 CGGGGGAGACGACCAAACAAGCCAGACTCCATCACCATGGTTATCACCAAGGT  
 AGCAGACAGCTACCCTGAGCCTGCCCGCCTACTAACAGGGTCCAAGTCTGTGT  
 GTGAAATAAGCAACAACCTGGTTCCAGTCCCTCTACCTTGGGGCCACGTTCTCC  
 TTGGAAGAAGGAGACAGACTAATGGTAAACGTCAGTGACATCTCCTTGGTGGA  
 TTACACAAAAGAAGATAAACTTTCTTTGGAGCTTTCTTGCTATAAGGAGGAG  
 AAAACCATCATTCCAAGGGGGCTCCCCTGCCTCCTACTTTCCAATTTCTTTTC  
 TCATATGGATCTATAAACAGGGGGCTTTAGAGGGATCAGGGAAGGGGACAGTGG  
 TTTAGCTATATAATTTAGGAACCCCAATATTGATCCGTATATGCCTTATGGACT  
 AAAATAGTAAATGGAAAACCCAGTACAGCTCATGTTTGATAGAGACCTGCTGG  
 GTTTTAAAAATTGAAACACGCCTCATCCAATGGCACAATCTACTGATTTTCAGG  
 ACAGAACCTTTCCACAGTGCCCTCTGTCCAAGTCCTTTCTGAATTCAGCAGTT  
 CAGTTAGAGCTGAATTCGACAATGAACTTACTCCAGATCAAGAGCTAAAGACA  
 GAATCCAAAGAAAGACTGAGAAAATGATGTTATTTCTCCAAGAGGCAATGCAT  
 TTCCACATTCTTTTGTGCCTAACCTAAAAAATAAGAAAGAAGAAAGGAAGGAA  
 GGAAGGAAGGAAGGAAGGAAGGAAGGAAGGAAGGAAGGAAGGAAGGAAGGAAG  
 GAAGGAAGGGACAAGAAAAGACAAGACAAGACAAGAAAAAAGAAAAAATGGTA  
 TTTCTCGTGAATATTCCCTAAAAGGAATTGGTTTTCTGCTGTGAAGGAGAAAC  
 CTCACCTTTCTTCTGATTGCATCCTTTAGTATCCAAACATACAAGTGGGAATT  
 CCAAATGCACATGGAACATAGAACACTTTTATTATTGTGAGAACATGTTTATT  
 GAGTACCTACTATGCTCTGGGCACTCAGCCCACAGGACCATGAAGAGAAAGTC  
 AAATTTTCTTAAAACTAAATGAATCCTCAATACATACTTCCTGATCAACTAC  
 CACTCAAAATGTATAACTTCCAAAGTATAACTTCAAGTCAGCCATCTAGGTGG  
 TTTCTTGGGTAAAGGTGCTTGTCATTAAGCCTGACACCTGGGTTTGACCTCCC  
 AGAACCCAAAAGCTGGAAGGAGAGAATTGGTTCCCAAAATTATCCTCAAACC  
 CCCATACAAATGATGTGGCATGCACACATGTAATAAATAAATAAGTGTA  
 CAAAAACAAAAACAAAATTTTAAAGAAAAATTTCAAGTCCTGAAAGACAGCAT  
 TCCTGAGAATGTTGTCTCCATCGTTGTCCAGTATAGGCTAACCAGCTGATAGA  
 GACACTGAAGGAATTTAAAGACAGACATCAAGTGAAATGGAGCACTGTAGAAA  
 CACTTGATTCATGCCAGGAGTCAATGTACTATGAAGACCAACAACAAAGTGTC

AGTCATCAAAATCCAGAGAGAGGTGTTTATCTAGATCTGCTTTTCAAGTTTGGTTTGCAC  
GCCTTTATATAGTCTCTATTACAAATGCTCGTGTTCATGGTAGATGCCACAAGG  
AGTCAGAGGGTAAACTTAGCCCCAAACCACTGCTGAGCCATCTTCTAGGAAAC  
CTTCGAAGCAGAGCTGGGCAGCGTGACTCCCACACAATGACTGGGAAAGTAGT  
AGCTGATCAAAATTTGTTGAGTAATAATTTGTTAGAAAATTCATCTCCACTGC  
CTACTAAACCTAAGTTGTATACTATCTAGCTTCTGCTAAGCCAACCTTACATTG  
GCCACTTTTTCTGTCTTCAACTTCTTGAAGTATCACAGGTCTCAGTGAGAACA  
CAGGGAAAGGTGAGGTCGCCTTCCCCTGGTTCTTCATAGGGGAAACCACACCT  
GAAAGAAGATGAGCAGCCTGAGGTGACCTGGAGGAAGGGCTGTCTCAGAAGAA  
GGACTTATTTTTTTGGCTTAGGTCTAAAACCTTGAGAGTAATGCTCACTGGTCA  
ATTGAGGATGCTTTATCAATGACTCCAGTCTGACTCCAAGGTCAGAAAGGAGA  
GTGAGATGCTCTCTCTGCCTGCATATATCTTCATGGAACATGAGAATATTGAG  
CAACATAGACTTATAGGAAAACACTTGCCCCAAAAGTAGCCAGAGTGACCTGGT  
CATCCCCTCTACTAAACCCAAGCTTTGTGTCAAGGGCCTTCAAAGCTGCCCAG  
AAGTGATCTGGATGGCTTGGGAATTTATCCAAGACAGGAATTCCTGACAGCC  
AAAGATGCTTGAGTCCTTGTGCCTGACATGCATTTATTTTGCCCCTGTTTATT  
GAAGACTGTAACCTGTTGATTTGTGGGTATACATACATACATACATACAT  
ACATACATACATACATATGCTGTCATGAAGGCAGCATCAAACATTACTAATTG  
GACTCAAACCAGCATTCTGTTTCCAAGATACTAAGTATTCCCATGCAAACAG  
GAGCATGCTATTTTTTCTAAAGCAAAATGAAAAAAATAGTTTTTGAAAGTATATA  
TATGATGGAGTCAAGTGTAATGGCATACATCTGTAAACCCAGCACATGGGATG  
CTGAGCCAGGAGGATTGCCGTGAGTTTGAGGAGAACAGGGGGCTAAATAGTAAT  
TTTCAGGAAAGCCTTGCCTATATAACAAGACCTTGTCTCAAATGAAAAAAA  
AAAAAAATAGACCCCAGGCTGGTCCTTGGAGATAAGGTAATATATTCATTGGG  
TGAGGGGGGTGTGTGTTTTTGAAAATAGTTAATTTAGTGAGAAATGCTTTTTCGG  
TCAAATGCATCTCAAAGGCTGCTGAATTCAAATCGGGTCTGTAAATGCTTACC  
TAGTGCTTGCTTGCCCTGGGGACAGAGACATAAATTACTTTAGTCTCAGATCC  
ACTCGTTCTAACAGATTGGCATCTCCATCGTCTGTGGAGCTTTTAATCACTCT  
GTTTGTATTAGCTAATTAATTAGCTAACTTGAGACACACTGATATTTTCTTAT  
TATAAACATGGGTGCCATTTGATAAAAGACAATCATTAAACAAAATGGTTTCGAA  
TTTCCGCTTAAGTGATCTTCTTTTTTTCCTTTTTCATTTTTTTTAACTAGCTAAT  
CAAAGGTAGTTTCCCAAAAATAAATGCAAAGGGAGTATAAAGAAAAAATTCCC  
TGTGGTGGGAGCTAGTATTGAAACAACAGTATCAAAGAGGCTGTTACCTACTG  
GCCTCAAATTTTGGCAGGAACGCCTTTGAAAATGTTAGAACTTTACGGACAGC  
CTAGAGGTGCTTTGAAAAGTCTCTGTTGCCAACAAAAGCCATTAATCAGCATG  
CGGCACAGGTTACTCAAATTTTGACCTTGACTGTTTTTTAGATCTGTTACACA  
GAACACAACCTTCTGGGCTGTAATCTCTGATGTGGATTTGGTGATTTACTAAGG  
TACCGTGGGAAACAAGGAAAGTGTAAGTGTACTTGTACCACATCGTTTCTCAGTGCATG  
TCAGAGTCTACTCAACAGCAGGGCATGCCAGAGCCTTGGATACATTCCGGGAC  
AACTATGTCACTCCTAAGGAAATTCCAAGTGTGTGCCTGTCAAGCACTCTGG  
ATCATAGAAGCCCACGAGTTCAGTGTGCACAAGGCACAGCCATGGCCAGCACT  
CTCTTGCAATGGTATTTCTCTTAAAGCTCTTACTCAATCACGGTCCCATGATTGT  
GACATTGGGGATTAATTGCTTGAGCAGGTTTATTTACAGTCTGTTCCCTTGCAA  
AATACATGCAGATATGTCTGGCCTCAAATCCCCTGATTGTTTTAGGGCTTAG  
AGAATACTGGGGATGTTTTTGTCTGTTTTTCAGATGTACTTTATTTAAGCTTGCA  
GAATTACCCTGAATATTAACAGTGTTCTAAGATATTGCCTGCTAGCTTCTGGC  
TAATTTACTAGTGGTGACAGTATCAGATCAGAGTATCTATATTTATGTCTTGC  
TATTATAGTTAAACTTCCTGATCTCTGTAAACACACTCACCCCTACCTCATCT  
ATCTACCCATCTTGTGGATGTAGCTGTGAGAAGACTCACAAAGCCCGAGTTGCA  
GTTACTTTTTCTGAAGCAACATAGTATGTTAATGGAATGGCCAGAAGCTCTACTC

TTGGCATTGACCTGAATTTGACCTAAAGAAATTTGAAGGCAGAAA  
TATTTTTTACTATGCATGGGACAACGTAGAAAGAGCAAGGAGACTGCTTACACA  
TGGTGGTCACATCTCTGGCTTCATCCCTAAACCAATTTTCTGACCCCAAGTCG  
ATTTTTTTTCATGTAGTTATTGTTTCATTTTCTGGAAAGAGTCAAGCAAAAAGA  
GAGTTTTATAGAAACCATTGCATCATGGAGGTCAGGGGAGGGATTAAGCCAAA  
GAATTCCTTCTCCAAATCTATAGCCATATGGCCACCCTTTGGTGTACTTCTAT  
TTGATCATGACAAACCTGAGAGCCCTGCCCAGAGTTCAGTGGATCCTAATGAA  
CTCCAAGAGTAATTCATTCCCTCACCAACTCTAGGGGGCTTGGCCAGTGCAGAA  
AATGTCATGGGATTTTAAAGTTAACATGAGCTGCTATCCAAACTTATGTCTCT  
TTAAGAATGGAGAGACACAGGCCAGGAGAGGTAACATATGAAGCCTGGTATTG  
GGCAGTAGCTTGATGGAGTATTGAGGCTAAAAGTAGACTTCCTGCCCCTGACC  
ATACACAACACCCTTTCAGTTTGATCCATGGTGGTCTTATTCTACTTTATTTT  
GAGCACCTGTCACACCTAGTTACTGTCATGCCAAGAAGGTCCATAACAGGCAA  
ATCCTACTCTGCTGTGTGCACACAAGAGGAAGGAGGCTCACAGTAGCAAGTAA  
ACAGATAAGCAAACGTACACGATTTTCGTCTTAAAGTCATTAAGACACACGCG  
TACCCCTCTTTTGTTCAGAGGGTATACAGGCTGAACAGATGTCAGTGTTTAC  
CTATTCTTATTGATAAGCCCCATGTGCTTTCATTGGTTGAATGTTTTATGTTA  
AAACGTCATATTGCCATCGTAAAATGCATATTGTATGTTGTTGGGTATATAAT  
TAACTAATATGCATCGCATGTATGAATTCTAATCTCTGTAAATGAAAACCTTAT  
ATATGTTAACATATGTAATAGTTATAATTTAATAAACTGACACTGGAGACTAC (SEQ ID  
NO: 135) >NP\_796345.4 tumor necrosis factor ligand superfamily member 15  
[*Mus musculus*]

MAEELGLGFGEVPEVLPEGCRHRPEARAGLAARSKACLALTCCLLSFPILA  
GLSTLLMAGQLRVPKGKDCMLRAITEERSESPQQVYSPPRGKPRAHLTIKKQT  
PAPHLKNQLSALHWEHDLGMAFTKNGMKYINKSLVIPESGDYFIYSQITFRGT  
TSVCGDISRGRPRPNKPDSITMVITKVADSYPEPARLLTGSKSVCEISNNWFQS  
LYLGATFSLEEGDRLMVNVSDISLVDYTKEDKTFFGAFLL (SEQ ID NO: 136)  
Human CD80 >NM\_005191.4 *Homo sapiens* CD80 molecule (CD80), mRNA  
AAACCCTCTGTAAAGTAACAGAAGTTAGAAGGGGAAATGTCGCCTCTCTGAAG  
ATTACCCAAAGAAAAAGTGATTTGTCATTGCTTTATAGACTGTAAGAAGAGAA  
CATCTCAGAAGTGGAGTCTTACCCTGAAATCAAAGGATTTAAAGAAAAAGTGG  
AATTTTTCTTCAGCAAGCTGTGAAACTAAATCCACAACCTTTGGAGACCCAGG  
AACACCCTCCAATCTCTGTGTGTTTTGTAAACATCACTGGAGGGTCTTCTACG  
TGAGCAATTGGATTGTCATCAGCCCTGCCTGTTTTGCACCTGGGAAGTGCCCT  
GGTCTTACTTGGGTCCAAATTGTTGGCTTTCACTTTTGACCCTAAGCATCTGA  
AGCCATGGGCCACACACGGAGGCAGGGAACATCACCATCCAAGTGTCCATACC  
TCAATTTCTTTCAGCTCTTGGTGCTGGCTGGTCTTTCTCACTTCTGTTCAGGT  
GTTATCCACGTGACCAAGGAAGTGAAAGAAGTGGCAACGCTGTCCTGTGGTCA  
CAATGTTTCTGTTGAAGAGCTGGCACAACTCGCATCTACTGGCAAAAGGAGA  
AGAAAATGGTGCTGACTATGATGTCTGGGGACATGAATATATGGCCCGAGTAC  
AAGAACCGGACCATCTTTGATATCACTAATAACCTCTCCATTGTGATCCTGGC  
TCTGCGCCCATCTGACGAGGGCACATACGAGTGTGTTGTTCTGAAGTATGAAA  
AAGACGCTTTCAAGCGGGAACACCTGGCTGAAGTGACGTTATCAGTCAAAGCT  
GACTTCCCTACACCTAGTATATCTGACTTTGAAATTCCAACCTTCTAATATTAG  
AAGGATAATTTGCTCAACCTCTGGAGGTTTTCCAGAGCCTCACCTCTCCTGGT  
TGGAATAATGGAGAAGAATTAATGCCATCAACACAACAGTTTCCCAAGATCCT  
GAAACTGAGCTCTATGCTGTTAGCAGCAAACCTGGATTTCAATATGACAACCAA  
CCACAGCTTCATGTGTCTCATCAAGTATGGACATTTAAGAGTGAATCAGACCT  
TCAACTGGAATACAACCAAGCAAGAGCATTTTCCTGATAACCTGCTCCCATCC  
TGGGCCATTACCTTAATCTCAGTAAATGGAATTTTTGTGATATGCTGCCTGAC



AAGTTAAAGCTTGCTTAAAGCTTGAAGCTTCCATCAAAAGCTGACTTCTCTACC  
CCCAACATAACTGAGTCTGGAAACCCATCTGCAGACACTAAAAGGATTACCTG  
CTTTGCTTCCGGGGGTTTCCCAAAGCCTCGCTTCTCTTGGTTGGAAAATGGAA  
GAGAATTACCTGGCATCAATACGACAATTTCCCAGGATCCTGAATCTGAATTG  
TACACCATTAGTAGCCAACTAGATTTCAATACGACTCGCAACCACACCATTAA  
GTGTCTCATTAATATGGAGATGCTCACGTGTCAGAGGACTTCACCTGGGAAA  
AACCCCCAGAAGACCCTCCTGATAGCAAGAACACACTTGTGCTCTTTGGGGCA  
GGATTCCGGCGCAGTAATAACAGTCGTCGTCATCGTTGTCATCATCAAATGCTT  
CTGTAAGCACAGAAGCTGTTTCAGAAGAAATGAGGCAAGCAGAGAAACAAACA  
ACAGCCTTACCTTTCGGGCCTGAAGAAGCATTAGCTGAACAGACCGTCTTTCCTT  
TAGTTCTTCTCTGTCCATGTGGGATACATGGTATTATGTGGCTCATGAGGTAC  
AATCTTTCTTTCAGCACCGTGCTAGCTGATCTTTCGGACAACCTTGACACAAGA  
TAGAGTTAACTGGGAAGAGAAAGCCTTGAATGAGGATTTCTTTCCATCAGGAA  
GCCTACGGGCAAGTTTGCTGGGCCTTTGATTGCTTGATGACTGAAGTGGAAAG  
GCTGAGCCCCTGTGGGTGGTGCTAGCCCTGGGCAGGGGCAGGTGACCCTGGG  
TGGTATAAGAAAAAGAGCTGTCACTAAAAGGAGAGGTGCCTAGTCTTACTGCA  
ACTTGATATGTCATGTTTGGTTGGTGTCTGTGGGAGGCCTGCCCTTTTCTGAA  
GAGAAGTGGTGGGAGAGTGGATGGGGTGGGGGCAGAGGAAAAGTGGGGGAGAG  
GGCCTGGGAGGAGAGGAGGGAGGGGGACGGGGTGGGGGTGGGGAAAACCTATGG  
TTGGGATGTAAAAACGATAATAATATAAATATTAAATAAAAAGAGAGTATTGA GCAA  
(SEQ ID NO: 139) >NP\_033985.3 T-lymphocyte activation antigen CD80  
precursor [*Mus musculus*]

MACNCQLMQDTPLLKFPCPRLILLFVLLIRLSQVSSDVDEQLSKSVKDKVLLP  
CRYNPHEDESEDRIYWQKHDKVVL SVIAGKLKVWPEYKNRTLYDNTTYSLII  
LGLVLSDRGTYSVVQKKERGTYEVKHLALVKLSIKADFSTPNITESGNPSAD  
TKRITCFASGGFPKPRFSWLENGRELPGINTTISQDPESELYTISSQLDFNTT  
RNHTIKCLIKYGDAHVSEDFWEKPPEDPPDSKNTLVLFAGFGAVITVVVIV  
VIKCFCKHRSCFRNEASRETNNSLTFGP EEALAEQTVFL (SEQ ID NO: 140)  
Human CD86 >NM\_175862.5 *Homo sapiens* CD86 molecule (CD86), transcript  
variant 1, mRNA

AGTCATTGCCGAGGAAGGCTTGCACAGGGTGAAAGCTTTGCTTCTCTGCTGCT  
GTAACAGGGACTAGCACAGACACACGGATGAGTGGGGTCAATTCAGATATTA  
GGTCACAGCAGAAGCAGCCAAAATGGATCCCCAGTGC ACTATGGGACTGAGTA  
ACATTCTCTTTGTGATGGCCTTOCTGOTCTCTGGTGCTGCTCCTCTGAAGATT  
CAAGCTTATTTCAATGAGACTGCAGACCTGCCATGCCAATTTGCAA ACTCTCA  
AAACCAAAGCCTGAGTGAGCTAGTAGTATTTTGGCAGGACCAGGAAA ACTTG  
TTCTGAATGAGGTATACTTAGGCAAAGAGAAATTTGACAGTGTTCA TTCCAAG  
TATATGGGCCGCACAAGTTTGTATTCCGACAGTTGGACCCTGAGACTTCACAA  
TCTTCAGATCAAGGACAAGGGCTTGTATCAATGTATCATCCATCACAAAAGC  
CCACAGGAATGATTCGCATCCACCAGATGAATTCTGAACTGTCAGTGCTTGCT  
AACTTCAGTCAACCTGAAATAGTACCAATTTCTAATATAACAGAAAATGTGTA  
CATAAATTTGACCTGCTCATCTATACACGGTTACCCAGAACCTAAGAAGATGA  
GTGTTTTGCTAAGAACCAAGAATTCAACTATCGAGTATGATGGTGTTATGCAG  
AAATCTCAAGATAATGTCACAGAACTGTACGACGTTTCCATCAGCTTGTCTGT  
TTCATTCCCTGATGTTACGAGCAATATGACCATCTTCTGTATTCTGGAACTG  
ACAAGACGCGGCTTTTATCTTCACCTTTCTCTATAGAGCTTGAGGACCCTCAG  
CCTCCCCCAGACCACATTCCTTG GATTACAGCTGTACTTCCAACAGTTATTAT  
ATGTGTGATGGTTTTCTGTCTAATTCTATGGAAATGGAAGAAGAAGAGCGGC  
CTCGCAACTCTTATAAATGTGGAACCAACACAATGGAGAGGGAAGAGAGTGAA  
CAGACCAAGAAAAGAGAAAAAATCCATATACCTGAAAGATCTGATGAAGCCCA

GCGTGTGTTTTTGAAGTTCATGCGACACAAAAGTGATACATGTT  
TTTAATTAAAGAGTAAAGCCCCATACAAGTATTCATTTTTTCTACCCTTTCCTT  
TGTAAGTTCCTGGGCAACCTTTTTTGATTTCTTCCAGAAGGCCAAAAGACATTA  
CCATGAGTAATAAGGGGGCTCCAGGACTCCCTCTAAGTGGAATAGCCTCCCTG  
TAACTCCAGCTCTGCTCCGTATGCCAAGAGGAGACTTTAATTCTCTTACTGCT  
TCTTTTCACTTCAGAGCACACTTATGGGCCAAGCCCAGCTTAATGGCTCATGA  
CCTGGAAATAAAATTTAGGACCAATACCTCCTCCAGATCAGATTCTTCTCTTA  
ATTTCATAGATTGTGTTTTTTTTTTTAAATAGACCTCTCAATTTCTGGAAAAC  
GCCTTTTATCTGCCCAGAATTCTAAGCTGGTGCCCCACTGAATTTTGTGTGTA  
CCTGTGACTAAACAACCTACCTCCTCAGTCTGGGTGGGACTTATGTATTTATGA  
CCTTATAGTGTTAATATCTTGAAACATAGAGATCTATGTACTGTAATAGTGTG  
ATTACTATGCTCTAGAGAAAAGTCTACCCCTGCTAAGGAGTTCTCATCCCTCT  
GTCAGGGTCAGTAAGGAAAACGGTGGCCTAGGGTACAGGCAACAATGAGCAGA  
CCAACCTAAATTTGGGGAAATTAGGAGAGGCAGAGATAGAACCTGGAGCCACT  
TCTATCTGGGCTGTTGCTAATATTGAGGAGGCTTGCCCCACCCAACAAGCCAT  
AGTGAGAGAACTGAATAAACAGGAAAATGCCAGAGCTTGTGAACCCTGTTTC  
TCTTGAAGAACTGACTAGTGAGATGGCCTGGGGAAGCTGTGAAAGAACCAAAA  
GAGATCACAATACTCAAAAGAGAGAGAGAGAGAGAAAAAAGAGAGATCTTGATCC  
ACAGAAATACATGAAATGTCTGGTCTGTCCACCCCATCAACAAGTCTTGAAAC  
AAGCAACAGATGGATAGTCTGTCCAAATGGACATAAGACAGACAGCAGTTTCC  
CTGGTGGTCAGGGAGGGGTTTTGGTGATACCCAAGTTATTGGGATGTCATCTT  
CCTGGAAGCAGAGCTGGGGAGGGAGAGCCATCACCTTGATAATGGGATGAATG  
GAAGGAGGCTTAGGACTTTCCACTCCTGGCTGAGAGAGGAAGAGCTGCAACGG  
AATTAGGAAGACCAAGACACAGATCACCCGGGGCTTACTTAGCCTACAGATGT  
CCTACGGGAACGTGGGCTGGCCCAGCATAGGGCTAGCAAATTTGAGTTGGATG  
ATTGTTTTTGTCTCAAGGCAACCAGAGGAACTTGCATACAGAGACAGATATAC  
TGGGAGAAATGACTTTGAAAACCTGGCTCTAAGGTGGGATCACTAAGGGATGG  
GGCAGTCTCTGCCCAAACATAAAGAGAACTCTGGGGAGCCTGAGCCACAAAAA  
TGTTCCTTTATTTTATGTAAACCCTCAAGGGTTATAGACTGCCATGCTAGACA  
AGCTTGTCATGTAATATTTCCATGTTTTTACCCTGCCCTGCCTTGATTAGA  
CTCCTAGCACCTGGCTAGTTTCTAACATGTTTTGTGCAGCACAGTTTTTAATA  
AATGCTTGTTACATTCA (SEQ ID NO: 141) >NP\_787058.5 T-lymphocyte

activation antigen CD86 isoform 1 precursor [*Homo sapiens*]  
MDPQCTMGLSNILFVMAFLLSGAAPLKIQAYFNETADLPCQFANSQNQSLSEL  
VFWQDQENLVLNEVYLGKEKFDVSHSKYMGRTSFDSDSWTLRLHNLQIKDKG  
LYQCIIHHKKPTGMIRIHQMNSELSVLANSFSQPEIVPISNITENVYINLTCSS  
IHGYPEPKKMSVLLRKTSTIEYDGVMQKSQDNVTELYDVSISLSVSFPDVTS  
NMTIFCILETDKTRLLSPFSIELEDPPPPDHIPWITAVLPTVIICVMVFCL  
ILWKWKKKKRPRNSYKCGTNTMERESEQTKKREKIHIPERSDEAQRVFKSSK  
TSSCDKSDTCF (SEQ ID NO: 142) Mouse CD86 >NM\_019388.3 *Mus musculus*  
CD86 antigen (Cd86), mRNA

ATTGCTGAGGAAGAAAGAGGAGCAAGCAGACGCGTAAGAGTGGCTCCTGTAGG  
CAGCACGGAAGTGAACAACCAGACTCCTGTAGACGTGTTCCAGAACTTACGGA  
AGCACCCACGATGGACCCAGATGCACCATGGGCTTGGCAATCCTTATCTTTG  
TGACAGTCTTGCTGATCTCAGATGCTGTTTCCGTGGAGACGCAAGCTTATTTT  
AATGGGACTGCATATCTGCCGTGCCCATTTACAAAGGCTCAAAACATAAGCCT  
GAGTGAGCTGGTAGTATTTTGGCAGGACCAGCAAAAAGTTGGTTCTGTACGAGC  
ACTATTTGGGCACAGAGAACTTGATAGTGTGAATGCCAAGTACCTGGGCCGC  
ACGAGCTTTGACAGGAACAACCTGGACTCTACGACTTCACAATGTTTCAGATCAA  
GGACATGGGCTCGTATGATTGTTTTATACAAAAAAGCCACCCACAGGATCAA



TTATCCTCCAACAGACATTAACAGAACTGTTCAGTGATCGCCAACTTCAGTGAA  
CCTGAAATAAAACTGGCTCAGAATGTAACAGGAAATTCTGGCATAAATTTGAC  
CTGCACGTCTAAGCAAGGTCACCCGAAACCTAAGAAGATGTATTTTCTGATAA  
CTAATTCAACTAATGAGTATGGTGATAACATGCAGATATCACAAGATAATGTC  
ACAGAACTGTTTCACTATCTCCAACAGCCTCTCTCTTTTCATTCCCGGATGGTGT  
GTGGCATATGACCGTTGTGTGTGTCTGGAACGGAGTCAATGAAGATTTCCCT  
CCAAACCTCTCAATTTCACTCAAGAGTTTCCATCTCCTCAAACGTATTGGAAG  
GAGATTACAGCTTCAGTTACTGTGGCCCTCCTCCTTGTGATGCTGCTCATCAT  
TGTATGTCACAAGAAGCCGAATCAGCCTAGCAGGCCAGCAACACAGCCTCTA  
AGTTAGAGCGGGATAGTAACGCTGACAGAGAGACTATCAACCTGAAGGAACTT  
GAACCCCAAATTGCTTCAGCAAAACCAAATGCAGAGTGAAGGCAGTGAGAGCC  
TGAGGAAAGAGTTAAAAATTGCTTTGCCTGAAATAAGAAGTGCAGAGTTTCTC  
AGAATTCAAAAATGTTCTCAGCTGATTGGAATTCTACAGTTGAATAATTAAAG  
AACAAAATACACAACAGTGTCCATATTTTATCCTGTTTCCTTTCCAAGTTTTT  
GGGCAATGTCAATTGTGTCCCCTATGCCAGGAGCAGACATCTATTTTGTCTTG  
CTTTGTTTAACTCAGTGCACACTCATAGGCCAAGAGCACTGAAATGGCTTCTT  
TCCCAGGAATAACATTTTGGATCAATCTCTCCTACTTGAGATCAGATTCTTCT  
TCTAATTTTGCATAGTGTGTTTTTATATGGAACCTCCTTGTTGTAGGAATACTG  
GCTTTTATCTGTCTTGCACACTTGCATACTTATATACTTATACCTGGACAGCT  
ACCTCTTCAGTCAGGATGGGAGTGGTATATTTGGTGATGTTATTTGATGTGTT  
CGTGTTGCTATCTTAAAACAGCAAAGAGCATATACTATAGTAGCTCAACTACA  
ATGATCTAGAGAAAGACCCAGCACTTATAAGAAACACTGTCCCTCCATCAGGG  
TCAATAATGAATACAATGACCTAAGTAATATACAGGTGACAGCAACAGCACAG  
AGTTCTCAGTGCTGGCAAATCAAGAAACACAAATATGGAACCATCTCTAGATC  
CAAGAGCCACTCCTACCTGGGCTGCCACAGATACTGGAAGAATCCACCTGCCT  
GGCCAGCAAGTCACAACCTTAGCAGGCAGCACTGAAGAAAGCAAGATGTACTGT  
ATGCCCTTTTAAGAAAATGCCTGGAAAGGTCTGGAGAATGCTGTGCAAGGATA  
AGACAGCCAAGCACTCAAAACCAGGAGACATCACTAGAATCCAACCAACAAAT  
GTTTATGGAAGGACTGATCTGCCCAGTCCATTGAAAAGTCAAGAGGTCAGAGA  
TAGACCAGTGTGTGTCTCAATGGATGTAGATATCAGCCACCTCGGTGCTCAAC  
AGGTATTTTATGATCTCCTTGTTTCAAATTCATCTAGATGTAGAACTAGGGAG  
AGAGCAGTCACATTGATGAAAGGCTAGGACTCTTTCAGCTCATGGCTTGTGTG  
GAAGGAGGGAAAGCAGAAATCACAACACTCTGAGACTACTGTAGTCTGCAGAT  
ACCTGAGTGGGTGTGGCTTGGCCTTTCAAAGGACAAAGAGCAACTAATGCTGA  
AAGCACATAGTGTATCTATACGGCATGGAATAGTCATCACCCAGACTTAAAGA  
GAACTTTGGCAGGTCTGAGCAGCAAAATATTGTTGTTTCCATTTTACATAAAG  
GGCCCTGGAGGGCTATAGACTATTCCGCTGGCAGGGCTCATGCTTGTAATGTG  
TCCATCTTGATTCACCCTGTGCAGACTCTTAAGATCTGGCCAGTTACCAACAT  
GTTCTGTACAGAGTGGATTTCAATAAAGTTTTCTTGAATTTTTTCAAG (SEQ ID NO:  
143) >NP\_062261.3 T-lymphocyte activation antigen 0D86 precursor [*Mus musculus*]  
MDPRCTMGLAILIFVTVLLISDAVSVETQAYFNGTAYLPCPFTKAQNISLSEL  
VFWQDQQLVLYEHLGTEKLDVNAKYLGRTSFDNRNNWTLRLHNVQIKDMG  
SYDCFIQKKPPTGSILQQTLTELSVIANFSEPEIKLAQNVTGNSGINLTCTS  
KQGHPPKPKMYFLITNSTNEYGDNMQISQDNVTELFISISNSLSLSPDGVWHM  
TVVVCVLETESMKISSKPLNFTQEFSPQTYWKEITASVTVALLLVMLLIIVCH  
KKPNQPSRPSNTASKLERDSNADRETINLKELEPQIASAKPNAE (SEQ ID NO: 144)  
Human LFA-3 >NM\_001779.3 *Homo sapiens* CD58 molecule (CD58), (CD58)  
transcript variant 1, mRNA  
GAACTTAGGGCTGCTTGTGGCTGGGCACTCGCGCAGAGGCCGGCCCGACGAGC  
CATGGTTGCTGGGAGCGACGCGGGGCGGGCCCTGGGGGTCTCAGCGTGGTCT

GGCTGCTGCTGCTGCTGCTTTTCTTCCCAACATAATATGGT  
GTTGTGTATGGGAATGTAACCTTTCCATGTACCAAGCAATGTGCCTTTAAAAGA  
GGTCCTATGGAAAAACAAAAGGATAAAGTTGCAGAACTGGAAAATTCTGAAT  
TCAGAGCTTTCTCATCTTTTAAAAATAGGGTTTATTTAGACACTGTGTCAGGT  
AGCCTCACTATCTACAACCTTAACATCATCAGATGAAGATGAGTATGAAATGGA  
ATCGCCAAATATTACTGATACCATGAAGTTCCTTTCTTTATGTGCTTGAGTCTC  
TTCCATCTCCCACACTAACTTGTGCATTGACTAATGGAAGCATTGAAGTCCAA  
TGCATGATACCAGAGCATTACAACAGCCATCGAGGACTTATAATGTACTCATG  
GGATTGTCCTATGGAGCAATGTAAACGTAACCTCAACCAGTATATATTTTAAGA  
TGGAAAATGATCTTCCACAAAAAATACAGTGTACTCTTAGCAATCCATTATTT  
AATACAACATCATCAATCATTTTGTACAACCTGTATCCCAAGCAGCGGTCATTC  
AAGACACAGATATGCACTTATACCCATACCATTAGCAGTAATTACAACATGTA  
TTGTGCTGTATATGAATGGTATTCTGAAATGTGACAGAAAACCAGACAGAACC  
AACTCCAATTGATTGGTAACAGAAGATGAAGACAACAGCATAACTAAATTATT  
TTAAAACTAAAAAGCCATCTGATTTCTCATTTGAGTATTACAATTTTGAAC  
AACTGTTGAAATGTAACCTGAAGCAGCTGCTTTAAGAAGAAATACCCACTAA  
CAAAGAACAAGCATTAGTTTTGGCTGTCATCAACTTATTATATGACTAGGTGC  
TTGCTTTTTTTTGTGTCAGTAAATTGTTTTTACTGATGATGTAGATACTTTTGTA  
ATAATGTAAATATGTACACAAGTGA (SEQ ID NO: 145) >NP\_001770.1  
lymphocyte function-associated antigen 3 isoform 1 [*Homo sapiens*]  
MVAGSDAGRALGVLSVVCLLHCFGFISCFSSQQIYGVVYGNVTFHVPSNVPLKE  
VLWKKQKDKVAELENSEFRAFSSFKNRVYLDTVSGSLTIYNLTSSDEDEYEME  
SPNITDTMKFFLYVLESLPSPTLTLCALTNGSIEVQCMIPHYNSHRGLIMYSW  
DCPMEQCKRNSTSIYFKMENDLPQKIQCTLSNPLFNTTSSIILTTICIPSSGHS  
RHRYALIPIPLAVITTCIVLYMNGILKCDRKPDRNTSN (SEQ ID NO: 146) Human  
SLAM >NM\_003037.5 *Homo sapiens* signaling lymphocytic (CD150) activation  
molecule family member 1 (SLAMF1), transcript variant 1, mRNA  
AGACAGCCTCTGCTGCATGACACGAAGCTTGCTTCTGCCTGGCATCTGTGAGC  
AGCTGCCAGGCTCCGGCCAGGATCCCTTCCTTCTCCTCATTGGCTGATGGATC  
CCAAGGGGCTCCTCTCCTTGACCTTCGTGCTGTTTCTCTCCCTGGCTTTTGGG  
GCAAGCTACGGAACAGGTGGGCGCATGATGAACTGCCCAAAGATTCTCCGGCA  
GTTGGGAAGCAAAGTGCTGCTGCCCCTGACATATGAAAGGATAAATAAGAGCA  
TGAACAAAAGCATCCACATTGTGCGTCACAATGGCAAATCACTGGAGAACAGT  
GTCGAGAACAAAATAGTGTCTCTTGATCCATCCGAAGCAGGCCCTCCACGTTA  
TCTAGGAGATCGCTACAAGTTTTATCTGGAGAATCTCACCTGGGGATACGGG  
AAAGCAGGAAGGAGGATGAGGGATGGTACCTTATGACCCTGGAGAAAAATGTT  
TCAGTTCAGCGCTTTTGCCTGCAGTTGAGGCTTTATGAGCAGGTCTCCACTCC  
AGAAATTAAAGTTTTAAACAAGACCCAGGAGAACGGGACCTGCACCTTGATAC  
TGGGCTGCACAGTGGAGAAGGGGGACCATGTGGCTTACAGCTGGAGTGAAAAG  
GCGGGCACCCACCCACTGAACCCAGCCAACAGCTCCCACCTCCTGTCCCTCAC  
CCTCGGCCCCCAGCATGCTGACAATATCTACATCTGCACCGTGAGCAACCCTA  
TCAGCAACAATTCCCAGACCTTCAGCCCGTGGCCCGGATGCAGGACAGACCCC  
TCAGAAACAAAACCATGGGCAGTGTATGCTGGGCTGTTAGGGGGTGTCATCAT  
GATTCTCATCATGGTGGTAATACTACAGTTGAGAAGAAGAGGTAAAACGAACC  
ATTACCAGACAACAGTGGAACAAAAAAGCCTTACGATCTATGCCCAAGTCCAG  
AAACCAGGTCCTCTTCAGAAGAACTTGACTCCTTCCCAGCTCAGGACCCTTG  
CACCACCATATATGTTGCTGCCACAGAGCCTGTCCCAGAGTCTGTCCAGGAAA  
CAAATTCCATCACAGTCTATGCTAGTGTGACACTTCCAGAGAGCTGACACCAG  
AGACCAACAAAGGGACTTTCTGAAGGAAAATGGAAAAACCAAAATGAACACTG  
AACTTGGCCACAGGCCCCAAGTTTCCTCTGGCAGACATGCTGCACGTCTGTAC

CCTTCTCATGATCAATCCCTGCTTCCATGCTTCCACATGCTGATGTTT  
AACAAAGGAAGTGAGGCTTCCCAAGAATTTAGCTTGCTGTGCAGTGGCTGCAGG  
CGCAGAACAGAGCGTTACTTGATAACAGCGTTCCATCTTTGTGTTGTAGCAGA  
TGAAATGGACAGTAATGTGAGTTCAGACTTTGGGCATCTTGCTCTTGGCTGGA  
ACTGGATAATAAAAATCAGACTGAAAGCCAGGACATCTGAGTACCTATCTCAC  
ACACTGGACCACCAGTCACAAAGTCTGGAAAAGTTTACATTTTGGCTATCTTT  
ACTTTGTTCTGAGGCTGATCATGATAACCTGCAGACCTGATCAAGCCTCTGT  
GCCTCAGTTTCTCTCTCAGGATAAAGAGTGAATAGAGGCTGAAGGGTGAATTT  
CTTATTATACATAAAACACTCTGATATTATTGTATAAAGGAAGCTAAGAATAT  
TATTTTATTTGCAAAACCCAGAAGCTAAAAAGTCAATAAACAGAAAGAATGAT  
TTTGAGATCTCTGAGTTTTGAACAGTGGACTGGAAACCATGTAAGAGCCTTAA  
AAGTACAGTTCTGTGCAAATGGCATTCAAGTTTTAAAGAAAAACGTAGCAAATG  
TTTGATGGTGCTGTTACAAAGGAGCTTGGAATACTCAGAGGAAGTTGTCCCAT  
GGTGATTTTTCACTTCTCAAATGATGTTTAAATCCCAGTTCTCTGTTGATTC  
CCTTGAACAACAAACCTGGAACCTCAGCTAAGACTCTCTGTGACCAGATTCTG  
AACCTCTTATATCCAGGGCTTCAAGGGGTATTGCAGGTCAAGGTOTTTCCCTAG  
GCACCTTCTACTCCCTGCATACCTCTCCTCACACTAAATTTATCCTCTAGTAG  
AAAATTAAGTTATTTTGGTCTAACAGCTTCAAATCTTTGAATGCTCAATAACT  
TATTTTGCAAGCTGCAGGCAGAAAGAGACTTTTTTAAGTAAAGTCCTTTGTTTT  
TTCCTATTCTCTGCTTTTAGACAGGCTGTCCTCAATTTAAGCCCTGCTTTTTTC  
TTATTGTTTCTTATATAAACTTGGTAAGTACTGTAAGAAACAGCCACTATCAT  
ACCATTGCATAATAAGGAGCACCAACTTCCCAGCTCAAAACTCAGGTCCCTTAT  
TGCCTTGATCTTACCTCCTCTATGAGGTCAATTCACATTGTAAGCCTGTTGC  
TTAGTGCATCTCGTTTTCTGGTACCAGCTTCTTTAATAGAGTTCTTAGTTGCA  
ATCAACAGAAGCTGGCTTTGGCTTTTTTATGTAGAAAAGGAACCTATTGAAAA  
GATACTGATTGGTTCCAATAACTGCTAGAAGTTTCTGCAAAACCATGCTTTGA  
AAGTGAGCAGGAAAAGAAGAGACTAGGCTGTGGCTGGGAGCACAGCCAAAATT  
ACAAAACCAGCCCAGGGATGATGATCCTGTTTCATGCACAGCCACTGTCCCCAG  
CACTAGGCACAGACTCTACCACTGCCTCACTGTCTCTGOTGGACTTGAAACT  
TGATATTACTGTTACTGCTGCACTGTCTGCCATGAAAATGAATTCTCCAGGGT  
CCCTTCTTCATCCTTTTCATCTCTAGCTTATAATTCAAAGTCTGGGATTGAGTG  
GCCAATCCTAGGTCACATGTCCATGTCCTATCTCCAAGGGGGGCTGGGAATTG  
AATATCTGGCATTTTCCACTTTCCTTCTTATGAATTAAGGAATTCTACAAAT  
AATAGAAGTGGGATTCAGGTGGTAGGCAGACAAAAAAGCCTCACAATTATCCA  
CTACGCCACCCTTGATAACCTTACCCTCATTCACTGTCTACTCTCAAAACTG  
TGGAGCTACTAATGAAGATTTGTAAACCCGGGCTTATGAGCACCCATTCTTTT  
ACTACAACTCAGATTGCTCTAGAAGCTCAGTTCCCAGCACTTGGATTTTTCCA  
GTAGCTGAATTCTACCTGAAGGAAGGGCAGAAACAAAGGGTGAAGAAGAGGCT  
ATCACTTCCAAGTATCCTGCACCCCTGGGCTCAAGACCTCACTGGGGAGGGAG  
TCTTTTGGGCCACCCACCAAACAGCACTGGCATTATGCCTCTCACCCCTAGACC  
ATGGTTACACGTGGTAAAACAACCCCTTCTGGTGATACATTCACAACTCTCTA  
GTTTCCCCCAAATGGCACTATGGGGAGCGGGAGCTTGCCTTTTCTCAGACTT  
AAAACAATAAGTTTTCCCCGTGTTTCCCCTCTAATGCTGTTTTCTTTTGACCA  
AGCATGTCTGAATTCTAGAGAAGTCAGGAGGAACACACCCATTCTCGGTTTGA  
AGGGACTGATGTTCTGAAGTACAACCTGGGCACAGTCCCAGGCTCTTCAGGACG  
CTTCCTCCATTCACACAGCGGGGATGTGATTGTTACAGCGGGTGGTGTGTGCT  
GGCTGAGAAGCCACTGTGAATTGATTCTTCTTCTGAAGTTTATGTTTCTACTT  
TTTGGAATGAATAAATTACAGCCAGTCCATCAAGGAAA (SEQ ID NO: 147)

>NP\_003028.1 signaling lymphocytic activation molecule isoform b precursor  
[Homo sapiens]

MPDKGLLSLTFVLFSLAFGASYGTGGRMMNCPKILRQLGSKVLLPLTYERIN  
KSMNKSIHIVVTMAKSLENSVENKIVSLDPSEAGPPRYLGDRYKFYLENLTLG  
IRESRKEDEGWYLMTLEKNVSVQRFCLQLRLYEQVSTPEIKVLNKTQENGCTCT  
LILGCTVEKGDHVAYSWSEKAGTHPLNPANSSHLLSLTLGPQHADNIYICTVS  
NPISNNSQTFSPWPGCRTDPSETKPWAVYAGLLGGVIMILIMVVILQLRRRGK  
TNHYQTTVEKKSLTIYAQVQKPGPLQKKLDSFPAQDPCTTIYVAATEPVPEVS  
QETNSITVYASVTLPESE (SEQ ID NO: 148) Mouse SLAM >NM\_013730.4 *Mus musculus* signaling lymphocytic (CD150) activation molecule family member 1 (Slamf1), transcript variant 1, mRNA  
GAGCTTCTTCCTTGGGGGTAACAGTAAGCAGCTGTCCTGCCGAGCTGAGCTGA  
GCTGAGCTCACAGCTGGGGACCCTGTCTGCGATTGCTGGCTAATGGATCCCAA  
AGGATCCCTTTCCTGGAGAATACTTCTGTTTCTCTCCCTGGCTTTTGAGTTGA  
GCTACGGAACAGGTGGAGGTGTGATGGATTGCCCAGTGATTCTCCAGAAGCTG  
GGACAGGACACGTGGCTGCCCCTGACGAATGAACATCAGATAAATAAGAGCGT  
GAACAAAAGTGTCGCGATCCTCGTCACCATGGCGACGTCCCCAGGAAGCAAAT  
CCAACAAGAAAATTGTGTCTTTTGATCTCTCTAAAGGGAGCTATCCAGATCAC  
CTGGAGGATGGCTACCACTTTCAATCAAAAAACCTGAGCCTGAAGATCCTCGG  
GAACAGGCGGGAGAGTGAAGGATGGTACTTGGTGAGCGTGGAGGAGAACGTTT  
CTGTTTCAGCAATTCTGCAAGCAGCTGAAGCTTTATGAACAGGTCTCCCCTCCA  
GAGATTAAAGTGCTAAACAAAACCCAGGAGAACGAGAATGGGACCTGCAGCTT  
GCTGTTGGCCTGCACAGTGAAGAAAGGGGACCATGTGACTTACAGCTGGAGTG  
ATGAGGCAGGCACCCACCTGCTGAGCCGAGCCAACCGCTCCCACCTCCTGCAC  
ATCACTCTTAGCAACCAGCATCAAGACAGCATCTACAACCTGCACCGCAAGCAA  
CCCTGTCAGCAGTATCTCTAGGACCTTCAACCTATCATCGCAAGCATGCAAGC  
AGGAATCCTCCTCAGAATCGAGTCCATGGATGCAATATACTCTTGTACCACTG  
GGGGTCGTTATAATCTTCATCCTGGTTTTACGGCAATAATAATGATGAAAAG  
ACAAGGTAAATCAAATCACTGCCAGCCACCAGTGGAAGAAAAAAGCCTTACTA  
TTTATGCCCAAGTACAGAAATCAGGGCCTCAAGAGAAGAAACTTCATGATGCC  
CTAACAGATCAGGACCCCTGCACAACCATTTATGTGGCTGCCACAGAGCCTGC  
CCCAGAGTCTGTCCAGGAACCAAACCCACCACAGTTTATGCCAGTGTGACAC  
TGCCAGAGAGCTGACCCATATAACCCAGTGAAAGGACTTTTTGAAGGAGGATAG  
AAGAACC AAAATCCACACTGAACTGGACCCCGGGTCCCAAGTTCTCTGTGACA  
GAAACTGCACATCTGTAACCTTCTCCAATCAGTTCCCTGGTGACGGATCTGCA  
CAGGCGTGCTTATGAAGTAGATGAGAAGTGAGGCTTCCCTGGGCATGCAACCTG  
CTCTGCTGCTGACACAGATATGAAGCAGAGATCCCGTGGTACAGTGTACCATC  
TTTGCTGTAGCAGATAATGTGGGTTTAGGCATCTCACTCTTTGCTGGACTGGA  
TAACAGAACTCAAAAAAAAACCAACAAGCCAAAGACATAGACTCCATCTCAGA  
TGGCTGAGCACAAAGTATAAAAGCCATTTTGGCTCTCTGGACTTTATTCTGGA  
AGCTGATCCTGATCACCTCAAGGCCAAGGGCTCCATGCCTCAGTTTCTCTCTC  
ACCCTCTAGATGAAGAGGGAACAAAGCATAAAGAGTGAAATCCTTGTTGTCTG  
AGATCATTCTATAAACGAACTGACATTTTATTTGCAAAACTCAAGCTAGTAAT  
TCAGTAGACTTGAAGATGATTTTAGAGCCTCTTATGCTTCAAACAACAGAATG  
AAATCCATCCAATGTTCTTCAAAGTGTTGTTCTCTGATTAAGTCAAAGCAACA  
CTGTTTGGCAATGCTGCTGTAAAGTTGCCTGGAATACTCAGAGGAACTTGTCC  
CAGGGAGGTTTTTTTCACTTCTTCAAAGAACTTTTGAATTTAAGTTCTCTGTT  
TATTCCTTGAGCAAACTCTGGAACCTCAAGAGTCTCTCTCCGTTGGTTCTG  
AGGCCATTTTATAGCCTAGGCCTCCTGTGGATCTACATGTGTATCACCCACTT  
CCTATCTCACTGCATACCTCTGTGTAGTAGTAAATTTAACCTCAAGTAGAAAA  
TTAAATTATTTTGGATGATCAGTTCCAAATGATTAGATGTTTAGTCTCTTATA  
ATAGGATGTAGGTAGAGTCTATATAAAGTCCTATATTCTTCACGTTGTCTGTC

>NM\_001250.6 *Homo sapiens* CD40 molecule (CD40), transcript variant 1, mRNA  
AGTGGTCCTGCCGCTGGTCTCACCTCGCTATGGTTTCGTCTGCCTCTGCAGTG  
CGTCCTCTGGGGCTGCTTGCTGACCGCTGTCCATCCAGAACCACCCACTGCAT  
GCAGAGAAAAACAGTACCTAATAAACAGTCAGTGOTGTTCTTTGTGCCAGCCA  
GGACAGAAACTGGTGAGTGACTGCACAGAGTTCACTGAAACGGAATGCCTTCC  
TTGCGGTGAAAGCGAATTCCTAGACACCTGGAACAGAGAGACACACTGCCACC  
AGCACAAATACTGCGACCCCAACCTAGGGCTTCGGGTCCAGCAGAAGGGCACC  
TCAGAAACAGACACCATCTGCACCTGTGAAGAAGGCTGGCACTGTACGAGTGA  
GGCCTGTGAGAGCTGTGTCCTGCACCGCTCATGCTCGCCCGGCTTTGGGGTCA  
AGCAGATTGCTACAGGGGTTTCTGATAACCATCTGCGAGCCCTGCCCAGTCGGC  
TTCTTCTCCAATGTGTCATCTGCTTTTCGAAAAATGTCACCCTTGGACAAGCTG  
TGAGACCAAAGACCTGGTTGTGCAACAGGCAGGCACAAACAAGACTGATGTTG  
TCTGTGGTCCCCAGGATCGGCTGAGAGCCCTGGTGGTGATCCCCATCATCTTC  
GGGATCCTGTTTGCCATCCTCTTGGTGCTGGTCTTTATCAAAAAGGTGGCCAA  
GAAGCCAACCAATAAGGCCCCCCCACCCCAAGCAGGAACCCCAGGAGATCAATT  
TTCCCGACGATCTTCCTGGCTCCAACACTGCTGCTCCAGTGCAGGAGACTTTA  
CATGGATGCCAACCGGTCACCCAGGAGGATGGCAAAGAGAGTCGCATCTCAGT  
GCAGGAGAGACAGTGAGGCTGCACCCACCCAGGAGTGTGGCCACGTGGGCAAA  
CAGGCAGTTGGCCAGAGAGCCTGGTGCTGCTGCTGCTGTGGCGTGAGGGTGAG  
GGGCTGGCACTGACTGGGCATAGCTCCCCGCTTCTGCCTGCACCCCTGCAGTT  
TGAGACAGGAGACCTGGCACTGGATGCAGAAACAGTTCACCTTGAAGAACCTC  
TCACTTCACCCTGGAGCCCATCCAGTCTCCCAACTTGTATTAAAGACAGAGGC  
AGAAGTTTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGT  
TTCCGATCCAGCAGTTTGGTGCCAGAGAGGCATCATGGTGGCTTCCCTGCGC  
CCAGGAAGCCATATACACAGATGCCCATTCAGCATTGTTTGTGATAGTGAAC  
AACTGGAAGCTGCTTAACTGTCCATCAGCAGGAGACTGGCTAAATAAAATTAG  
AATATATTTATACAACAGAATCTCAAAAACACTGTTGAGTAAGGAAAAAAAGG  
CATGCTGCTGAATGATGGGTATGGAACCTTTTTTAAAAAAGTACATGCTTTTATG  
TATGTATATTGCCTATGGATATATGTATAAATACAATATGCATCATATATTGA  
TATAACAAGGGTTCTGGAAGGGTACACAGAAAACCCACAGCTCGAAGAGTGGT  
GACGTCTGGGGTGGGGAAGAAGGGTCTGGGGGAGGGTTGGTTAAAGGGGAGATT

TGGCTTCCATTCATGCTTCATATTTTCCCAAAGGAGAGTGGAATTCACAT  
AATGCTTATGTAATTAATAAATCATCAAACATGTAAAAA (SEQ ID NO: 151)  
>NP\_001241.1 tumor necrosis factor receptor superfamily member 5 isoform 1  
precursor [*Homo sapiens*]  
MVRLPLQCVLWGCLLTAVHPEPPTACREKQYLINSQCCSLCQPGQKLVSDCTE  
FTETECLPCGESEFLDTWNRETHCHQHKEYCDPNLGLRVQQKGTSETDTICTCE  
EGWHCTSEACESCVLHRSCSPGFGVKQIATGVSDTICEPCPVGFFSNVSSAFE  
KCHPWTSCETKDLVVQQAGTNKTDVVCGPQDRLRALVVIPIIFGILFAILVL  
VFIKKVAKKPTNKAPHPKQEPQEINFPDDLPGSNTAAPVQETLHGCQPVTQED  
GKESRISVQERQ (SEQ ID NO: 152) Mouse CD40 >NM\_170703.2 *Mus musculus*  
CD40 antigen (Cd40), transcript variant 2, mRNA  
AGCAGGGACTTTGGAGTGAAGTGTGGCTTCAGCAGGAGCCCTGTGATTTGGCT  
CTTCTGATCTCGCCCTGCGATGGTGTCTTTGCCTCGGCTGTGCGCGCTATGGG  
GCTGCTTGTGACAGCGGTCCATCTAGGGCAGTGTGTTACGTGCAGTGACAAA  
CAGTACCTCCACGATGGCCAGTGCTGTGATTTGTGCCAGCCAGGAAGCCGACT  
GACAAGCCACTGCACAGCTCTTGAGAAGACCCAATGCCACCCATGTGACTCAG  
GCGAATTCTCAGCCCAGTGGAACAGGGAGATTCGCTGTCACCAGCACAGACAC  
TGTGAACCCAATCAAGGGCTTCGGGTAAAGAAGGAGGGCACCGCAGAATCAGA  
CACTGTCTGTACCTGTAAGGAAGGACAACACTGCACCAGCAAGGATTGCGAGG  
CATGTGCTCAGCACACGCCCTGTATCCCTGGCTTTGGAGTTATGGAGATGGCC  
ACTGAGACCACTGATACCGTCTGTCTATCCCTGCCAGTCGGCTTCTTCTCCAA  
TCAGTCATCACTTTTCGAAAAGTGTTATCCCTGGACAAGGTTTAAAGTCCCGG  
ATGCGAGCCCTGCTGGTCATTCTGTCTGATGGGCATCCTCATCACCATTTT  
CGGGGTGTTTCTCTATATCAAAAAGGTGGTCAAGAAACCAAAGGATAATGAGA  
TCTTACCCCCTGCGGCTCGACGGCAAGATCCCCAGGAGATGGAAGATTATCCC  
GGTCATAACACCGCTGCTCCAGTGCAGGAGACGCTGCACGGGTGTCAGCCTGT  
CACACAGGAGGATGGTAAAGAGAGTCGCATCTCAGTGCAGGAGCGGCAGGTGA  
CAGACAGCATAGCCTTGAGGCCCTGGTCTGAACCCTGGAAGTCTTTGGAGG  
CGATGGCTCGGCTCGGGAGCAGGGGCCTGGCTCTGAGGACTGCTTGCTGACCT  
TTGAAGTTTGAGATGAGCCAAGACAGAGCCCAGTGCAGCTAACTCTCATGCCT  
GCCCCCTATCATTCTCAACTTGCTTTTAAAGGATGGAGGGAGAGCTCGGGCA  
TCGGGGGTCCACAGTGATACCTACCAAGTGCAGCAGTGCAGGACCCAGAGTCG  
TCTTGCTGCGGCGTTCAGTGTAAAGGAGTCATGGACACAGGAGTCCGTGGCCCA  
CAGCTTGTGCTGCTAGAGGGCACCTGGTTGCCCATCAGCAGGGTACTGGCTAA  
ATAAATCTGTAATTATTTATACAATGACATCTCAGAACTCTAGCAGGTGGGG  
CAGAAAACAGGTAGTAGAATGATGGGTAGAGAAATAGCTTTTAAACACATTC  
CAAGGCAGGTAAGATGGCTTTTGTGAGTAAAGGAGCTTGCTGCCCAAACCCGG  
TTACCTGATTTTGATCCCTGGGACTTCATGGTAAAAGGGAGAGAACCAAATCC  
AGAGGGTTGTCAATTTGACCTCCATGTGTGCTCTGTGGTAATGTACCCCGTGTG  
TGCACATGTGCACATATCCTAAAATGGATGTGGTGGTGTATTGTAGAAATTAT  
TTAATCCCGCCCTGGGGTTTCTACCTGTGTGTTACCATTAGTTCTTGAATAA  
AAGACACACTCAACCTTTATATTTACAATAA (SEQ ID NO: 153) >NP\_733804.1  
tumor necrosis factor receptor superfamily member 5 isoform 2 precursor [*Mus  
musculus*]  
MVSLPRLCALWGCLLTAVHLGQCVTCSDKQYLHDGQCCDLCPGSRSLTSHCTA  
LEKTQCHPCDSGEFSAQWNREIRCHQHRHCEPNQGLRVKKEGTAESDTVCTCK  
EGQHCTSKDCEACAQHTPCIPGFGVMEMATETTDTVCHPCPVGFFSNQSSLFE  
KCYPWTRFKVPDASPAGHSCRDGHPHHHFRGVSLYQKGGQETKG (SEQ ID NO: 154)  
Human CD28 >NM\_006139.4 *Homo sapiens* CD28 molecule (CD28), transcript  
variant 1, mRNA  
ACACTTCGGGTTCTCTCGGGGAGGAGGGGCTGGAACCCTAGCCCATCGTCAGGA

CAAGAGATGCTCAGGCTCTGCTCTTATTCCTCTTCAATTGAGT  
ACAGGAAACAAGATTTTGGTGAAGCAGTCGCCCATGCTTGTAGCGTACGACAA  
TGCGGTCAACCTTAGCTGCAAGTATTCTTCTCAAGGGAGTTCC  
GGGCATCCCTTCACAAAGGACTGGATAGTGCTGTGGAAGTCTGTGTTGTATAT  
GGGAATTACTCCCAGCAGCTTCAGGTTTACTCAAAAACGGGGTTCAACTGTGA  
TGGGAAATTGGGCAATGAATCAGTGACATTCTACCTCCAGAATTTGTATGTTA  
ACCAAACAGATATTTACTTCTGCAAAATTGAAGTTATGTATCCTCCTCCTTAC  
CTAGACAATGAGAAGAGCAATGGAACCATTATCCATGTGAAAGGGAAACACCT  
TTGTCCAAGTCCCCTATTTCCCGGACCTTCTAAGCCCTTTTGGGTGCTGGTGG  
TGGTTGGTGGAGTCCTGGCTTGCTATAGCTTGCTAGTAACAGTGGCCTTTATT  
ATTTTCTGGGTGAGGAGTAAGAGGAGCAGGCTCCTGCACAGTGACTACATGAA  
CATGACTCCCCGCCGCCCGGGCCCCACCCGCAAGCATTACCAGCCCTATGCCC  
CACCACGCGACTTCGCAGCCTATCGCTCCTGACACGGACGCCTATCCAGAAGC  
CAGCCGGCTGGCAGCCCCCATCTGCTCAATATCACTGCTCTGGATAGGAAATG  
ACCGCCATCTCCAGCCGCCACCTCAGGCCCCTGTTGGGCCACCAATGCCAAT  
TTTTCTCGAGTGACTAGACCAAATATCAAGATCATTTTGAGACTCTGAAATGA  
AGTAAAAGAGATTTCTGTGACAGGCCAAGTCTTACAGTGCCATGGCCACAT  
TCCAACTTACCATGTACTTAGTGACTTGACTGAGAAGTTAGGGTAGAAAACAA  
AAAGGGAGTGGATTCTGGGAGCCTCTTCCCTTTCTCACTCACCTGCACATCTC  
AGTCAAGCAAAGTGTGGTATCCACAGACATTTTAGTTGCAGAAGAAAGGCTAG  
GAAATCATTCTTTTGGTTAAATGGGTGTTTAATCTTTTGGTTAGTGGGTAA  
ACGGGGTAAGTTAGAGTAGGGGGAGGGATAGGAAGACATATTTAAAACCAT  
AAAACACTGTCTCCCCTCATGAAATGAGCCACGTAGTTCCTATTTAATGCTG  
TTTTCTTTAGTTTAGAAATACATAGACATTGTCTTTTATGAATTCTGATCAT  
ATTTAGTCATTTTGACCAAATGAGGGATTTGGTCAAATGAGGGATTCCCTCAA  
AGCAATATCAGGTAAACCAAGTTGCTTTCCTCACTCCCTGTCATGAGACTTCA  
GTGTTAATGTTTACAATATACTTTCGAAAGAATAAAATAGTTCTCCTACATGA  
AGAAAGAATATGTCAGGAAATAAGGTCACTTTATGTCAAATTTATTTGAGTAC  
TATGGGACCTGGCGCAGTGGCTCATGCTTGTAATCCCAGCACTTTGGGAGGCC  
GAGGTGGGCAGATCACTTGAGATCAGGACCAGCCTGGTCAAGATGGTGAACT  
CCGTCTGTACTAAAAATACAAAATTTAGCTTGGCCTGGTGGCAGGCACCTGTA  
ATCCCAGCTGCCCAAGAGGCTGAGGCATGAGAATCGCTTGAACCTGGCAGGCG  
GAGGTTGCAGTGAGCCGAGATAGTGCCACAGCTCTCCAGCCTGGGCGACAGAG  
TGAGACTCCATCTCAAACAACAACAACAACAACAACAACAACAACAACCACA  
AAATTATTTGAGTACTGTGAAGGATTATTTGTCTAACAGTTCATTCCAATCAG  
ACCAGGTAGGAGCTTTCCTGTTTCATATGTTTCAGGGTTGCACAGTTGGTCTC  
TTTAATGTCGGTGTGGAGATCCAAAGTGGGTGTGGAAAGAGCGTCCATAGGA  
GAAGTGAGAATACTGTGAAAAGGGATGTTAGCATTATTAGAGTATGAGGAT  
GAGTCCCAAGAAGGTTCTTTGGAAGGAGGACGAATAGAATGGAGTAATGAAAT  
TCTTGCCATGTGCTGAGGAGATAGCCAGCATTAGGTGACAATCTTCCAGAAGT  
GGTCAGGCAGAAAGGTGCCCTGGTGAGAGCTCCTTTACAGGGACTTTATGTGGT  
TTAGGGCTCAGAGCTCCAAAACCTCTGGGCTCAGCTGCTCCTGTACCTTGGAGG  
TCCATTCACATGGGAAAGTATTTTGGAAATGTGTCTTTTGAAGAGAGCATCAGA  
GTTCTTAAGGGACTGGGTAAGGCCTGACCCTGAAATGACCATGGATATTTTTC  
TACCTACAGTTTGAGTCAACTAGAATATGCCTGGGGACCTTGAAGAATGGCCC  
TTCAGTGGCCCTCACCATTTGTTTCATGCTTCAGTTAATTCAGGTGTTGAAGGA  
GCTTAGGTTTTAGAGGCACGTAGACTTGGTTCAAGTCTCGTTAGTAGTTGAAT  
AGCCTCAGGCAAGTCACTGCCCACCTAAGATGATGGTTCTTCAACTATAAAAT  
GGAGATAATGGTTACAAATGTCTCTTCCTATAGTATAATCTCCATAAGGGCAT  
GGCCCAAGTCTGTCTTTGACTCTGCCTATCCCTGACATTTAGTAGCATGCCCCG

ACATACAACTGTTGTTATGTTGCTATATAGTATAAATTATGTATATAAA  
AATTAAACTGGGCAATAGCCTAAGAAGGGGGGAATATTGTAACACAAATTTAA  
ACCCACTACGCAGGGATGAGGTGCTATAATATGAGGACCTTTTAACTTCCATC  
ATTTTCCTGTTTCTTGAAATAGTTTATCTTGTAATGAAATATAAGGCACCTCC  
CACTTTTATGTATAGAAAGAGGTCTTTTAAATTTTTTTTTTAATGTGAGAAGGAA  
GGGAGGAGTAGGAATCTTGAGATTCCAGATCGAAAATACTGTACTTTGGTTGA  
TTTTTAAGTGGGCTTCCATTCCATGGATTTAATCAGTCCCAAGAAGATCAAAC  
TCAGCAGTACTTGGGTGCTGAAGAACTGTTGGATTACCTTGGCACGTGTGCC  
ACTTGCCAGCTTCTTGGGCACACAGAGTTCTTCAATCCAAGTTATCAGATTGT  
ATTTGAAAATGACAGAGCTGGAGAGTTTTTTTGAAATGGCAGTGGCAAATAAAT  
AAATACTTTTTTTTAAATGGAAAGACTTGATCTATGGTAATAAATGATTTTGT  
TTTCTGACTGGAAAAATAGGCCTACTAAAGATGAATCACACTTGAGATGTTTC  
TTACTCACTCTGCACAGAAACAAAGAAGAAATGTTATACAGGGAAGTCCGTTT  
TCACTATTAGTATGAACCAAGAAATGGTTCAAAAACAGTGGTAGGAGCAATGC  
TTTCATAGTTTCAGATATGGTAGTTATGAAGAAAACAATGTCATTTGCTGCTA  
TTATTGTAAGAGTCTTATAATTAATGGTACTCCTATAATTTTTTGATTGTGAGC  
TCACCTATTTGGGTAAAGCATGCCAATTTAAAGAGACCAAGTGTATGTACATT  
ATGTTCTACATATTCAGTGATAAAATTACTAACTACTATATGTCTGCTTTAA  
ATTTGTACTTTAATATTGTCTTTTGGTATTAAGAAAGATATGCTTTCAGAATA  
GATATGCTTCGCTTTGGCAAGGAATTTGGATAGAACTTGCTATTTAAAAGAGG  
TGTGGGGTAAATCCTTGTATAAATCTCCAGTTTAGCCTTTTTTTGAAAAAGCTA  
GACTTTCAAATACTAATTTCACTTCAAGCAGGGTACGTTTCTGGTTTGTTC  
TTGACTTCAGTCACAATTTCTTATCAGACCAATGGCTGACCTCTTTGAGATGT  
CAGGCTAGGCTTACCTATGTGTTCTGTGTCATGTGAATGCTGAGAAGTTTGAC  
AGAGATCCAACCTTCAGCCTTGACCCCATCAGTCCCTCGGGTTAACTAACTGAG  
CCACCGGTCCTCATGGCTATTTTAATGAGGGTATTGATGGTTAAATGCATGTC  
TGATCCCTTATCCCAGCCATTTGCACTGCCAGCTGGGAACTATAACCAGACCTG  
GATACTGATCCCAAAGTGTTAAATTCAACTACATGCTGGAGATTAGAGATGGT  
GCCAATAAAGGACCCAGAACCAGGATCTTGATTGCTATAGACTTATTAATAAT  
CCAGGTCAAAGAGAGTGACACACACTCTCTCAAGACCTGGGGTGAGGGAGTCT  
GTGTTATCTGCAAGGCCATTTGAGGCTCAGAAAGTCTCTCTTTCTTATAGATA  
TATGCATACTTTCTGACATATAGGAATGTATCAGGAATACTCAACCATCACAG  
GCATGTTCTACCTCAGGGCCTTTACATGTCCTGTTTACTCTGTCTAGAATGT  
CCTTCTGTAGATGACCTGGCTTGCCTCGTCACCCTTCAGGTCCTTGCTCAAGT  
GTCATCTTCTCCCCTAGTTAAACTACCCACACCCTGTCTGCTTTCTTGCTT  
ATTTTTCTCCATAGCATTTTACCATCTCTTACATTAGACATTTTTCTTATTTA  
TTTGTAGTTTATAAGCTTCATGAGGCAAGTAACTTTGCTTTGTTTCTTGCTGT  
ATCTCCAGTGCCCAGAGCAGTGCCTGGTATATAATAAATATTTATTGACTGAG TGAA

(SEQ ID NO: 155) >NP\_006130.1 T-cell-specific surface glycoprotein CD28 isoform 1 precursor [*Homo sapiens*]

MLRLLALNLFPSIQVTGNKILVKQSPMLVAYDNAVNLSCKYSYNLFSREFRA  
SLHKGLDSAVEVCVVYGNYSQQLQVYSKTGFNC DGKLGNESVTFY LQNLVYNQ  
TDIYFCKIEVMYPPPYLDNEKSNGTIIHVKGKHLCPSP LFPGPSKPFWVLVVV  
GGVLACYSLLVTVAFIIFWVR SKRSRL LHS DYMNMTPRRPGPTRKHYPYAPP  
RDFAAYRS (SEQ ID NO: 156) Mouse CD28 >NM\_007642.4 *Mus musculus*  
CD28 antigen (Cd28), mRNA

AGACCTTGGCAGATGTGACTTCAGTTCACACCACACTCTGCCTTGCTCACAGA  
GGAGGGGCTGCAGCCCTGGCCCTCATCAGAACAAATGACACTCAGGCTGCTGTT  
CTTGGCTCTCAACTTCTTCTCAGTTCAAGTAACAGAAAACAAGATTTTGGTAA  
AGCAGTCGCCCCTGCTTGTGGTAGATAGCAACGAGGTCAGCCTCAGCTGCAGG



TATTCCTACAACCTTCTCTCGAAGGCGGATCCGCTGTACCGGCTGTAACCGG  
GAACAGCGACGTGGAAGTCTGTGTCGGGAATGGGAATTTTACCTATCAGCCCC  
AGTTTCGCTCGAATGCCGAGTTCAACTGCGACGGGGATTTTCGACAACGAAACA  
GTGACGTTCCGTCTCTGGAATCTGCACGTCAATCACACAGATATTTACTTCTG  
CAAAATTGAGTTCATGTACCCTCCGCCTTACCTAGACAACGAGAGGAGCAATG  
GAACTATTATTCACATAAAAGAGAAACATCTTTGTGCATACTCAGTCATCTCCT  
AAGCTGTTTTTGGGCACTGGTCGTGGTTGCTGGAGTCCTGTTTTTGTATGGCTT  
GCTAGTGACAGTGGCTCTTTGTGTTATCTGGACAAATAGTAGAAGGAACAGAC  
TCCTTCAAAGTGACTACATGAACATGACTCCCCGGAGGCCTGGGCTCACTCGA  
AAGCCTTACCAGCCCTACGCCCTGCCAGAGACTTTGCAGCGTACCGCCCCCTG  
ACAGGGACCCCTATCCAGAAGCCCGCCGGCTGGTACCCGTCTACCTGCTCATC  
ATCACTGCTCTGGATAGGAAAGGACAGCCTCATCTTCAGCCGGCCACTTTGGA  
CCTCTACTGGGCCACCAATGCCAACTATTTTAGAGTGTCTAGATCTAACATCA  
TGATCATCTTGAGACTCTGGAATGAATGACAGAAGCTTCTATGGCAGGATAAA  
GTCTGTGTGGCTTGACCCAACTCAAGCTTAATACATTTATTGACTTGATTGG  
GGAAGTTAGAGTAGAGCAATCAAAAAGATCATTCAATCAGCCTTGGGAAAGTCA  
ATTTGCAGGCTCCTGGATGAGCCCTGCCCCGTTTTTCACTTGCCAGCACATTC  
AGTCATGTGGTGTGATAGCCAAAGATGTTTTTGGACAGAGAAGAAAGGATAGAA  
AAACCTTCTCTTTGGCTAAGTTGGTGTTTGGGGTGGGGATAGGTTAGAGTATA  
GTACTTAACTATTTGAAAAATAATGAAAACACTTTTTTCACTCATGAAATGAG  
CCACTTAGCTCCTAAATAGTGTTTTCTCTGTTAGTTTAGAAAGTTGTGGACATA  
TTTTTTTAATGATTTCTGACCATTTTAAATCACATTGACTCATGGAATGGCCT  
CAAAGCACCCCCCAGTGCTTCTTTCCTCATTCCCGGTCATGGGAAGTCAAGTAT  
TATTAATAGTCACAACATGATTTCAGAACTAGATAGCCCTCCCACACCAAGAA  
GAATGTGAGAGGAAGTAAGGTCACCTTTATGTAAAAAAGGATAGGATAGGATAG  
GTACACATATGTATGTATACATACATACCTATGTGCACACACACACATATA  
CATACACACAAAATGCTATGAAGAGTTATCTGTTTAGTAGCCTGTTATAGTCA  
AATCATTTTAAAGTTTCAACTTCTTACAGTTGGGCCACTTGTTGTCCTTTGTGG  
ATGGATATCTGAAATTGTGTCTATATATTGCTAGTCATGATACTGTGAACAAA  
AAGGGTAGTGTTAGTATTTGTCAGGGTGGTAAGGATGCATTCCAGGAAGCTTC  
CTCTGAGGAAGGGAATGAGGTCATTCTTGCCATGTATGAAAGACATAGATGTT  
TTCCAGAAGGCACCATTGGGAGCCCCAGTATAAGTTCCCTTAGACTCTACAGT  
TTAGAGGGATTTTATATGTCCTAGGACTCAGGACTCCAGAACTTTGTGGGCTC  
AGCTGCTTCATACCATGGGGATACATTGACATGAACAATTATTTTGGAAATGTG  
TCTTTAGGGACGACATCAAAGTTCTCAAGTACCTACAAGACCTGATACTGGAA  
TGAAGGTGGACTTTCTTTTTTGTCTTCCAGTTCGGATCAACTGGAATGTATCTG  
GGGACCTTGAAGAACGGCTGTCCAGCTGTCTTCACCATTTGTATAGTGCTTTG  
AATTATTCAGAGGTTTAAAGTCAGGAAGACCTGGTTTAAAAAACATTTCAAT  
ATGAGTTAAATGGCCTCAGGCAAGTCACTGTTTCATCCAAGTCTATGACTCCTC  
AACTGTAAGATGGCCACACTGAAACTTGCTAAGATCCTCTGGCCTCTGCCTCC  
CAAGAGTTGGGATTTCAAGGAGTGCACAATCATGACCCAACTCGTGATAATCT  
CTCAGCTTCAATAACTTTCCAGCTAATTGGAATATCCTGTAATCAAACATGAG  
GCATTTCCCCTCCCCCCTGTTTTTGTGTATAAAGAGATCTTTAAACTTTTT  
TTTTAATATGAGGGGTAAGAAAAGATAGGAATCTTTTAATTCTAGACAGAAGA  
TATTGTGCTTTGGTTTTTTTTTTTTTTAATGGCTTCTATTCTGTGCTTTTAAT  
TAAACCAGAGAAGGCCAAGATTAGCCCTACTTGTGTGATAAAGAATGCTGGC  
CCTTGTGATTGCAGTCAGCCTCTTGACACATAGAGTTCTTGAATCTAAGTTAT  
AAAATTATATTTGAAAATGACAGAGCTGGAGAATTTATAGAAAGGGTCATAGC  
AAATAACAAACCATTTTTTTTTTAAACGGAAAGATTTGGTCTTTGGCAATCAAT  
AACTTTGTTTTCTAACTGGAAAAGGAGGTTTACTGGAGATGAATCACACCTGA

AGTGTTCCTCTTCACTGACGATACGAGTGTCCAGCTTT  
CGCTCTCGGTATGAACCAACAGGCGGGTTAAAAACACTGGGTCAGAGTAAAGC  
TTTTGCAGTTTCAGATGTAGTGTGTATGAAGAAAACATATGTCACCTTGCTGCTA  
TTATTGTAAGAGTCTAAGAACTAAAGGTGTGCCTGTAATTTCTAATTATGAGC  
TCACCTATTTGGTACCGAGCATGCCAATTTTAAAGAGACCCGGTGTACCTTAT  
AGCTACATCCAATGATAAAATTACCACACTAGCACATGCCTGTGTTTAAACTC  
GTGCTTTAATGTTTTTCTTAGGGCAGGTATGCACCCCTTTGCAGTGAGTTGG  
GAGAGATTTTGAAAAAGTGTATGACAAACATTTTAAACACCTTTGGTTTCCTC  
TCTCTGTGTCTCTTTGTCTCTGTCTCTCTCTTTCTCTCCTGTGCATATGTCTC  
CCCTCCCTCACTTCTCTGTCTCTTCCTCTCTCCCTCTCTCTGTCTTTCTCTGT  
GTGTCTCTCTGTCTCTGTGTATCTCTCTGTCTGTCTCTTTCTCTGCAGATTTT  
CAAAACGTTGTTTTTCTATGGAAGAAATACAAGCTGTGGTTGGTTTGCTACGA  
GTCAGTAGCAGTTTATCAGTAGGCCAATGTTTTATCTCTTGGAGATTTTCAGTC  
TGGGTTTACCCAATGTATTCTCTGTAATGTGACTGCTGGGGACAGATATAACT  
TGATTGAGCCTTCAAATCATTAGGTCTTCAATCATTAGTCAACGGAGTGAG  
CCACTAATCTGCAATGGCTATTTTAATATGCATACTGATGGTCAAATGGATGT  
CTGATCTCTCATCCCAGCTTTCTGTACTACCATATGGGAACCTATATGTAACCT  
GTATACTTACCTGAATATGTTAAATTCAACTACATGGTAAGATGGACCAGAAA  
TTGCAATGTTTCATGTCCATATAGCCACCATTAAACCCAAGTTAAGCACAGTAGT  
GTGGGTTCTCTCAGGACTTGTGAATGAGTTTATGCTCTCTACAAAGACAGGTG  
AAGCTTAAATCTCTCTTGCCTGCTATGTTTATGCAAATATCAAGATTGTTTC  
TGTACCAGGGACTTAACACATTCTATTCTACTATTTTCCCTGTCTACAATGT  
TATTTTCATAGATATCTACTTGGTTTGTCTTACTTCCTTGACATATTTGCCCA  
AATGCCACCTTCAACTGTAGTTAATTACCTGTACAACCTGTCTCCATGCCTTG  
TTTTATTTTCTCTATAACTCTACTAATAGGTATTTTCTTATTTATTGGTTTA  
TTGCCTGTTTTTTTTTCTAAATCTACACCGGATCTCCAAAGGGAAAGAACTCC  
ATTTGCTTTGATTTTATTGCTGTATCCCCAGTGCCTAGAATAATGCTTAGCCT  
GCAATAAATATTTATTTCATTGACT (SEQ ID NO: 157)>NP\_031668.3 T-cell-  
specific surface glycoprotein CD28 precursor [*Mus musculus*]  
MTLRLFLALNFFSVQVTENKILVKQSPLLVDSDNEVSLSCRYSYNLLAKEFR  
ASLYKGVNSDVEVCVGNGNFTYQPQFRSNAEFNCDGDFDNETVTFRLWNLHVN  
HTDIYFCKIEFMYPYPYLDNERSNGTIIHIKEKHLCHTQSSPKLFWALVVVAG  
VLFCYGLLVTVALCVIWTNSRRNRLQLSDYMNMTPRRPGLTRKPYQPYAPARD FAAYRP  
(SEQ ID NO: 158) Human>NM\_144615.2 *Homo sapiens* transmembrane and  
CD28H immunoglobulin domain containing 2 (TMIGD2), transcript variant 1, mRNA  
GGAAGTCTGTCAACTGGGAGGGGGGAGAGGGGGGTGATGGGCCAGGAATGGGGT  
CCCCGGGCATGGTGCTGGGCCTCCTGGTGCAGATCTGGGCCCTGCAAGAAGCC  
TCAAGCCTGAGCGTGCAGCAGGGGCCCAACTTGCTGCAGGTGAGGCAGGGCAG  
TCAGGCGACCCTGGTCTGCCAGGTGGACCAGGCCACAGCCTGGGAACGGCTCC  
GTGTTAAGTGGACAAAGGATGGGGCCATCCTGTGTCAACCGTACATCACCAAC  
GGCAGCCTCAGCCTGGGGGTCTGCGGGCCCCAGGGACGGCTCTCCTGGCAGGC  
ACCCAGCCATCTCACCCTGCAGCTGGACCCTGTGAGCCTCAACCACAGCGGGG  
CGTACGTGTGCTGGGCGGCCGTAGAGATTCCTGAGTTGGAGGAGGCTGAGGGC  
AACATAACAAGGCTCTTTGTGGACCCAGATGACCCACACAGAACAGAAACCG  
GATCGCAAGCTTCCCAGGATTCCTCTTCGTGCTGCTGGGGGTGGGAAGCATGG  
GTGTGGCTGCGATCGTGTGGGGTGCCTGGTTCTGGGGCCGCCGCAGCTGCCAG  
CAAAGGGACTCAGGTAACAGCCCAGGAAATGCATTCTACAGCAACGTCCTATA  
CCGGCCCCGGGGGGCCCCAAAGAAGAGTGAGGACTGCTCTGGAGAGGGGAAGG  
ACCAGAGGGGCCAGAGCATTTATTCAACCTCCTTCCCGCAACCGGCCCCCGC  
CAGCCGCACCTGGCGTCAAGACCCTGCCCCAGCCCGAGACCCTGCCCCAGCCC

CCAGCCGGCCCGCCGAGGCCAAAAGGGTTCCTAGAGGAGCCCA  
CCCAGCAGCCGAGGCCAAAAGGGTTCCTAGAGGAGTGGGAGAGGAGTGAGAGATC  
CCAGGAGACCTCAACAGGACCCCATAGGTACACACAAAAAAGGGGGGAT  
CGAGGCCAGACACGGTGGCTCACGCCTGTAATCCCAGCAGTTTGGGAAGCCGA  
GGCGGGTGGAACTTGAGGTCAGGGGTTTGAGACCAGCCTGGCTTGAACCTG  
GGAGGCGGAGGTTGCAGTGAGCCGAGATTGCGCCACTGCACTCCAGCCTGGGC  
GACAGAGTGAGACTCCGTCTCAAAAAAAACAAAAAGCAGGAGGATTGGGAGCC  
TGTCAGCCCCATCCTGAGACCCCGTCCTCATTTCTGTAATGATGGATCTCGCT  
CCCCTTTCCCCCAAGAACCTAATAAAGGCTTGTGAAGAAAAAGCAAAAAAA  
AAAAA (SEQ ID NO: 159) >NP\_653216.2 transmembrane and  
immunoglobulin domain- containing protein 2 isoform 1 precursor [*Homo sapiens*]  
MGSPGMVLGLLVQIWALQEASSLSVQQGPNLLQVRQGSQATLVCQVDQATAWE  
RLRVKWKTDGAILCQPYITNGSLSLGVCGPQGRLSWQAPSHLTLQLDPVSLNH  
SGAYVCWAAVEIPELEEAEGNITRLFVDPDDPTQNRNRIASFPGFLFVLLGVG  
SMGVAAIVWGAWFWGRRSCQQRDSGNSPGNAFYSNVLYRPRGAPKKSEDCSGE  
GKDQRGQSIYSTSFPQPAPRQPHLASRPCSPRCPSPRPGHPVSMVRVSPRP  
SPTQQPRPKGFVKVGE (SEQ ID NO: 160) Human CD2 >NM\_001328609.2  
*Homo sapiens* CD2 molecule (CD2), transcript variant 1, mRNA  
AGTCTCACTTCAGTTCCTTTTGCATGAAGAGCTCAGAATCAAAGAGGAAACC  
AACCCCTAAGATGAGCTTTCCATGTAAATTTGTAGCCAGCTTCCTTCTGATTT  
TCAATGTTTCTTCCAAAGGTGCAGTCTCCAAAGAGATTACGAATGCCTTGGAA  
ACCTGGGGTGCCTTGGGTCAGGACATCAACTTGGACATTCTAGTTTTCAAAT  
GAGTGATGATATTGACGATATAAAATGGGAAAAAACTTCAGACAAGAAAAAGA  
TTGCACAATTCAGAAAAGAGAAAGAGACTTTCAAGGAAAAAGATACATATAAG  
CTATTTAAAAATGGAACCTCTGAAAATTAAGCATCTGAAGACCGATGATCAGGA  
TATCTACAAGGTATCAATATATGATACAAAAGGAAAAAAATGTGTTGGAAAAA  
TATTTGATTTGAAGATTCAAGAGAGGGTCTCAAACCAAAGATCTCCTGGACT  
TGTATCAACACAACCCTGACCTGTGAGGTAATGAATGGAACCTGACCCCGAATT  
AAACCTGTATCAAGATGGGAAACATCTAAACTTTCTCAGAGGGTTCATCACAC  
ACAAGTGGACCACCAGCCTGAGTGCAAAATTCAAGTGCACAGCAGGGAACAAA  
GTCAGCAAGGAATCCAGTGTCGAGCCTGTCAGCTGTCCAGGAGGCAGCATCCT  
TGGCCAGAGTAATGGGCTCTCTGCCTGGACCCCTCCCAGCCATCCCCTTCTC  
TTCCTTTTGCAGAGAAAGGTCTGGACATCTATCTCATCATTGGCATATGTGGA  
GGAGGCAGCCTCTTGATGGTCTTTGTGGCACTGCTCGTTTTCTATATCACCAA  
AAGGAAAAAACAGAGGAGTCGGAGAAATGATGAGGAGCTGGAGACAAGAGCCC  
ACAGAGTAGCTACTGAAGAAAGGGGGCCGGAAGCCCCACCAAATTCCAGCTTCA  
ACCCCTCAGAATCCAGCAACTTCCCAACATCCTCCTCCACCACCTGGTCATCG  
TTCCCAGGCACCTAGTCATCGTCCCCCGCCTCCTGGACACCGTGTTTCAGCACC  
AGCCTCAGAAGAGGCCTCCTGCTCCGTCGGGCACACAAGTTCACCAGCAGAAA  
GGCCCGCCCCCTCCCCAGACCTCGAGTTCAGCCAAAACCTCCCCATGGGGCAGC  
AGAAAACCTCATTGTCCCCTTCTCTAATTAAGATAAGAACTGTCTTTTT  
CAATAAAAAGCACTGTGGATTTCTGCCCTCCTGATGTGCATATCCGTACTTCC  
ATGAGGTGTTTTCTGTGTGCAGAACATTGTCACCTCCTGAGGCTGTGGGCCAC  
AGCCACCTCTGCATCTTCGAACTCAGCCATGTGGTCAACATCTGGAGTTTTTG  
GTCTCCTCAGAGAGCTCCATCACACCAGTAAGGAGAAGCAATATAAGTGTGAT  
TGCAAGAATGGTAGAGGACCGAGCACAGAAATCTTAGAGATTTCTTGTCCCT  
CTCAGGTCATGTGTAGATGCGATAAATCAAGTGATTGGTGTGCCTGGGTCTCA  
CTACAAGCAGCCTATCTGCTTAAGAGACTCTGGAGTTTCTTATGTGCCCTGGT  
GGACACTTGCCACCATCCTGTGAGTAAAGTGAAATAAAAGCTTTGACTAGA (SEQ  
ID NO: 161) >NP\_001315538.1 T-cell surface antigen CD2 isoform 1 precursor

[*Homo sapiens*]

MSFPCKFVASFLIFNVSSKGAVSKEITNALETWGALGQDINLDIPSFQMSDD  
IDDIKWEKTSDDKKKIAQFRKEKETFEKEDTYKLFKNGTLKIKHLKTDDQDIYK  
VSIYDTKGKNVLEKIFDLKIQERSKPKISWTCINTTLTCEVMNGTDPENLY  
QDGKHLKLSQRVITHKWTTSLSAKFKCTAGNKVSKESSVEPVSCPGGGSILGQS  
NGLSAWTPPSHPTSLPFAEKGLDIYLIIGICGGGSLLMVFVALLVFYITKRKK  
QRSRRNDEELETRAHRVATEERGRKPHQIPASTPQNPATSQHPPPPPGHRSQA  
PSHRPPPPGHRVQHQPQKRPPAPSGTQVHQKGPPLPRPRVQPKPPHGAAENS LSPSSN  
(SEQ ID NO: 162) Mouse CD2 >NM\_013486.2 *Mus musculus* CD2 antigen  
(Cd2), mRNA

GCCTCACACAGTCCTGACAGAAAGAACTCAGAGTCACCCCTGGGAAAAGAAC  
TCTAAAGATGAAATGTAAATTCCTGGGTAGCTTCTTTCTGCTCTTCAGCCTTT  
CCGGCAAAGGGGCGGACTGCAGAGACAATGAGACCATCTGGGGTGTCTTGGGT  
CATGGCATCACCTGAACATCCCCAACTTTCAAATGACTGATGATATTGATGA  
GGTGCGATGGGTAAAGGAGGGGCACCCTGGTCGCAGAGTTTAAAAGGAAGAAGC  
CACCTTTTTTGTATATCAGAAACGTATGAGGTCTTAGCAAACGGATCCCTGAAG  
ATAAAGAAGCCGATGATGAGAAACGACAGTGGCACCTATAATGTAATGGTGTA  
TGGCACAAATGGGATGACTAGGCTGGAGAAGGACCTGGACGTGAGGATTCTGG  
AGAGGGTCTCAAAGCCCATGATCCACTGGGAATGCCCCAACACAACCCTGACC  
TGTGCGGTCTTGCAAGGGACAGATTTTGAAGTGAAGCTGTATCAAGGGGAAAC  
ACTACTCAATAGTCTCCCCCAGAAGAACATGAGTTACCAGTGGACCAACCTGA  
ACGCACCATTCAAGTGTGAGGCGATAAACCCGGTCAGCAAGGAGTCTAAGATG  
GAAGTTGTAACTGTCCAGAGAAAGGTCTGTCCTTCTATGTCACAGTGGGGGT  
CGGTGCAGGAGGACTCCTCTTGGTGCTCTTGGTGGCGCTTTTTTATTTTCTGTA  
TCTGCAAGAGGAGAAAACGGAACAGGAGGAGAAAAGATGAAGAGCTGGAAATA  
AAAGCTTCCAGAACAAGCACTGTGGAAAGGGGCCCCAAGCCGCACTCAACCCC  
AGCCGCAGCAGCGCAGAATTCAGTGGCGCTCCAAGCTCCTCCTCCACCTGGCC  
ATCACCTCCAGACACCTGGCCATCGTCCCTTGCCCTCCAGGCCACCGTACCCGT  
GAGCACCAGCAGAAGAAGAGACCTCCTCCATCAGGCACACAGATTCACCAGCA  
GAAAGGCCCTCCTTTACCCAGACCCCGAGTTCAGCCAAAACCTCCCTGTGGGA  
GTGGAGATGGTGTTCCTGCTGCGCCCCCTAATTAAGAAGGCAGAGTTCGTCAT  
TTCCAATAAAAAGCTGTGTGGATTTATCTTC (SEQ ID NO: 163) >NP\_038514.1 T-  
cell surface antigen CD2 precursor [*Mus musculus*]

MKCKFLGSFLLFSLSGKGADCRDNETIWGVLGHGITLNPNFQMTDDIDEVR  
WVRRGTLVAEFKRKKPPFLISETYEVLANGSLKIKKPMRNDSTYNNVMVYGT  
NGMTRLEKDLDRILERSKPMIHWECNNTTLTCAVLQGTDFELKLYQGETLL  
NSLPQKNMSYQWTNLNAPFKCEAINPVSKESKMEVVNCPEKGLSFYVTVGVGA  
GGLLLVLLVALFIFCICKRRKRNRRRRKDEELEIKASRTSTVERGPKPHSTPAA  
AAQNSVALQAPPPPGHHLQTPGHRPLPPGHRTREHQKQKRPPPSGTQIHQQKG  
PPLPRPRVQPKPPCGSGDGVSLPPN (SEQ ID NO: 164) Human LFA-3  
>NM\_001779.3 *Homo sapiens* CD58 molecule (CD58), (CD58) transcript variant  
1, mRNA

GAAGTTAGGGCTGCTTGTGGCTGGGCACTCGCGCAGAGGCCGGCCCGACGAGC  
CATGGTTGCTGGGAGCGACGCGGGGCGGGGCCCTGGGGGTCTCAGCGTGGTCT  
GCCTGCTGCACTGCTTTGGTTTCATCAGCTGTTTTTCCCAACAAATATATGGT  
GTTGTGTATGGGAATGTAACCTTCCATGTACCAAGCAATGTGCCTTTAAAAGA  
GGTCCTATGGAAAAACAAAAGGATAAAGTTGCAGAACTGGAAAATTCTGAAT  
TCAGAGCTTTCTCATCTTTTAAAAATAGGGTTTATTAGACACTGTGTCAGGT  
AGCCTCACTATCTACAACCTTAACATCATCAGATGAAGATGAGTATGAAATGGA  
ATCGCCAAATATTACTGATACCATGAAGTTCTTTCTTTATGTGCTTGAGTCTC

TTCCATCTCCACACACTAATCTTGTCATTGACTAATGGAAGCATTGAAGTCCAA  
TGCATGATACCAGAGCATTACAACAGCCATCGAGGACTTATAATGTACTCATG  
GGATTGTCCTATGGAGCAATGTAAACGTA ACTCAACCAGTATATATTTTAAGA  
TGGAAAATGATCTTCCACAAAAAATACAGTGTACTCTTAGCAATCCATTATTT  
AATACAACATCATCAATCATTTTTGACAACCTGTATCCCAAGCAGCGGTCATTC  
AAGACACAGATATGCACTTATACCCATAACCATTAGCAGTAATTACAACATGTA  
TTGTGCTGTATATGAATGGTATTCTGAAATGTGACAGAAAACCAGACAGAACC  
AACTCCAATTGATTGGTAACAGAAGATGAAGACAACAGCATAACTAAATTATT  
TTAAAAACTAAAAAGCCATCTGATTTCTCATTTGAGTATTACAATTTTTGAAC  
AACTGTTGGAAATGTA ACTTGAAGCAGCTGCTTTAAGAAGAAATACCCACTAA  
CAAAGAACAAGCATTAGTTTTTGGCTGTCATCAACTTATTATATGACTAGGTGC  
TTGCTTTTTTTTGTCTAGTAAATTGTTTTTACTGATGATGTAGATACTTTTTGTAA  
ATAAATGTAAATATGTACACAAGTGA (SEQ ID NO: 165) >NP\_001770.1  
lymphocyte function-associated antigen 3 isoform 1 [*Homo sapiens*]  
MVAGSDAGRALGVLSVCLLHCFGFISCFSSQIYGVVYGNVTFHVP SNVPLKE  
VLWKKQKDKVAELENSEFRAFSSFKNRVYLDTVSGSLTIYNLTSSDEDEYEME  
SPNITDTMKFFLYVLES LPSPTLTALTN GSI EVQCM IPEHYNSHRGLIMYSW  
DCPMEQCKRNSTSIYFKMENDLPQKIQCTLSNPLFNTTSSIILTTCIPSSGHS  
RHR YALIP IPLAVITTCIVLYMNGILKCDRKPDR TNSN (SEQ ID NO: 166) Human  
CD48 >NM\_001778.4 *Homo sapiens* CD48 molecule (CD48), transcript variant 1,  
mRNA CTTTTTCTAGCCAGGCTCTCAACTGTCTCCTGCGTTGCTGGGAAGTTCTGGAA  
GGAAGCATGTGCTCCAGAGGTTGGGATTCGTGTCTGGCTCTGGAATTGCTACT  
GCTGCCTCTGTCACTCCTGGTGACCAGCATTCAAGGTC ACTTGGTACATATGA  
CCGTGGTCTCCGGCAGCAACGTGACTCTGAACATCTCTGAGAGCCTGCCTGAG  
AACTACAAACA ACTAACCTGGTTTTTATACTTTTCGACCAGAAGATTGTAGAATG  
GGATTCCAGAAAATCTAAGTACTTTGAATCCAAATTTAAAGGCAGGGTCAGAC  
TTGATCCTCAGAGTGGCGCACTGTACATCTCTAAGGTCCAGAAAGAGGACAAC  
AGCACCTACATCATGAGGGTGTTGAAAAAGACTGGGAATGAGCAAGAATGGAA  
GATCAAGCTGCAAGTGCTTGACCCTGTACCCAAGCCTGTCATCAAAATTGAGA  
AGATAGAAGACATGGATGACA ACTGTTATCTGAAACTGTCATGTGTGATACCT  
GGCGAGTCTGTAAACTACACCTGGTATGGGGACAAAAGGCCCTTCCCAAAGGA  
GCTCCAGAACAGTGTGCTTGAAACCACCCTTATGCCACATAATTACTCCAGGT  
GTTATACTTGCCAAGTCAGCAATTCTGTGAGCAGCAAGAATGGCACGGTCTGC  
CTCAGTCCACCCTGTACCCTGGCCCGGTCCTTTGGAGTAGAATGGATTGCAAG  
TTGGCTAGTGGTCACGGTGCCCACTTCTTGGCCTGTTACTTACCTGAGATG  
AGCTCTTTTAACTCAAGCGAAACTTCAAGGCCAGAAGATCTTGCCTGTTGGTG  
ATCATGCTCCTCACCAGGACAGAGACTGTATAGGCTGACCAGAAGCATGCTGC  
TGAATTATCAACGAGGATTTTCAAGTTAACTTTTAAATACTGGTTATTATTTA  
ATTTTATATCCCTTTGTTGTTTTCTAGTACACAGAGATATAGAGATACACATG  
CTTTTTTCCCACCCAAAATTGTGACAACATTATGTGAATGTTTTATTATTTTT  
TAAAATAAACATTTGATATAATTGTCAATTA ACTGAA (SEQ ID NO: 167)  
>NP\_001769.2 CD48 antigen isoform 1 precursor [*Homo sapiens*]  
MCSRGWDSCLALELLLLPLSLLVTSIQGHLVHMTVVSGSNVTLNISESLPENY  
KQLTW FYTFDQKIVEWDSR KSKYFESKFKGRVRLDPQSGALYISKVQKEDNST  
YIMRVLKKTGNEQEWKIKLQVLDPVPKPVIKIEKIEDMDDNCYLKLSCVIPGE  
SVNYTWYGD KRPFPKELQNSVLETTLM PHNYSRCYTCQVSNSVSSKNGTVCLS  
PPCTLARSFGVEWIASWL VVTVP TILGLLLT (SEQ ID NO: 168) Mouse CD48  
>NM\_007649.5 *Mus musculus* CD48 antigen (Cd48), transcript variant 1, mRNA  
ATACGACTTCCGGTTTTTGGGTTTTGCTTCCTGATTGAAGGGCAGGCGCCCTGA  
CTTCTCTTACAGTTGTCTCCAGTGTCTGGGGAAGCTTCTCTAAGTATTATGT



GGTCTAGCTTAGATACTAGTGTGCTGTACTTTGATGATAAGTAAATA  
CTTCATTACCTAAAATCATTTTTTCATAGAACTGTTTCAGAAAACCTGACTCTA  
ACTGGTTTATATACAAAAGAAAACCTTACTGTATCATATAACAGAATGATCCAG  
GGGAGATTAAAGCTTTGGGCAAGGGCTATTTACCAGGGCTTAAATGTTGTGTCT  
AGAATTAAGTATGGGCATAAACTGGCTTCTGAATCCCTTTCCAGAGTGTTGGA  
TCCATTTCCCTGGTCTTGGCCTCACTCTCATGCAGGCTTTCCTCTTGTGTTGG  
CAAGATGGCTGCCAACTCTTGGCAATTCATACATCCTTGTTTCTGTCTGGTAG  
AGAGTTTGCTTCTCAAATGGAGCAAACAAATTTGATTATTTTTTTCATTGTTAA  
ATAGGCAACATGACCAGAAAGGATGGAATGGCTTAAGTAAACTAAGGGTTCAC  
TTCTAGAGCTGAGAAGCAGGGTCAAAGCACATACTGGGCAATTCAGAGCATG  
GTTAGAAGAGGAAAGGGGAGTCTCAAAGCTGGAGAGTTTACCAACAAATATTG  
ACTGCAGTGATTAACCAAGACATTTTTGTAACTAAAAAGTGAAATATGGGAT  
GGATTCTAGAAATGGGGTATCTCTGTCCATACTTCTAGAATCCACTCTATCAG  
CATAGTCCAGAAGAATACCTGGCAGTAGAAGAAATGAATATTCAAGAGGAAGA  
TAAATGCGAGAGGGCAATCCTTTACTATTCTCATATTTATTTATCTCTCATTC  
TGTATAGAATTCTTGCCGCCATCCCAGGTCTAGCCTTAGGAGCAAATGTAGTA  
GATAGTCGAATAATAAATACTTAATGTTTTGGACATATTTTGTCTACTTTTG  
AGAATTATTTTTAATATGTAAATTCTCTCAAAGGGTCAGGCACCTAGTTATT  
ATTTTTTAATGATTATGTGAAAGTTGAATATAATATACCACTAAAAGTGACAG  
TTGAAAGTGGTGGCATAGGACGGTAGGGTAGAAATTTGGGAGGGGAAAAAAGAA  
ATTGGGAGGGTACAGGCAACAGGAGAAAGGAATCAAACCACAGAAAAATACAA  
AGGGAAACTTCTGCTTCACTATTCAGACAAAGACAGCCCTAATGACATCACCA  
ACAGTCAAAGCAATTAGAGACCATACCTAATATTGTTTAAATTCTAGATGTAG  
GCTAACAATGAAAAGTATTTGCCAAACTGAATAAAACTGTCATGGTTACCTTG  
AAAGGACAATGGTTATTGTTAAATATAGTGATCATTCATGTCTAAAAGATTCA  
TTATTTATCTCTAAAGATTTCTAAAGACCACCATCTAGAAAAGATTCAATTATG  
AAGGCTGTATTTAAATATCAAAGTTGTGGACTTCATGATAATCTTAAATAAAG  
CAAATCCAAATTCTCCTGTTGCCTAGACAGATTCTAAGATGTAATTTACACTT  
TTAAGCTAATTAGTGAGTATTTTATGATTTTAGCCTTAAACACCATGTATGCC  
AAATAATGCACTTGTTTTGTGAATTACAGAAATGGTAAGTGCCACATTTCTG  
TGAATTATAAAATTTGTGAGTTTCTTTTAACCCTTTTCAGGAGTGAAAAATA  
AAAACGACCATTTCCCTGGTTGTGCTTAAGTATATGCAAGAAGGGTAAACTCTC  
ATTTTTATTATGTTTGCTTAAAGATCTTTTTATACCTGGATTCATGAAATGTT  
TCCACAAATATATTAGTGTAACAAACTTGAAAGGCAGTTTACAAGAAAGCACT  
CTACTATCAGATCAATCAAAGATTCTGTGAGTGAATTTATTGGTTTGCATGGT  
GAAGCAAGCTTAGCATCAATTAAGGTAATAATTTCTTTTCTGAATGGTAA  
AGACAATCAAATATTACTTTCTGGAAAACCTCCAATAACCAAATTCTCAATGA  
TTAGTGTATGTGAGCAGGAAAACATTTTTACAGTTGTAGTATGGGGAAATATA  
AATCCAATTTTAAGAGAGAAAATTATGACTGGGTGTGGAAGGGACAGTATAGT  
CAGATACCAATTGTCATGGTGGTTTTTACTGGGAACTTCATGAAAGACTTTTGT  
AGCAAACCACTGCAGTATTGCAAAGCCTCCAGAACATTTGGAACCTTGTCTCTT  
TTTCCTTGTGTGTGTTTGTGTTTTTGGTCTCTCATTCAAATATTGATGAGAA  
CTATTTACTCTGTCCTTTCTTCTCTATATATTCTTCCTCTACAGAGTGTAGGG  
TTTTTTCAGGAATTTGGAGCCATCTGAAGTCCTCCCAAAAATTCTCTGACGTC  
TTCTGATGCTCCTGTTATACCCTCAGGGGTAATGCTTGTGAAATTCCATTCAT  
TCATTTTCTTTCTCTGGACATCTTTACTTACCAAAGCACTTTCATTGTCATCT  
TTTTAACATCATTCTTAATTCGTGATAGTTTTGGGACTCTCCCTAGTGATGT  
TTCTCCCCCTCTACTCTTTTGCACCTATGATTCTGATTGTTACTAAGAAAGCA  
GATGAAAAACAGATCCACAGAATAAACGATCAGAATTCCAGTAAATTCTATTT  
TAAATACAGATACTTTTTACAAGTTGCTGCTTTGGAAGCAAATGCTTCTTAA

[illegible]



ATGGTGAACACCCCTGTCTTAGTAATAAATAGTAAGAAATAGGCCAGGCAGGCCTGGTGGCA  
 CGCACCTGTAGTCCCAGCTACTTGGGAGGGCTGAGGCAGAAGAATCGCTTGAAC  
 CCGGGAGGCAGAGGTTGCAGTGAGCCAAGATCGCACCACTGCACTCCACCCTG  
 GGCAACAGAGCGAGACTCCGTCTCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA  
 GTTAAGTATTTGAACATAGGGGTGGCTCATAGAATTCCCAGGACACCCGATGG  
 AGTAGGCTTGCAAAACACAACATGTGGCAACTCCAGTGGGAAACGAGGCAGGA  
 AACACTCGTTTCCTGCGAGAAAGCAACAATTTGGGCTTCGATACCCTCCCTAGA  
 ACACAGGGCAGTGAATCTGAGCAGCATCAGTACCCACGTTTCGGATGAGTCCT  
 GAGCCCCCTATTTTTATTCACTGACTTATTCCAAAATCAGTGTCTCTTAAATAT  
 ATCTGGAAGGCAGCAGCTTGTATCTCCCCCTTCAGCTTCCATAGTGGCAGTCA  
 GGGTACAACCTTACTTTCCAAACAGAACACACTGCGACATTCCCTCCAGGCTCG  
 TTGAAGAACTTCAACTGACAAATGTCCCTCCTCGACCAGATGATAGTTTTCTT  
 AAAGGCAGGGTTTAATATACCCTTTTATAAATGTTTCAAGGCCCTGTGTAATA  
 CCTGAGTTTATTCCAGATGTAATAAATATATCCAAGATTGTTTTAAAATAAA  
 TTGCTGAAAAAACAATAAATACAGTTAGTATCTATATCAATATTCTCAGTTG  
 GCAGTTTTGCAATAATGGCCGATAGTTCATTTTTAGTAACACTATTGACATTG  
 CATTTGGATATTAGGGTTTACTAATCATCCGCATGTATACATTGCATATTTTT  
 CTAGACTTTAACTTTATTCAAATCTATTGATTTTTTAAACCTGCAACTTATGTC  
 TAGACACAGGTATACCTTTACAAGAATACTACCATTTTTTTTTGGTAACATACTAC  
 CTCCAAAATTTCAAGTAAGAAGTTGATTTTTTGTCCATTTTTTAAATGGAAAAC  
 TGTAAATCAAATGCCACAAAATTATACTGTGTATCATTTGACCTATAGAAACC  
 AATATTATTACAGGAAGAAAGCAGAGCCAATCTTCTACCTGTGGTCAAATAAG  
 TGGAGGCCCTTTCTAGACTAAGTTCTCATGAGTTTAAAATACCAAGCATAAGT  
 TCTCCAAATTCCTGAAAAGGAAGCCTTGTGTTGTATTGCCCAGCCATATTTGT  
 AAGACATAAAAATAAAACTTGAGAAGAAGCTATGATAACTTACTTTCTTCATT  
 CTTCAAATTTTACATAATCTCAACTGATTTTTATGTTTTTATGAAAATGCATTC  
 TTAAGATATATCCTTATTCAATCATGTATTCATTACATCCTTTATGCCAGGTA  
 TCCAAAAGTACTTACAGTGACTAAGACCATTATTCTTTGATCAGCTGCCTGAG  
 TAAGACTTTGAGCTCTCCAATATACTCTCAGTGATACTAAGTTTTCTGAGTAA  
 CAGCTTTGGATGTGGCTTCAGTTGAGCTGATTTATCCCACACTTTATTTTTAT  
 CGTATAATGGTCCTCAGAAGCAAATTTTGATTTTAGCTCACATAAAAAATGTA  
 CAAAGAAATGTAATGGCTCAGTAGCTTCTAGAGATAGAGATTACTCTTCTAAC  
 CTTTCTGTAATTTTGTATGTCTATTTTTATAATTCTTTCAATGTCTAATGAATA  
 GCTATCTTTTTTTGAGACGGAGTCTCGCTCTGTGCGCCAGGCTGGAGTGCAGT  
 GGTGCGACCTCGGCTCACCGCAAGCTGCGTCTTCCAGGTTACAGCCATTCTCC  
 TGCCTCAGCCTCCCGAGTAGCTGGGACTTCAGGCGCCCACCACCATGCCCAGC  
 TAATTTTTTTGTATTTTTAGTAGAGACGGGGTTTCACCGTGTTAGCCAGGGTG  
 GTCTCGATCTCCTGACCTCGTGATCCGCCCCGCTCGGCCTCCCAACGTGCTGG  
 GATTACAGGAGTGAGCCACCGCGCCCCGGCCTCCTTAGTTTCTTAAGGTGGAAG  
 CCTAGATTATTGATTTTATATGTTGTTTTCTTTTCCAATAGTGGCACTTAATG  
 CTATAAATTTCACTTTGTTCCACAAGTTTTGGTAAGCTCTATTTTTATTTTCA  
 TTTAGTCCAAAATATTTTAAAATTTCTTTTGATATTTCTTCTTTGAGCCATGA  
 ATTATTTACAATGTGTTGTTTAATCTCTATATATTTTGGGATTTTTCTACTTT  
 ATATCTCTTACAGATTTCTAACTTAATTTTCATCATGTTTTTAAAACATTCTTT  
 GTATAATTTCTATTCTTTTAAATTTTTTCAGGTGTATTTTATGGCCCAGAATAT  
 GGTCTATCTTGTAGAATGTTTCATGTGATCTTAAGAAGAATGTTTCATTCTGCT  
 GTTGAGTGTAATATTCTACAAATGTCCATTAGATTAAACTGATTGATACCACC  
 GTTCAGATTATCTATATCCTTTCTGATTTTCCCTCTTCTTGATCTATCACATA  
 CTGACAGATCAAGTGATCAAGTCTCGTTAAAGACTGCAAGTAAAATAGTGGAT  
 TTTTCTATTTCTCCTTGCAGTTTTTGTTAGTTTTTGTCTCATGTATCTTGATAC

TCTTGTAGTACATATAATGATGTTTCTTGGAGGTTTCTTGGAGGTTGACC  
CCTTTACCACATGTAATGTCCCTTTTATTCTTGATAATCTTTCTTGTTCTGTC  
TGCTTTTTCTGATATTAACATAACTTTCAGTTTTTTAAAAAATTAACATTAGC  
ATCTCACATCTTTATCCTTTTAATTTTAAATTATCTAAATATTTATATTTAAT  
GTGCCTTTCTTATAGACAATGTATAGTTGCGTCTATTTGTAATTTCCCCACTT  
TTCTTACTTAAAAATGTTGTAGATATATAGGAGTTGTATATATTTGGGGGGTA  
CATGTGATGTTTTGATACCTGTATACAATATGTAATGATCATATTGGGTAATC  
GTGATATCTGTCACCTCTAACATTCATCTTTTTTGTGTGTTTAAACCCACCAC  
TTCTAATTGGTACATTTAGATTATTCAAATTTAAGTGATTATTGATATAGTTG  
GATTAATATCTACTATGTTTGTAACTTTTCTATCCTTGCCTCGTTCTTTCTT  
TTTTATCCTCCTTTTTCTGTGTTCTCTGATTTTAACTGGGGTTTTTACATGAT  
TTAATTTTCTCTCGTGGCATATCTTTCATTGATCAACCTAGGTTTTTCTCCTT  
TTCCCCTCTTTTTTTTTGGTATTTATTCTATTTAGTGTTATCTGAGCTACCTGA  
GTTGGTGTCTATCACTAATTTTGGCAAGTTCCCAGACGTTATTACTTCTAACA  
TTCTTTTGCTCCATTCTTTCTTCTTCTTCAATTATTCCATAGTCTTGAATATT  
CTGGGTTTTTCCCCTCTTTGAATTTTAGTTTGAAAAGTTTCTATTGGCCTAG  
CTTCAAAGTCATTCACTTCTTCTTCCGTTTCCAAGTCAACTGATAATTGCAT  
CAAAGATATCCTTCTTCTTCTATTACTATGTTTTTTATTGCTACCATTTCTTTT  
TTATTCCTTCTTAGTGTTTCCATCTTTCTTCTTACATTATCCATCTGTTGTCT  
ATTTTTTTTCATGAGAGCTCTTAACATATTAATGATAAGTTCCATGTCTGATAA  
TTCTGACACGTGTCATGTCTCTATCTGGTTCCAATGATTGCTTTATCTCTTCA  
GACCATGACTTTTCTTGCCTTTTGACGTTCTTTGACATTTTTTTTTGAATTTT  
TGTTGCAAGCCAGATCTGGTGTGTTATGTAATAGGAACAGGTAAATAAGTCTT  
TAGCTTGCAGACTTATCTTAATCTGACTAACTATTAGACTGTGTTTAAAGTCT  
GTTATAACCATAGGTGCTAAATTTCTTCAAATTCCTCTAGTGCTTTTGTTTTG  
TTTGTTTCATGTGTTTTTCCCCTTCTTGAGTTCAGGCTTCCCTAAGTGCTCCTC  
TTCAGAGAGACTTTCTGTCTTTCAGCTCTTTCCTCTGCAATTCACTGTTACTA  
TACTGGAGCCCTGTTGGTGTAGTACTAAGCTGTGGGAAAGGAGAGTGCTCTGT  
AATCTTACAGTGAAATCTCAGTOTTTTAGTGGGTCTGTGTCTGGGACATTAC  
AGAGCTTCTCCAGTGGTATTGCTTCCCTCATCCTCAACTCTCTTTCCTGGCTGC  
AGCATTCCCAATGTATTTCTTTGAAGGCCTGCCCCCTGTTGACTGTTATTTTC  
CCTCTTTCCTTAAGTGGGACAGGGAGACTTCAGGGGCTGGGATGAGGTTTGGG  
AATTGTGCTTGGCAGAGTCCTTTCCATCTTTGTTACCAAGAAGGTTTCATGGCT  
TATTTCTCAATGGATGTCCCTCTCTATCTGTTGCCAGAGCCACGAGGAAATTT  
TTCTTGGATCCTCATAATGAGAACCTTGGAGTTTCCTACTGGAAAAGCCCTTG  
AATGTGTGGAGTGCCTCAAGAGCACAGCCCCCATGGGTTTTCTTGCTCACACCA  
GTCCACAAACAGATGCCAGCAATTCACCCAACCTTACCATATAAAGGCTCATA  
TAGTTTATGGCTCCAGTGCTTTGACTCCAGATAAATGGCTATTGGTTGCGTAT  
CTCTCTGGATGTATCTGTATCTCCAGATTTTGGGGTGGCAGTTTGCTCAGGAC  
CTTGGTTCTCTAATAGGTCTAATAAGAAAAGTCATTGATTTTCAGCTTTCCAA  
CTTTCAGCTTTGTCTTGTTATAAGCATGGCAGCAACATCTTCCATGCCTTAA  
CATGATGACACTAAAGGCAGAAGTCGATCTCCATGTATAAACATTTTAACACA  
TATGTTTTTTGTTATCGTGGTTTCTGACCTGTCTCTTTGCCCTGACTTTCTGA  
TACTGCACTAGGGTTCCTGTTGCTGGACTCCATTCCATATGACTTGCTCTCGT  
CTAGGCTGCTCTTTGGCTCATCTTTATAAATCATGATCCAAAATGAAGCACAT  
ATTTATTTTTTAAATAAATATGAAATGAAGTATAGACATCAAACCTGAAGATGA  
GTAGATCATACTGAGTTTCACTGTCTGTGCTTGGATCAACATCAGGCCTTATA  
CAAATATTCAAGTCCAGAGGCCAAAAGGTAATAAGGAAAATTTGTAGCACAAGC  
CACAAGGAGATAACATGTCAAGTCTATGCGATTGGAAATAAACTAAAGATGAA  
CTGCTGGGGATGCTCACTCATCACAGAGCTCAGTCTAAAGCACCAGATTTAC

[illegible]

GATGAAGGACCTAGASUAGTTGACTTAAAGACTTTGAGTGCGAAGAT  
ATTACTCATTAACCAGATAACAAGTGAATATGCTGTAGCATCAACATAATTCA  
AAAGAGTAAAGAAATGGCTAGGAATGAGGTAGTTGTGTAATTATTTCTTCTCT  
TACTAGTTTCAAATAAATTCATCTCTAATTCTATAGAGAATTCTTGCCTCCCA  
TTCAGGACTGGCCTTCTATACAGTGAGATGGTCCAGTAAGAAATAATTTTTAT  
TAGTGTTTTTTCTATTTTGAGAATTATTTTAATATATATTTTAATATATAAAC  
TTGTGAGTTAAATTTTTTTTTTTTGCAAAATTAGCACATGAAAAGAGATTGATGG  
TTTTAAGTAGTAGAACACAGTAGTGTAGGAATCTGAGAGCAGAGAGTTTGGGA  
GGGGGTGAAGAGAAAACAACATCACCAAATAGTGATATATAAGAGAAAATCTG  
TGCTTCAGAGTTTGATCAGGGCCATCTCTCCCAACTCTGCTGGAAGTGAAGAGA  
ATGCACCTGATGTTGTCTCCATTTTAGATAGAGAAAAAAAACCCGAATATT  
TATAAACTAAATAAACTATAGTTACCTCAAACTATGGGGATCACTATAAC  
ATAGAATAGAATAGAATAGAATAGAATAGAATAGAATAGAATAGAATAG (SEQ ID  
NO: 173) >NP\_848802.2 CD226 antigen isoform a precursor [*Mus musculus*]  
MAYVTWLLAILHVHKALCEETLWDTTVRLSETMTLECVYPLTHNLTQVEWTKN  
TGTKTVSIAVYNPNHNMHIESNYLHRVHFLNSTVGFRNMSLSFYNAASEADIGI  
YSCLFHAFPNGPWEEKKIKVWSDSFEIAAPSDSYLSAEPGQDVT LTCQLPRTW  
PVQQVIWEKVQPHQVDILASCNLSQETRYTSKYLRQTRSNCSQGSMSIL IIP  
NAMAADSGLYRCRSEAITGKNKSFVIRLIITDGGTNKHFILPIVGGLVSLLLV  
ILIIIIFILYNRKRRRQVRIPLKEPRDKQSKVATNCRSPTSPIQSTDDEKEDI  
YVNYPTFSRRPKPRL (SEQ ID NO: 174) Human DR3 >NM\_003790.3 *Homo  
sapiens* TNF receptor superfamily member 25 (TNFRSF25), transcript variant 2,  
mRNA

GAAGGCGGAACCACGACGGGCAGAGAGCACGGAGCCGGGAAGCCCCTGGGCGC  
CCGTCGGAGGGCTATGGAGCAGCGGCCGCGGGGCTGCGCGGCGGTGGCGGGCGG  
CGCTCCTCCTGGTGCTGCTGGGGGGCCCGGGCCAGGGCGGCACTCGTAGCCCC  
AGGTGTGACTGTGCCGGTGACTTCCACAAGAAGATTGGTCTGTTTTGTTGCAG  
AGGCTGCCCAGCGGGGCACTACCTGAAGGCCCTTGACGAGCCCTGCGGGCA  
ACTCCACCTGCCTTGTGTGTCCCCAAGACACCTTCTTGCCCTGGGAGAACCAC  
CATAATTCTGAATGTGCCCCGCTGCCAGGCCTGTGATGAGCAGGCCTCCCAGGT  
GGCGCTGGAGAACTGTTTCAGCAGTGGCCGACACCCGCTGTGGCTGTAAGCCAG  
GCTGGTTTGTGGAGTGCCAGGTCAGCCAATGTGTTCAGCAGTTCACCCTTCTAC  
TGCCAACCATGCCTAGACTGCGGGGGCCCTGCACCGCCACACACGGCTACTCTG  
TTCCCGCAGAGATACTGACTGTGGGACCTGCCTGCCTGGCTTCTATGAACATG  
GCGATGGCTGCGTGTCCTGCCCCACGAGCACCCCTGGGGAGCTGTCCAGAGCGC  
TGTGCCGCTGTCTGTGGCTGGAGGCAGATGTTCTGGGTCCAGGTGCTCCTGGC  
TGGCCTTGTGGTCCCCCTCCTGCTTGGGGCCACCCTGACCTACACATAACCGCC  
ACTGCTGGCCTCACAAGCCCCTGGTTACTGCAGATGAAGCTGGGATGGAGGCT  
CTGACCCCAACACCGGCCACCCATCTGTACCCCTTGGACAGCGCCACACCCT  
TCTAGCACCTCCTGACAGCAGTGAGAAGATCTGCACCGTCCAGTTGGTGGGTA  
ACAGCTGGACCCCTGGCTACCCCGAGACCCAGGAGGCGCTCTGCCCCGAGGTG  
ACATGGTCCTGGGACCAGTTGCCAGCAGAGCTCTTGGCCCCGCTGCTGCGCC  
CACACTCTCGCCAGAGTCCCCAGCCGGCTCGCCAGCCATGATGCTGCAGCCGG  
GCCCCGAGCTCTACGACGTGATGGACGCGGTCCAGCGCGGCGCTGGAAGGAG  
TTCGTGCGCACGCTGGGGCTGCGCGAGGCAGAGATCGAAGCCGTGGAGGTGGA  
GATCGGCCGCTTCCGAGACCAGCAGTACGAGATGCTCAAGCGCTGGCGCCAGC  
AGCAGCCCGCGGGCCTCGGAGCCGTTTACGCGGCCCTGGAGCGCATGGGGCTG  
GACGGCTGCGTGGAAGACTTGCGCAGCCGCCTGCAGCGCGGGCCCGTGACACGG  
CGCCCACTTGCCACCTAGGCGCTCTGGTGGCCCTTGACAGAAGCCCTAAGTACG  
GTTACTTATGCGTGTAGACATTTTATGTCACCTATTAAGCCGCTGGCACGGCC

CTGCTAGCAGCAGCAGCCGCGCTTCGCCCCCTATCGCCCTCCAGCC  
AAGGCGAAGAAGCACGAACGAATGTCGAGAGGGGGGTGAAGACATTTCTCAACT  
TCTCGGCCGGAGTTTGGCTGAGATCGCGGTATTAAATCTGTGAAAGAAAACAA  
AACAAAACAAAAACGGCTTCTTGGCGTTTCTGCGGGGGCTGGGGTGTAAAGTGG  
ACTGGACTTTTCTCGAGGGATTCTGAAGGGGACGGGAATCTTGTCACCCCGGGA  
TCTGGCACCCATGGTGGAGTCCAGTGTGGCCTTAGCTCCCAAGCCTGCCCTC  
CCGAGTCCACTCTGGCTCAATTACCCCGAGAAGGAGAGAGCAAGTCGCGGCCA  
CAGCGAGTGAGTGAACCGGAGCCCAGATGAGAGCGCTTTAATGGGGCTGCGAG  
GTGGCGGAGACAGGGTTCGGGATGGGGTGCAGCAGTTGGAGACACAGGGTCAGG  
GCCCCTCATCCTCTATTCACTCCACCGGGGCAGTGAAAGGGTCCCGGCAGCGA  
GTGGGTC (SEQ ID NO: 175)>NP\_003781.1 tumor necrosis factor receptor  
superfamily member 25 isoform 2 precursor [*Homo sapiens*]  
MEQRPRGCAAVAAALLVLLGARAQGGTRSPRCDGDFHKKIGLFCRCRCPA  
GHYLKAPCTEPCGNSTCLVCPQDTFLAWENHHNSECARCQACDEQASQVALEN  
CSAVADTRCGCKPGWFVECVSQVSSSPFYCQPCLDCGALHRHTRLLCSRRD  
TDCGTCLPGFYEHDGCVSCPTSTLGSCPERCAAVCGWRQMFVWQVLLAGLVV  
PLLLGATLTYTYRHCWPHKPLVTADAEAGMEALTPPPATHLSPLDSAHTLLAPP  
DSSEKICTVQLVGNSWTPGYPETQEALCPQVTWSWDQLPSRALGPAAAPTLS  
P  
ESPAGSPAMMLQPGPQLYDVM DAVPARRWKEFVRTLGLREAEIEAVEVEIGRF  
RDQQYEM LKRWRQQPAGLGAVYAALERMGLDGCVEDLRSRLQRGP (SEQ ID NO:  
176) Mouse DR3 >NM\_033042.4 *Mus musculus* tumor necrosis factor receptor  
superfamily, member 25 (Tnfrsf25), transcript variant 2, mRNA  
CTGCGTGGAGGGGAAATGGGCCAGAGGCTGCTGGCAGGGGGCCTCCTCTGCTG  
TACACAAGCTGGTTTTGTAGACAGTGAGAGGGAAGCTGATCCCAGTCCCCTAA  
CCCTGTTCTGCCCAGGAGCCTGAGAACTGAGCTTACTCGGGCAAATGCTAGGG  
CTTCAGAAATGGAGGAGCTGCCTAGGAGGGAGAGGTCACCTCCTGGGGCAGCC  
ACACCAGGGTCAACTGCACGTGTTCTCCAGCCTCTGTTCTTACCCTGCTGCT  
GCTGCTGCTGCTGCTGCTTGGTGGCCAGGGCCAGGGCGGCATGTCTGGCAGGT  
GTGACTGTGCCAGTGAGTCCCAGAAGAGGTATGGCCCGTTTTTGTGTCAGGGGC  
TGCCCAAAGGGACACTACATGAAGGCCCCCTGCGCAGAACCCTGTGGCAACTC  
CACCTGCCTTCCCTGTCCCTCGGACACCTTCTTGACCAGAGACAACCACTTTA  
AGACTGACTGTACCCGCTGCCAAGTCTGTGATGAAGAGGCCCTTCAAGTGACC  
CTTGAGAACTGCTCGGCAAAGTCGGACACCCACTGTGGCTGCCAGTCAGGCTG  
GTGTGTTGACTGCTCCACCGAGCCATGTGGGAAAAGCTCACCTTTCTCTTGTG  
TCCCATGCGGGGCTACAACACCAGTCCATGAGGCTCCAACCCCCCTGTTTTGG  
GTCCAGGTGCTTCTAGGAGTCGCGTTTCTTTTTGGGGCTATCCTGATCTGTGC  
ATATTGTCGATGGCAGCCTTGTAAGGCCGTGGTCACTGCAGACACAGCTGGGA  
CGGAGACCCTGGCCTCACCACAGACTGCCCATCTCTCAGCCTCAGACAGCGCC  
CACACCCTCCTGGCACCTCCAAGCAGTACTGGGAAAATCTGTACCACTGTOCA  
GTTGGTAGGCAACAACCTGGACCCCTGGCTTATCCCAGACTCAGGAGGTGGTCT  
GCGGACAGGCCTCACAACCCTGGGATCAGCTGCCAAACAGAACTCTTGGAAC  
CCTCTGGCATCTCCGCTCTCGCCAGCGCCCCCTGCGGGCTCTCCGGCTGCTGT  
GCTCCAGCCTGGCCCGCAGCTCTACGATGTGATGGATGCGGTCCCAGCACGAA  
GGTGGAAGGAGTTCGTGCGCACGCTGGGGCTGCGGGAAGCGGAAATTGAAGCC  
GTGGAGGTGGAATCTGCCGCTTCCGAGACCAGCAGTATGAGATGCTCAAGCG  
CTGGCGTCAGCAGCAGCCTGCAGGCCTCGGTGCCATCTATGCGGCTCTGGAGC  
GCATGGGTCTGGAAGGCTGTGCCGAGGACCTGCGCAGCCGCCTGCAGCGTGGC  
CCGTGATGGAAGGTCCATCAGCCACTTTGACACCCTAGTGACCCTTGAAGGAG  
CCTTAAGTATTGTTACTTATGCGTGTAGACATTTTATGTCAATTACTAACCC  
CTGCCGTGGTCCTGCGTAGCAGGGCTGGCTGCCTCACTTTTGCTTATCTGCAG

[illegible]

(SEQ ID NO: 181)>NP\_000630.1 tumor necrosis factor ligand superfamily member 6 isoform 1 [*Homo sapiens*]

LVNFEESEQTFFGLYKL (SEQ ID NO: 182) Mouse FasL>NM\_010177.4 *Mus musculus* Fas ligand (TNF superfamily, member 6) (FasL), transcript variant 1, mRNA TGAGGCTTCTCAGCTTCAGATGCAAGTGAGTGGGTGTCTCACAGAGAAGCAAA GAGAAGAGAACAGGAGAAAGGTGTTTCCCTTGACTGCGGAAACTTTATAAAGA AAACCTTAGCTTCTCTGGAGCAGTCAGCGTCAGAGTTCTGTCCTTGACACCTGA GTCTCCTCCACAAGGCTGTGAGAAGGAAACCCTTTCCCTGGGGCTGGGTGCCAT GCAGCAGCCCATGAATTACCCATGTCCCCAGATCTTCTGGGTAGACAGCAGTG CCACTTCATCTTGGGCTCCTCCAGGGTCAGTTTTTCCCTGTCCATCTTGTGGG CCTAGAGGGCCGGACCAAAGGAGACCGCCACCTCCACCACCACCTGTGTCACC ACTACCACCGCCATCACAACCACTCCCACTGCCGCCACTGACCCCTCTAAAGA AGAAGGACCACAACACAAATCTGTGGCTACCGGTGGTATTTTTTCATGGTTCTG GTGGCTCTGGTTGGAATGGGATTAGGAATGTATCAGCTCTTCCACCTGCAGAA GGAAGTGGCAGAACTCCGTGAGTTCACCAACCAAAGCCTTAAAGTATCATCTT TTGAAAAGCAAATAGCCAACCCCAGTACACCCTCTGAAAAAAAAAGAGCCGAGG AGTGTGGCCCATTTAACAGGGAACCCCCACTCAAGGTCCATCCCTCTGGAATG GGAAGACACATATGGAACCGCTCTGATCTCTGGAGTGAAGTATAAGAAAGGTG

CGCTTGTGACGAAACTGGGTTGTACTTTCGTGTATTCCAAAGTATACTTC  
CGGGGTCAGTCTTGCAACAACCAGCCCCTAAACCACAAGGTCTATATGAGGAA  
CTCTAAGTATCCTGAGGATCTGGTGCTAATGGAGGAGAAGAGGTTGAACTACT  
GCACTACTGGACAGATATGGGCCACAGCAGCTACCTGGGGGCAGTATTCAAT  
CTTACCAGTGCTGACCATTTATATGTCAACATATCTCAACTCTCTCTGATCAA  
TTTTGAGGAATCTAAGACCTTTTTTCGGCTTGTATAAGCTTTAAAAGAAAAAGC  
ATTTTAAAATGATCTACTATTCTTTATCATGGGCACCAGGAATATTGTCTTGA  
ATGAGAGTCTTCTTAAGACCTATTGAGATTAATTAAGACTACATGAGCCACAA  
AGACCTCATGACCGCAAGGTCCAACAGGTCAGCTATCCTTCATTTTCTCGAGG  
TCCATGGAGTGGTCCTTAATGCCTGCATCATGAGCCAGATGGAAGGAGGTCTG  
TGACTGAGGGACATAAAGCTTTGGGCTGCTGTGTGACAATGCAGAGGCACAGA  
GAAAGAAGTGTCTGATGTTAAATGGCCAAGAGAATTTTAACCATTGAAGAAGA  
CACCTTTACACTCACTTCCAGGGTGGGTCTACTTACTACCTCACAGAGGCCGT  
TTTTGAGACATAGTTGTGGTATGAATATACAAGGGTGAGAAAGGAGGCTCATT  
TGACTGATAAGCTAGAGACTGAAAAAAGACAGTGTCTCATTGGCACCATCTT  
TACTGTTACCTAATGTTTTCTGAGCCGACCTTTGATCCTAACGGAGAAGTAAG  
AGGGATGTTTGAGGCACAAATCATTCTCTACATAGCATGCATACCTCCAGTGC  
AATGATGTCTGTGTGTTTGTATGTATGAGAGCAAACAGATTCTAAGGAGTCAT  
ATAAATAAAATATGTACATTATGGAGTACATATTAGAAACCTGTTACATTTGA  
TGCTAGATATCTGAATGTTTCTTGGCAATAAACTCTAATAGTCTTCAAATCT  
TTTATTATCAGCTACTGATGCTGTTTTTCTTTAATACAACTAGTATTTATGCT  
CTGAACATCCTAATGAGGAAAAGACAAATAAAATTATGTTATAGAATACAGAA  
ATGCCTTAAGGACATAGACTTTGGAAA (SEQ ID NO: 183) >NP\_034307.1 tumor  
necrosis factor ligand superfamily member 6 isoform 1 [*Mus musculus*]  
MQQPMNYPCPQIFWVDSSATSSWAPPGSVFPCPSCGPRGPDQRRPPPPPPV  
PLPPPSQPLPLPPLTPLKKKDHNTNLWLPVVFVFMVLVALVGMGLGMYQLFHLQ  
KELAEFREFTNQSLKVSSFEEKQIANPSTPSEKKEPRSVAHLTGNPHSRISPLE  
WEDTYGTALISGVKYYKKGGLVINETGLYFVYSKVYFRGQSCNNQPLNHKVYMR  
NSKYPEDLVLMEEKRLNYCTTGQIWAHSSYLGAVFNLTSADHLYVNISQLSLI  
NFEESKTFFGLYKL (SEQ ID NO: 184) Human TIM-1 >NM\_012206.3 *Homo  
sapiens* hepatitis A virus cellular (CD365) receptor 1 (HAVCR1), transcript  
variant 1, mRNA  
GACCAGGAGTCAGTTTGGCGGTTATGTGTGGGGAAGAAGCTGGGAAGTCAGGG  
GCTGTTTCTGTGGACAGCTTTCCTGTCTTTTGGGAAGGCACAGAGCTCTCAGC  
TGCAGGGAACAAACAGAGCTCTGAAGCCGTTATATGTGGTCTTCTCTCATTTT  
CAGCAGAGCAGGCTCATATGAATCAACCAACTGGGTGAAAAGATAAGTTGCAA  
TCTGAGATTTAAGACTTGATCAGATAACCATCTGGTGGAGGGTACCAACCAGCC  
TGTCTGCTCATTTTCTTCAGGCTGATCCCATATGCATCCTCAAGTGGTCAT  
CTTAAGCCTCATCCTACATCTGGCAGATTCTGTAGCTGGTTCTGTAAAGGTTG  
GTGGAGAGGCAGGTCCATCTGTCACACTACCCTGCCACTACAGTGGAGCTGTC  
ACATCCATGTGCTGGAATAGAGGCTCATGTTCTCTATTACATGCCAAAATGG  
CATTGTCTGGACCAATGGAACCCACGTCACCTATCGGAAGGACACACGCTATA  
AGCTATTGGGGGACCTTTCAAGAAGGGATGTCTCTTTGACCATAGAAAATACA  
GCTGTGTCTGACAGTGGCGTATATTGTTGCCGTGTTGAGCACCGTGGGTGGTT  
CAATGACATGAAAATCACCGTATCATTGGAGATTGTGCCACCCAAGGTCACGA  
CTACTCCAATTGTCACAACCTGTTCCAACCGTCACGACTGTTCTGAACGAGCACC  
ACTGTTCCAACGACAACGACTGTTCCAATGACGACTGTTCCAACGACAACCTGT  
TCCAACAACAATGAGCATTCCAACGACAACGACTGTTCTGACGACAATGACTG  
TTTCAACGACAACGAGCGTTCCAACGACAACGAGCATTCCAACAACAACAAGT  
GTTCCAGTGACAACAACCTGTCTCTACCTTTGTTCTCTCCAATGCCTTTGCCCAG



GCAGAACTACGACACTGCAGGGAGCAATAAGGAGAGAACCCACCAGCTCACCA  
CCCACCCTACGACACTGCAGGGAGCAATAAGGAGAGAACCCACCAGCTCACCA  
TTGTACTCTTACACAACAGATGGGAATGACACCGTGACAGAGTCTTCAGATGG  
CCTTTGGAATAACAATCAAACCTCAACTGTTCTAGAACATAGTCTACTGACGG  
CCAATACCACTAAAGGAATCTATGCTGGAGTCTGTATTTCTGTCTTGGTGCTT  
CTTGCTCTTTTGGGTGTCATCATTGCCAAAAAGTATTTCTTCAAAAAGGAGGT  
TCAACAACCTAAGTGTTTCATTTAGCAGCCTTCAAATTAAGCTTTGCAAAATG  
CAGTTGAAAAGGAAGTCCAAGCAGAAGACAATATCTACATTGAGAATAGTCTT  
TATGCCACGGACTAAGACCCAGTGGTGCTCTTTGAGAGTTTACGCCCATGAGT  
GCAGAAGACTGAACAGACATCAGCACATCAGACGTCTTTTAGACCCCAAGACA  
ATTTTTCTGTTTCAGTTTCATCTGGCATTCCAACATGTCAGTGATACTGGGTA  
GAGTAACTCTCTCACTCCAACTGTGTATAGTCAACCTCATCATTAAATGTAGT  
CCTAATTTTTTATGCTAAAACTGGCTCAATCCTTCTGATCATTGCAGTTTTCT  
CTCAAATATGAACACTTTATAATTGTATGTTCTTTTTTAGACCCCATAAATCCT  
GTATACATCAAAGAGAA (SEQ ID NO: 185) >NP\_036338.2 hepatitis A virus  
cellular receptor 1 isoform a precursor [*Homo sapiens*]

MHPQVVILSLILHLADSVAGSVKVGGEAGPSVTLPCHYSGAVTSMCWNRGSCS  
LFTCQNGIVWTNGTHVTYRKDTRYKLLGDLSSRDVSLTIENTAVSDSGVYCCR  
VEHRGWFNDMKITVSLEIVPPKVTTTPIVTTVPTVTTVRTSTTVPTTTTVPMT  
TVPTTTVPPTTMSIPTTTTTLTMTVSTTTSVPTTTSIPTTTSVPVTTTVSTFV  
PPMPLPRQNHEPVATSPSSPQPAETHPTTLQGAIIRREPTSSPLYSYTTDGNDT  
VTESSDGLWNNNQTLFLEHSLTANTTKGIYAGVCISVLVLLALLGVIIAKK  
YFFKKEVQQLSVSFSSLQIKALQNAVEKEVQAEDNIYIENSLYATD (SEQ ID NO: 186)  
Mouse TIM-1 >NM\_134248.2 *Mus musculus* hepatitis A virus cellular receptor 1  
(Havcr1), transcript variant 1, mRNA

GTCAGTACCATGAATCAGATTCAAGTCTTCATTTCAGGCCTCATACTGCTTCT  
CCCAGGCGCTGTGGATTCTTATGTGGAAGTAAAGGGGGTGGTGGGTCACCCTG  
TCACACTTCCATGTACTTACTCAACATATCGTGGAATCACAACGACATGTTGG  
GGCCGAGGGCAATGCCCATCTTCTGCTTGTCAAAATACACTTATTTGGACCAA  
TGGACATCGTGTCACCTATCAGAAGAGCAGTCGGTACAACCTTAAAGGGGCATA  
TTTCAGAAGGAGATGTGTCCTTGACGATAGAGAACTCTGTTGAGAGTGACAGT  
GOTCTGTATTGTTGTCGAGTGGAGATTOCTGOATGGTTTAATGATCAGAAAGT  
GACCTTTTTCATTGCAAGTTAAACCAGAGATTCCCACACGTCCTCCAAGAAGAC  
CCACAACCTACAAGGCCACAGCTACAGGAAGACCCACGACTATTTCAACAAGA  
TCCACACATGTACCAACATCAACCAGAGTCTCTACCTCCACTCCTCCAACATC  
TACACACACATGGACTCACAAACCAGAACCCACTACATTTTGTCCCCATGAGA  
CAACAGCTGAGGTGACAGGAATCCCATCCCATACTCCTACAGACTGGAATGGC  
ACTGTGACATCCTCAGGAGATACCTGGAGTAATCACACTGAAGCAATCCCTCC  
AGGGAAGCCGCAGAAAAACCCTACTAAGGGCTTCTATGTTGGCATCTGCATCG  
CAGCCCTGCTGCTACTGCTCCTTGTGAGCACCGTGGCTATCACCAGGTACATA  
CTTATGAAAAGGAAGTCAGCATCTCTAAGCGTGGTTGCCTTCCGTGTCTCTAA  
GATTGAAGCTTTGCAGAACGCAGCGGTTGTGCATTCCCGAGCTGAAGACAACA  
TCTACATTGTTGAAGATAGACCTTGAGGGGCAGAATGAGTACCAGTGGCCCTC  
TGAGGGACCTTCTGCCTGAGATTTATAGAGACTGTCACTGATGTCATAGAGTC  
ACACCCATTACAGCGCCAAGGCGATTTTCTGTGTTGGTTCTTCCAGCTGCAGC  
AGAGAGGGTAACCCTCTACTGTGTATACTCAAACTCAGATTAACATCATCCT  
AATTTTGGTATCTGCACCACCTCCGTGTCTCTGCTCACTACAGAGATTCTCTC  
AAACATGAACGTTTTAGAAAGTTTGTGTTTCCCTTAGTCAATGTAATCATTGGT  
AATACTATTCTATTCTTGGTTACTAAAACCATTACTAAGAGAGGGATAGGAAT  
TAAAAGTTGGTGTGAGGGGCCTCCTGAATTTAGAAGCACTTGATTCTGTTTTA



[illegible]

AAGGGGATGATACCAAGCTTACCGGTTAGGCTGGGTGTTAGAAATCTCCTCA  
TTCACCTCTGTCCTGGAGTTGGGTTCTGGCTCTCCTTCACACCTAGGACTCTGA  
AATGAGCAAGCACTTCAGACAGTCAGGGTAGCAAGAGTCTAGCTGTCTGGTGG  
GCACCCAAAATGACCAGGGCTTAAGTCCCTTTTCTTTGGTTTAAGCCCGTTAT  
AATTAAATGGTACCAAAAAGCTTTAA (SEQ ID NO: 191) >NP\_032824.1  
programmed cell death protein 1 precursor [*Mus musculus*]  
MWVRQVPWSFTWAVLQLSWQSGWLLEVPNGPWRSLTFYPAWLTVSEGANATFT  
CSLSNWSIDLMLNWNRLSPSNQTEKQAAFCNGLSQPVQDARFQIIQLPNRHDF  
HMNILDTRRNDSGIYLCGAISLHPKAKIEESPGAELVVTERILETSTRYSPSPS  
PKPEGRFQGMVIGIMSALVGIPVLLLLLAWALAVFCSTSMSEARGAGSKDDTLK  
EEPSAAPVPSVAYEELDFQGREKTPELPTACVHTEYATIVFTEGLGASAMGRR  
GSADGLQGPRPPRHEDGHCSWPL (SEQ ID NO: 317) mScarlet >KY021423.1  
Synthetic construct mScarlet gene, partial cds, mRNA  
ATGGTGAGCAAGGGCGAGGCAGTGATCAAGGAGTTCATGCGGTTCAAGGTGCA  
CATGGAGGGCTCCATGAACGGCCACGAGTTCGAGATCGAGGGCGAGGGCGAGG  
GCCGCCCCCTACGAGGGCACCCAGACCGCCAAGCTGAAGGTGACCAAGGGTGGC  
CCCCTGCCCTTCTCCTGGGACATCCTGTCCCCTCAGTTCATGTACGGCTCCAG  
GGCCTTCACCAAGCACCCCGCCGACATCCCCGACTACTATAAGCAGTCCTTCC  
CCGAGGGGCTTCAAGTGGGAGCGCGTGATGAAGTTCGAGGACGGCGGGCGCCGTG  
ACCGTGACCCAGGACACCTCCCTGGAGGACGGCACCCCTGATCTACAAGGTGAA  
GCTCCGCGGCACCAACTTCCCTCCTGACGGCCCCGTAATGCAGAAGAAGACAA  
TGGGCTGGGAAGCGTCCACCGAGCGGTTGTACCCCGAGGACGGCGTGCTGAAG  
GGCGACATTAAGATGGCCCTGCGCCTGAAGGACGGCGGGCCGCTACCTGGCGGA  
CTTCAAGACCACCTACAAGGCCAAGAAGCCCGTGCAGATGCCCCGGCGCCTACA  
ACGTCGACCGCAAGTTGGACATCACCTCCCACAACGAGGACTACACCGTGGTG  
GAACAGTACGAACGCTCCGAGGGCCGCGCCACTCCACCGGCGGCATGGACGAGCT  
GTACAAG (SEQ ID NO: 192) >APD76535.1 mScarlet, partial [synthetic  
construct] MVSKGEAVIKEFMRFKVHMEGSMNGHEFEIEGEGEGRPHYEGTQTAKLKVTKGG  
PLPFSWDILSPQFMYGSRAFTKHPADIPDYKQSFPEGFKWERMNFEDGGAV  
TVTQDTSLEDGTLIYKVKLRGTNFPDGPVMQKKTMGWEASTERLYPEDGVLK  
GDIKMALRLKDGGRYLADFKTTYKAKKPVQMPGAYNVDRKLDITSHNEDYTVV  
EQYERSEGRHSTGGMDELYK (SEQ ID NO: 193) Nanoluciferase >JQ513379. 1  
NanoLuc reporter vector pNL1.1.CMV[Nluc/CMV], complete sequence, mRNA  
GGCCTAACTGGCCTCAATATTGGCCATTAGCCATATTATTCATTGGTTATATA  
GCATAAATCAATATTGGCTATTGGCCATTGCATACGTTGTATCTATATCATAA  
TATGTACATTTATATTGGCTCATGTCCAATATGACCGCCATGTTGGCATTGAT  
TATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCA  
TATATGGAGTTCCGCGTTACATAACTTACGGTAAATGGCCCCGCCTGGCTGACC  
GCCCAACGACCCCCGCCCATTGACGTCAATAATGACGTATGTTCCCATAGTAA  
CGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACT  
GCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTCCGCCCCCTATTGA  
CGTCAATGACGGTAAATGGCCCCGCCTGGCATTATGCCAGTACATGACCTTAC  
GGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATG  
GTGATGCGGTTTTTGGCAGTACACCAATGGGCGTGGATAGCGGTTTTGACTCACG  
GGGATTTCCAAGTCTCCACCCCATGACGTCAATGGGAGTTTTGTTTTGGCACC  
AAAATCAACGGGACTTTCCAAAATGTCGTAATAACCCCGCCCCGTTGACGCAA  
ATGGGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCGTTTAGT  
GAACCGTCAGATCACTAGAAGCTTTATTGCGGTAGTTTATCACAGTTAAATTG  
CTAACGCAGTCAGTGGGCCTCGGCGGCCAAGCTTGGCAATCCGGTACTGTTGG  
TAAAGCCACCATGGTCTTCACACTCGAAGATTTCTGTTGGGGACTGGCGACAGA

CAGCGGCTGACCAACTGCAAGTCCAGGACCTGAGGAGCTTGTGTTG  
TTTCAGAATCTCGGGGTGTCCGTAACCTCCGATCCAAAGGATTGTCCTGAGCGG  
TGAAAATGGGCTGAAGATCGACATCCATGTCATCATCCCGTATGAAGGTCTGA  
GCGGCGACCAAATGGGCCAGATCGAAAAAATTTTTAAGGTGGTGTACCCTGTG  
GATGATCATCACTTTAAGGTGATCCTGCACTATGGCACACTGGTAATCGACGG  
GGTTACGCCGAACATGATCGACTATTTTCGGACGGCCGTATGAAGGCATCGCCG  
TGTTTCGACGGCAAAAAGATCACTGTAACAGGGACCCTGTGGAACGGCAACAAA  
ATTATCGACGAGCGCCTGATCAACCCCGACGGCTCCCTGCTGTTCCGAGTAAC  
CATCAACGGAGTGACCGGCTGGCGGCTGTGCGAACGCATTCTGGCGTAATTCT  
AGAGTCGGGGCGGCCGGCCGCTTCGAGCAGACATGATAAGATAACATTGATGAG  
TTTGGACAAACCACAACCTAGAAATGCAGTGAAAAAAATGCTTTATTTGTGAAAT  
TTGTGATGCTATTGCTTTATTTGTAACCATTATAAGCTGCAATAAACAAAGTTA  
ACAACAACAATTGCATTCATTTTATGTTTCAGGTTTCAGGGGGAGGTGTGGGAG  
GTTTTTTTAAAGCAAGTAAAACCTCTACAAATGTGGTAAAATCGATAAGGATCC  
GTCGACCGATGCCCTTGAGAGCCTTCAACCCAGTCAGCTCCTTCCGGTGGGCG  
CGGGGCATGACTATCGTCGCCGCACTTATGACTGTCTTCTTTATCATGCAACT  
CGTAGGACAGGTGCCGGCAGCGCTCTTCCGCTTCCTCGCTCACTGACTCGCTG  
CGCTCGGTCGTTTCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAAT  
ACGGTTATCCACAGAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAG  
GCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCAT  
AGGCTCCGCCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTG  
GCGAAACCCGACAGGACTATAAAGATAACCAGGCGTTTCCCCCTGGAAGCTCCC  
TCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGCCTTT  
CTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCACGCTGTAGGTATCTCAG  
TTCGGTGTAGGTCGTTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTTC  
AGCCCGACCGCTGCGCCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGGTA  
AGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGC  
GAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCT  
ACACTAGAAGAACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTC  
GGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAGCGG  
TGGTTTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAAGGATCTCAAG  
AAGATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGGAACGAAAACCTCA  
CGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCTAGATCCT  
TTTAAATTAAAAATGAAGTTTTTAAATCAATCTAAAGTATATATGAGTAACTT  
GGTCTGACAGCGGCCGCAAATGCTAAACCACTGCAGTGGTTACCAGTGCTTGA  
TCAGTGAGGCACCGATCTCAGCGATCTGCCTATTTTCGTTTCGTCCATAGTGGCC  
TGACTCCCCGTCGTGTAGATCACTACGATTCGTGAGGGCTTACCATCAGGCCC  
CAGCGCAGCAATGATGCCGCGAGAGCCGCGTTACCGGGCCCCCGATTGTGTCAG  
CAATGAACCAGCCAGCAGGGAGGGCCGAGCGAAGAAGTGGTCCTGCTACTTTG  
TCCGCCTCCATCCAGTCTATGAGCTGCTGTCGTGATGCTAGAGTAAGAAGTTC  
GCCAGTGAGTAGTTTCCGAAGAGTTGTGGCCATTGCTACTGGCATCGTGGTAT  
CACGCTCGTCGTTTCGGTATGGCTTCGTTCAACTCTGGTTCCCAGCGGTCAAGC  
CGGGTCACATGATCACCCATATTATGAAGAAATGCAGTCAGCTCCTTAGGGCC  
TCCGATCGTTGTCAGAAGTAAGTTGGCCGCGGTGTTGTCGCTCATGGTAATGG  
CAGCACTACACAATTCTCTTACCGTCATGCCATCCGTAAGATGCTTTTTCCGTG  
ACCGGCGAGTACTCAACCAAGTCGTTTTTGTGAGTAGTGTATACGGCGACCAAG  
CTGCTCTTGCCCGGCGTCTATACGGGACAACACCGCGCCACATAGCAGTACTT  
TGAAAGTGCTCATCATCGGGAATCGTTCTTTCGGGGCGGAAAGACTCAAGGATC  
TTGCCGCTATTGAGATCCAGTTCGATATAGCCCCTTTCGACCCAGTTGATC  
TTCAGCATCTTTTACTTTTACCAGCGTTTCGGGGTGTGCAAAAACAGGCAAGC

AAAATGCCGCAAGGAATGAGTGCGACACGAAAATGTTGGATGCTCATA  
CTCGTCCTTTTTCAATATTATTGAAGCATTATCAGGGTTACTAGTACGTCTC  
TCAAGGATAAGTAAGTAATATTAAGGTACGGGAGGTATTGGACAGGCCGCAAT  
AAAATATCTTTATTTTCATTACATCTGTGTGTTGGTTTTTTGTGTGAATCGAT  
AGTACTAACATACGCTCTCCATCAAAACAAAACGAAACAAAACAAACTAGCAA  
AATAGGCTGTCCCCAGTGCAAGTGCAGGTGCCAGAACATTTCTCT (SEQ ID NO:  
194) >AFJ15599.1 NanoLuc luciferase [NanoLuc reporter vector  
pNL1.1.CMV[Nluc/CMV]]  
MVFTLEDFVGDWRQTAGYNLDQVLEQGGVSSLFQNLGVSVTPIQRIVLSGENG  
LKIDIHVIIPYEGLSGDQMGQIEKIFKVVPVDDHHFKVILHYGTLVIDGVTP  
NMIDYFGRPYEGIAVFDGKKITVTGTLWNGNKIIDERLINPDGSLLFRVTING  
VTGWRLCERILA (SEQ ID NO: 195)

(65) The polypeptides provided in Table 1 above are involved in a range of biological processes, including but not limited to, suppressing the adaptive arm of the immune system (e.g., PD-L1); cellular adhesion (e.g., nectin), immune activation (e.g., HVEM), and the like. The POI domains can also be used to track, purify, or identify the engineered EVs from native EVs (e.g., mScarlet and nanoluciferase). The genes, transcripts, polypeptides, variants, and fragments thereof can be used in any combination from Table 1 to be expressed by an engineered EV provided herein. In some embodiments, the POI domain is the human polypeptide. In some embodiments, the POI domain is a homologue of the human polypeptide (e.g., mouse).

(66) In some embodiments of any of the aspects, the engineered cell or EV provided herein comprises an exogenous nucleic acid encoding one or more exogenous polypeptide(s) selected from the group consisting of: the polypeptides listed in Table 1.

(67) In some embodiments of any of the aspects, the POI domain is PD-L1 or a fragment thereof. In some embodiments of any of the aspects, the POI domain is PD-L2 or a fragment thereof. In some embodiments of any of the aspects, the POI domain is FGL1 or a fragment thereof. In some embodiments of any of the aspects, the POI domain is 4-1BBL or a fragment thereof. In some embodiments of any of the aspects, the POI domain is CTLA or a fragment thereof.

(68) In some embodiments of any of the aspects, the POI domain substantially binds to one or more of a target polypeptide. In some embodiments of any of the aspects, the target polypeptide is a cellular receptor. In some embodiments of any of the aspects, the target polypeptide is an immunosuppressive polypeptide. In some embodiments of any of the aspects, the target polypeptide is an immunostimulatory polypeptide. The engineered exosomes provided herein can be designed to activate, block, or modulate a given target polypeptide with the appropriate POI domain that binds to or modulates the function or expression of the target polypeptide. Non-limiting examples of target polypeptides include those listed in Table 2 (below).

(69) TABLE-US-00002 TABLE 2 Exemplary Target Polypeptides PD-1 VISTA LAG-3 CD44 CD80 BTLA CD112 IL10RA CD86 CD160 CD200R IL10RB CD28 HVEM CD200 Tim-3 ICOS CD2 Galectin 9 TNFRSF25 CD28H SLAM CD150 TIM-3 TNFRSF6B PD-L1 CD58 CD226 CD113 CTLA-4 TIM-1 CD155 CD27 4-1BB (CD137) TIM-4 CD112 CD30 GITR CD40 DR3 LFA-3 (CD58) CD27L CD30L GITRL CD40L CD48 CD244 DcR3 CD28H LFA-3 (CD58) CD98 TNF Receptor TNF receptor Superfamily associated factor members (TRAF) family members Butyrophilin family PD-L2 Nectin TIM family members members B7/CD28 family SLAM family Nectin-like Collagen members members binding receptors family proteins LAIR-1 (CD305)

(70) The EVs provided herein further comprise at least one fusion protein comprising a vesicle targeting domain. In various embodiments, the vesicle targeting domain provided herein is capable of binding or anchoring the fusion polypeptide provided herein to an extracellular vesicle, e.g., via targeting of the phospholipid bilayer membrane. In various embodiments, the vesicle targeting domain is a GPI domain (i.e., GPI linker, GPI anchor), fatty acylation site, or prenylation site. One of skill in the art can appreciate that the aforementioned refer to peptide or protein sites, wherein

covalent lipid attachment supports embedding of the lipid in a cell membrane (i.e., phospholipid bilayer).

(71) Biochemical forces that anchor EV targeting domains to the EV phospholipid bilayer may include, but are not limited to, electrostatic forces, affinity for EVs through protein-protein interactions with natively resident proteins (e.g., CD81, CD63, CD9, ALIX, TSG101, CD98, CD298, MARCKS, PTGFRN, Lactadherin (MFG8)), association or affinity for negatively or positively curved phospholipids, association or affinity for negatively or positively charged domains of resident membrane associated proteins, etc., or the like.

(72) Additional non-limiting examples of membrane targeting domains that can be used and their properties are further described in detail, e.g., Alberts B, Johnson A, Lewis J, et al., *Molecular Biology of the Cell*, 4th edition, New York: Garland Science, 2002. Membrane Proteins; Marilyn D. Resh, Fatty acylation of proteins: new insights into membrane targeting of myristoylated and palmitoylated proteins. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research*. Volume 1451, Issue 1, 12 Aug. 1999, Pages 1-16, doi: 10.1016/S0167-4889(99)00075-0; Ann Apolloni, et al., H-ras but Not K-ras Traffics to the Plasma Membrane through the Exocytic Pathway, *Molecular and Cellular Biology* April 2000, 20 (7) 2475-2487, doi: 10.1128/MCB.20.7.2475-2487.2000; Rosie Dawaliby et. al., Phosphatidylethanolamine Is a Key Regulator of Membrane Fluidity in Eukaryotic Cells, *Membrane Biology*, VOLUME 291, ISSUE 7, doi: 10.1074/jbc.M115.706523; R. J. Deschenes, Protein Palmitoylation, *Encyclopedia of Biological Chemistry (Second Edition)*, Academic Press, 2013, Pages 645-647, ISBN 9780123786319, doi: 10.1016/B978-0-12-378630-2.00022-0; Charuta C. Palsuledesai and Mark D. Distefano, Protein Prenylation: Enzymes, Therapeutics, and Biotechnology Applications, *ACS Chemical Biology* 2015 10 (1), 51-62, doi: 10.1021/cb500791f; Hung M E, Leonard J N. Stabilization of exosome-targeting peptides via engineered glycosylation, *J Biol Chem*, 2015 Mar. 27; 290 (13): 8166-72, doi: 10.1074/jbc.M114.621383; Udenwobele Daniel Ikenna, et. al., Myristoylation: An Important Protein Modification in the Immune Response, *Frontiers in Immunology*, Vol: 8, 2017, doi: 10.3389/fimmu.2017.00751; Kinoshita Taroh 2020 Biosynthesis and biology of mammalian GPI-anchored proteins *Open Biol.* 10190290, doi: 10.1098/rsob.190290, the contents of which are incorporated herein by reference in their entireties.

(73) In some embodiments, the fusion polypeptide comprises one or more, two or more, three or more, four or more, five or more, or six or more vesicle targeting domains on the same polypeptide or nucleic acid construct encoding said polypeptide. For example, the fusion polypeptides provided herein can comprise PD-L1 and Glycosylphosphatidylinositol (GPI).

(74) In some embodiments, the vesicle targeting domain is a prenylated protein. Prenylated proteins are proteins that have at least one prenylation site. Prenylation occurs when a 15-carbon or 20-carbon, farnesyl or geranylgeranyl isoprenoid, respectively, is covalently bound via a thioether bond to a cysteine at or near the carboxy terminus of a protein. In general, a prenylation site comprises an amino acid sequence CAAX, wherein C represents cysteine, A represents an aliphatic amino acid (glycine, alanine, valine, leucine, or isoleucine), and X represents alanine, methionine, serine, leucine, or glutamine.

(75) In some embodiments, the vesicle targeting domain is a fatty acylated protein. Fatty acylated proteins are proteins that have been modified post-translationally by covalent attachment of one or more fatty acids, generally with a saturated fatty acid that comprises 14-carbon (e.g., myristic acid) via myristoylation or 16-carbons (e.g., palmitic acid) via palmitoylation. For example, proteins destined to become myristoylated begin with the amino acids Met-Gly-X-X-X followed by a serine or threonine at position 6 and lysine or arginine at position 7 and/or 8 wherein X can be any amino acid. The methionine is removed and a myristate is linked to the glycine via an amide bond. Palmitoylation herein means a posttranslational covalent attachment of fatty acids (e.g., palmitic acid) to cysteine (S-palmitoylation), serine and/or threonine (O-palmitoylation), and to the amino group of lysine (N-palmitoylation) of proteins.

(76) Palmitoylated proteins may be acylated by attachment of a thioester linkage to a sulfhydryl group of cysteine, or via a palmitate linked to the amino group of an N-terminal cysteine. Palmitoylation sites may be present near the N- or C-terminus of a protein.

(77) In some embodiments, the vesicle targeting domain is a glycosylphosphatidylinositol (GPI) anchor. A glycosylphosphatidylinositol (GPI) anchor (“GPI anchor”) or “GPI sticky binder” are used interchangeably and refer to a means of stably anchoring a protein to an outer leaflet (e.g., exterior layer of a phospholipid bilayer) of a cell membrane. A GPI anchor comprises a glycan, a phosphoethanolamine linker, a phospholipid tail, and may be modified by various glycan sidechains. The glycan core comprises phosphoinositol, glucosamine, and mannose residues wherein said mannose residues may be modified for example with phosphoethanolamine or carbohydrates. The phosphoethanolamine is amide-bonded to the carboxyl terminus of a protein during the process of GPI attachment. In some embodiments, the vesicle targeting domain may have affinity to EV resident proteins, e.g., CD81, CD63, CD9, ALIX, TSG101, CD98, CD298, MARCKS, PTGFRN, Lactadherin (MFG8).

(78) Sticky binders can include a sequence for one or more myristoylation and/or palmitoylation (Myr/Palm) sites fused to a transmembrane domain from 4F2 (CD98). For example, the myristoylation sequence from the MARCKS protein may be modified to encode for one or more myristoylation and palmitoylation sites, wherein the modified MARCKS protein sequence is fused to a protein sequence of the transmembrane domain from 4F2 via a covalent peptide bond. A Myr/Palm followed by the 4F2 transmembrane domain can improve loading of the fusion proteins provided herein when compared with 4F2 transmembrane domain alone or Myr/Palm alone.

(79) Non-limiting examples of vesicle targeting domains that enhance fusion polypeptide structure and function on the extracellular vesicles are provided in Table 3 (below).

(80) TABLE-US-00003 TABLE 3 Exosome Targeting Domain Exosome Targeting Domain/Sticky Nucleic Acid Sequence (SEQ ID NO:) Binder Amino Acid Sequence (SEQ ID NO:) Human CD55 >NM\_000574.5 *Homo sapiens* CD55 molecule (Cromer blood (DAF) group) (CD55), transcript variant 1, mRNA Glycosylphosphatidylinositol

CTGCTTACTGCAACTCGCTCCGGCCGCTGGGCGTAGCTGCGACTCGGCGGAGTCCCG (GPI)  
GCGGCGCGTCCTTGTTCTAACCCGGCGCGCCATGACCGTCGCGCGGCGGAGCGTGCC  
CGCGGCGCTGCCCCCTCCTCGGGGAGCTGCCCCGGCTGCTGCTGCTGGTGCTGTTGTG  
CCTGCCGGCCGTGTGGGGTGACTGTGGCCTTCCCCCAGATGTACCTAATGCCAGCC  
AGCTTTGGAAGGCCGTACAAGTTTTCCCGAGGATACTGTAATAACGTACAAATGTGA  
AGAAAGCTTTGTGAAAATTCCTGGCGAGAAGGACTCAGTGATCTGCCTTAAGGGCAG  
TCAATGGTCAGATATTGAAGAGTTCTGCAATCGTAGCTGCGAGGTGCCAACAAGGCT  
AAATTCTGCATCCCTCAAACAGCCTTATATCACTCAGAATTATTTTCCAGTCGGTAC  
TGTTGTGGAATATGAGTGCCGTCCAGGTTACAGAAGAGAACCTTCTCTATCACCAA  
ACTAATTGCCTTCAGAATTTAAAATGGTCCACAGCAGTCGAATTTTGTAAAAAGAA  
ATCATGCCCTAATCCGGGAGAAATACGAAATGGTCAGATTGATGTACCAGGTGGCAT  
ATTATTTGGTGCAACCATCTCCTTCTCATGTAACACAGGGTACAAATTATTTGGCTC  
GACTTCTAGTTTTTGTCTTATTTTCAAGGAGCTCTGTCCAGTGAGTGACCCGTTGCC  
AGAGTGCAGAGAAATTTATTGTCCAGCACCACCACAAATTGACAATGGAATAATTCA  
AGGGGAACGTGACCATTATGGATATAGACAGTCTGTAACGTATGCATGTAATAAAGG  
ATTCACCATGATTGGAGAGCACTCTATTTATTGTACTGTGAATAATGATGAAGGAGA  
GTGGAGTGGCCCACCACCTGAATGCAGAGGAAAATCTCTAACTTCCAAGGTCCCACC  
AACAGTTCAGAAACCTACCACAGTAAATGTTCCAACCTACAGAAGTCTCACCAACTTC  
TCAGAAAACCACCACAAAAACCACCACACCAATGCTCAAGCAACACGGAGTACACC  
TGTTTCCAGGACAACCAAGCATTTTTCATGAAACAACCCCAAATAAAGGAAGTGGAAC  
CACTTCAGGTACTACCCGTCTTCTATCTGGGCAGACGTGTTTCACGTTGACAGGTTT





ATTAGCTTGAGCAAGGCTTGAAGGCTTGGGAGAGCACTTGGGAGAG  
CCAGTCCTGTTTTGCAGGGAAGCCCCACTTGAAGGAAGAAGTCTAAGAGTGAA  
GTAGGTGTGACTTGAAGTAGATTGCATGCTTCCTCCTTTGCTCTTGGGAAGA  
CCAGCTTTGCAGTGACAGCTTGAGTGGGTTCTCTGCAGCCCTCAGATTATTT  
TTCCTCTGGCTCCTTGGATGTAGTCAGTTAGCATCATTAGTACATCTTTGGA  
GGGTGGGGCAGGAGTATATGAGCATCCTCTCTCACATGGAACGCTTTCATAA  
ACTTCAGGGATCCCGTGTTGCCATGGAGGCATGCCAAATGTTCCATATGTGG  
GTGTCAGTCAGGGACAACAAGATCCTTAATGCAGAGCTAGAGGACTTCTGGC  
AGGGAAGTGGGGAAGTGTTCCAGATAGCAGGGCATGAAAACCTTAGAGAGGTA  
CAAGTGGCTGAAAATCGAGTTTTTTCCTCTGTCTTTAAATTTTATATGGGCTT  
TGTTATCTTCCACTGGAAAAGTGTAATAGCATACATCAATGGTGTGTAAAG  
CTATTTCCCTTGCCTTTTTTTTTTATTGGAATGGTAGGATATCTTGGCTTTGCCA  
CACACAGTTACAGAGTGAACACTCTACTACATGTGACTGGCAGTATTAAGTG  
TGCTTATTTTAAATGTTACTGGTAGAAAGGCAGTTCAGGTATGTGTGTATAT  
AGTATGAATGCAGTGGGGACACCCTTTGTGGTTACAGTTTGAGACTTCCAAA  
GGTCATCCTTAATAACAACAGATCTGCAGGGGTATGTTTTACCATCTGCATC  
CAGCCTCCTGCTAACTCCTAGCTGACTCAGCATAGATTGTATAAAATACCTT  
TGTAACGGCTCTTAGCACACTCACAGATGTTTGAGGCTTTCAGAAGCTCTTC  
TAAAAAATGATACACACCTTTCACAAGGGCAAACCTTTTTCCCTGTG  
TATTCTAGTGAATGAATCTCAAGATTCAGTAGACCTAATGACATTTGTATTT  
TATGATCTTGGCTGTATTTAATGGCATAGGCTGACTTTTGCAGATGGAGGAA  
TTTCTTGATTAATGTTGAAAAAAAACCCTTGATTATACTCTGTTGGACAAAC  
CGAGTGCAATGAATGATGCTTTTCTGAAAATGAAATATAACAAGTGGGTGAA  
TGTGGTTATGGCCGAAAAGGATATGCAGTATGCTTAATGGTAGCAACTGAAA  
GAAGACATCCTGAGCAGTGCCAGCTTTCCTTCTGTTGATGCCGTTCCCTGAAC  
ATAGGAAAATAGAAACTTGCTTATCAAAACTTAGCATTACCTTGGTGCTCTG  
TGTTCTCTGTTAGCTCAGTGTCTTTCCTTACATCAATAGGTTTTTTTTTTTT  
TTTTTGGCCTGAGGAAGTACTGACCATGCCACAGCCACCGGCTGAGCAAAG  
AAGCTCATTTTCATGTGAGTTCTAAGGAATGAGAAACAATTTTGATGAATTTA  
AGCAGAAAATGAATTTCTGGGAACCTTTTTTGGGGGCGGGGGGGTGGGGAATT  
CAGCCACACTCCAGAAAGCCAGGAGTCGACAGTTTTTGGAAGCCTCTCTCAGG  
ATTGAGATTCTAGGATGAGATTGGCTTACTGCTATCTTGTGTCATGTACCCA  
CTTTTTGGCCAGACTACACTGGGAAGAAGGTAGTCCTCTAAAGCAAAATCTG  
AGTGCCACTAAATGGGGAGATGGGGCTGTTAAGCTGTCCAAATCAACAAGGG  
TCATATAAATGGCCTTAAACCTTTGGGGTTGCTTTCCTGCAAAAAGTTGCTGTG  
ACTCATGCCATAGACAAGGTTGAGTGCCTGGACCCAAAGGCAATACTGTAAT  
GTAAAGACATTTATAGTACTAGGCAAACAGCACCCCAGGTA CTCCAGGCCCT  
CCTGGCTGGAGAGGGCTGTGGCAATAGAAAATTAGTGCCAACTGCAGTGAGT  
CAGCCTAGGTTAAATAGAGAGTGTAAGAGTGCTGGACAGGAACCTCCACCCT  
CATGTCACATTTCTTCAATGTGACCCTTCTGGCCCCCTCTCCTCCTGACAGCG  
GAACAATGACTGCCCCGATAGGTGAGGCTGGAGGAAGAATCAGTCCTGTCCCT  
TGGCAAGCTCTTCACTATGACAGTAAAGGCTCTCTGCCTGCTGCCAAGGCCT  
GTGACTTTCTAACCTGGCCTCACGCTGGGTAAAGCTTAAGGTAGAGGTGCAGG  
ATTAGCAAGCCCACCTGGCTACCAGGCCGACAGCTACATCCTCCA ACTGACC  
CTGATCAACGAAGAGGGATTCATGTGTCTGTCTCAGTTGGTTCCAAATGAAA  
CCAGGGAGCAGGGGAGTTAGGAATCGAACACCAGTCATGCCTACTGGCTCTC  
TGCTCGAGAGCCAATACCCTGTGCCCTCCACTCATCTGGATTTACAGGAACT  
GTCATAGTGTT CAGTATTGGGTGGTGATAAGCCCATTTGGATTGTCCCCTTGG  
GGGGATGAGCTAGGGGTGCAAGGAACACCTGATGAGTAGATAAGTGAGGCTC  
ATGGTATTTCCCTGAAAGATGCTAATCTATTTGCCAAACTTGGTCTTGAATGT

ACTGGGCTTCAAGGTTAGTATATTTGTGTCCTTTGGTCACTTTGGCAGTTAGCC  
CCCATGTCTTATGTGTGTCCTGAAAAAATAAGAGCCTGCCCAAGACTTTTGGG  
CCTCTTGACAGAATTAACCACTTTTATACATCTGAGTTCTCTTGGTAAGTTC  
TTTAGCAGTGTTCAAAGTCTACTAGCTCGCATTAGTTTCTGTTGCTGCCAAC  
AGATCTGAACTAATGCTAACAGATCCCCCTGAGGGATTCTTGATGGGCTGAG  
CAGCTGGCTGGAGCTAGTACTGACTGACATTCATTGTGATGAGGGCAGCTTT  
CTGGTACAGGATTCTAAGCTCTATGTTTTTATATACATTTTCATCTGTACTTG  
CACCTCACTTTACACAAGAGGAACTATGCAAAGTTAGCTGGATCGCTCAAG  
GTCACTTAGGTAAGTTGGCAAGTCCATGCTTCCCCTCAGCTCCTCAGGTCA  
GCAAGTCTACTTCTCTGCCTATTTTGTATACTCTCTTTAATATGTGCCTAGC  
TTTGGAAGTCTAGAATGGGTCCCTGGTGCCTTTTTACTTTGAAGAAATCAG  
TTTCTGCCTCTTTTTTGGAAGAAACAAAGTGCAATTGTTTTTTACTGGAA  
AGTTACCCAATAGCATGAGGTGAACAGGACGTAGTTAGGCCTTCCTGTAAAC  
AGAAAATCATATCAAAACACTATCTTCCCCTCTGTTTCTCAATGCCTGCTAC  
TTCTTGATAGATATTTCAATTCAGGAGAGCAGCAGTTAAACCCGTGGATTTTG  
TAGTTAGGAACCTGGGTCAAACCCCTCTTCCACTAATTGGCTATGTCTCTGG  
ACAAGTTTTTTTTTTTTTTTTTTTTTAAACCCTTTCTGAACCTTCACTTTCT  
ATGTCTACCTCAAAGAATTGTTGTGAGGCTTGAGATAATGCATTTGTAAAGG  
GTCTGCCAGATAGGAAGATGCTAGTTATGGATTACAAGGTTGTTAAGGCTG  
TAAGAGTCTAAAACCTACAGTGAATCACAATGCATTTACCCCCACTGACTTG  
GACATAAGTGAAAACCTAGCCAGAAGTCTCTTTTTCAAATTACTTACAGGTTA  
TTCAATATAAAATTTTTGTAATGGATAATCTTATTTATCTAAACTAAAGCTT  
CCTGTTTATACACACTCCTGTTATTCTGGGATAAGATAAATGACCACAGTAC  
CTTAATTTCTAGGTGGGTGCCTGTGATGGTTCATTGTAGGTAAGGACATTTT  
CTCTTTTTTCAGCAGCTGTGTAGGTCCAGAGCCTCTGGGAGAGGAGGGGGTA  
GCATGCACCCAGCAGGGGACTGAACTGGGAACTCAAGGTTCTTTTTACTGT  
GGGGTAGTGAGCTGCCTTTCTGTGATCGGTTTCCCTAGGGATGTTGCTGTTT  
CCCTCCTTGCTATTCGCAGCTACATAACAAGTGGCCAACCCCAAGTAGGCTGA  
TCCTATATATGATCAGTGCTGGTGCTGACTCTCAATAGCCCCACCCAAGCTG  
GCTATAGGTTTACAGATACATTAATTAGGCAACCTAAAATATTGATGCTGGT  
GTTGGTGTGACATAATGCTATGGCCAGAAGTGAAGCTTAGAGTTATAATTCA  
TGTATTAGGGTTCTCCAGAGGGACAGAATTAGTAGGATATATGTATATATGA  
AAGGGAGGTTATTAGGGAGAAGTGGCTCCCACAGTTAGAAGGCGAAGTCGCA  
CAATAGGCCGTCTGCAAGCTGGGTAGAGAGAAGCCAGTAGTGGCTCAGCCT  
GAGTTCAAAAACCTCAAACTGGGGAAGCTGACAGTGCAGCCAGCCTTCAGT  
CTGTGGCCAAAGGCCCAAGAGCCCCTGGCAACCAACCCACTGGTGCAAGTCC  
TAGATTCAAAGGCTGAAGAACCTGGAGTCTGATGTCCAAGAGCAGGAAGAG  
TGGAAGAAAGCCAGAAGACTCAGCAAACAAGGTAGACAGTGTCTACCACCAT  
AGTGGCCATACCAAAGAGGCTACCGATTCTTCCCTGCTACCTGGATCCCTGA  
AGTTGCCCTGGTCTCTGCACCTTCTAAACCTAGTTCTTAAGAGCTTTCCATT  
ACATGAGCTGTCTCAAAGCCCTCCAATAAATTCTCAGTGTAAGCTTCTGTTG  
CTTGTGGACAGAAAATTCTGACAGACCTACCCTATAAGTGTTACTGTCAGGA  
TAACATGAGAACGCACAACAGTAAGTGGTCACTAAGTGTTAGCTACGGTTAT  
TTTGCCCAAGGTAGCATGGCTAGTTGATGCCGGTTGATGGGGCTTAAACCCA  
GCTCCCTCATCTTCCAGGCCTCTGTACTCCCTATTCCACTAACTACCTCTC  
AGGTTTATTTTTTTTAAATTCTTACTCTGCAAGTACATAGGACCACATTTACC  
TGGGAAAACAAGAATAAAGGCTGCTCTGCATTTTTTTAGAACTTTTTTGAAA  
GGGAGATGGGAATGCCTGCACCCCCAAGTCCAGACCAACACAATGGTTAATT  
GAGATGAATAATAAAGGAAAGACTGTTCTGGGCTTCCCAGAATAGCTTGGTC  
CTTAAATTGTGGCACAAACAACCTCCTGTCAGAGCCAGCCTCCTGCCAGGAA

GAGGGTAGGAGACTAGAGGACCCCTGTGTGCAGCCTTGCCCTGAAGGCTAGGG  
TGACAATTTGGAGGCTGTCCAAACACCCTGGCCTCTAGAGCTGGCCTGTCTA  
TTTGAAATGCCGGCTCTGATGCTAATCGGCGACCCTCAGGCAAGTTACTTAA  
CCTTACATGCCTCAGTTTTTCTCATCTGGAAAATGAGAACCCTAGGTTTAGGG  
TTGTTAGAAAAGTTAAATGAGTTAAGACAAGTGCCTGGGACACAGTAGCCTC  
TTGTGTGTGTTTATCATTATGTCCTCAGCAGGTCGTAGAAGCAGCTTCTCAG  
GTGTGAGGCTGGCGCGATTATCTGGAGTGGGTGGGTCTTAGGATGGACC  
CCCTGCTGCATTTTCCTCATTATCCACCAGGGCTTAATGGGGAATCAAGGA  
ATCCATGTGTAAGTGTATAATACTGTAGCCACACTCCAATGACCACCTACT  
AGTTGTCCCTGGCACTGCTTATACATATGTCCATCAAATCAATCCTATGAAG  
TAGATACTGTCTTCATTTTATAGATCAGAGACAATTGGGGTTCAGAGAGCTG  
ATGTGATTTTCCCAGGGTCACAGAGAGTCCCAGATTCAGGCACAACCTCTTGT  
ATTCCAAGACACAACCCTACATGTCCAAAGGCTGCCCAGAGCCACCGGGCA  
CGGCAAATTGTGACATATCCCTAAAGAGGCTGAGCACCTGGTCAGGATCTGA  
TGGCTGACAGTGTGTCCAGATGCAGAGCTGGAGTGGGGGAGGGGAAGGGGGG  
CTCCTTGGGACAGAGAAGGCTTTCTGTGCTTTCTCTGAAGGGAGCAGTCTGA  
GGACCAAGGGAACCCGGCAAACAGCACCTCAGGTACTCCAGGCCCTCCTGGC  
TGGAGAGGGCTGTGGCAATGGAAAATTAGTGCCAACTGCAATGAGTCAGCCT  
CGGTTAAATAGAGAGTGAAGAATGCTGGACAGGAACCTCCACCCTCATGTCA  
CATTTCTTCAGTGTGACCCTTCTGGCCCCCTCTCCTCCTGACAGCGGAACAAT  
GACTGCCCCGATAGGTGAGGCTGGAGGAAGAATCAGTCCTGTCCTTGGCAAG  
CTCTTCACTATGACAGTAAAGGCTCTCTGCCTGCTGCCAAGGCCTGTGACTT  
TCTAACCTGGCCTCACGCTGGGTAAGCTTAAGGTAGAGGTGCAGGATTAGCA  
AGCCCACCTGGCTACCAGGCCGACAGCTACATCTTTCAACTGACCCTGATCA  
ACGAAGAGGGACTTGTGTCTCTCAGTTGGTTCCAAATGAAACCAGGGAGCAG  
GGGCGTTAGGAAGCTCCAACAGGATGGTACTTAATGGGGCATTTGAGTGGAG  
AGGTAGGTGACATAGTGCTTTGGAGCCCAGGGAGGGGAAAGGTTCTGCTGAAG  
TTGAATTCAAGACTGTTCTTTCATCACAACTTGAGTTTCCTGGACATTTGT  
TTGCAGAAACAACCGTAGGGTTTTGCCTTAACCTCGTGGGTTTATTATTACC  
TCATAGGGACTTTGCCTCCTGACAGCAGTTTATGGGTGTTTATTGTGGCACT  
TGAGTTTTCTTGCATACTTGTTAGAGAAACCAAGTTTGTGCATCAACTTCTTA  
TTTAACCCCTGGCTATAACTTCATGGATTATGTTATAATTAAGCCATCCAG  
AGTAAATCTGTTTAGATTATCTTGGAGTAAGGGGGGAAAAAATCTGTAATTT  
TTTCTCCTCAACTAGATATATACATAAAAAAATGATTGTATTGCTTCATTAA  
AAAATATAACGCAAAATCTCTTTTCCTTCTAAAAAAAAAAAAAAAAAAAAA (SEQ ID  
NO: 321)>NP\_976075.1 CD59 glycoprotein preproprotein [*Homo sapiens*]  
MGIQGGSVLFLGLLLVLAVFCHSGHSLQCYNCPNPTADCKTAVNCSSDFDACL  
ITKAGLQVYNKCWKFEHCNFDVTTTRLRENELTYYCCKKDLCNFNEQLENGG  
**TSLSEKTVLLLVTPLAAWSLHP** (SEQ ID NO: 198) Human C1C2 from  
NM\_005928.4 *Homo sapiens* milk fat globule-EGF factor 8 MFGE8 protein  
(MFGE8), transcript variant 1, mRNA  
AGAACCCCGCGGGGTCTGAGCAGCCCAGCGTGCCCATTCAGCGCCCGCGTCCCCGC  
AGCATGCCGCGCCCCCGCCTGCTGGCCGCGCTGTGCGGCGCGCTGCTCTGCGCCCCC  
AGCCTCCTCGTCGCCCTGGATATCTGTTCCAAAAACCCCTGCCACAACGGTGTTTA  
TGCAGAGGAGATTTCCCAAGAAGTGCGAGGAGATGTCTTCCCCTCGTACACCTGCACG  
TGCCTTAAGGGCTACGCGGGCAACCACTGTGAGACGAAATGTGTCGAGCCACTGGGC  
**CTGGAGAATGGGAACATTGCCAACTCACAGATCGCCGCCTCGTCTGTGCGTGTGACC**  
**TTCTTGGGTTTGCAGCATTGGGTCCCGGAGCTGGCCCCGCCTGAACCGCGCAGGCATG**  
**GTCAATGCCTGGACACCCAGCAGCAATGACGATAACCCCTGGATCCAGGTGAACCTG**  
**CTGCGGAGGATGTGGGTAACAGGTGTGGTGACGCAGGGTGCCAGCCGCTTGGCCAGT**

CATGAGTCTGACGAGGCCTTCAAGGCTTACAGGCTTAATGACACGCAATTCGAT  
TTCATCCATGATGTTAATAAAAAACACAAGGAGTTTGTGGGTAACTGGAACAAAAAC  
GCGGTGCATGTCAACCTGTTTGAGACCCCTGTGGAGGCTCAGTACGTGAGATTGTAC  
CCCACGAGCTGCCACACGGCCTGCACTCTGCGCTTTGAGCTACTGGGCTGTGAGCTG  
AACGGATGCGCCAATCCCCTGGGCCTGAAGAATAACAGCATCCCTGACAAGCAGATC  
ACGGCCTCCAGCAGCTACAAGACCTGGGGCTTGCATCTCTTCAGCTGGAACCCCTCC  
TATGCACGGCTGGACAAGCAGGGCAACTTCAACGCCTGGGTTGCGGGGAGCTACGGT  
AACGATCAGTGGCTGCAGGTGGACCTGGGCTCCTCGAAGGAGGTGACAGGCATCATC  
ACCCAGGGGGGCCGTAACCTTTGGCTCTGTCCAGTTTGTGGCATCCTACAAGGTTGCC  
TACAGTAATGACAGTGCGAACTGGACTGAGTACCAGGACCCCAGGACTGGCAGCAGT  
AAGATCTTCCCTGGCAACTGGGACAACCACTCCCACAAGAAGAAGTGTGTTGAGACG  
CCCATCCTGGCTCGCTATGTGCGCATCCTGCCTGTAGCCTGGCACAACCGCATCGCC  
CTGCGCCTGGAGCTGCTGGGCTGTTAGTGGCCACCTGCCACCCCCAGGTCTTCCTGC  
TTTCCATGGGCCCCGCTGCCTCTTGGCTTCTCAGCCCCCTTTAAATCACCATAGGGCTG  
GGGACTGGGGAAGGGGAGGGTGTTTCAGAGGCAGCACCACCACACAGTCACCCCTCCC  
TCCCTCTTTCCCACCCTCCACCTCTCACGGGGCCCTGCCCCAGCCCCTAAGCCCCGTC  
CCCTAACCCCCAGTCCTCACTGTCCTGTTTTCTTAGGCACTGAGGGATCTGAGTAGG  
TCTGGGATGGACAGGAAAGGGCAAAGTAGGGCGTGTGGTTTCCCTGCCCCCTGTCCGG  
ACCGCCGATCCCAGGTGCGTGTGTCTCTGTCTCTCCTAGCCCCCTCTCTCACACATCA  
CATTCCCATGGTGGCCTCAAGAAAGGCCCGGAAGCGCCAGGCTGGAGATAACAGCCT  
CTTGCCCGTCGGCCCTGCGTCGGCCCTGGGGTACCATGTGGCCACAACCTGCTGTGGC  
CCCCTGTCCCAAGACACTTCCCCTTGTCTCCCTGGTTGCCTCTCTTGCCCCCTTGTC  
CTGAAGCCCAGCGACACAGAAGGGGGTGGGGCGGGTCTATGGGGAGAAAGGGAGCGA  
GGTCAGAGGAGGGCATGGGTTGGCAGGGTGGGCGTTTGGGGCCCTCTATGCTGGCTT  
TTCACCCCAGAGGACACAGGCAGCTTCCAAAATATATTTATCTTCTTCACGGGAA  
(SEQ ID NO: 199)>NP\_005919.2 lactadherin isoform a preproprotein [*Homo sapiens*]

MPRPRLLAALCGALLCAPSLLLVALDICKSNPCHNGGLCEEISQEVRGDVFPSTCTC  
LKGYAGNHCETKCV EPLGLENGNIANSQIAASSVRVTFGLGLQHWVPELARLN RAGMV  
NAWTPSSNDDNPWIQVNLLRRMWVTGVVTQ GASRLASHEY LKAFKVAYS LNHGHEFDF  
IHDVNKKHKEFVGNWNKNAVHVNL FETPVEAQYVRLYPT SCHTACTLR FELLGCELN  
GCANPLGLKNNSIPDKQITASSSYKTWGLHLFSWNPSYARLDKQGNFN AWVAGSYGN  
DQWLQVDLGSSKEVTGIITQ GARNEFSVQFVASYKVAYSND SANWTEYQDPRTGSSK  
IFPGNWDN HSHKKNLFETPILARYVRILPVAWHNR IALRLELLGC (SEQ ID NO:  
200) Human 4F2 (CD98)>NM\_002394.6 *Homo sapiens* solute carrier family 3  
member 2 (SLC3A2), transcript variant 3, mRNA  
GCATTGCGGCTTGGTTTTCTCACCCAGTGCATGTGGCAGGAGCGGTGAGATC  
ACTGCCTCACGGCGATCCTGGACTGACGGTCACGACTGCCTACCCTCTAACC  
CTGTTCTGAGCTGCCCCCTTGCCCACACACCCCCAAACCTGTGTGCAGGATCCG  
CCTCCATGGAGCTACAGCCTCCTGAAGCCTCGATCGCCGTCGTGTCGATTCC  
GCGCCAGTTGCCTGGCTCACATTCGGAGGCTGGTGTCCAGGGTCTCAGCGCG  
GGGGACGACTCAGAGTTGGGGTCTCACTGTGTTGCCCAGACTGGTCTCGAAC  
TCTTGGCCTCAGGTGATCCTCTTCCCTCAGCTTCCCAGAATGCCGAGATGAT  
AGAGACGGGGTCTGACTGTGTTACCCAGGCTGGTCTTCAACTCTTGGCCTCA  
AGTGATCCTCCTGCCTTAGCTTCCAAGAATGCTGAGGTTACAGGCACCATGA  
GCCAGGACACCGAGGTGGATATGAAGGAGGTGGAGCTGAATGAGTTAGAGCC  
CGAGAAGCAGCCGATGAACGCGGCGTCTGGGGCGGGCATGTCCCTGGCGGGA  
GCCGAGAAGAATGGTCTGGTGAAGATCAAGGTGGCGGAAGACGAGGCGGAGG  
CGGCAGCCGCGGCTAAGTTCACGGGCCTGTCCAAGGAGGAGCTGCTGAAGGT  
GGCAGGCAGCCCCGGCTGGGTACGCACCCGCTGGGCACTGCTGCTGCTCTTC

>NP\_002385.3 4F2 cell-surface antigen heavy chain isoform c [*Homo sapiens*]  
**MELQPPEASIAVVSIPRQLPGSHSEAGVQGLSAGDDSELGSHCVAQTGLELL**  
**ASGDPLPSASQNAEMIETGSDCVTQAGLQLLASSDPPALASKNAEVTGTMSQ**  
**DTEVDMKEVELNELEPEKQPMNAASGAAMSLAGAEEKNLVKIKVAEDEAEAA**  
**AAAKFTGLSKEELLKVAGSPGWVRTRWALLLLFWLGWLGMLAGAVVIIVRAP**  
RCRELPAQKWWHTGALYRIGDLQAFQGHGAGNLAGLKGRLDYLSSLKVKGLV  
LGPIHKNQKDDVAQTDLLQIDPNFGSKEDFDSLLQSAKKKSIRVILDLT  
PNYRGNSWFTQVDTVATKVKDALEFWLQAGVDGFQVRDIENLKDASSFLAEWQ  
NITKGFSEDRLLIAGTNSSDLQQILSLLESNKDLLLTSSYLSDSGSTGEHTK  
SLVTQYLNATGNRWCSWSLSQARLLTSFLPAQLLRLYQLMLFTLPGTPVFSY  
GDEIGLDAAALPGQPMAPVMLWDESSFPDIPGAVSANMTVKGQSEDPGSL  
LLSLFRRLSDQRSKERSLLHGDFHAFSAGPGLFSYIRHWDQNERFLVVLNFGDV  
GLSAGLQASDLPASASLPAKADLLLSTQPGREEGSPLELERLKLEPHEGLLL RFPYAA  
(SEQ ID NO: 202) Human TFR2 >NM\_003227.4 *Homo sapiens* transferrin  
receptor 2 (TFR2), transcript variant 1, mRNA

CTGATCTTCACTAGGGGCCCTTCTACTAGTGGGCTACGTCGCGCTTCCGAGGGGTCC  
TGCCAGGCGTGCGGAGACTCTGTGTTGGTGGTCAGTGAGGATGTCAACTATG  
AGCCTGACCTGGATTTCACCAGGGCAGACTCTACTGGAGCGACCTCCAGGC  
CATGTTCTGTCAGTTCTTGGGGGAGGGGCGCCTGGAGGACACCATCAGGCAA  
ACCAGCCTTCGGGAACGGGTGGCAGGCTCGGCCGGGATGGCCGCTCTGACTC  
AGGACATTTCGCGCGGCGCTCTCCCGCCAGAAGCTGGACCACGTGTGGACCGA  
CACGCACTACGTGGGGCTGCAATTCCCGGATCCGGCTCACCCCAACACCCTG  
CACTGGGTTCGATGAGGCCGGGAAGGTCGGAGAGCAGCTGCCGCTGGAGGACC  
CTGACGTCTACTGCCCTACAGCGCCATCGGCAACGTCACGGGAGAGCTGGT  
GTACGCCCACTACGGGCGGCCCCGAAGACCTGCAGGACCTGCGGGGCCAGGGGC  
GTGGATCCAGTGGGCGCCTGCTGCTGGTGCGCGTGGGGGTGATCAGCTTCG  
CCCAGAAGGTGACCAATGCTCAGGACTTCGGGGCTCAAGGAGTGCTCATATA  
CCCAGAGCCAGCGGACTTCTCCAGGACCCACCCAAGCCAAGCCTGTCCAGC  
CAGCAGGCAGTGTATGGACATGTGCACCTGGGAACCTGGAGACCCCTACACAC  
CTGGCTTCCCTTCCTTCAATCAAACCCAGTTCCCTCCAGTTGCATCATCAGG  
CCTTCCCAGCATCCCAGCCCAGCCCATCAGTGCAGACATTGCCTCCCGCCTG  
CTGAGGAAGCTCAAAGGCCCTGTGGCCCCCAAGAATGGCAGGGGAGCCTCC  
TAGGCTCCCCTTATCACCTGGGCCCCGGGCCACGACTGCGGCTAGTGGTCAA  
CAATCACAGGACCTCCACCCCATCAACAACATCTTCGGCTGCATCGAAGGC  
CGCTCAGAGCCAGATCACTACGTTGTCATCGGGGGCCAGAGGGATGCATGGG  
GCCAGGAGCAGCTAAATCCGCTGTGGGGACGGCTATACTCCTGGAGCTGGT  
GCGGACCTTTTCTCCATGGTGAGCAACGGCTTCCGGCCCCGCAGAAGTCTC  
CTCTTCATCAGCTGGGACGGTGGTGACTTTGGAAGCGTGGGCTCCACGGAGT  
GGCTAGAGGGCTACCTCAGCGTGCTGCACCTCAAAGCCGTAGTGTACGTGAG  
CCTGGACAACGCAGTGCTGGGGGATGACAAGTTTCATGCCAAGACCAGCCCC  
CTTCTGACAAGTCTCATTGAGAGTGTCTGAAGCAGGTGGATTCTCCCAACC  
ACAGTGGGCAGACTCTCTATGAACAGGTGGTGTTCACCAATCCCAGCTGGGA  
TGCTGAGGTGATCCGGCCCCCTACCCATGGACAGCAGTGCCTATTCCTTCACG  
GCCTTTGTGGGAGTCCCTGCCGTCGAGTTCTCCTTTATGGAGGACGACCAGG  
CCTACCCATTCTCTGCACACAAAGGAGGACACTTATGAGAACCTGCATAAGGT  
GCTGCAAGGCCGCTGCCCGCCGTGGCCCAGGCCGTGGCCCAGCTCGCAGGG  
CAGCTCCTCATCCGGCTCAGCCACGATCGCCTGCTGCCCTCGACTTCGGCC  
GCTACGGGGACGTCGTCCTCAGGCACATCGGGAACCTCAACGAGTTCTCTGG  
GGACCTCAAGGCCCGCGGGCTGACCCTGCAGTGGGTGTACTCGGCGCGGGGG  
GACTACATCCGGGCGGCGGAAAAGCTGCGGCAGGAGATCTACAGCTCGGAGG  
AGAGAGACGAGCGACTGACACGCATGTACAACGTGCGCATAATGCGGGTGGA  
GTTCTACTTCTTTCCAGTACGTGTCGCCAGCCGACTCCCCGTTCCGCCAC  
ATCTTCATGGGCCGTGGAGACCACACGCTGGGCGCCCTGCTGGACCACCTGC  
GGCTGCTGCGCTCCAACAGCTCCGGGACCCCCGGGGCCACCTCCTCCACTGG  
CTTCCAGGAGAGCCGTTTCCGGCGTCAGCTAGCCCTGCTCACCTGGACGCTG  
CAAGGGGCAGCCAATGCGCTTAGCGGGGATGTCTGGAACATTGATAACAAC  
TCTGAGGCCCTGGGGATCCTCACATCCCCGTCCCCCAGTCAAGAGCTCCTCT  
GCTCCTCGCTTGAATGATTCAGGGTCAGGGAGGTGGCTCAGAGTCCACCTCT  
CATTGCTGATCAATTTCTCATTACCCCTACACATCTCTCCACGGAGCCCAGA  
CCCCAGCACAGATATCCACACACCCAGCCCTGCAGTGTAGCTGACCCTAAT  
GTGACGGTCATACTGTCGGTTAATCAGAGAGTAGCATCCCTTCAATCACAGC  
CCCTTCCCCTTTCTGGGGTCTCCATACCTAGAGACCACTCTGGGAGGTTTG  
CTAGGCCCTGGGACCTGGCCAGCTCTGTTAGTGGGAGAGATCGCTGGCACCA  
TAGCCTTATGGCCAACAGGTGGTCTGTGGTGAAAGGGGCGTGGAGTTTCAAT  
ATCAATAAACCACCTGATATCAATAA (SEQ ID NO: 203) >NP\_003218.2

transferrin receptor protein 2 isoform 1 [*Homo sapiens*]  
**MERLWGLFQRAQQLSRPSSTVYQRVEGPRKGHLEEEEDGEEGAETLAHFC**  
**PMELRGPEPLGSRPRQPNLIPWAAAGRRAAPYLVLTAALLIFTGAFLLG YVAF**  
**RGSCQACGDSVLVSEDEVNYEPDLDFHQGRLYWSDLQAMFLQFLGEGRLEDT**  
IRQTSLRERVAGSAGMAALTQDIRAALSRQKLDHVWTDTHYVGLQFPDPAHP  
NTLHWVDEAGKVGEQLPLEDPDVYCPYSAIGNVTGELVYAHYGRPEDLQDLR  
ARGVDPVGRLLLVRVGVISFAQKVTNAQDFGAQGVLIYPEPADFSQDPPKPS  
LSSQQAVYGHVHLGTGDPYTPGFPSFNQTQFPPVASSGLPSIPAQPISADIA  
SRLLRKLKGPVAPQEWFQGSLLGSPYHLGPGPRLRLVVNNHRTSTPINNIFGC  
IEGRSEPDHYVVIGAQRDAWGPGAAKSAVGTAILELVRVTFSSMVSNGFRPR  
RSLLFISWDGGDFGSGVSTEWLEGYLSVLHLKAVVYVSLDNAVLGDDKFHAK  
TSPLLTSLIESVLKQVDSPNHSGQTLYEQVVFTNPSWDAEVIRPLPMDSSAY  
SFTAFVGVPAVEFSFMEDDQAYPFLHTKEDTYENLHKVLQGRLPAVAQAVAQ  
LAGQLLIRLSHDRLLPLDFGRYGDVVLRHIGNLNEFSGDLKARGLTQLQWVYS  
ARGDYIRAAEKLQRQEIYSSEERDERLTRMYNVRIMRVEFYFLSQYVSPADSP  
FRHIFMGRGDHTLGALLDHLRLRLRSNSSGTPGATSSTGFQESRFRRLALLT  
WTLQGAANALSGDVWNIDNNF (SEQ ID NO: 204) Human ADAM10  
>NM\_001110.4 *Homo sapiens* ADAM metalloproteinase domain 10 (ADAM10),  
transcript variant 1, mRNA

GTTGCCGGCCCCCTGAAGTGGAGCGAGAGGGAGGTGCTTCGCCGTTTCTCCTG  
CCAGGGGAGGTCCCGGCTTCCCGTGGAGGCTCCGGACCAAGCCCCTTCAGCT  
TCTCCCTCCGGATCGATGTGCTGCTGTTAACCCGTGAGGAGGCGGCGGCGGC  
GGCAGCGGCAGCGGAAGATGGTGTGCTGAGAGTGTTAATTCTGCTCCTCTC  
CTGGGCGGCGGGGATGGGAGGTCAGTATGGGAATCCTTTAAATAAATATATC  
AGACATTATGAAGGATTATCTTACAATGTGGATTCATTACACCAAAAACACC  
AGCGTGCCAAAAGAGCAGTCTCACATGAAGACCAATTTTTACGTCTAGATTT  
CCATGCCCATGGAAGACATTTCAACCTACGAATGAAGAGGGACACTTCCCTT  
TTCAGTGATGAATTTAAAGTAGAAACATCAAATAAAGTACTTGATTATGATA  
CCTCTCATATTTACACTGGACATATTTATGGTGAAGAAGGAAGTTTTAGCCA  
TGGGTCTGTTATTGATGGAAGATTTGAAGGATTCATCCAGACTCGTGGTGGC  
ACATTTTATGTTGAGCCAGCAGAGAGATATATTAAAGACCGAACTCTGCCAT  
TTCACTCTGTCAATTTATCATGAAGATGATATTAACCTATCCCCATAAATACGG  
TCCTCAGGGGGGCTGTGCAGATCATTGAGTATTTGAAAGAATGAGGAAATAC  
CAGATGACTGGTGTAGAGGAAGTAACACAGATACCTCAAGAAGAACATGCTG  
CTAATGGTCCAGAACTTCTGAGGAAAAAACGTACAACCTCAGCTGAAAAAAA  
TACTTGTCAGCTTTATATTCAGACTGATCATTTGTTCTTTAAATATTACGGA  
ACACGAGAAGCTGTGATTGCCCAGATATCCAGTCATGTTAAAGCGATTGATA  
CAATTTACCAGACCACAGACTTCTCCGGAATCCGTAACATCAGTTTCATGGT  
GAAACGCATAAGAATCAATACAACTGCTGATGAGAAGGACCCTACAAATCCT  
TTCCGTTTCCCAAATATTGGTGTGGAGAAGTTTCTGGAATTGAATTCTGAGC  
AGAATCATGATGACTACTGTTTGGCCTATGTCTTCACAGACCGAGATTTTGA  
TGATGGCGTACTTGGTCTGGCTTGGGTTGGAGCACCTTCAGGAAGCTCTGGA  
GGAATATGTGAAAAAAGTAACTCTATTCAGATGGTAAGAAGAAGTCCTTAA  
ACACTGGAATTATTACTGTTTCAAGAACTATGGGTCTCATGTACCTCCCAAAGT  
CTCTCACATTACTTTTGCTCACGAAGTTGGACATAACTTTGGATCCCCACAT  
GATTCTGGAACAGAGTGCACACCAGGAGAATCTAAGAATTTGGGTCAAAAAG  
AAAATGGCAATTACATCATGTATGCAAGAGCAACATCTGGGGACAACTTAA  
CAACAATAAATTCTCACTCTGTAGTATTAGAAATATAAGCCAAGTTCTTGAG  
AAGAAGAGAAACAACCTGTTTTGTTGAATCTGGCCAACCTATTTGTGGAAATG  
GAATGGTAGAACAAGGTGAAGAATGTGATTGTGGCTATAGTGACCAGTGTA



AGATGAATGATGCTGCTTCCGATGCAAAATCAACACAGAGGAGGAAGAAAGAAACTG  
AAACCTGGGAAACAGTGCAGTCCAAGTCAAGGTCCTTGTTGTACAGCACAGT  
GTGCATTCAAGTCAAAGTCTGAGAAGTGTCGGGATGATTCAGACTGTGCAAG  
GGAAGGAATATGTAATGGCTTCACAGCTCTCTGCCCAGCATCTGACCCTAAA  
CCAAACTTCACAGACTGTAATAGGCATACACAAGTGTGCATTAATGGGCAAT  
GTGCAGGTTCTATCTGTGAGAAATATGGCTTAGAGGAGTGTACGTGTGCCAG  
TTCTGATGGCAAAGATGATAAAGAATTATGCCATGTATGCTGTATGAAGAAA  
ATGGACCCATCAACTTGTGCCAGTACAGGGTCTGTGCAGTGGAGTAGGCACT  
TCAGTGGTCGAACCATCACCTGCAACCTGGATCCCCTTGCAACGATTTTAG  
AGGTTACTGTGATGTTTTTCATGCGGTGCAGATTAGTAGATGCTGATGGTCCT  
CTAGCTAGGCTTAAAAAAGCAATTTTTTAGTCCAGAGCTCTATGAAAACATTG  
CTGAATGGATTGTGGCTCATTGGTGGGCAGTATTACTTATGGGAATTGCTCT  
GATCATGCTAATGGCTGGATTTATTAAGATATGCAGTGTTTCATACTCCAAGT  
AGTAATCCAAAGTTGCCTCCTCCTAAACCACTTCCAGGCACTTTAAAGAGGA  
GGAGACCTCCACAGCCCATTTCAGCAACCCCAGCGTCAGCGGCCCCCGAGAGAG  
TTATCAAATGGGACACATGAGACGCTAACTGCAGCTTTTGCCTTGGTTCTTC  
CTAGTGCCTACAATGGGAAAACCTTCACTCCAAAGAGAAACCTATTAAGTCAT  
CATCTCCAAACTAAACCCTCACAAGTAACAGTTGAAGAAAAAATGGCAAGAG  
ATCATATCCTCAGACCAGGTGGAATTACTTAAATTTTAAAGCCTGAAAATTC  
CAATTTGGGGGTGGGAGGTGGAAAAGGAACCCAATTTTCTTATGAACAGATA  
TTTTTAACTTAATGGCACAAGTCTTAGAATATTATTATGTGCCCCGTGTTC  
CCTGTTCTTCGTTGCTGCATTTTCTTCACTTGCAGGCCAACTTGGCTCTCAA  
TAACTTTTACCACAAATTGAAATAAATATATTTTTTTTCAACTGCCAATCAA  
GGCTAGGAGGCTCGACCACCTCAACATTGGAGACATCACTTGCCAATGTACA  
TACCTTGTTATATGCAGACATGTATTTCTTACGTACACTGTACTTCTGTGTG  
CAATTGTAAACAGAAATTGCAATATGGATGTTTCTTTGTATTATAAAATTTT  
TCCGCTCTTAATTAAAAATTACTGTTTAATTGACATACTCAGGATAACAGAG  
AATGGTGGTATTCAGTGGTCCAGGATTCTGTAATGCTTTACACAGGCAGTTT  
TGAAATGAAAATCAATTTACCTTTCTGTTACGATGGAGTTGGTTTTGATACT  
CATTTTTTCTTTATCACATGGCTGCTACGGGCACAAGTGACTATACTGAAGA  
ACACAGTTAAGTGTTGTGCAAACTGGACATAGCAGCACATACTACTTCAGAG  
TTCATGATGTAGATGTCTGGTTTCTGCTTACGTCTTTTAACTTTCTAATTC  
AATTCCATTTTTCAATTAATAGGTGAAATTTTATTCATGCTTTGATAGAAAT  
TATGTCAATGAAATGATTCTTTTTTATTTGTAGCCTACTTATTTGTGTTTTTC  
ATATATCTGAAATATGCTAATTATGTTTTCTGTCTGATATGGAAGAAAG  
CTGTGTCTTTATCAAAATATTTAAACGGTTTTTTTCAGCATATCATCACTGAT  
CATTGGTAACCACTAAAGATGAGTAATTTGCTTAAGTAGTAGTTAAAATTGT  
AGATAGGCCTTCTGACATTTTTTTTCTTAAATTTTTTAAACAGCATTGAAGGT  
GAAACAGCACAAATGTCCCATTCCAAATTTATTTTTGAAACAGATGTAAATAA  
TTGGCATTTTTAAAGAGAAAGCAAAAACATTTAATGTATTAACAGGCTTATTG  
CTATGCAGGAAATAGAAGGGGCATTACAAAAATTGAAGCTTGTGACATATTT  
ATTGCTTCTGTTTTTCCAACCTACATCACTTCAACTAGAAGTAAAGCTATGATT  
TTCCTGACTTCACATAGGAGGCAAATTTAGAGAAAGTTGTAAAGATTTCTAT  
GTTTTGGGTTTTTTTTTTTTCCTTTTTTTTTTTAAGAGTATAAGGTTTACACA  
ATCATTCTCATAATGTGACGCAAGCCAGCAAGGCCAAAAATGCTAGAGAAAA  
TAACGGGATCTCTTCCTTGTAACCTTGTACAGTATGTGGTGACTTTTTTCAA  
ATACAGCTTTTTGTACATGATTTAGAGACAAATTTGTACATGAAACCCAG  
ATAGACTATAAATAATTCTAAACAAACAAGTAGGTAGATATGTATGTAATTG  
CTTTTAAATCATTAAATGCCTTTGTTTTTGGACTGTGCAAAGGTTGGAAGT  
GGGTTTGCATTTCTAAATGGTGACTTTTATTCTGCAAGAGTTCTTAGTAAC

TTCTTGAGTGTGGTAGGATTTTGGCAGGTACACACACTGCCCTATTTTATTTTGAGT  
TCCTGTGGTAGGATTTTGGCAGGTACACACACTGCCCTATTTTATTTTGAGT  
CTAAGTTAAATGTTTTCTGAAAAGAGATACATGCACTGAACTCTTTCCACTG  
CGAATCAAGATGTGGTAATATAAAAGGATCAAGACAAATGAGATCTAATACT  
ACTGTCAGTTTTAATGTCCACTGTGTTTTATACAGTATCTTTTTTTTGTTCAC  
TTTGGAATTTTTACTAAAAATTGCAAAAAATAAAGTATTGTGCAAAGATGT  
AAGGTTTTTTGAACTTGAAATGCATTAATAAATAGACGATTAAATCAACTT  
GAAGGTTCTATACTCTTTGAACTCTGAGAACTATCACAAGAAGCTTCCCACA  
AGGCAGTGTTTTCTTACAGTTGTCTCTTCCCTACAAAAGTATAGATTATCTTT  
ATTCTTAATACTTTGGAATCCATGTAGAAAATTTCCAGTTAGATACTCTGCG  
TACACACAATAAACCTTTTTTAAAACACCCCACTAATCTCAACTGCATTACAT  
TGTTTCTAATCAATATTCAGTGCTTGTCTTGGTGGAAGAGGTGAGTCATTTT  
GAAAACCTTATGGTCTTGTTTTTATGTGTTTTTCAAAGTTTTGAATGCTAAGT  
ACCTCATTTATTTTAAAAAGCCTAGTTTAATGATAAGTTTTGTTTAAAAATTTT  
GAGCCATCATTTTTTCTCTTCATAGCAAATAAGGAGAGAATTGACATTTTCAGT  
GTTACCTAGAAAAGGAATTGTAAGCCCAGAATAATTCCTGCGATGAGGTAAT  
CTGCTTCAAATTCTTTTTTTTAGTCAAGGTTAGCTATAAGTAATACTTGTTAA  
ATGAGTAAATATGTAATACTTTGTGAATTACTTTGTTAATTTAGGAGCATCA  
AATGTATATTATGTTTAGTTATTTATGAAACTCTCAATATTGATTGATTTGG  
GTAATTATAAATTAGTTATTTTTTACTTGTAATTGAATGCTTAAATTCTGTTT  
ACAGTCCGTCCTCTCTCCCTCCATCCCTCCCTCCCCAGTTTTATAAATTCAG  
GTACCAATTCACAAACAAAATCAGAAATAAAATAAATTTATTGACTGCTTCT  
GGATTTAGCATTCCTGTAGTGTCAGCAATGTCATGCAGTTTGGGGAAGCA  
TTTATTTAAGGAAATGACAACCTTCTCTGATCAGTCTTGTTTTGTGAGGTGT  
CTTCAACACTTTTATGCTTTGGGTACTTCGTGTTTGTGACAGTCTTAGGATAG  
TGAAATCTGATTTGTCCAAGCGGAGCAAACACTACTCGACCCTCAGTCCTTGTA  
TTTGTCCCTGTAGTAAGACCTAATTATTATTATTTCTTAAAGATGGGATTGG  
TGTCCTTGGCAACTATGAAATTTTCGGGGCTTGTGCATGAGAAGGCATTTCTT  
ATTAAGTATTTCTAATTGAAGGTATCAGAGTGTCAGCATTACAAACCTGGA  
CAGTTCACCTGGAGGAGTACAAGAAGAGATATTCATTATCCATATTTAAAGG  
GTCAAGGTTTCCCAAACCAAGGGTGCAAGCCAGATGTAGTTTTAAAGCAGCT  
GCCAGGGACAGTTCATCTTTAGAGAAGTCACTAAAGTTGTAAGAAATTTTAG  
TTTCCCCAAAACCACTTTCAACTTCTTAGAACTAGAAAGACAATTGGTTTTG  
CCCCACAGAGGACAACCTTCAGTTTCAGCATCTCTCATGTTGTGTTCTTGATT  
AAAAACAACCTTCCATTTGATATACTTTTCCGTTTATTACCAGTTTAGTTTTT  
TCACTATTGTTTCTGTATTCAACTCTTTATATGATTAGGATAGAAATTTAGC  
CCTTCTGTTTTATATTACTATATTGTTTGTGTGTCTTAGATATATACATGTA  
TGTAATTTTTTTCAGTAGAAATTCATGTATTTTATAATTGGTAAGTTCTTCAG  
AGCATCTCTTCTATAAAAAGCAACAGGATGCTAGGTAAAACGGAGCATTGAG  
CAAAATACTGATTAGTTTTTTGCTTTTTTCTGAAATCTACACTAAAGTGATAG  
GGTGTGGGGTAATCCAACAAGGACAAGGTGAATTGAACAAGAACGAAATCTG  
GAAGCAGATGAAGGAGTACTATTGATTGGGCAGACCCAGGGAAGTCAAATCC  
TAAACCAGCAGTGGGAACACAACAGAATGGTGTAGTTTGCAGTGGTAAGATT  
TGGGTACCTGGCAGGGCTGGGTGCGGTGGCTCACACCTGTAATCCCAGCACT  
TTGGGAGGCCAAGGCGGGTGGATCACTTGAGGTCAGGAGTTCGAGACCACCC  
TGGCCAACATGGTGAAACCCCGTCTCTACTAAAAATAACAGCTGGGCGTGGTG  
GCACATGCTTGTAATCCCAGCTACTCGGGAGGCAGAGGCAGGAGATTTGCTT  
GAACCCGGGAGGCAGAGGCAGGAGATTTGCTTGAACCCAGGAGGCAGAGGCT  
GCAGTGAGCCGCGATTGCGCCATTGCACTCCAGCCTGGGTGACAGAGCGAGA  
CTCTGTCTCAA

AATAATTGAGTACCTGGCCTTTGTTACTTTTTTTCTATGTGTGTGACAAAA  
 ACATAATATGCACACTTTTGTAAACCCACCTTTCTCATTTAATGGTACATTGA  
 TAATGTATATCACATTAATACTACTCTAAATATTTCTGTGGATGTATGTTTTT  
 TTTTTCTTAACCAATTTCCCATTGTTTGGACATGTAGGTTCCACATTGTTTA  
 TTATTTTAAACAATTCTAAAGAATTTTAAACAATTCTTAGGAAAATCCTCAG  
 CCTAATAATGAAATTAATTCCTAAAAGTGGAATTGTTGGGGTAAAGGTTTTT  
 TGAGGGACATTGATAAAAATTATGGTACTGTCTCCCAGATAGATGTACCAAG  
 TTATACTACCACGATTTAATATATATATATATATATATTAATCAGAGTCCC  
 ATCCTTAGAAATCCACATATATGCAGCCACATGAATGTATTAGAAACAATAA  
 TAGAAGACTCATGCTTAATTCAGTTGATTAGCTTTAGACATAATTCAAATGC  
 AAGTCAAATTGAGTGGCCTAATTGTGGTCTCTTAAGTACCATTTTTCTTCAA  
 GGAACCAGACTCCTTTGGATAAATCACTAATTCCACCTGTAAGAAAGAAAT  
 GTACAAGAAGAACCTAGGAAACATTGTTTTGTACCAGATCAGAAAGATTGAG  
 GAGGCACCTTAGAAGTTACCACTGGCCAAGGCTGAGATAACTTTAGCATCAG  
 CAAGGATAATATCTGCAAGAGATTGAAACTCATAGTATTGTATTTAACTCTG  
 TGAGTTAATGATGGTAGTGGACAGAATTATAGTTACCTTTGGGATACGCTTT  
 TAAAGAAATTCCAGGTAATAAGAGAAATGATAGAATTAGGATATCACCATT  
 TACCCCCCAACAATTTATGGATCTAGACAATAATCGCCAGTGAAGTCTAAC  
 CTCACAAAGTGAGAGCAATCAGATTTTGTGCCTCCTAATGGAAGTACATATA  
 CCACCTATGAAGCAGTTCTGCCAAAAGTCACATCTCATCATGATGAAGCCTC  
 CTGATCTAACTACCCCTTCATTAGAAATACAGGGGACAGAGGGACAAATAAT  
 ATACAAGGGACTCAATCAGCAAAATCCAGACTCTGGAAAACCTACAAGACATA  
 TGGTCCTGCTTCAACAACAGAAATGCAAAGAGAAAAGACAACGATGGGTAA  
 AGGAGACTTAAGAGCTACATCTATCAAGACAATTTATGGACTTATTTGGATA  
 CTGATTTGAACAACTGTTGAGACCATTGGAAAAATGTGAAAAGTGGATATT  
 TGATATTAAGGTTTTTAAATTATTTTATAGGTGTGATAATGGTATTGTTACATT  
 TTTTAAAGGACCCCTTTTAGAGATGCAAATTGAAACACTTAAAAAATGAAAT  
 GATACGATGTATAAGTTTTTGTCTTAAAAATAAGGATTGAAGTTGGCTGGTGT  
 GTGTGGATATAGTTGAAACAAGATTGGCTGTGAGTTGATAATTATTGAAGCT  
 GGGTGATGGGCACTTGGGGATTTATTATACTATTTTCTCTACCTGTGTTTAT  
 ATTTGAAATTTTTTCATAGAAGTTTTTAAATGTGGCCAGTTGTGATGGCTCAT  
 ACCTGTAATCCCAACACTTTGGGAGGCCAAGGTGGGAGGATCACTTGAGCTC  
 AGGAGTTAGAGACCAGCCTGGGCAAAATAGTGAGACTCCATCTCAAAGAAAA  
 AAAAAAAGTGTTTTAAATGTGAATCAAATTCCTATAGAAGCTGATTCATTAC  
 TGTTTTTATTTTAGCAGTAATTCATGATAATGACCTGTATTCATAATGATTT  
 TCATAATGATTGTTTTAGTGGAATTAACTTGAACCAGTCAAGCTAACATAA  
 TTATATTCTGCTCCAGTTACAATGAATAATTAATTGATTTCAACTGCTAGGG  
 TGAAGTCTTGAAGCTATCAGTCATCCAGCAATCTTAGCAAGCAGGCCATTGG  
 GTCCTGTGTTGCTCTGTCTCTCTCTCTCTCTCTCACTGTTGAAGGGCTTAGC  
 TAACTACTTAAGTAAAATATTTGTTCTCTGTAAACATGTCAAGGAGTATGG  
 TCAGCTTATCCACATTAAGCCTGTGTGTCCCACGTTGGAGTAAATGTTAAGT  
 AGCTCACTACAATAAACTAGATTCTTCTGCCCTCTCTTGTTTAAATGATCAT  
 GTTCCCTGGAGGTGGAAATAGATCTTTAAAAAGATATTCTGTAGTTGTTTGT  
 TCTCAGTGTAAAAAAATGAGAATAATTTGATAAGAGTGTAGGTTGTCTTATA  
 TAAAAAGTGTTCCATTTGCATGAATTTTAGAAAAATCATTTTGGAAAAATG  
 AAGGCTATGTGGTTATACTGAACACATTAAGCAATTTTATTCTTTATTTTAA  
 ATGAATATTTTATTATCGTTTTCTTCCCTTGCCCTTTGGGTATGGGAGTTAG  
 CCTTTGTGTTTCTAAATACAACAGGCCGGTTTTTATAAATTAAGGTGTCAAT  
 ATATTCTTCATTATTTAGTTTTGTGATTGTGGTTAGTTTTCATTTTTCTTAA  
 GTATCTGCTAGTAGCATCTGTAATTAAGTGAAGTGACCTGTTAACCATTTTC

disintegrin and metalloproteinase domain-containing protein 10 isoform 1  
preproprotein [*Homo sapiens*]

**IQQPQRQRPRESYQMGHMRR** (SEQ ID NO: 206) Transmembrane >NM\_001769.4  
*Homo sapiens* CD9 molecule (CD9), domain 2 or transcript variant 1, mRNA  
transmembrane

AGCCGCCTGCATCTGTATCCAGCGCCAGGTCCCGCCAGTCCCAGCTGCGCGC domain  
3 from GCCCCCAGTCCCGCACCCGTTCTGGCCCAGGCTAAGTTAGCCCTCACCATGC  
Human CD9

CGGTCAAAGGAGGCACCAAGTGCATCAAATACCTGCTGTTTCGGATTTAACCTT  
CATCTTCTGGCTTGCCGGGATTGCTGTCCTTGCCATTGGACTATGGCTCCGA  
TTCGACTCTCAGACCAAGAGCATCTTCGAGCAAGAACTAATAATAAATT  
CCAGCTTCTACACAGGAGTCTATATTCTGATCGGAGCCGGCGCCCTCATGAT  
**GCTGGTGGGCTTCCTGGGCTGCTGCGGGGCTGTGCAGGAGTCCCAGTGCATG**  
CTGGGACTGTTCTTCGGCTTCCTCTTGGTGATATTCGCCATTGAAATAGCTG  
CGGCCATCTGGGGATATTCCCACAAGGATGAGGTGATTAAGGAAGTCCAGGA  
GTTTTACAAGGACACCTACAACAAGCTGAAAACCAAGGATGAGCCCCAGCGG  
GAAACGCTGAAAGCCATCCACTATGCGTTGAACTGCTGTGGTTTGGCTGGGG  
GCGTGGAACAGTTTATCTCAGACATCTGCCCCAAGAAGGACGTACTCGAAAC  
CTTCACCGTGAAGTCCTGTCCTGATGCCATCAAAGAGGTCTTCGACAATAAA  
TTCCACATCATCGGCGCAGTGGGCATCGGCATTGCCGTGGTCATGATATTTG  
GCATGATCTTCAGTATGATCTTGTGCTGTGCTATCCGCAGGAACCGCGAGAT  
GGTCTAGAGTCAGCTTACATCCCTGAGCAGGAAAGTTTACCCATGAAGATTG  
GTGGGATTTTTTGTGTTTGTGTTTGTGTTTGTGTTTGTGTTTGTGTTTGTGTTT  
TTTTGCCACTAATTTTAGTATTCATTCTGCATTGCTAGATAAAAGCTGAAGT  
TACTTTATGTTTGTCTTTTAATGCTTCATTCAATATTGACATTTGTAGTTGA  
GCGGGGGGTTTGGTTTGTCTTTGGTTTATATTTTTTCAGTTGTTTGTGTTTGC  
TTGTTATATTAAGCAGAAATCCTGCAATGAAAGGTACTATATTTGCTAGACT  
CTAGACAAGATATTGTACATAAAAGAATTTTTTTGTCTTTAAATAGATACAA  
ATGTCTATCAACTTTAATCAAGTTGTAACCTTATATTGAAGACAATTTGATAC  
ATAATAAAAAATTATGACAATGTCCTGGA (SEQ ID NO: 207) >NP\_001760.1

CD9 antigen isoform 1 [*Homo sapiens*]

MPVKGGTKCIKYLFLGFNFIFWLAGIAVLAIGLWLRFDSQTKSIFEQETNNN  
NSSFYTGVIILIGAGALMMLVGFLGCCGAVQESQCMLGLFFGFLLVIFAIEI  
AAAIWGYSHKDEVIKEVQEFYKDTYNKLKTKDEPQRETLKAIHYALNCCGLA  
GGVEQFISDICPKKDVLETFTVKSCPDAIKEVFDNKFHIIGAVGIGIAVVM

FGMIFSMILCCAIRRNREMV (SEQ ID NO: 208) Human CD298 >NM\_001679.4

*Homo sapiens* ATPase Na<sup>+</sup>/K<sup>+</sup> transporting subunit beta 3 (ATP1B3), mRNA  
AGTCGGCTCGAGTACTCCCCGTAACGAGGAGGTGTTCTCGGCCGTCCCACCC  
TTCAGTCCCGTCTCCGGGCTGCGCCGCCGGAGCCGGGACGCGCCTCCGCAGC  
CCTCGCCGCCTCCATCCCCGCGGCCGCAGCTCCTCTCGCCGTCCGCGCGCAC  
ACCATGACGAAGAACGAGAAGAAGTCCCTCAACCAGAGCCTGGCCGAGTGGA  
AGCTCTTCATCTACAACCCGACCACCGGAGAATTCCTGGGGCGCACCGCCAA  
GAGCTGGGGTTTGATCTTGCTCTTCTACCTAGTTTTTTATGGGTTCCTGGCT  
GCACTCTTCTCATTACGATGTGGGTTATGCTTCAGACTCTCAACGATGAGG  
TTCCAAAATACCGTGACCAGATTCCTAGCCCAGGACTCATGGTTTTTCCAAA  
ACCAGTGACCGCATTGGAATATACATTCAGTAGGTCTGATCCAACCTTCGTAT  
GCAGGGTACATTGAAGACCTTAAGAAGTTTCTAAAACCATATACTTTAGAAG  
AACAGAAGAACCTCACAGTCTGTCCTGATGGAGCACTTTTTGAACAGAAGGG  
TCCAGTTTATGTTGCATGTCAGTTTCCTATTTCACTTCAAGCATGCAGT  
GGTATGAATGATCCTGATTTTGGCTATTCTCAAGGAAACCCTTGATTTCTTG  
TGAAAATGAACAGAATAATTGGATTAAAGCCTGAAGGAGTGCCAAGGATAGA  
TTGTGTTTCAAAGAATGAAGATATACCAAATGTAGCAGTTTATCCTCATAAT

GGAAATGATAGATAGCTTAAATAATTTCCCATATTATGGGAAAAAACCTGCATGTTG  
GGTATCTACAGCCATTGGTTGCTGTTTCAGGTCAGCTTTGCTCCTAACAACAC  
TGGGAAAGAAGTAACAGTTGAGTGCAAGATTGATGGATCAGCCAACCTAAAA  
AGTCAGGATGATCGTGACAAGTTTTTTGGGACGAGTTATGTTCAAAATCACAG  
CACGTGCATAGTATGAGTAGGATATCTCCACAGAGTAAATGTTGTGTTGTCT  
GTCTTCATTTTGTAAACAGCTGGACCTTCCATTCTAGAATTATGAGACCACCT  
TGGAGAAAGGTGTGTGGTACATGACATTGGGTTACATCATAACGTGCTTCCA  
GATCATAGTGTTTCAGTGTCTCTGAAGTAACTGCCTGTTGCCTCTGCTGCCC  
TTTGAACCAGTGACAGTCGCCAGATAGGGACCGGTGAACACCTGATTCCAA  
ACATGTAGGATGGGGGTCTTGTCCTCTTTTTTATGTGGTTTAATTGCCAAGTG  
TCTAAAGCTTAATATGCCGTGCTATGTAAATATTTTATGGATATAACAACCTG  
TCATATTTTGATGTCAACAGAGTTTTAGGGATAAAATGGTACCCGGCCAACA  
TCAAGTGACTTTATAGCTGCAAGAAATGTGGTATGTGGAGAAGTTCTGTATG  
TGAGGAAGGAAAAAAGAAAAATAAAAGTGTGTTTGAAAAATATTATCTTGGG  
TTCTTTGTAAAATTTATTTTTTACATGCTGAATTAGCCTCGATCTTTTTTGAT  
TAAGAGCACAACTTTTTTTGTAAACATGTAAAAAATAAACTGGGATTA  
ATTTTTAGTGTTGGAAGTGCCTCTTATTTTAGGCTGTAGATAAAATAGCATT  
TTTAGGTTAGCCAGTGTGACTATGCACCTAATTTTTTATGAGATTAAATTCA  
TAAGACTTAATTTGTACAATAGTTTGTGAAATATCTTGTTACTGCTTTTATT  
TAGCAGACTGTGGACTGTAATAAAGTATATAAATTGTGAAATATAAAAACTT  
GGAAGTTATTCAAAGCTTCAAAGCAAA (SEQ ID NO: 209) >NP\_001670.1  
sodium/potassium-transporting ATPase subunit beta-3 [*Homo sapiens*]  
MTKNEKKS LNQSLAEWKLFIYNPTTGEFLGR TAKSWGLILLFYLVFYGFLAA  
LFSFTMWV MLQTLNDEV PKYRDQIPSPGLMVFPKPVTALEYTF SRSDPTSYA  
GYIEDLKKFLKPYTLEEQKNLTVCPD GALFEQKGPVYVACQFPISLLQACSG  
MNDPDFGYSQGNPCILVKMNRIIGLKPEGVPRIDCVSKNEDIPNVAVYPHNG  
MIDLKYFPYYGKKLHVGYLQPLVAVQVSFAPNNTGKEVTVECKIDGSANLKS  
QDDRDKFLGRVMFKITARA (SEQ ID NO: 210) Lipid affinity tag >NM\_004985.5  
*Homo sapiens* KRAS proto-oncogene, modified from GTPase (KRAS), transcript  
variant b, mRNA Human KRAS  
CTAGGCGGCGGCCGCGGCGGCGGAGGCAGCAGCGGCGGCGGCAGTGGCGGCG  
GCCAAGGTGGCGGCGGCTCGGCCAGTACTCCCGGCCCCCGCCATTTCGGACT  
GGGAGCGAGCGCGGCGCAGGCACTGAAGGCGGCGGCGGGGCCAGAGGCTCAG  
CGGCTCCCAGGTGCGGGAGAGAGGCCTGCTGAAAATGACTGAATATAAACTT  
GTGGTAGTTGGAGCTGGTGGCGTAGGCCAAGAGTGCCTTGACGATACAGCTAA  
TTCAGAATCATTTTGTGGACGAATATGATCCAACAATAGAGGATTCCTACAG  
GAAGCAAGTAGTAATTGATGGAGAAACCTGTCTCTTGATATTCTCGACACA  
GCAGGTCAAGAGGAGTACAGTGCAATGAGGGACCAGTACATGAGGACTGGGG  
AGGGCTTTCTTTGTGTATTTGCCATAAATAATACTAAATCATTTGAAGATAT  
TCACCATTATAGAGAACAAATTAAGAGTTAAGGACTCTGAAGATGTACCT  
ATGGTCCTAGTAGGAAATAAATGTGATTTGCCTTCTAGAACAGTAGACACAA  
AACAGGCTCAGGACTTAGCAAGAAGTTATGGAATTCCTTTTATTGAAACATC  
AGCAAAGACAAGACAGGGTGTTGATGATGCCTTCTATACATTAGTTTCGAGAA  
ATTGAAAACATAAAGAAAAGATGAGCAAAGATGGTAAAAAGAAGAAAAAGA  
**AGTCAAAGACAAAGTGTGTAATTATGTAAATACAATTTGTACTTTTTTCTTA**  
AGGCATACTAGTACAAGTGGTAATTTTTGTACATTACACTAAATTATTAGCA  
TTTGTTTTAGCATTACCTAATTTTTTTCTGCTCCATGCAGACTGTTAGCTT  
TTACCTTAAATGCTTATTTTAAAATGACAGTGGAAGTTTTTTTTTCTCTAA  
GTGCCAGTATTCCCAGAGTTTTGGTTTTTGAAGTACCAATGCCTGTGAAAAA  
GAAACTGAATACCTAAGATTTCTGTCTTGGGGCTTTTGGTGCATGCAGTTGA

TTACTTCTTATTCTTTCTTACCAATTGTAAGATGTTGGTGTGAAACAAATTAAT  
GAAGCTTTTGAATCATCCCTATTCTGTGTTTTATCTAGTCACATAAATGGAT  
TAATTACTAATTTTCAGTTGAGACCTTCTAATTGGTTTTTACTGAAACATTGA  
GGGAACACAAATTTATGGGCTTCCTGATGATGATTCTTCTAGGCATCATGTC  
CTATAGTTTGTTCATCCCTGATGAATGTAAAGTTACACTGTTTACAAAGGTTT  
TGTCTCCTTTCCACTGCTATTAGTCATGGTCACTCTCCCCAAAATATTATAT  
TTTTTCTATAAAAAGAAAAAAATGGAAAAAAATTACAAGGCAATGGAAACTA  
TTATAAGGCCATTTTCCTTTTTCACATTAGATAAATTACTATAAAGACTCCTAA  
TAGCTTTTCCCTGTTAAGGCAGACCCAGTATGAAATGGGGATTATTATAGCAA  
CCATTTTGGGGCTATATTTACATGCTACTAAATTTTTTATAATAATTGAAAAG  
ATTTTAACAAGTATAAAAAAATTCTCATAGGAATTAAATGTAGTCTCCCTGTG  
TCAGACTGCTCTTTCATAGTATAACTTTAAATCTTTTCTTCAACTTGAGTCT  
TTGAAGATAGTTTTAATTCTGCTTGTGACATTAAAAGATTATTTGGGCCAGT  
TATAGCTTATTAGGTGTTGAAGAGACCAAGGTTGCAAGGCCAGGCCCTGTGT  
GAACCTTTGAGCTTTCATAGAGAGTTTTCACAGCATGGACTGTGTCCCCACGG  
TCATCCAGTGTTGTCATGCATTGGTTAGTCAAATGGGGAGGGACTAGGGCA  
GTTTGGATAGCTCAACAAGATACAATCTCACTCTGTGGTGGTCTCTGCTGACA  
AATCAAGAGCATTGCTTTTGTTCCTTAAGAAAACAACTCTTTTTTAAAAAT  
TACTTTTAAATATTAACCTCAAAGTTGAGATTTTGGGGTGGTGGTGTGCCAA  
GACATTAATTTTTTTTTTAAACAATGAAGTGAAAAAGTTTTACAATCTCTAG  
GTTTGGCTAGTTCTCTTAACACTGGTTAAATTAACATTGCATAAACACTTTT  
CAAGTCTGATCCATATTTAATAATGCTTTAAAATAAAAAACAATCCT  
TTTGATAAATTTAAAATGTTACTTATTTTAAAATAAATGAAGTGAGATGGCA  
TGGTGAGGTGAAAGTATCACTGGACTAGGAAGAAGGTGACTTAGGTTCTAGA  
TAGGTGTCTTTTAGGACTCTGATTTTGAGGACATCACTTACTATCCATTTCT  
TCATGTTAAAAGAAGTCATCTCAAACCTCTTAGTTTTTTTTTTTACAACAT  
GTAATTTATATTCCATTTACATAAGGATACACTTATTTGTCAAGCTCAGCAC  
AATCTGTAAATTTTAAACCTATGTTACACCATCTTCAGTGCCAGTCTTGGGC  
AAAATTGTGCAAGAGGTGAAGTTTATATTTGAATATCCATTCTCGTTTTAGG  
ACTCTTCTTCCATATTAGTGTCATCTTGCCCTCCCTACCTTCCACATGCCCCA  
TGACTTGATGCAGTTTTAATACTTGTAATTTCCCCTAACCATAAGATTTACTG  
CTGCTGTGGATATCTCCATGAAGTTTTCCCCTGAGTCACATCAGAAATGCC  
CTACATCTTATTTCCCTCAGGGCTCAAGAGAATCTGACAGATACCATAAAGGG  
ATTTGACCTAATCACTAATTTTCAGGTGGTGGCTGATGCTTTGAACATCTCT  
TTGCTGCCCCAATCCATTAGCGACAGTAGGATTTTTCAAACCTGGTATGAATA  
GACAGAACCCTATCCAGTGGAAGGAGAATTTAATAAAGATAGTGCTGAAAGA  
ATTCCTTAGGTAATCTATAACTAGGACTACTCCTGGTAACAGTAATACATTC  
CATTGTTTTAGTAACCAGAAATCTTCATGCAATGAAAAATACTTTAATTCAT  
GAAGCTTACTTTTTTTTTTGGTGTGAGAGTCTCGCTCTTGTACCCAGGCT  
GGAATGCAGTGGCGCCATCTCAGCTCACTGCAACCTCCATCTCCCAGGTTCA  
AGCGATTCTCGTGCCTCGGCCTCCTGAGTAGCTGGGATTACAGGCGTGTGCC  
ACTACACTCAACTAATTTTTGTATTTTAGGAGAGACGGGGTTTCACCCTGT  
TGGCCAGGCTGGTCTCGAACTCCTGACCTCAAGTGATTACCCACCTTGGCC  
TCATAAACCTGTTTTGCAGAACTCATTATTTCAGCAAATATTATTGAGTGC  
CTACCAGATGCCAGTCACCACACAAGGCACTGGGTATATGGTATCCCCAAC  
AAGAGACATAATCCCGGTCCTTAGGTAGTGCTAGTGTGGTCTGTAATATCTT  
ACTAAGGCCTTTGGTATACGACCCAGAGATAACACGATGCGTATTTTAGTTT  
TGCAAAGAAGGGGTTTGGTCTCTGTGCCAGCTCTATAATTGTTTTGCTACGA  
TTCCACTGAAACTCTTCGATCAAGCTACTTTATGTAAATCACTTCATTGTTT  
TAAAGGAATAAACTTGATTATATTGTTTTTTTATTTGGCATAACTGTGATTC

TTTATGACCAATTAGTATGTTATGTCAGATATTCTAGTATGTA  
CCCAAATGTGTAATATTCCAGTTTTTCTCTGCATAAGTAATTAATAATACTT  
AAAAATTAATAGTTTTATCTGGGTACAAATAAACAGGTGCCTGAACTAGTTC  
ACAGACAAGGAACTTCTATGTAAAAATCACTATGATTTCTGAATTGCTATG  
TGAAACTACAGATCTTTGGAACACTGTTTAGGTAGGGTGTTAAGACTTACAC  
AGTACCTCGTTTTCTACACAGAGAAAGAAATGGCCATACTTCAGGAACTGCAG  
TGCTTATGAGGGGATATTTAGGCCTCTTGAATTTTTTGATGTAGATGGGCATT  
TTTTTAAGGTAGTGGTTAATTACCTTTATGTGAACTTTGAATGGTTTAAACA  
AAGATTTGTTTTTGTAGAGATTTTAAAGGGGGAGAATTCTAGAAATAAATGT  
TACCTAATTATTACAGCCTTAAAGACAAAAATCCTTGTTGAAGTTTTTTTTAA  
AAAAAGCTAAATTACATAGACTTAGGCATTAAACATGTTTGTGGAAGAATATA  
GCAGACGTATATTGTATCATTTGAGTGAATGTTCCCAAGTAGGCATTCTAGG  
CTCTATTTAACTGAGTCACACTGCATAGGAATTTAGAACCTAACTTTTATAG  
GTTATCAAAACTGTTGTCACCATTGCACAATTTTGTCTAATATATACATAG  
AACTTTGTGGGGCATGTTAAGTTACAGTTTGCACAAGTTCATCTCATTGT  
ATTCCATTGATTTTTTTTTTCTTCTAAACATTTTTTCTTCAAACAGTATATA  
ACTTTTTTTTAGGGGATTTTTTTTTTAGACAGCAAAAACCTATCTGAAGATTCC  
ATTTGTCAAAAAGTAATGATTTCTTGATAATTGTGTAGTAATGTTTTTTAGA  
ACCCAGCAGTTACCTTAAAGCTGAATTTATATTTAGTAACCTTCTGTGTTAAT  
ACTGGATAGCATGAATTCTGCATTGAGAACTGAATAGCTGTCATAAAATGA  
AACTTTCTTTCTAAAGAAAGATACTCACATGAGTTCTTGAAGAATAGTCATA  
ACTAGATTAAGATCTGTGTTTTAGTTTAAATAGTTTGAAGTGCCTGTTTGGGA  
TAATGATAGGTAATTTAGATGAATTTAGGGGAAAAAAAAGTTATCTGCAGAT  
ATGTTGAGGGCCCATCTCTCCCCCACACCCCCACAGAGCTAACTGGGTAC  
AGTGTTTTATCCGAAAGTTTCCAATTCCACTGTCTTGTGTTTTCATGTTGAA  
AATACTTTTGCATTTTTTCCTTTGAGTGCCAATTTCTTACTAGTACTATTTCT  
TAATGTAACATGTTTACCTGGAATGTATTTTAACTATTTTTGTATAGTGTA  
ACTGAAACATGCACATTTTGTACATTGTGCTTTCTTTTGTGGGACATATGCA  
GTGTGATCCAGTTGTTTTCCATCATTTGGTTGCGCTGACCTAGGAATGTTGG  
TCATATCAAACATTAAAAATGACCACTCTTTTAAATTGAAATTAACTTTTAAA  
TGTTTATAGGAGTATGTGCTGTGAAGTGATCTAAAATTTGTAATATTTTTGT  
CATGAAGTGTACTACTCCTAATTATTGTAATGTAATAAAAATAGTTACAGTG AC (SEQ  
ID NO: 211)>NP\_004976.2 GTPase KRas isoform b [*Homo sapiens*]  
MTEYKLVVVGAGGVGKSALTIQLIQNHVDEYDPTIEDSYRKQVVIDGETCL  
LDILDTAGQEEYSAMRDQYMRTGEGFLCVFAINNTKSFEDIHHYREQIKRVK  
DSEDVPMVLVGNKCDLPSRTVDTKQAQDLARSYGIPFIETSAKTRQGVDDAF  
YTLVREIRKHKEKMSKD**GKKKKKKSKTKCVIM** (SEQ ID NO: 212)>Lipid

affinity tag nucleotide sequence  
AAAAAGAAGAAAAAGAAGAAGAAGACAAAGTGTGTAATTATG (SEQ ID NO:  
213)>Lipid affinity tag peptide sequence KKKKKKK**KTKCVIM** (SEQ ID NO:  
214) Myr/Palm tag >NM\_002356.7 *Homo sapiens* myristoylated alanine rich  
modified from protein kinase C substrate (MARCKS), mRNA Human MARCKS  
GCACTTGGGCGTTGGACCCCGCATCTTATTAGCAACCAGGGAGATTTCTCCA  
TTTTCTCTTGTCTACAGTGCGGCTACAAATCTGGGATTTTTTTTATTACTTC  
TTTTTTTTTTCGAACTACACTTGGGCTCCTTTTTTTTGTGCTCGACTTTTCCAC  
CCTTTTTCCCTCCCTCCTGTGCTGCTGCTTTTTTGATCTCTTCGACTAAAATT  
TTTTTATCCGGAGTGTATTTAATCGGTTCTGTTCTGTCCTCTCCACCACCCC  
CACCCCCCTCCCTCCGGTGTGTGTGCCGCTGCCGCTGTTGCCGCCGCCGCTG  
CTGCTGCTGCTCGCCCCGTCGTTACACCAACCCGAGGCTCTTTGTTTCCCCT  
CTTGATCTGTTGAGTTTCTTTGTTGAAGAAGCCAGCAT**GGGTGCCCAGTTC**



TCCAAGCCACGACCGAGGAGAGCGCGGAGGCGCGGAGGCGCGG  
CTGTGGCCTCGTCGCCTTCCAAAGCGAACGGACAGGAGAATGGCCACGTGAA  
GGTAAACGGCGACGCTTCGCCCCGCGGCCGCCGAGTCGGGCGCCAAGGAGGAG  
CTGCAGGCCAACGGCAGCGCCCCGGCCGCCGACAAGGAGGAGCCCCGCGGCCG  
CCGGGAGCGGGGCGGCGTCGCCCTCCGCGGCCGAGAAAGGTGAGCCGGCCGC  
CGCCGCTGCCCCCGAGGCCGGGGCCAGCCCGGTAGAGAAGGAGGCCCCCGCG  
GAAGGCGAGGCTGCCGAGCCCGGCTCGCCACGGCCGCGGAGGGAGAGGCCG  
CGTCGGCCGCCTCCTCGACTTCTTCGCCCAAGGCCGAGGACGGGGCCACGCC  
CTCGCCAGCAACGAGACCCCGAAAAAAGAGCGCTTTTCCTTCAAG  
AAGTCTTTCAAGCTGAGCGGCTTCTCCTTCAAGAAGAACAAGAAGGAGGCTG  
GAGAAGGCGGTGAGGCTGAGGCGCCCGCTGCCGAAGGCGGCAAGGACGAGGC  
CGCCGGGGGCGCAGCTGCGGCCGCCGCCGAGGCGGGCGCGGCCTCCGGGGAG  
CAGGCAGCGGCGCCGGGCGAGGAGGCGGCAGCGGGCGAGGAGGGGGCGGGCGG  
GTGGCGACCCGCAGGAGGCCAAGCCCCAGGAGGCCGCTGTGCGCCAGAGAA  
GCCGCCCGCCAGCGACGAGACCAAGGCCGCCGAGGAGCCCAGCAAGGTGGAG  
GAGAAAAAGGCCGAGGAGGCCGGGGCCAGCGCCGCCGCCTGCGAGGCCCCCT  
CCGCCGCCGGGCCCGGCGCGCCCCCGGAGCAGGAGGCAGCCCCCGCGGAGGA  
GCCCCGCGGCCGCCGCAGCCTCGTCAGCCTGCGCAGCCCCCTCACAGGAGGCC  
CAGCCCGAGTGCAAGTCCAGAAGCCCCCCCCAGCGGAGGCGGCAGAGTAAAGA  
GCAAGCTTTTGTGAGATAATCGAAGAACTTTTCTCCCCCGTTTGTGTTGG  
AGTGGTGCCAGGTACTGGTTTTTGAGAACTTGTCTACAACCAGGGATTGATT  
TTAAAGATGTCTTTTTTTTATTTTACTTTTTTTTAAGCACCAAATTTTGTGTT  
TTTTTTTTTTTTCTCCCTCCCCACAGATCCCATCTCAAATCATTCTGTTAAC  
CACCATTTCCAACAGGTCGAGGAGAGCTTAAACACCTTCTTCCTCTGCCTTGT  
TTCTCTTTTATTTTTTATTTTTTTCGCATCAGTATTAATGTTTTTGCATACTT  
TGCATCTTTATTCAAAAGTGTAAGCTTTCTTTGTCAATCTATGGACATGCCC  
ATATATGAAGGAGATGGGTGGGTCAAAAAGGGATATCAAATGAAGTGATGGG  
GTCACAATGGGGAAATTGAAGTGGTGATAACATTGCCAAAATAGTGTGCCA  
CTAGAAATGGTGTAAGGCTGTCTTTTTTTTTTTTTTTTAAAGAAAAGTTATT  
ACCATGTATTTTGTGAGGCAGGTTTACAACACTACAAGTCTTGAGTTAAGAA  
GGAAAGAGGAAAAAAGAAAAAACACCAATACCCAGATTTAAAAAAGGAGGAG  
CGATCATAGTCTTAGGAGTTCATTTAAACCATAGGAACCTTTTCACTTATCTC  
ATGTTAGCTGTACCAGTCAGTGATTAAGTAGAACTACAAGTTGTATAGGCTT  
TATTGTTTATTGCTGGTTTATGACCTTAATAAAGTGTAATTATGTATTACCA  
GCAGGGTGTTTTTAACTGTGACTATTGTATAAAAACAAATCTTGATATCCAG  
AAGCACATGAAGTTTGCAACTTTCCACCCTGCCCATTTTTGTAAACTGCAG  
TCATCTTGGACCTTTTAAACACAAATTTTAAACTCAACCAAGCTGTGATAA  
GTGGAATGGTTACTGTTTATACTGTGGTATGTTTTTGATTACAGCAGATAAT  
GCTTTCTTTTCCAGTCGTCTTTGAGAATAAAGGAAAAAAATCTTCAGATGC  
AATGGTTTTGTGTAGCATCTTGTCTATCATGTTTTGTAAATACTGGAGAAGC  
TTTGACCAATTTGACTTAGAGATGGAATGTAACCTTTGCTTACAAAATTGCT  
ATTAAACTCCTGCTTAAGGTGTTCTAATTTTCTGTGAGCACACTAAAAGCGA  
AAAATAAATGTGAATAAAATGTACAAATTTGTTGTGTTTTTTTATGTTCTAA  
TAATACTGAGACTTCTAGGTCTTAGGTAAATTTTAGGAAGATCTTGCATGC  
CATCAGGAGTAAATTTTATTGTGGTTCTTAATCTGAAGTTTTCAAGCTCTGA  
AATTCATAATCCGCAGTGTGAGATTACGTAGAGGAAGATCTTACAACATTTT  
CATGTCAAATCTGTTACCATTTATTGGCATTTAGTTTTTCATTTAAGAATTGA  
ACATAATTATTTTTATTGTAGCTATATAGCATGTCAGATTAAATCATTTACA  
ACAAAAGGGGTGTGAACCTAAGACTATTTAAATGTCTTATGAGAAAATTTCA  
TAAAGCCATTCTCTTGTCAATTCAGGTCCAGAAACAAATTTTAAACTGAGTGA

GTGCTCTATAGTAACTACTGCAGATGCAATGTGACCATGTAATGTGCTTAGTGCTTGGCT  
TTCAAAAATTGATGGTGTATTACCTGCTATTGTAATTGCTTAGTGCTTGGCT  
AATTTCCAAATTATTGCATAATATGTTCTACCTTAAGAAAACAGGTTTATGT  
AACAAAGTAATGGTGTGGAATGGATGATGTCAGTTCATGGGCCTTTAGCATA  
GTTTTAAGCATCCTTTTTTTTTTTTTTTTTTTTGAAGTGTGTTAGCATCTTGT  
TACTCAAAGGATAAGACAGACAATAATACTTCACTGAATCTTAATAATCTTT  
ACTAGTTTACCTCCTCTGCTCTTTGCCACCCGATAACTGGATATCTTTTCCT  
TCAAAGGACCCTAAACTGATTGAAATTTAAGATATGTATCAAAAACATTATT  
TCATTTAATGCACATCTGTTTTGCTGTTTTTGAGCAGTGTGCAGTTTAGGGT  
TCATGATAAATCATTGAACCACATGTGTAACAACCTGAATGCCAAATCTTAAA  
CTCATTAGAAAAATAACAAATTAGGTTTTGACACGCATTCTTAATTGGAATA  
ATGGATCAAAAATAGTGGTTCATGACCTTACCAAACACCCTTGCTACTAATA  
AAATCAAATAACACTTAGAAGGGTATGTATTTTTTAGTTAGGGTTTCTTGATC  
TTGGAGGATGTTTGAAAGTTAAAAATTGAATTTGGTAACCAAAGGACTGATT  
TATGGGTCTTTCCTATCTTAACCAACGTTTTCTTAGTTACCTAGATGGCCAA  
GTACAGTGCCTGGTATGTAGTAAGACTCAGTAAAAAAGTGGATTTTTAAAAA  
TAACTCCCAAAGTGAATAGTCAAAAATCCTGTTAGCAAACCTGTTATATATTG  
CTAAGTTTGTTCTTTTAACAGCTGGAATTTATTAAGATGCATTATTTTGATT  
TTATTCAGTGCCTAAAACACTTTGGGTGGTATTGATGGAGTTGGTGGATTTT  
CCTCCAAGTGATTAAATGAAATTTGACGTATCTTTTCATCCAAAGTTTTGTA  
CATCATGTTTTCTAACGGAAAAAAATGTTAATATGGCTTTTTTTGTATTACTA  
AAAATAGCTTTGAGATTAAGGAAAAAATAAATACTCTTGTACAGTTCAGTAT  
TGTCTATTAAATCTGTATTGGCAGTATGTATAATGGCATTGCTGTGGTTAC  
AAAATACTTCCTCTGGGTATAATAATCATTGATCCAATTCCTATTGCTTG  
TAAAATAAAGTTTTACCAGTTGATATAATCAA (SEQ ID NO: 215)>NP\_002347.5  
myristoylated alanine-rich C-kinase substrate [*Homo sapiens*]

**MGAFQSKTAAKGEAAAERPGEAAVASSPSKANGQENGHVKVNGDASPAAAES**  
GAKEELQANGSAPAADKEEPAAAGSGAASPSAAEKGEPAAAAAPEAGASPVE  
KEAPAEGEAAEPGSPTAAEGEAASAASTSSPKAEDGATPSPSNETPKKKKK  
RFSFKKSFKLSGFSFKKNKKEAGEGGEAEAPAAEGGKDEAAGGAAAAAAEAG  
AASGEQAAAPGEEAAAGEEGAAGGDPQEAKPQEAAVAPEKPPASDETKAAEE  
PSKVEEKKAEEAGASAAACEAPSAAGPGAPPEQEAPAEPA AAAAASSACAA  
PSQEAQPECSPEAPPAEAAE (SEQ ID NO: 216)>Myr/Palm tag modified from  
Human MARCKS, nucleotide sequence ATGGGTTGCTGTTTCTCCAAGACC (SEQ  
ID NO: 217)>Myr/Palm tag modified from Human MARCKS, peptide sequence  
MGCCFSKT (SEQ ID NO: 218)

(81) In some embodiments of any of the aspects provided herein, the fusion polypeptide further comprises a peptide linker. The linker may be flexible, rigid, or cleavable. Further, the linker can be linked directly or via another linker (e.g., a peptide of one, two, three, four, five, six, seven, eight, nine, ten or more amino acids) to the fusion polypeptides described herein. Linkers can be configured according to a specific need, e.g., based on at least one of the following characteristics. In some embodiments of any of the aspects, linkers can be configured to have a sufficient length and flexibility such that it can allow for a cleavage at a target site. In some embodiments of any of the aspects, linkers can be configured to allow multimerization of the fusion polypeptides provided herein. In some embodiments of any of the aspects, linkers can be configured to facilitate expression and purification of the fusion proteins or engineered extracellular vesicles provided herein.

(82) In some embodiments of any of the aspects, a linker can be configured to have any length in a form of a peptide, peptidomimetic, an aptamer, a protein, a nucleic acid (e.g., DNA or RNA), or any combinations thereof. For example, in one embodiment, the linker may be a polypeptide linker

such as Gly-Ser-Ser-Gly (SEQ ID NO: 319) or a variation thereof as known by one of ordinary skill in the art. In another embodiment the linker may be a protein sequence for a self-cleavable peptide. For example, 2A sequences such as P2A, E2A, F2A, and T2A code for self-cleavable peptides by inducing ribosomal slippage on the mRNA at the 2A site which prevents peptide bond formation. The slippage will result in two separate peptides after translation. This allows the expression of two separate proteins from one promoter region. Any combination of the proteins described herein may be expressed with a self-cleavable peptide as known by one of ordinary skill in the art.

(83) In some embodiments of any of the aspects, the polypeptide linker is a non-cleavable linker. In some embodiments of any of the aspects, a linker can be a chemical linker of any length.

(84) In some embodiments of any of the aspects, the linker is an Fc linker. An exemplary nucleic acid sequence encoding an Fc polypeptide is:

(85) TABLE-US-00004 >KY053479.1 Synthetic construct Fc-adiponectin gene, complete cds (SEQ ID NO: 219)

```
ATGTACAGGATGCAACTCCTGTCTTGCAATTGCACTAAGTCTTGCACTTGTC
ACGAACTCGATATCGGCCATGGTTAGATCTGACAAAACCTCACACATGCCCA
CCGTGCCCAGCACCTGAACTCCTGGGGGGGACCGTCAGTCTTCCTCTTCCCC
CCAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACATGC
GTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTAC
GTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAG
TACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGAC
TGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCA
GCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCA
CAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTC
AGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAG
TGGGAGAGCAATGGGCAGCCGGAGAACAACCTACAAGACCACGCCTCCCGTG
CTGGACTCCGACGGCTCCTTCTTCCTCTACAGCAAGCTCACCGTGGACAAG
AGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCACGAGGCT
CTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAAGCC
AGCGGAAGTGGCGGAGGAGGCGGTCCTGGAGAAGGTGCCTATGTATACCGC
TCAGCATTCAAGTGTGGGATTGGAGACTTACGTTACTATCCCCAACATGCCC
ATTGCTTTACCAAGATCTTCTACAATCAGCAAAACCACTATGATGGCTCC
ACTGGTAAATTCCACTGCAACATTCCTGGGCTGTACTACTTTGCCTACCAC
ATCACAGTCTATATGAAGGATGTGAAGGTCAGCCTCTTCAAGAAGGACAAG
GCTATGCTCTTCACCTATGATCAGTACCAGGAAAATAATGTGGACCAGGCC
TCCGGCTCTGTGCTCCTGCATCTGGAGGTGGGCGACCAAGTCTGGCTCCAG
GTGTATGGGGAAGGAGAGCGTAATGGACTCTATGCTGATAATGACAATGAC
TCCACCTTCACAGGCTTTCTTCTCTACCATGACACCAACTCTAGAAAGCTT
CCTGGAGAAGGTGCCTATGTATACCGCTCAGCATTCAAGTGTGGGATTGGAG
ACTTACGTTACTATCCCCAACATGCCCATTGCTTTACCAAGATCTTCTAC
AATCAGCAAAACCACTATGATGGCTCCACTGGTAAATTCCACTGCAACATT
CCTGGGCTGTACTACTTTGCCTACCACATCACAGTCTATATGAAGGATGTG
AAGGTCAGCCTCTTCAAGAAGGACAAGGCTATGCTCTTCACCTATGATCAG
TACCAGGAAAATAATGTGGACCAGGCCTCCGGCTCTGTGCTCCTGCATCTG
GAGGTGGGCGACCAAGTCTGGCTCCAGGTGTATGGGGAAGGAGAGCGTAAT
GGACTCTATGCTGATAATGACAATGACTCCACCTTCACAGGCTTTCTTCTC
TACCATGACACCAACACTAGTCCTGGAGAAGGTGCCTATGTATACCGCTCA
GCATTCAAGTGTGGGATTGGAGACTTACGTTACTATCCCCAACATGCCCATT
CGCTTTACCAAGATCTTCTACAATCAGCAAAACCACTATGATGGCTCCACT
GGTAAATTCCACTGCAACATTCCTGGGCTGTACTACTTTGCCTACCACATC
```

ACAGTATGATGAGGATGAGTCAAGTCAAGTCAAGGACAAGGCT  
ATGCTCTTTCACCTATGATCAGTACCAGGAAAATAATGTGGACCAGGCCTCC  
GGCTCTGTGCTCCTGCATCTGGAGGTGGGCGACCAAGTCTGGCTCCAGGTG  
TATGGGGAAGGAGAGCGTAATGGACTCTATGCTGATAATGACAATGACTCC  
ACCTTCACAGGCTTTCTTCTCTACCATGACACCAACTAA.

The amino acid sequence of the Fc linker is:

(86) TABLE-US-00005 >Fc Translation (SEQ ID NO: 220)

DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDP  
EVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCK  
VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFY  
PSDIAVEWESNGQPENNYKTTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFS  
CSVMHEALHNHYTQKSLSLSPGK.

(87) In some embodiments of any of the aspects, the linker is a P2A peptide linker. P2A is a self-cleaving peptide sequence allowing for expression of two proteins from one promoter. In some embodiments, the P2A linker is encoded by the nucleic acid sequence:

GCTACTAACTTCAGCCTGCTGAAGCAG (SEQ ID NO: 221). The amino acid sequence of P2A is ATNFSLKQAGDVENPGP (SEQ ID NO: 222).

(88) In some embodiments of any of the aspects, the linker provides a multimerization (e.g., dimerization) domain wherein one fusion polypeptide may connect with another fusion polypeptide at each fusion polypeptide's respective multimerization domain. Multimerization of multiple fusion polypeptides will provide multiple fusion polypeptides within close proximity to one another to one or more a target receptor on the target cell, wherein the multiple fusion peptides will enhance receptor clustering on the target cell. Clustering receptors on a target cell will result in enhanced signal transduction. Without receptor clustering a signal may be weaker or not occur all together. For example, Fc domain sequences presented herein dimerize resulting in two fusion polypeptides connected by a covalent bond via the two Fc domains on their respective fusion polypeptide. One preferred embodiment of an Fc domain is from IgG4, herein labeled 4Fc. In other embodiments Fc may be from IgG1, herein labeled Fc. In certain embodiments Fc from other immunoglobulin, (e.g., IgG2, IgG3, etc.) may be used.

(89) Additional non-limiting examples of linkers that can be used and their properties are further described in detail, e.g., in Chen X, Zaro J L, Shen W C. Fusion protein linkers: property, design and functionality. *Adv Drug Deliv Rev.* 2013; 65 (10): 1357-1369. doi: 10.1016/j.addr.2012.09.039; O'Shea E K, Lumb K J, Kim P S. Peptide 'Velcro': design of a heterodimeric coiled coil. *Curr Biol.* 1993 Oct. 1; 3 (10): 658-67. doi: 10.1016/0960-9822 (93) 90063-t. PMID: 15335856; and Müller K M, Arndt K M, Alber T. Protein fusions to coiled-coil domains. *Methods Enzymol.* 2000; 328:261-82. doi: 10.1016/s0076-6879 (00) 28402-4. PMID: 11075350, the contents of which are incorporated herein by reference in their entireties.

(90) The engineered extracellular vesicle compositions provided herein can comprise variations in the configuration of the POI domain, linkers, and/or vesicle targeting domain. The specific combination and localization of these domains can enhance fusion polypeptide anchoring, function, or therapeutic effect, e.g., modulating inflammation.

(91) Thus, in one aspect, provided herein is an engineered extracellular vesicle comprising: at least one fusion polypeptide comprising: (i) at least one protein of interest (POI) domain or a fragment thereof; and (ii) at least one vesicle targeting domain, wherein the POI domain is in an extracellular position relative to a lipid membrane of the extracellular vesicle.

(92) In some embodiments, the POI domain or a fragment thereof is a N-terminal domain of the fusion polypeptide. In some embodiments, the vesicle targeting domain or a fragment thereof is a C-terminal domain of the fusion polypeptide.

(93) In another aspect, provided herein is an engineered extracellular vesicle comprising: at least one fusion polypeptide comprising: (i) at least one protein of interest (POI) domain or a fragment

thereof; and (ii) at least one vesicle targeting domain, wherein the POI domain is in an extracellular position relative to a lipid membrane of the extracellular vesicle, and wherein the vesicle targeting domain is a transmembrane domain relative to a lipid membrane of the extracellular vesicle.

(94) In some embodiments, the POI domain or a fragment thereof is a C-terminal domain of the fusion polypeptide. In some embodiments, the vesicle targeting domain or a fragment thereof is a N-terminal domain of the fusion polypeptide. In some embodiments, the vesicle targeting domain is in a luminal position relative to the lipid membrane of the extracellular vesicle.

(95) In some embodiments, the linker is in an exterior position relative to the lipid membrane of the extracellular vesicle. In some embodiments, the linker is a transmembrane linker. In some embodiments, the linker is in a luminal position relative to the lipid membrane of the extracellular vesicle.

(96) The engineered extracellular vesicle compositions provided herein can comprise one or more of the following fusion polypeptide sequences in Table 4.

(97) TABLE-US-00006 TABLE 4 Full Length Constructs Nucleic Acid Sequence (SEQ ID NO:) Fusion Polypeptide Amino Acid Sequence (SEQ ID NO:) hCTLA4-Fc-GPI >Artificial sequence; **hCTLA4-Fc-GPI**, DNA

**ATGGCTTGCTTGATTTCAGCGGCACAAGGCTCAGCTGAACCTGGCTACCAGGACC**  
**TGGCCCTGCACTCTCCTGTTTTTTCTTCTCTTCATCCCTGTCTTCTGCAAAGCAATG**  
**CACGTGGCCCAGCCTGCTGTGGTACTGGCCAGCAGCCGAGGCATCGCCAGCTTTGTG**  
**TGTGAGTATGCATCTCCAGGCAAAGCCACTGAGGTCCGGGTGACAGTGCTTCGGCAG**  
**GCTGACAGCCAGGTGACTGAAGTCTGTGCGGCAACCTACATGATGGGGAATGAGTTG**  
**ACCTTCCTAGATGATTCCATCTGCACGGGCACCTCCAGTGGAATCAAGTGAACCTC**  
**ACTATCCAAGGACTGAGGGCCATGGACACGGGACTCTACATCTGCAAGGTGGAGCTC**  
**ATGTACCCACCGCCATACTACCTGGGCATAGGCAACGGAACCCAGATTTATGTAATT**  
**GATCCAGAACCGTGCCCAGATTCTGACATCGATGACAAAACCTCACACATGCCCACCG**  
**TGCCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCTCTTCCCCCCTCAAAACCC**  
**AAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGTGGACGTG**  
**AGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCAT**  
**AATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGC**  
**GTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTC**  
**TCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAG**  
**CCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAAC**  
**CAGGTGAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAG**  
**TGGGAGAGCAATGGGCAGCCGGAGAACAACCTACAAGACCACGCCTCCCGTGCTGGAC**  
**TCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAG**  
**CAGGGGAACGTCTTCTCATGCTCCGTGATGCACGAGGCTCTGCACAACCACTACACG**  
**CAGAAGAGCCTCTCCCTGTCTCCGGGTAAAATCGATCCAAATAAAGGAAGTGGAACC**  
**ACTTCAGGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACAGGTTT**  
**CTTGGGACGCTAGTAACCATGGGCTTGCTGACTTAG** (SEQ ID NO: 223) > Artificial  
sequence; **hCTLA4-Fc-GPI**, Amino Acid

**MACLGFRHKAQLNLATRTWPCTLLFFLLFIPVFCKAMHVAQPAVVLASSRGIASFV**  
**CEYASPGKATEVRVTVLRLQADSQVTEVCAATYMMGNELTFLDDSICTGTSSGNQVNL**  
**TIQGLRAMDTGLYICKVELMYPPPYLGLGNGTQIYVIDPEPCPDSIDDKTHTCPP**  
**CPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVH**  
**NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQ**  
**PREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD**  
**SDGSFFLYSKLTVDKSRWQQGNVFCSSVMHEALHNHYTQKSLSLSPGKIDPNKSGT**  
**TSGTTRLLSGHTCFTLTGLLGLVTMGLLT** (SEQ ID NO: 224) hPDL1-GPI > Artificial  
sequence; **hPDL1-GPI**, DNA

**ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATTT**

ACTGTCACGGTACCGTACCGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT  
GAATGCAAATTCCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG  
GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT  
CAGCATAGTAGCTACAGACAGAGGGGCCCGGCTGTTGAAGGACCAGCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGAGAGGAGAAGCTT  
TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGAA  
CTACCTCTGGCACATCCTCCAAATGAAAGGCCAAATAAAGGAAGTGGAACCACTTCA  
GGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACAGGTTTGCTTGGG

ACGCTAGTAACCATGGGCTTGCTGACTTAG (SEQ ID NO: 225) >Amino Acid  
Sequence; **hPDL1-GPI**, Amino Acid

MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVYW  
EMEDKNIIQFVHGEEDLKVQHSSYRQRARLLKDQLSLGNAALQITDVKLQDAGVYRC  
MISYGGADYKRITVKVNAPYNKINQRILVVDPTSEHELTCQAEGYPKAEVIWTSSD  
HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPE  
LPLAHPNERPNKSGTTS GTTRLLSGHTCFTLTGLLGLVTMGLLT (SEQ ID NO: 226)

hPDL1-C1C2 >Artificial Sequence; **hPDL1-C1C2**, DNA

ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATTT  
ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT  
GAATGCAAATTCCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG  
GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT  
CAGCATAGTAGCTACAGACAGAGGGGCCCGGCTGTTGAAGGACCAGCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGAGAGGAGAAGCTT  
TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGAA  
CTACCTCTGGCACATCCTCCAAATGAAAGGATCGATGTCGAGCCACTGGGCATGGAG  
AATGGGAACATTGCCAACTCACAGATCGCCGCCTCATCTGTGCGTGTGACCTTCTTG  
GGTTTGACGATTGGGTCCCGGAGCTGGCCCCGCCTGAACCGCGCAGGCATGGTCAAT  
GCCTGGACACCCAGCAGCAATGACGATAACCCCTGGATCCAGGTGAACCTGCTGCGG  
AGGATGTGGGTAAACAGGTGTGGTGACGCAGGGTGCCAGCCGCTTGGCCAGTCATGAG  
TACCTGAAGGCCTTCAAGGTGGCCTACAGCCTTAATGGACACGAATTCGATTTTCATC  
CATGATGTTAATAAAAAACACAAGGAGTTTGTGGGTAACTGGAACAAAAACGCGGTG  
CATGTCAACCTGTTTGAGACCCCTGTGGAGGCTCAGTACGTGAGATTGTACCCACG  
AGCTGCCACACGGCCTGCACTCTGCGCTTTGAGCTACTGGGCTGTGAGCTGAACGGA  
TGCGCCAATCCCCTGGGCCTGAAGAATAACAGCATCCCTGACAAGCAGATCACGGCC  
TCCAGCAGCTACAAGACCTGGGGCTTGCATCTCTTCAGCTGGAACCCCTCCTATGCA  
CGGCTGGACAAGCAGGGCAACTTCAACGCCTGGGTTGCGGGGAGCTACGGTAACGAT  
CAGTGGCTGCAGATCTTCCCTGGCAACTGGGACAACCACTCCCACAAGAAGAACTTG  
TTTGAGACGCCCATCCTGGCTCGCTATGTGCGCATCCTGCCTGTAGCCTGGCACAAC  
CGCATCGCCCTGCGCCTGGAGCTGCTGGGCTGTTAG (SEQ ID NO: 227) >Artificial  
Sequence, **hPDL1-C1C2**, Amino Acid

MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVYW

EMEDKNIIQFVHGEEDLKVQHSSYRQRRARLLKDQLSLGNAALQITDVKLQDAGVYRC  
MISYGGADYKRITVKVNAPYNKINQRILVVD PVTSEHELTCQAEGYPKAEVIWTSSD  
HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPE  
LPLAHPPNERIDVEPLGMENGNIANSQIAASSVRVTFGLGLQHWVPELARLN RAGMVN  
AWTPSSNDDNPWIQVNLLRRMWVTGVVTQGASRLASHEYLKAFKVAYSLNGHEFDFI  
HDVNNKKHKEFVGNWNKNNAVHVNLFETPVEAQYVRLYPTSCHTACTLRFELLGCELNG  
CANPLGLKNNSIPDKQITASSSYKTWGLHLFSWNPSYARLDKQGNFNAWVAGSYGND  
QWLQIFPGNWDNHSHKKNLFETPILARYVRILPVAWHNRIALRLELLGC (SEQ ID NO:

228) hPDL1-Fc-GPI >Artificial Sequence; **hPDL1-Fc-GPI**, DNA  
ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATTT  
ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT  
GAATGCAAATTCCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG  
GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT  
CAGCATAGTAGCTACAGACAGAGGGGCCCGGCTGTTGAAGGACCAGCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGAGAGGAGAAGCTT  
TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGAA  
CTACCTCTGGCACATCCTCCAAATGAAAGGATCGATGACAAAACCTCACACATGCCCA  
CCGTGCCCAGCACCTGAACTCCTGGGGGGGACCGTCAGTCTTCTCTTCCCCCCAAAA  
CCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGTGGAC  
GTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTG  
CATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTC  
AGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAG  
GTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGG  
CAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAG  
AACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTG  
GAGTGGGAGAGCAATGGGCAGCCGGAGAACAACTACAAGACCACGCCTCCCGTGCTG  
GACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGG  
CAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCACGAGGCTCTGCACAACCACTAC  
ACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAAATCGATCCAAATAAAGGAAGTGGA  
ACCACTTCAGGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACAGGT  
TTGCTTGGGACGCTAGTAACCATGGGCTTGCTGACTTAG (SEQ ID NO: 229) >

Artificial Sequence; **hPDL1-Fc-GPI**, Amino Acid  
MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVYW  
EMEDKNIIQFVHGEEDLKVQHSSYRQRRARLLKDQLSLGNAALQITDVKLQDAGVYRC  
MISYGGADYKRITVKVNAPYNKINQRILVVD PVTSEHELTCQAEGYPKAEVIWTSSD  
HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPE  
LPLAHPPNERIDDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVD  
VSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCK  
VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAV  
EWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNV FSCSVMHEALHNHY  
TQKSLSLSPGKIDPNKGS GTTSGTTRLLSGHTCFTLTGLLGLVTMGLLT (SEQ ID NO:

230) hPDL2-C1C2 >Artificial Sequence; **hPDL2-C1C2**, DNA  
ATGATCTTCTCCTGCTAATGTTGAGCCTGGAATTGCAGCTTCACCAGATAGCAGCT  
TTATTCACAGTGACAGTCCCTAAGGAAGTGTACATAATAGAGCATGGCAGCAATGTG  
ACCCTGGAATGCAACTTTGACACTGGAAGTCATGTGAACCTTGGAGCAATAACAGCC





GTACGCGTCTCACCGTCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGC  
AAGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAGCCAAA  
GGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACC  
AAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCC  
GTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCCGTG  
CTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGG  
TGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCACGAGGCTCTGCACAACCAC  
TACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAAATCGATCCAAATAAAGGAAGT  
GGAACCACTTCAGGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACA  
GGTTTGCTTGGGACGCTAGTAACCATGGGCTTGCTGACTTAG (SEQ ID NO: 233)

>Artificial Sequence; **hPDL2-Ec-GPI**, Amino Acid

**MIFLLMLSLLEQLHQIAALFTVTVPKELYIIHGSNVTLECNFDTGSHVNLGAITA**  
**SLQKVENDTSPHRERATLLEEQLPLGKASFHIPQVQVRDEGQYQCIIIYGVAWDYKY**  
**LTLKVKASYRKINTHILKVPETDEVELTCQATGYPLAEVSWPNVSVPANTSHSRTPE**  
**GLYQVTSVLRLKPPPGRNFSCVFWNTHVRELTLASIDLQSQMEPRTHPTIDDKTHTC**  
**PPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVE**  
**VHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAK**  
**GQPREPQVYITLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPV**  
**LDSDGSFFLYSKLTVDKSRWQQGNVFSQVMHEALHNHYTQKSLSLSPGKIDPNKGS**  
**GTTSGTTRLLSGHTCFTLTGLLGLVTMGLLT** (SEQ ID NO: 234) **4F2-h41BBL**

>Artificial Sequence; **4F2-41BBL**, DNA

**ATGAGCCAGGACACCGAGGTGGATATGAAGGAGGTGGAGCTGAATGAGTTAGAGCCC**  
**GAGAAGCAGCCGATGAACGCGGCGTCTGGGGCGGCCATGTCCCTGGCGGGAGCCGAG**  
**AAGAATGGTCTGGTGAAGATCAAGGTGGCGGAAGACGAGGCGGAGGCGGCAGCCGCG**  
**GCTAAGTTCACGGGCTGTCCAAGGAGGAGCTGCTGAAGGTGGCAGGCAGCCCCGGC**  
**TGGGTACGCACCCGCTGGGCACTGCTGCTGCTCTTCTGGCTCGGCTGGCTCGGCATG**  
**CTTGCTGGTGCCGTGGTCATAATCGTGGCCTGCCCCCTGGGCGGTGTCCGGGGGCTCGC**  
**GCCTCGCCCGGCTCCGCGGCCAGCCCGAGACTCCGCGAGGGTCCCGAGCTTTTCGCCC**  
**GACGATCCCGCCGGCCTCTTGACCTGCGGCAGGGCATGTTTGCGCAGCTGGTGGCC**  
**CAAATGTTCTGCTGATCGATGGGCCCCCTGAGCTGGTACAGTGACCCAGGCCTGGCA**  
**GGCGTGTCCCTGACGGGGGGCCTGAGCTACAAAGAGGACACGAAGGAGCTGGTGGTG**  
**GCCAAGGCTGGAGTCTACTATGTCTTCTTTCAACTAGAGCTGCGGGCGCGTGGTGGCC**  
**GGCGAGGGCTCAGGCTCCGTTTCACTTGCGCTGCACCTGCAGCCACTGCGCTCTGCT**  
**GCTGGGGCCGCCGCCCTGGCTTTGACCGTGGACCTGCCACCCGCCTCCTCCGAGGCT**  
**CGGAACTCGGCCTTCGGTTTCCAGGGCCGCTTGCTGCACCTGAGTGCCGGGCCAGCGC**  
**CTGGGCGTCCATCTTCACACTGAGGCCAGGGCACGCCATGCCTGGCAGCTTACCCAG**  
**GGCGCCACAGTCTTGGGACTCTTCCGGGTGACCCCCGAAATCCCAGCCGGACTCCCT**  
**TCACCGAGGTCGGAATAA** (SEQ ID NO: 235) >Artificial Sequence; **4F2-**

**h41BBL**, Amino Acid

**MSQDTEVDMKEVELNELEPEKQPMNAASGAAMSLAGAEKNGLVKIKVAEDEAEAAAA**  
**AKFTGLSKEELLKVAGSPGWVRTRWALLLLFWLGWLGMLAGAVVIIIVACPWAVSGAR**  
**ASPGSAASPRLREGPELSPDDPAGLLDLRQGMFAQLVAQNVLLIDGPLSWYSDPGLA**  
**GVSLTGGLSYKEDTKELVVAKAGVYYVFFQLELRRVVAGEGSGSVSLALHLQPLRSA**  
**AGAAALALTVDLPPASSEARNSAFGFQGRLLHLSAGQRLGVHLHTEARARHAWQLTQ**  
**GATVLGLFRVTPEIPAGLPSRSE** (SEQ ID NO: 236) **hPDL1-4Fc-GPI** >Artificial

Sequence; **hPDL1-4Fc-GPI**, DNA

**ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATTT**  
**ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT**  
**GAATGCAAATTCCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG**  
**GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT**

CAGCATAGTAGACAGGACGAGGACCGGCTGTGTGAAGGACGACCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGAGAGGAGAAGCTT  
TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGAA  
CTACCTCTGGCACATCCTCCAAATGAAAGGGAGTCCAAATATGGTCCCCCATGCCCA  
TCATGCCCAGCACCTGAGTTCCTGGGGGGACCATCAGTCTTCCTGTTCCCCCAAAA  
CCCAAGGACACTCTCATGATCTCCCGGACCCCTGAGGTCACGTGCGTGGTGGTGAC  
GTGAGCCAGGAAGACCCCGAGGTCCAGTTCAACTGGTACGTGGATGGCGTGGAGGTG  
CATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTTCAACAGCACGTACCGTGTGGTC  
AGGGTCCTCACCGTCCTGCACCAGGACTGGCTGAACGGTAAGGAGTACAAGTGCAAG  
GTCTCCAACAAAGGCCTCCCGTCCTCCATCGAGAAAACCATCTCCAAAGCCAAAGGG  
CAGCCCCGAGAGCCACAGGTGTACACCCTGCCCCCATCCCAGGAGGAGATGACCAAG  
AACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTACCCCAGCGACATCGCCGTG  
GAGTGGGAGAGCAATGGGCAGCCGGAGGACAACCTACAAGACCACGCCTCCCGTGCTG  
GACTCCGACGGCTCCTTCTTCTCTACAGCAGGCTAACCGTGGACAAGAGCAGGTGG  
CAGGAGGGGAATGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTAC  
ACACAGAAGAGCCTCTCCCTGTCTCCGGGTAAACCAAATAAAGGAAGTGGAACCACT  
TCAGGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACAGGTTTGCTT  
GGGACGCTAGTAACCATGGGCTTGCTGACTTAG (SEQ ID NO: 237) >Artificial  
Sequence; **hPDL1-4Fc-GPI**, Amino Acid

**MRIFAVFI****MTYWHLLNAFTVTVPKDLYVVEYGSNM****TIECKFPVEKQLDLAALIVYW**  
**EMEDKNIIQFVHGEEDLKVQHSSYRQ****RARLLKDQLSLGNAALQITDVKLQDAGVYRC**  
**MISYGGADYKRITVKVNAPYNKINQRILV****VD****PVTSEHELTCQAEGYPKAEVIWTSSD**  
**HQVLSGKTTTTNSKREEKLFNVTSTLRINTT****NEIFYCTFRRLDPEENHTAELVIPE**  
**LPLAHPPNERESKYGPPCPSCPAPEFLGGPSV****FLFPPKPKD****TL****MISRTPEVTCVVVD**  
**VSQEDPEVQFNWYVDGVEVHNAKTKPREEQ****FN****STYRVVRVLTVLHQDWLNGKEYKCK**  
**VSNKGLPSSIEKTISKAKGQPREPQVYTLPP****SQEEMTKNQVSLTCLVKGFYPSDIAV**  
**EWESNGQPEDNYKTTPPVLDSDGSFFLYSRL****TVDKSRWQEGNVFSCSVMHEALHNHY**  
**TQKSLSLSPGKPNKSGTTS****GT****TRLLSGHTCFTLTGLLGLVTMGLLT** (SEQ ID NO: 238)

hPDL1-GPI-P2A- >Artificial Sequence; **hPDL1-GPI-P2A-hFGL1-GPI**, DNA hFGL1-GPI  
**ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATT****TGCTGAACGCATTT**  
**ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT**  
**GAATGCAAATTTCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG**  
**GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT**  
**CAGCATAGTAGCTACAGACAGAGGGCCCGGCTGTTGAAGGACCAGCTCTCCCTGGGA**  
**AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC**  
**ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCA**  
**TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA**  
**CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC**  
**CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGAGAGGAGAAGCTT**  
**TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC**  
**ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGAA**  
**CTACCTCTGGCACATCCTCCAAATGAAAGGCCAAATAAAGGAAGTGGAACCACTTCA**  
**GGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACAGGTTTGCTTGGG**  
**ACGCTAGTAACCATGGGCTTGCTGACTGGAAGCGGAGCTACTAACTTCAGCCTGCTG**  
**AAGCAGGCTGGCGACGTGGAGGAGAACCCTGGACCTATGGCAAAGGTGTTCAGTTTC**

ATCTTGTATTACCACCGCTCTGACAATGGGGCAGGGAAATTTCCGGCGCTCGAGGACTGT  
GCCCAGGAGCAGATGCGGCTCAGAGCCCAGGTGCGCCTGCTTGAGACCCGGGTCAAA  
CAGCAACAGGTCAAGATCAAGCAGCTTTTGCAGGAGAATGAAGTCCAGTTCCTTGAT  
AAAGGAGATGAGAATACTGTCATTGATCTTGGAAGCAAGAGGCAGTATGCAGATTGT  
TCAGAGATTTTCAATGATGGGTATAAGCTCAGTGGATTTTACAAAATCAAACCTCTC  
CAGAGCCCAGCAGAATTTTCTGTTTATTGTGACATGTCCGATGGAGGAGGATGGACT  
GTAATTCAGAGACGATCTGATGGCAGTGAAAACCTTTAACAGAGGATGGAAAGACTAT  
GAAAATGGCTTTGGAAATTTTGTCCAAAAACATGGTGAATATTGGCTGGGCAATAAA  
AATCTTCACTTCTTGACCACTCAAGAAGACTACACTTTAAAAATCGACCTTGCAGAT  
TTTGAAAAAATAGCCGTTATGCACAATATAAGAATTTCAAAGTTGGAGATGAAAAG  
AATTTCTACGAGTTGAATATTGGGGAATATTCTGGAACAGCTGGAGATTCCCTTGCG  
GGGAATTTTTCATCCTGAGGTGCAGTGGTGGGCTAGTCACCAAAGAATGAAATTCAGC  
ACGTGGGACAGAGATCATGACAACTATGAAGGGAACTGCGCAGAAGAAGATCAGTCT  
GGCTGGTGGTTTAAACAGGTGTCACCTCTGCAAACCTGAATGGTGTATACTACAGCGGC  
CCCTACACGGCTAAAACAGACAATGGGATTGTCTGGTACACCTGGCATGGGTGGTGG  
TATTCTCTGAAATCTGTGGTTATGAAAATTAGGCCAAATGATTTTATTCCAAATGTA  
ATTCCAAATAAAGGAAGTGGAACCACTTCAGGTACTACCCGTCTTCTATCTGGGCGAC  
ACGTGTTTCACGTTGACAGGTTTGCTTGGGACGCTAGTAACCATGGGCTTGCTGACT TAG  
(SEQ ID NO: 239) >Artificial Sequence; hPDL1-GPI-P2A-hFGL1-GPI, Amino  
Acid

MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVYW  
EMEDKNIIQFVHGEEDLKVQHSSYRQRARLLKDQLSLGNAALQITDVKLQDAGVYRC  
MISYGGADYKRITVKVNAPYNKINQRILVVDPTSEHELTCQAEGYPKAEVIWTSSD  
HQVLSGKTTTTNSKREEKLEFNVSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPE  
LPLAHPNERPNKGSGTTS GTTRLLSGHTCFTLTGLLGTLVTMGLLTGSGATNFSLL  
KQAGDVEENPGPMAKVFSFILVTTALTMGREISALEDCAQEQMRLRAQVRLLETRVK  
QQQVKIKQLLQENEVQFLDKGDENTVIDLGSKRQYADCSEIFNDGYKLSGFYKIKPL  
QSPAEFVYCDMSDGGGWTVIQRRSDGSENFNRGWKDYENGFGNFVQKHGEYWLGNK  
NLHFLTQEDYTLKIDLADFEKNSRYAQYKNFKVGDEKNFYELNIGEYSGTAGDSLA  
GNFHPEVQWWASHQRMKFSTWDRDHDNYEGNCAEEDQSGWWFNRCHSANLNGVYYSG  
PYTAKTDNGIVWYTW HGWWSLKSVMKIRPNDFIPNVIPNKGSGTTS GTTRLLSGH  
TCFTLTGLLGTLVTMGLLT (SEQ ID NO: 240) Myr-mScarlet >Artificial Sequence;  
Myr-mScarlet, DNA

ATGGGTTGCTGTTTCTCCAAGACCGGCTCGAGCGGCGTGAGCAAGGGGCGAGGCAGTG  
ATCAAGGAGTTCATGCGGTTCAAGGTGCACATGGAGGGGCTCCATGAACGGCCACGAG  
TTCGAGATCGAGGGGCGAGGGGCGAGGGCCGCCCTACGAGGGGCACCCAGACCGCCAAG  
CTGAAGGTGACCAAGGGTGGCCCCCTGCCCTTCTCCTGGGACATCCTGTCCCCTCAG  
TTCATGTACGGCTCCAGGGCCTTCACCAAGCACCCCGCCGACATCCCGACTACTAT  
AAGCAGTCCTTCCCCGAGGGGCTTCAAGTGGGAGCGCGTGATGAACTTCGAGGACGGC  
GGCGCCGTGACCGTGACCCAGGACACCTCCCTGGAGGACGGCACCCCTGATCTACAAG  
GTGAAGCTCCGCGGCACCAACTTCCCTCCTGACGGCCCCGTAATGCAGAAGAAGACA  
ATGGGCTGGGAAGCGTCCACCGAGCGGTTGTACCCCGAGGACGGCGTGCTGAAGGGC  
GACATTAAGATGGCCCTGCGCCTGAAGGACGGCGGCCGCTACCTGGCGGACTTCAAG  
ACCACCTACAAGGCCAAGAAGCCCGTG CAGATGCCCGGGCGCCTACAACGTGACCGC  
AAGTTGGACATCACCTCCCACAACGAGGACTACACCGTGGTGGAAACAGTACGAACGC  
TCCGAGGGGCCGCACTCCACCGGCGGCATGGACGAGCTGTACAAG (SEQ ID  
NO: 241) >Artificial Sequence; Myr-mScarlet, Amino Acid  
MGCCFSKTGSSGVSKGEAVIKEFMRFKVHMEGSMNGHEFEIEGEGEGRPYEGTQTAK  
LKVTKGGPLPFSWDILSPQFMYGSRAFTKHPADIPDYYKQSFPEGFKWERVMNFEDG  
GAVTVTQDTSLEDGTLIYKVKLRGTNFPDPGPMQKKTMGWEASTERLYPEDGVLKG

**DIKMALRDLKADRGYFADFKTTYKAKKPVQMPGAYNVDRKLDITSHNEDYTVVQEYER**  
**SEGRHSTGGMDELYK** (SEQ ID NO: 242) Myr-NanoLuc > Artificial Sequence;

*Myr-NanoLuc*, DNA Luciferase

ATGGGTTGCTGTTTCTCCAAGACCGGCTCGAGCGGCGTCTTCACACTCGAAGATTTC  
GTTGGGGACTGGCGACAGACAGCCGGCTACAACCTGGACCAAGTCCTTGAACAGGGA  
GGTGTGTCCAGTTTGTTCAGAATCTCGGGGTGTCCGTAACCTCCGATCCAAAGGATT  
GTCCTGAGCGGTGAAAATGGGCTGAAGATCGACATCCATGTCATCATCCCGTATGAA  
GGTCTGAGCGGCGACCAAATGGGCCAGATCGAAAAAATTTTAAAGGTGGTGTACCCT  
GTGGATGATCATCACTTTAAGGTGATCCTGCACTATGGCACACTGGTAATCGACGGG  
GTTACGCCGAACATGATCGACTATTTTCGGACGGCCGTATGAAGGCATCGCCGTGTTT  
GACGGCAAAAAGATCACTGTAACAGGGACCCTGTGGAACGGCAACAAAATTATCGAC  
GAGCGCCTGATCAACCCCGACGGCTCCCTGCTGTTCCGAGTAACCATCAACGGAGTG  
ACCGGCTGGCGGCTGTGCGAACGCATTCTGGCGTAA (SEQ ID NO: 243)

>Artificial Sequence; *Myr-NanoLuc*, Amino Acid

MGCCFSKTGSSGVFTLEDVFGDWRQTAGYNLDQVLEQGGVSSLFQNLGVSVTPIQRI  
VLSENGLKIDIHVIIPYEGLSGDQMGQIEKIFKVVPVDDHHFKVILHYGTLVIDG  
VTPNMIDYFGRPYEGIAVFDGKKITVTGTLWNGNKIIDERLINPDGSLFRVTINGV  
TGWRLCERILA (SEQ ID NO: 244) hSecPDL1-GPI >Artificial Sequence; **hSecPDL1-**

*GPI*, DNA

ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATTT  
ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT  
GAATGCAAATTCCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG  
GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT  
CAGCATAGTAGCTACAGACAGAGGGGCCCGGCTGTTGAAGGACCAGCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGAGAGGAGAAGCTT  
TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGGT  
AATATTCTGAATGTGTCCATTAAAATATGTCTAACACTGTCCCCTAGCACCCCAAAT  
AAAGGAAGTGGAACCACTTCAGGTACTACCCGCTTCTATCTGGGCACACGTGTTTC  
ACGTTGACAGGTTTGCTTGGGACGCTAGTAACCATGGGCTTGCTGACTTAG (SEQ ID

NO: 245) >Artificial Sequence; **hSecPDL1-GPI**, Amino Acid

**MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVYW**  
**EMEDKNIIQFVHGEEDLKVQHSSYRQRRARLLKDQLSLGNAALQITDVKLQDAGVYRC**  
**MISYGGADYKRITVKVNAPYNKINQRILVVDPTSEHELTCQAEGYPKAEVIWTSSD**  
**HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPG**  
**NILNVSIIKICLTLSPTPNKGS GTTSGTTRLLSGHTCFTLTGLLGTLVTMGLLT** (SEQ ID

NO: 246) Tfr2-h41BBL >Artificial Sequence; **Tfr2-h41BBL**, DNA

ATGGAGCGGCTTTGGGGTCTATTCCAGAGAGCGCAACAACCTGTCCCCAAGATCCTCT  
CAGACCGTCTACCAGCGTGTGGAAGGCCCCCGGAAAGGGCACCTGGAGGAGGAAGAG  
GAAGACGGGGAGGAGGGGGCGGAGACATTGGCCCACTTCTGCCCCATGGAGCTGAGG  
GGCCCTGAGCCCCTGGGCTCTAGACCCAGGCAGCCAAACCTCATTCCCTGGGCGGCA  
GCAGGACGGAGGGCTGCCCCCTACCTGGTCCTGACGGCCCTGCTGATCTTCACTGGG  
GCCTTCCTACTGGGCTACGTCGCCTTCCGAGGGTCCGCCTGCCCTGGGCGGTGTCC  
GGGGCTCGCGCCTCGCCCGGCTCCGCGGCCAGCCCGAGACTCCGCGAGGGTCCCGAG  
CTTTCGCCCGACGATCCCGCCGGCCTCTTGACCTGCGGCAGGGCATGTTTGCGCAG  
CTGGTGGCCCAAATGTTCTGCTGATCGATGGGCCCCTGAGCTGGTACAGTGACCCA

GTGGCCTGCCCTGGGGCCGTGTCCGGGGGCTCGCGCCTCGCCCGGCTCCGCGGGCCAGC  
CCGAGACTCCGCGAGGGTCCCGAGCTTTCGCCCGACGATCCCGCCGGCCTCTTGGAC  
CTGCGGCAGGGCATGTTTGCGCAGCTGGTGGCCCAAATGTTCTGCTGATCGATGGG  
CCCCTGAGCTGGTACAGTGACCCAGGCCCTGGCAGGCGTGTCCCTGACGGGGGGCCTG  
AGCTACAAAGAGGACACGAAGGAGCTGGTGGTGGCCAAGGCTGGAGTCTACTATGTC  
TTCTTTCAACTAGAGCTGCGGCGCGTGGTGGCCGGCGAGGGCTCAGGCTCCGTTTCA  
CTTGCGCTGCACCTGCAGCCACTGCGCTCTGCTGCTGGGGCCGCCGCCCTGGCTTTG  
ACCGTGACCTGCCACCCGCCTCCTCCGAGGCTCGGAACCTCGGCCTTCGGTTTCCAG

GGCCGCTTGCTGCACCTGAGTGCGCGCCAGCGCCTGGGCGTCCATCTTCACACTGAG  
GCCAGGGCACGCCATGCCTGGCAGCTTACCCAGGGCGCCACAGTCTTGGGACTCTTC  
CGGGTGACCCCCGAAATCCCAGCCGGACTCCCTTCACCGAGGTTCGGAATAA  
(SEQ ID NO: 251) >Artificial Sequence; Myr/Palm-4F2-h41BBL, Amino Acid  
MGCCFSKTGSSGSQDTEVDMKEVELNELEPEKQPMNAASGAAMSLAGAEKNGLVKIK  
VAEDEAEAAAAAKFTGLSKEELLKVAGSPGWVRTRWALLLLFWLGLWGLMLAGAVVII  
VACPWAVSGARASPGSAASPRREGPELSPDDPAGLLDLRQGMFAQLVAQNVLLIDG  
PLSWYSDPGLAGVSLTGGLSYKEDTKELVVAKAGVYYVFFQLELRRVVAGEGSGSVS  
LALHLQPLRSAAGAAALALTVDLPPASSEARNSAFGFQGRLLHLSAGQRLGVHLHTE  
ARARHAWQLTQGATVLGLFRVTPEIPAGLPSRSE (SEQ ID NO: 252) Myr/Palm-  
Link- >Artificial Sequence; Myr/Palm-Link-41BBL, DNA 41BBL (41BBL  
ATGGGTTGCTGTTTCTCCAAGACCGGCTCGAGCGGCTGGGCCCTGGTCGCGGGGCTG  
transmembrane  
CTGCTGCTGCTGCTGCTCGCTGCCGCCTGCGCCGTCTTCCTCGCCTGCCCCCTGGGCC  
domain  
GTGTCCGGGGCTCGCGCCTCGCCCGGCTCCGCGGCCAGCCCGAGACTCCGCGAGGGT  
included)  
CCCGAGCTTTCGCCCCGACGATCCCGCCGGCCTCTTGACCTGCGGCAGGGCATGTTT  
GCGCAGCTGGTGGCCCAAATGTTCTGCTGATCGATGGGCCCCCTGAGCTGGTACAGT  
GACCCAGGCCTGGCAGGCGTGTCCCTGACGGGGGGGCTGAGCTACAAAGAGGACACG  
AAGGAGCTGGTGGTGGCCAAGGCTGGAGTCTACTATGTCTTCTTTCAACTAGAGCTG  
CGGCGCGTGGTGGCCGGCGAGGGCTCAGGCTCCGTTTCACTTGCGCTGCACCTGCAG  
CCACTGCGCTCTGCTGCTGGGGCCGCGCGCCCTGGCTTTGACCGTGGACCTGCCACCC  
GCCTCCTCCGAGGCTCGGAACTCGGCCTTCGGTTTCCAGGGCCGCTTGCTGCACCTG  
AGTGCCGGCCAGCGCCTGGGCGTCCATCTTCACACTGAGGCCAGGGCACGCCATGCC  
TGGCAGCTTACCCAGGGCGCCACAGTCTTGGGACTCTTCCGGGTGACCCCCGAAATC  
CCAGCCGGACTCCCTTCACCGAGGTTCGGAATAA (SEQ ID NO: 253) >Artificial  
Sequence; Myr/Palm-Link-41BBL, Amino Acid  
MGCCFSKTGSSGWALVAGLLLLLLLLAAACAVFLACPWAVSGARASPGSAASPRREG  
PELSPDDPAGLLDLRQGMFAQLVAQNVLLIDGPLSWYSDPGLAGVSLTGGLSYKEDT  
KELVVAKAGVYYVFFQLELRRVVAGEGSGSVSLALHLQPLRSAAGAAALALTVDLPP  
ASSEARNSAFGFQGRLLHLSAGQRLGVHLHTEARARHAWQLTQGATVLGLFRVTPEI  
PAGLPSRSE (SEQ ID NO: 254) hPDL1-Link-GPI >Artificial Sequence; hPDL1-  
Link-GPI, DNA  
ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATTT  
ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT  
GAATGCAAATTCCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG  
GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT  
CAGCATAGTAGCTACAGACAGAGGGGCCCGGCTGTTGAAGGACCAGCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCOAAGAGAGAGGAGAAGCTT  
TTCAANGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGGCTCGAGCGGC  
CCAAATAAAGGAAGTGGAACCACTTCAGGTACTACCCGTCTTCTATCTGGGCACACG  
TGTTTCACGTTGACAGGTTTGCTTGGGACGCTAGTAACCATGGGCTTGCTGACTTAG (SEQ  
ID NO: 255) >Artificial Sequence; hPDL1-Link-GPI, Amino Acid  
MRIFAVFIFMTYWHLNFAFTVTPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVYW

EMEDKNIIQFVHGEEDLKVQHSSYRQRRARLLKDQLSLGNAALQITDVKLQDAGVYRC  
MISYGGADYKRITVKVNAPYNKINQRILVVDVPTSEHELTCQAEGYPKAEVIWTSSD  
HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELGSSG  
PNKGS GTTSGTTRLLSGHTCFTLTGLLGLVTMGLLT (SEQ ID NO: 256) hSecPDL1-  
>Artificial Sequence; hSecPDL1-CD9tm2, DNA CD9tm2

ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATTT  
ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT  
GAATGCAAATTCCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG  
GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT  
CAGCATAGTAGCTACAGACAGAGGGGCCCGGCTGTTGAAGGACCAGCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGAGAGGAGAAGCTT  
TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGGT  
AATATTCTGAATGTGTCCATTAAAATATGTCTAACACTGTCCCCTAGCACCTTCTAC  
ACAGGAGTCTATATTCTGATCGGAGCCGGCGCCCTCATGATGCTGGTGGGCTTCCTG  
GGCTGCTGCGGGGCTGTGCAGGAGTCCCAGTGCTAG (SEQ ID NO: 257) >Artificial  
Sequence; hSecPDL1-CD9tm2, Amino Acid

MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNM TIECKFPVEKQLDLAALIVYW  
EMEDKNIIQFVHGEEDLKVQHSSYRQRRARLLKDQLSLGNAALQITDVKLQDAGVYRC  
MISYGGADYKRITVKVNAPYNKINQRILVVDVPTSEHELTCQAEGYPKAEVIWTSSD  
HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPG  
NILNVSIIKICLTLS PSTFYTG VYILIGAGALMMLVGFLGCCGAVQESQC (SEQ ID NO:  
258) hSecPDL1- >Artificial Sequence; hSecPDL1-CD9tm2-KRAS, DNA CD9tm2-  
ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATTT  
modified KRAS

ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT  
GAATGCAAATTCCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG  
GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT  
CAGCATAGTAGCTACAGACAGAGGGGCCCGGCTGTTGAAGGACCAGCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGAGAGGAGAAGCTT  
TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGGT  
AATATTCTGAATGTGTCCATTAAAATATGTCTAACACTGTCCCCTAGCACCTTCTAC  
ACAGGAGTCTATATTCTGATCGGAGCCGGCGCCCTCATGATGCTGGTGGGCTTCCTG  
GGCTGCTGCGGGGCTGTGCAGGAGTCCCAGTGCAAAAAGAAGAAAAGAAGTCAAAG  
ACAAAGTGTGTAATTATGTAA (SEQ ID NO: 259) >Artificial Sequence; hSecPDL1-  
CD9tm2-KRAS, Amino Acid

MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNM TIECKFPVEKQLDLAALIVYW  
EMEDKNIIQFVHGEEDLKVQHSSYRQRRARLLKDQLSLGNAALQITDVKLQDAGVYRC  
MISYGGADYKRITVKVNAPYNKINQRILVVDVPTSEHELTCQAEGYPKAEVIWTSSD  
HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPG  
NILNVSIIKICLTLS PSTFYTG VYILIGAGALMMLVGFLGCCGAVQESQCKKKKKKSK TKCVIM

(SEQ ID NO: 260) hSecPDL1- >Artificial Sequence; hSecPDL1-CD9tm4, DNA  
CD9tm4

ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATTT  
ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT  
GAATGCAAATTCCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG  
GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT  
CAGCATAGTAGCTACAGACAGAGGGGCCCGGCTGTTGAAGGACCAGCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGAGAGGAGAAGCTT  
TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGGT  
AATATTCTGAATGTGTCCATTAAAATATGTCTAACACTGTCCCCTAGCACCATCGGC  
GCAGTGGGCATCGGCATTGCCGTGGTCATGATATTTGGCATGATCTTCAGTATGATC  
TTGTGCTGTGCTATCCGCAGGAACCGCGAGATGGTCTAG (SEQ ID NO: 261)

>Artificial Sequence; hSecPDL1-CD9tm4, Amino Acid

MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVYW  
EMEDKNIIQFVHGEEDLKVQHSSYRQRRARLLKDQLSLGNAALQITDVKLQDAGVYRC  
MISYGGADYKRITVKVNAPYNKINQRILVVDPTSEHELTCQAEGYPKAEVIWTSSD  
HQVLSGKTTTTNSKREEKLFNVTSTIAINTTTNEIFYCTFRRLDPEENHTAELVIPG  
NILNVSIIKICLTLS PSTIGAVGIGIAVVMIFGMIFSMILCCAIRRNREMV (SEQ ID NO: 262)

hSecPDL1-CD81 >Artificial Sequence; hSecPDL1-CD81, DNA

ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATTT  
ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT  
GAATGCAAATTCCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG  
GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT  
CAGCATAGTAGCTACAGACAGAGGGGCCCGGCTGTTGAAGGACCAGCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGAGAGGAGAAGCTT  
TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGGT  
AATATTCTGAATGTGTCCATTAAAATATGTCTAACACTGTCCCCTAGCACCCCTGTAC  
CTCATCGGCATTGCTGCCATCGTGGTTCGCTGTGATCATGATCTTCGAGATGATCCTG  
AGCATGGTGCTGTGCTGTGGCATCCGGAACAGCTCCGTGTACTGA (SEQ ID NO: 263)

>Artificial Sequence; hSecPDL1-CD81, Amino Acid

MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVYW  
EMEDKNIIQFVHGEEDLKVQHSSYRQRRARLLKDQLSLGNAALQITDVKLQDAGVYRC  
MISYGGADYKRITVKVNAPYNKINQRILVVDPTSEHELTCQAEGYPKAEVIWTSSD  
HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPG  
NILNVSIIKICLTLS PSTLYLIGIAAIVVAVIMIFEMILSMVLCCGIRNSSVY (SEQ ID NO:

264) hCD200-Fc-GPI >Artificial Sequence; hCD200-Fc-GPI, DNA

ATGGAGAGGCTGGTGATCAGGATGCCCTTCTCTCATCTGTCTACCTACAGCCTGGTT  
TGGGTCATGGCAGCAGTGGTGCTGTGCACAGCACAAAGTGCAAGTGGTGACCCAGGAT  
GAAAGAGAGCAGCTGTACACACCTGCTTCCTTAAAATGCTCTCTGCAAAATGCCCAG  
GAAGCCCTCATTGTGACATGGCAGAAAAAGAAAGCTGTAAGCCCAGAAAAACATGGTC



ACCTCAGCAGCATGCGGGTGATGATCCAGCCTGCCTATAAGGACAAGATAAAC  
ATTACCCAGCTGGGACTCCAAAACCTCAACCATCACCTTCTGGAATATCACCCCTGGAG  
GATGAAGGGTGTTACATGTGTCTCTTCAATACCTTTGGTTTTGGGAAGATCTCAGGA  
ACGGCCTGCCTCACCGTCTATGTACAGCCCATAGTATCCCTTCACTACAAATTCTCT  
GAAGACCACCTAAATATCACTTGCTCTGCCACTGCCCCGCCAGCCCCCATGGTCTTC  
TGGAAGGTCCCTCGGTCAGGGATTGAAAATAGTACAGTGAAGTCTGTCTCACCCAAAT  
GGGACCACGTCTGTTACCAGCATCCTCCATATCAAAGACCCTAAGAATCAGGTGGGG  
AAGGAGGTGATCTGCCAGGTGCTGCACCTGGGGACTGTGACCGACTTTAAGCAAACC  
GTCAACAAAGGCATCGATGACAAAACCTCACACATGCCACCGTGCCAGCACCTGAA  
CTCCTGGGGGGGACCGTCAGTCTTCTCTTCCCCCAAAACCCAAGGACACCCTCATG  
ATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGTGGACGTGAGCCACGAAGACCCT  
GAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAG  
CCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTG  
CACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTC  
CCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAG  
GTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTCAGCCTGACC  
TGCCTGGTCAAAGGCTTCTATCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGG  
CAGCCGGAGAACAACCTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTC  
TTCCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTC  
TCATGCTCCGTGATGCACGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCC  
CTGTCTCCGGGTAAAATCGATCCAAATAAAGGAAGTGGAACCACTTCAGGTACTACC  
CGTCTTCTATCTGGGCACACGTGTTTCACGTTGACAGGTTTGCTTGGGACGCTAGTA  
ACCATGGGCTTGCTGACTTAG (SEQ ID NO: 265) >Artificial Sequence; **hCD200-Fc-**  
**GPI**, Amino Acid

MERLVIRMPFSLSTYSLVWVMAAVVLCTAQVQVVTQDEREQLYTPASLKCSLQNAQ  
EALIVTWQKKKAVSPENMVTFSENHGVVIQPAYKDKINITQLGLQNSTITFWNITLE  
DEGCYMCLFNTFGFGKISGTACLTVYVQPIVSLHYKFSEDHLNITCSATARPAPMVF  
WKVPRSGIENSTVTLSHPNGTTSVTSILHIKDPKNQVGKEVICQVLHLGTVTDFKQT  
VNKGIDDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDP  
EVKENWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKAL  
PAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG  
QPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVVFSCSVMHEALHNHYTQKSLS  
LSPGKIDPNKGSGTTSGTTRLLSGHTCFTLTGLLGLVTMGLLT (SEQ ID NO: 266)

hFGL1-GPI >Artificial Sequence; **hFGL1-GPI**, DNA  
ATGGCAAAGGTGTTTCAGTTTCATCCTTGTTACCACCGCTCTGACAATGGGCAGGGAA  
ATTTGCGCGCTCGAGGACTGTGCCAGGAGCAGATGCGGCTCAGAGCCCAGGTGCGC  
CTGCTTGAGACCCGGGTCAAACAGCAACAGGTCAAGATCAAGCAGCTTTTGCAGGAG  
AATGAAGTCCAGTTCCTTGATAAAGGAGATGAGAATACTGTCATTGATCTTGGAAGC  
AAGAGGCAGTATGCAGATTGTTCAAGATTTTCAATGATGGGTATAAGCTCAGTGGA  
TTTTACAAAATCAAACCTCTCCAGAGCCCAGCAGAATTTTCTGTTTATTGTGACATG  
TCCGATGGAGGAGGATGGACTGTAATTCAGAGACGATCTGATGGCAGTGAAAACCTT  
AACAGAGGATGAAAAGACTATGAAAATGGCTTTGGAAATTTTGTCCAAAAACATGGT  
GAATATTGGCTGGGCAATAAAAATCTTCACTTCTTGACCACTCAAGAAGACTACACT  
TTAAAAATCGACCTTGCAGATTTTGAAAAAAATAGCCGTTATGCACAATATAAGAAT  
TTCAAAGTTGGAGATGAAAAGAATTTCTACGAGTTGAATATTGGGGGAATATTCTGGA  
ACAGCTGGAGATTCCCTTGCGGGGAATTTTCATCCTGAGGTGCAGTGGTGGGCTAGT  
CACCAAAGAATGAAATTCAGCACGTGGGACAGAGATCATGACAACTATGAAGGGAAC  
TGCGCAGAAGAAGATCAGTCTGGCTGGTGGTTAACAGGTGTCACCTCTGCAAACCTG  
AATGGTGTATACTACAGCGGCCCTACACGGCTAAAACAGACAATGGGATTGTCTGG  
TACACCTGGCATGGGTGGTGGTATTCTCTGAAATCTGTGGTTATGAAAATTAGGCCA

ATGATTATTTCCAAATGTAATTCCTCCAAATAAAGGAAGTGGAAACCACTTCAGGTACT  
ACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACAGGTTTGCTTGGGACGCTA  
GTAACCATGGGCTTGCTGACTTAG (SEQ ID NO: 267) >Artificial Sequence; **hFGL1-GPI**, Amino Acid

MAKVFSFILVTTALTMGREISALEDCAQEQMRLRAQVRLLETRVKQQQVKIKQLLQE  
NEVQFLDKGDENTVIDLGSKRQYADCSEIFNDGYKLSGFYKIKPLQSPAEFVYCDM  
SDGGGWTVIQRRSDGSENFNRGWKDYENGFGNFVQKHGEYWLGNKNLHFLTQTQEDYT  
LKIDLADFEKNSRYAQYKNFKVGDEKNFYELNIGEYSGTAGDSLGNFHPVQWWAS  
HQRMKFSTWDRDHDNYEGNCAEEDQSGWWFNRCHSANLNGVYYSGPYTAKTDNGIVW  
YTWHGWWYSLKSVVMKIRPNDFIPNVIPNKGSGTTS GTTRLLSGHTCFTLTGLLGLT  
VTMGLLT (SEQ ID NO: 268) **hGal9-Fc-GPI** >Artificial Sequence; **hGal9-Fc-GPI**, DNA

ATGGCCTTCAGCGGTTCCCAGGCTCCCTACCTGAGTCCAGCTGTCCCCTTTTCTGGG  
ACTATTCAAGGAGGTCTCCAGGACGGACTTCAGATCACTGTCAATGGGACCGTTCTC  
AGCTCCAGTGGAACCAGGTTTGCTGTGAACCTTTCAGACTGGCTTCAGTGGAATGAC  
ATTGCCTTCCACTTCAACCCTCGGTTTGAAGATGGAGGGTACGTGGTGTGCAACACG  
AGGCAGAACGGAAGCTGGGGGGCCCGAGGAGAGGAAGACACACATGCCTTTCCAGAAG  
GGGATGCCCTTTGACCTCTGCTTCCTGGTGCAGAGCTCAGATTTCAAGGTGATGGTG  
AACGGGATCCTCTTCGTGCAGTACTTCCACCGCGTGCCCTTCCACCGTGTGGACACC  
ATCTCCGTCAATGGCTCTGTGCAGCTGTCCTACATCAGCTTCCAGAACCCCCGCACA  
GTCCCTGTTTCAGCCTGCCTTCTCCACGGTGCCGTTCTCCCAGCCTGTCTGTTTCCCA  
CCCAGGCCCAGGGGGCGCAGACAAAACCTCCCGGGCGTGTGGCCTGCCAACC CGGCT  
CCCATTACCCAGACAGTCATCCACACAGTGCAGAGCGCCCTGGACAGATGTTCTCT  
ACTCCCGCCATCCCACCTATGATGTACCCCCACCCCGCCTATCCGATGCCTTTCATC  
ACCACCATTCTGGGAGGGCTGTACCCATCCAAGTCCATCCTCCTGTCAGGCACTGTC  
CTGCCCAGTGCTCAGAGGTTCCACATCAACCTGTGCTCTGGGAACCACATCGCCTTC  
CACCTGAACCCCCGTTTTGATGAGAATGCTGTGGTCCGCAACACCCAGATCGACAAC  
TCCTGGGGGTCTGAGGAGCGAAGTCTGCCCCGAAAAATGCCCTTCGTCCGTGGCCAG  
AGCTTCTCAGTGTGGATCTTGTGTGAAGCTCACTGCCTCAAGGTGGCCGTGGATGGT  
CAGCACCTGTTTGAATACTACCATCGCCTGAGGAACCTGCCCACCATCAACAGACTG  
GAAGTGGGGGGCGACATCCAGCTGACCCATGTGCAGACAATCGATGACAAAACCTCAC  
ACATGCCCACCGTGCCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCCTCTTC  
CCCCCAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTG  
GTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGC  
GTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTAC  
CGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTAC  
AAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAA  
GCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAG  
ATGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGAC  
ATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCT  
CCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAG  
AGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCACGAGGCTCTGCAC  
AACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAAATCGATCCAAATAAA  
GGAAGTGGAACCACTTCAGGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACG  
TTGACAGGTTTGCTTGGGACGCTAGTAACCATGGGCTTGCTGACTTAG (SEQ ID NO:  
269) >Artificial Sequence; **hGal9-Fc-GPI**, Amino Acid

MAFSGSQAPYLSPAVPFSGTIQGGLQDGLQITVNGTVLSSSGTRFAVNFQTGFSGND  
IAFHFNPRFEDGGYVVCNTRQNGSWGPEERKTHMPFQKGMFDFLCFLVQSSDFKVMV  
NGILFVQYFHRVPFHRVDTISVNGSVQLSYISFQNPRTVPVQPAFSTVPFSQPVCFP  
PRPRGRRQKPPGVWPANPAPITQTVIHTVQSAPGQMFSTPAIPPMMPHPAYPMPFI

TTILGGLYPSKSLSTGLVLPSAQRFHNLCSGNHIAFHLNPRFDENAVVRNTQIDN  
SWGSEERSLPRKMPFVRGQSFSVWILCEAHCLKVAVDGQHLFEYYHRLRNLPTINRL  
EVGGDIQLTHVQTIDDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV  
VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEY  
KCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSD  
IAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALH  
NHYTQKSLSLSPGKIDPNKGSGTTSSGTTRLLSGHTCFTLTGLLGLVTMGLLT (SEQ ID

NO: 270) hCD200-GPI >Artificial Sequence; hCD200-GPI, DNA  
ATGGAGAGGCTGGTGATCAGGATGCCCTTCTCTCATCTGTCTACCTACAGCCTGGTT  
TGGGTCATGGCAGCAGTGGTGCTGTGCACAGCACAAAGTGCAAGTGGTGACCCAGGAT  
GAAAGAGAGCAGCTGTACACACCTGCTTCCTTAAAATGCTCTCTGCAAAATGCCCAG  
GAAGCCCTCATTGTGACATGGCAGAAAAAGAAAGCTGTAAGCCCAGAAAAACATGGTC  
ACCTTCAGCGAGAACCATGGGGTGGTGATCCAGCCTGCCTATAAGGACAAGATAAAC  
ATTACCCAGCTGGGACTCCAAAACCTCAACCATCACCTTCTGGAATATCACCTTGAG  
GATGAAGGGTGTTACATGTGTCTCTTCAATACCTTTGGTTTTGGGAAGATCTCAGGA  
ACGGCCTGCCTCACCGTCTATGTACAGCCCATAGTATCCCTTCACTACAAATTCTCT  
GAAGACCACCTAAATATCACTTGCTCTGCCACTGCCCGCCCAGCCCCCATGGTCTTC  
TGGAAGGTCCCTCGGTCAGGGATTGAAAATAGTACAGTGACTCTGTCTCACCCAAAT  
GGGACCACGTCTGTTACCAGCATCCTCCATATCAAAGACCCTAAGAATCAGGTGGGG  
AAGGAGGTGATCTGCCAGGTGCTGCACCTGGGGACTGTGACCGACTTTAAGCAAACC  
GTCAACAAAGGCCCAAATAAAGGAAGTGGAACCACTTCAGGTACTACCCGTCTTCTA  
TCTGGGCACACGTGTTTCACGTTGACAGGTTTGCTTGGGACGCTAGTAACCATGGGC  
TTGCTGACTTAG (SEQ ID NO: 271) >Artificial Sequence; hCD200-GPI, Amino  
Acid

MERLVIRMPFSLSTYSLVWVMAAVVLCTAQVQVVTQDEREQLYTPASLKCSLQNAQ  
EALIVTWQKKKAVSPENMVTFSENHGVVIQPAYKDKINITQLGLQNSTITFWNITLE  
DEGCYMCLFNTFGFGKISGTACLTVYVQPIVSLHYKFSEDHLNITCSATARPAPMVF  
WKVPRSGIENSTVTLSHPNGTTSVTSILHIKDPKNQVGKEVICQVLHLGTVTDFKQT  
VNKGPNKGSGTTS

SGTTRLLSGHTCFTLTGLLGLVTMGLLT (SEQ ID NO: 272) hGal9-  
GPI >Artificial Sequence; hGa19-GPI, DNA  
ATGGCCTTCAGCGGTTCCCAGGCTCCCTACCTGAGTCCAGCTGTCCCCTTTTCTGGG  
ACTATTCAAGGAGGTCTCCAGGACGGACTTCAGATCACTGTCAATGGGACCGTTCTC  
AGCTCCAGTGGAACCAGGTTTGCTGTGAACTTTCAGACTGGCTTCAGTGGAATGAC  
ATTGCCTTCCACTTCAACCCTCGGTTTGAAGATGGAGGGTACGTGGTGTGCAACACG  
AGGCAGAACGGAAGCTGGGGGGCCCGAGGAGAGGAAGACACACATGCCTTTCCAGAAG  
GGGATGCCCTTTGACCTCTGCTTCCTGGTGCAGAGCTCAGATTTCAAGGTGATGGTG  
AACGGGATCCTCTTCGTGCAGTACTTCCACCGCGTGCCCTTCCACCGTGTGGACACC  
ATCTCCGTCAATGGCTCTGTGCAGCTGTCTACATCAGCTTCCAGAACCCCCGCACA  
GTCCCTGTTTCAGCCTGCCTTCTCCACGGTGCCGTTCTCCCAGCCTGTCTGTTTCCCA  
CCCAGGCCCCAGGGGGCGCAGACAAAACCTCCCGGCGTGTGGCCTGCCAACCCGGCT  
CCCATTACCCAGACAGTCATCCACACAGTGCAGAGCGCCCTGGACAGATGTTCTCT  
ACTCCCGCCATCCACCTATGATGTACCCCCACCCCGCCTATCCGATGCCTTTTCATC  
ACCACCATTCTGGGAGGGCTGTACCCATCCAAGTCCATCCTCCTGTCAGGCACTGTC  
CTGCCCAGTGCTCAGAGGTTCCACATCAACCTGTGCTCTGGGAACCATCGCCTTC  
CACCTGAACCCCCGTTTTGATGAGAATGCTGTGGTCCGCAACACCCAGATCGACAAC  
TCCTGGGGGTCTGAGGAGCGAAGTCTGCCCCGAAAAATGCCCTTCGTCCGTGGCCAG  
AGCTTCTCAGTGTGGATCTTGTGTGAAGCTCACTGCCTCAAGGTGGCCGTGGATGGT  
CAGCACCTGTTTGAATACTACCATCGCCTGAGGAACCTGCCACCATCAACAGACTG  
GAAGTGGGGGGCGACATCCAGCTGACCCATGTGCAGACACCAAATAAAGGAAGTGGA  
ACCACTTCAGGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACAGGT

**TTGCTTGGGACGCTAGTAACCATGGGCTTGCTGACTTAG** (SEQ ID NO: 273)

>Artificial Sequence; **hGal9-GPI**, Amino Acid

**MAFSGSQAPYLSPAVPFSGTIQGGLQDGLQITVNGTVLSSSGTRFAVNFQTGFSGND  
IAFHFNPRFEDGGYVVCNTRQNGSWGPEERKTHMPFQKGMFDFLCFLVQSSDFKVMV  
NGILFVQYFHRVPFHRVDTISVNGSVQLSYISFQNPRTVPVQPAFSTVPFSQPVCFP  
PRPRGRRQKPPGVWPANPAPITQTVIHTVQSAPGQMPSTPAIPPMMPHPAYPMPFI  
TTILGGLYPSKSILLSGTVLPSAQRFHINLCSGNHIAFHLNPRFDENAVVRNTQIDN  
SWGSEERSLPRKMPFVRGQSFSVWILCEAHCLKVAVDGQHLFEYYHRLRNLPINRL  
EVGGDIQLTHVQTPNKGSGTTSGTTRLLSGHTCFTLTGLLGTLVTMGLLT** (SEQ ID NO:

274) **hHVEM-GPI** >Artificial Sequence; **hHVEM-GPI**, DNA

**ATGGAGCCTCCTGGAGACTGGGGGCCTCCTCCCTGGAGATCCACCCCCAAAACCGAC  
GTCTTGAGGCTGGTGCTGTATCTCACCTTCCTGGGAGCCCCCTGCTACGCCCCAGCT  
CTGCCGTCCTGCAAGGAGGACGAGTACCCAGTGGGCTCCGAGTGCTGCCCCAAGTGC  
AGTCCAGGTTATCGTGTGAAGGAGGCCTGCGGGGAGCTGACGGGCACAGTGTTGTGAA  
CCCTGCCCTCCAGGCACCTACATTGCCCACCTCAATGGCCTAAGCAAGTGTCTGCAG  
TGCCAAATGTGTGACCCAGCCATGGGCCTGCGCGCGAGCCGGAAGTGTCTCAGACA  
GAGAACGCCGTGTGTGGCTGCAGCCCAGGCCACTTCTGCATCGTCCAGGACGGGGAC  
CACTGCGCCGCGTGCCGCGCTTACGCCACCTCCAGCCCCGGGCCAGAGGGTGCAGAAG  
GGAGGCACCGAGAGTCAGGACACCCTGTGTGTCAGAACTGCCCCCGGGGACCTTCTCT  
CCCAATGGGACCCTGGAGGAATGTCAGCACCCAGACCAAGTGCAGCTGGCTGGTGACG  
AAGGCCGGAGCTGGGACCAGCAGCTCCCCTGGGTACCAAATAAAGGAAGTGGAACC  
ACTTCAGGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACAGGTTTG  
CTTGGGACGCTAGTAACCATGGGCTTGCTGACTTAG** (SEQ ID NO: 275) >Artificial

Sequence; **hHVEM-GPI**, Amino Acid

**MEPPGDWGGPPWRSTPKTDVLRVLVLYLTFLGAPCYAPALPSCKEDEYPVGSECCPKC  
SPGYRVKEACGELTGTVCEPCPPGTYIAHLNGLSKCLQCQMCDPAMGLRASRNCST  
ENAVCGCSPGHFCIVQDGDHCAACRAYATSSPGQRVQKGGTESQDTLCQNCPPGTFS  
PNGTLEECQHQTCSWLVTKAGAGTSSSHWVPNKGSGTTSGTTRLLSGHTCFTLTGL  
LGTTLVTMGLLT** (SEQ ID NO: 276) **hPDL2-GPI** >Artificial Sequence; **hPDL2-GPI**, DNA

**ATGATCTTCCTCCTGCTAATGTTGAGCCTGGAATTGCAGCTTCACCAGATAGCAGCT  
TTATTCACAGTGACAGTCCCTAAGGAACTGTACATAATAGAGCATGGCAGCAATGTG  
ACCCTGGAATGCAACTTTGACACTGGAAGTCATGTGAACCTTGGAGCAATAACAGCC  
AGTTTGCAAAAGGTGGAAAATGATACATCCCCACACCGTGAAAGAGCCACTTTGCTG  
GAGGAGCAGCTGCCCCCTAGGGAAGGCCTCGTTCCACATACCTCAAGTCCAAGTGAGG  
GACGAAGGACAGTACCAATGCATAATCATCTATGGGGTTCGCCTGGGACTACAAGTAC  
CTGACTCTGAAAGTCAAAGCTTCCTACAGGAAAATAAACACTCACATCCTAAAGGTT  
CCAGAAACAGATGAGGTAGAGCTCACCTGCCAGGCTACAGGTTATCCTCTGGCAGAA  
GTATCCTGGCCAAACGTCAGCGTTCCTGCCAACACCAGCCACTCCAGGACCCCTGAA  
GGCCTCTACCAGGTCACCAGTGTTCTGCGCCTAAAGCCACCCCCCTGGCAGAACTTC  
AGCTGTGTGTTCTGGAATACTCACGTGAGGGAACCTTACTTTGGCCAGCATTGACCTT  
CAAAGTCAGATGGAACCCAGGACCCATCCAACCTCAAATAAAGGAAGTGGAACCACT  
TCAGGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACAGGTTTGCTT  
GGGACGCTAGTAACCATGGGCTTGCTGACTTAG** (SEQ ID NO: 277) >Artificial

Sequence; **hPDL2-GPI**, Amino Acid

**MIFLLLMLSLELQLHQIAALFTVTVPKELYIIHGSNVTLECNFDTGSHVNLGAITA  
SLQKVENDTSPHRERATLLEEQLPLGKASFHIPQVQVRDEGQYQCIIYGVAWDYKY  
LTLKVKASYRKINTHILKVPETDEVELTCQATGYPLAEVSWPNVSVPANTSHSRTPE  
GLYQVTSVLRLLKPPPGRNFSCVFWNTHVRELTLASIDLQSQMEPRTHPTPNKGSGTT  
SGTTRLLSGHTCFTLTGLLGTLVTMGLLT** (SEQ ID NO: 278) **hTSG6-GPI** >Artificial

Sequence; **hTSG6-GPI**, DNA

ATGATCATCTTAATTTACTTATTTCTCTTGCTATGGGAAGACACTCAAGGATGGGGA  
TTCAAGGATGGAATTTTTTCATAACTCCATATGGCTTGAACGAGCAGCCGGTGTGTAC  
CACAGAGAAGCACGGTCTGGCAAATACAAGCTCACCTACGCAGAAGCTAAGGCGGTG  
TGTGAATTTGAAGGCGGCCATCTCGCAACTTACAAGCAGCTAGAGGCAGCCAGAAAA  
ATTGGATTTTCATGTCTGTGCTGCTGGATGGATGGCTAAGGGCAGAGTTGGATACCCC  
ATTGTGAAGCCAGGGCCCAACTGTGGATTTGGAAAACTGGCATTATTGATTATGGA  
ATCCGTCTCAATAGGAGTGAAAGATGGGATGCCTATTGCTACAACCCACACGCAAAG  
GAGTGTGGTGGCGTCTTTACAGATCCAAAGCAAATTTTTTAAATCTCCAGGCTTCCCA  
AATGAGTACGAAGATAACCAAATCTGCTACTGGCACATTAGACTCAAGTATGGTCAG  
CGTATTCACCTGAGTTTTTTTAGATTTTGACCTTGAAGATGACCCAGGTTGCTTGGCT  
GATTATGTTGAAATATATGACAGTTACGATGATGTCCATGGCTTTGTGGGAAGATAC  
TGTGGAGATGAGCTTCCAGATGACATCATCAGTACAGGAAATGTCATGACCTTGAAG  
TTTCTAAGTGATGCTTCAGTGACAGCTGGAGGTTTCCAAATCAAATATGTTGCAATG  
GATCCTGTATCCAAATCCAGTCAAGGAAAAAATACAAGTACTACTTCTACTGGAAAT  
AAAAACTTTTTAGCTGGAAGATTTAGCCACTTAATCGATCCAAATAAAGGAAGTGGA  
ACCACTTCAGGTACTACCCGTCTTCTATCTGGGACACAGTGTTTCACGTTGACAGGT  
TTGCTTGGGACGCTAGTAACCATGGGCTTGCTGACTTAG (SEQ ID NO: 279)

>Artificial Sequence; **hTSG6-GPI**, Amino Acid

MIILIYFLLLWEDTQGWGFKDGIHNSIWLERAAGVYHREARSGKYKLTAEAKAV  
CEFEGGHLATYKQLEAARKIGFHVCAAGWMAKGRVGYPIVKPGPNCGFGKTGIIDYG  
IRLNRSERWDAYCYNPHAKECGGVFTDPKQIFKSPGFPNEYEDNQICYWHIRLKYGQ  
RIHLSFLDFDLEDDPGCLADYVEIYDSYDDVHGFVGRYCGDELDDIISTGNVMTLK  
FLSDASVTAGGFQIKYVAMDPVSKSSQGKNTSTTSTGNKNFLAGRFSHLIDPNKGS  
TTSGTTRLLSGHTCFTLTGLLGLVTMGLLT (SEQ ID NO: 280) hHVEM-Fc-GPI

>Artificial Sequence; **hHVEM-Fc-GPI**, DNA

ATGGAGCCTCCTGGAGACTGGGGGCCTCCTCCCTGGAGATCCACCCCCAAAACCGAC  
GTCTTGAGGCTGGTGTGTATCTCACCTTCCTGGGAGCCCCCTGCTACGCCCCAGCT  
CTGCCGTCTGCAAGGAGGACGAGTACCCAGTGGGCTCCGAGTGCTGCCCAAGTGC  
AGTCCAGGTTATCGTGTGAAGGAGGCCTGCGGGGAGCTGACGGGCACAGTGTGTGAA  
CCCTGCCCTCCAGGCACCTACATTGCCACCTCAATGGCCTAAGCAAGTGTCTGCAG  
TGCCAAATGTGTGACCCAGCCATGGGCCTGCGCGCGAGCCGGAAGTGTCTCAGGACA  
GAGAACGCCGTGTGTGGCTGCAGCCCAGGCCACTTCTGCATCGTCCAGGACGGGGAC  
CACTGCGCCGCGTGCCGCGCTTACGCCACCTCCAGCCCGGGCCAGAGGGTGCAGAAG  
GGAGGCACCGAGAGTCAGGACACCCTGTGTGAGAAGTGGCCCCCGGGGACCTTCTCT  
CCCAATGGGACCCTGGAGGAATGTCAGCACCAGACCAAGTGCAGCTGGCTGGTGACG  
AAGGCCGGAGCTGGGACCAGCAGCTCCCCTGGGTAATCGATGACAAAACCTCACACA  
TGCCACCGTGCCAGCACCTGAACCTGGGGGGACCGTCAGTCTTCCTCTTCCCC  
CCAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTG  
GTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTG  
GAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGT  
GTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAG  
TGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCC  
AAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATG  
ACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAGCGACATC  
GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACTACAAGACCACGCCTCCC  
GTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGC  
AGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCACGAGGCTCTGCACAAC  
CACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAAATCGATCCAAATAAAGGA  
AGTGGAACCACTTCAGGTACTACCCGTCTTCTATCTGGGACACAGTGTTTCACGTTG

ACAGCTTTGCTTGACGACGATGAACCATGGGCTTGCTGACTTAG (SEQ ID NO: 281)

>Artificial Sequence; **hHVEM-Fc-GPI**, Amino Acid

**MEPPGDWGWPPPWRSTPKTDVLRRLVLYLTFLGAPCYAPALPSCKEDEYPVGSECCPKC  
SPGYRVKEACGELTGTVCPCPPGTIAHLNGLSKCLQCQMCDPAMGLRASRNCST  
ENAVCGCSPGHFCIVQDGDHCAACRAYATSSPGQRVQKGGTESQDTLCQNCPPGTFS  
PNGTLEECQHQTCKSWLVTKAGAGTSSSHWVIDDKTHTCPPCPAPPELLGGPSVFLFP  
PKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYR  
VVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEM  
TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKS  
RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGKIDPNKSGTTS GTTRLLSGHTCFTL**

**TGLLGTLVTMGLLT** (SEQ ID NO: 282) **hPDL1-GPI-P2A-** >Artificial Sequence;

**hPDL1-GPI-P2A-hHVEM-GPI**, DNA **hHVEM-GPI**

**ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATTT  
ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT  
GAATGCAAATTCCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG  
GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT  
CAGCATAGTAGCTACAGACAGAGGGGCCCGCTGTTGAAGGACCAGCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCGAAGAGAGAGGAGAAGCTT  
TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGAA  
CTACCTCTGGCACATCCTCCAAATGAAAGGCCAAATAAAGGAAGTGGAACCACTTCA  
GGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACAGGTTTGCTTGGG  
ACGCTAGTAACCATGGGCTTGCTGACTGGAAGCGGAGCTACTAACTTCAGCCTGCTG  
AAGCAGGCTGGCGACGTGGAGGAGAACCCTGGACCTATGGAGCCTCCTGGAGACTGG  
GGGCCTCCTCCCTGGAGATCCACCCCAAAACCGACGTCTTGAGGCTGGTGCTGTAT  
CTCACCTTCCTGGGAGCCCCCTGCTACGCCCCAGCTCTGCCGTCCTGCAAGGAGGAC  
GAGTACCCAGTGGGCTCCGAGTGCTGCCCCAAGTGCAGTCCAGGTTATCGTGTGAAG  
GAGGCCTGCGGGGAGCTGACGGGCACAGTGTGTGAACCCTGCCCTCCAGGCACCTAC  
ATTGCCACCTCAATGGCCTAAGCAAGTGTCTGCAGTGCCAAATGTGTGACCCAGCC  
ATGGGCCTGCGCGCGAGCCGGAAGTGTCTCCAGGACAGAGAACGCCGTGTGTGGCTGC  
AGCCCAGGCCACTTCTGCATCGTCCAGGACGGGGACCACTGCGCCGCGGTGCCGCGCT  
TACGCCACCTCCAGCCCCGGGCCAGAGGGTGCAGAAGGGAGGCACCGAGAGTCAGGAC  
ACCCTGTGTCAGAACTGCCCCCGGGGACCTTCTCTCCAATGGGACCCTGGAGGAA  
TGTCAGCACCAGACCAAGTGCAGCTGGCTGGTGACGAAGGCCGGAGCTGGGACCAGC  
AGCTCCCCTGAGTACCAAATAAAGGAAGTGGAACCACTTCAGGTACTACCCGTCTT  
CTATCTGGGCACACGTGTTTCACGTTGACAGGTTTGCTTGGGACGCTAGTAACCATG  
GGCTTGCTGACTTAG** (SEQ ID NO: 283) >Artificial Sequence; **hPDL1-GPI-P2A-**

**hHVEM-GPI**, Amino Acid

**MRIFAVFIFMTYWHLNAFTVTVPKDLYVVEYGSNMETIECKFPVEKQLDLAALIVYW  
EMEDKNIIQFVHGEEDLKVQHSSYRQRARLLKDQLSLGNAALQITDVKLQDAGVYRC  
MISYGGADYKRITVKVNAPYNKINQRILVVDPTSEHELTCQAEGYPKAEVIWTSSD  
HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPE  
LPLAHPNERPNKSGTTS GTTRLLSGHTCFTLTGLLGTLVTMGLLTGSGATNFSLL  
KQAGDVEENPGPMEPPGDWGWPPPWRSTPKTDVLRRLVLYLTFLGAPCYAPALPSCKED  
EYPVGSECCPKCSPGYRVKEACGELTGTVCPCPPGTIAHLNGLSKCLQCQMCDPA  
MGLRASRNCSTENAVCGCSPGHFCIVQDGDHCAACRAYATSSPGQRVQKGGTESQD**

**TLCQNCPPTGTEPCQHQTKCSWINTKAGTSSSHWVPNKGSGTTSGTTRL**

**LSGHTCFTLTGLLGTLVTMGLLT** (SEQ ID NO: 284) mCTLA4-Fc-GPI >Artificial

Sequence; **mCTLA4-Fc-GPI**, DNA

**ATGGCTTGTCTTGGACTCCGGAGGTACAAAGCTCAACTGCAGCTGCCTTCTAGGACT**  
**TGGCCTTTTGTAGCCCTGCTCACTCTTCTTTTCATCCCAGTCTTCTCTGAAGCCATA**  
**CAGGTGACCCAACCTTCAGTGGTGTGGCTAGCAGCCATGGTGTGCGCCAGCTTTCCA**  
**TGTGAATATTCACCATCACACAACACTGATGAGGTCCGGGTGACTGTGCTGCGGCAG**  
**ACAAATGACCAAATGACTGAGGTCTGTGCCACGACATTCACAGAGAAGAATACAGTG**  
**GGCTTCCTAGATTACCCCTTCTGCAGTGGTACCTTTAATGAAAGCAGAGTGAACCTC**  
**ACCATCCAAGGACTGAGAGCTGTTGACACGGGACTGTACCTCTGCAAGGTGGAACCTC**  
**ATGTACCCACCGCCATACTTTGTGGGCATGGGCAACGGGACGCAGATTTATGTCATT**  
**GATCCAGAACCATGCCCCGGATTCTGAATCGATGACAAAACCTCACACATGCCCCACCGT**  
**GCCCAGCACCTGAACTCCTGGGGGGGACCGTCAGTCTTCTCTTCCCCCCAAAACCCA**  
**AGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGTGGACGTGA**  
**GCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATA**  
**ATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCG**  
**TCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCT**  
**CCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGC**  
**CCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACC**  
**AGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGT**  
**GGGAGAGCAATGGGCAGCCGGAGAACTACAAGACCACGCCTCCCGTGCTGGACT**  
**CCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGC**  
**AGGGGAACGTCTTCTCATGCTCCGTGATGCACGAGGCTCTGCACAACCACTACACGC**  
**AGAAGAGCCTCTCCCTGTCTCCGGGTAAAATCGATCCAAATAAAGGAAGTGGAACCA**  
**CTTCAGGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACAGGTTTGC**  
**TTGGGACGCTAGTAACCATGGGCTTGCTGACTTAG** (SEQ ID NO: 285) >Artificial

Sequence; **mCTLA4-Fc-GPI**, Amino Acid

**MACLGLRRYKAQLQLPSRTWPFVALLTLLFIPVFSEAIQVTQPSVVLASSHGVASFP**  
**CEYSPSHNTDEVVRVTVLRQTNDQMTEVCATTFTEKNTVGFLDYPFCSGTFNESRVNL**  
**TIQGLRAVDLTGLYLCKVELMYPPPYFVGMGNGTQIYVIDPEPCPDS**  
**IDDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKF**  
**NWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPI**  
**EKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPN**  
**NYKTTTPVLDSDSGFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG**  
**KIDPNKGS GTTSGTTRL****LSGHTCFTLTGLLGTLVTMGLLT** (SEQ ID NO: 286) mPDL1-

C1C2 >Artificial Sequence; **mPDL1-C1C2**, DNA

**ATGAGGATATTTGCTGGCATTATATTCACAGCCTGCTGTCACTTGCTACGGGGCGTTT**  
**ACTATCACGGCTCCAAAGGACTTGACGTGGTGGAGTATGGCAGCAACGTCACGATG**  
**GAGTGCAGATTCCCTGTAGAACGGGAGCTGGACCTGCTTGCGTTAGTGGTGTACTGG**  
**GAAAAGGAAGATGAGCAAGTGATTCAGTTTGTGGCAGGAGAGGAGGACCTTAAGCCT**  
**CAGCACAGCAACTTCAGGGGGGAGAGCCTCGCTGCCAAAGGACCAGCTTTTGAAGGGA**  
**AATGCTGCCCTTCAGATCACAGACGTCAAGCTGCAGGACGCAGGCGTTTACTGCTGC**  
**ATAATCAGCTACGGTGGTGC GGACTACAAGCGAATCACGCTGAAAGTCAATGCCCCA**  
**TACCGCAAAATCAACCAGAGAATTTCCGTGGATCCAGCCACTTCTGAGCATGAACTA**  
**ATATGTCAGGCCGAGGGTTATCCAGAAGCTGAGGTAATCTGGACAAACAGTGACCAC**  
**CAACCCGTGAGTGGGAAGAGAAGTGTCACCACTTCCCGGACAGAGGGGGATGCTTCTC**  
**AATGTGACCAGCAGTCTGAGGGTCAACGCCACAGCGAATGATGTTTTCTACTGTACG**  
**TTTTGGAGATCACAGCCAGGGCAAACACACAGCGGAGCTGATCATCCAGAAGCTG**  
**CCTGCAACACATCCTCCACAGAACAGGACTATCGATGTCGAGCCACTGGGCATGGAG**  
**AATGGGAACATTGCCAACTCACAGATCGCCGCCTCATCTGTGCGTGTGACCTTCTTG**

GGTTGCTGACCTGAGGATGGTCCCGCCCTGACCGCGCAGGCATGGTCAAT  
GCCTGGACACCCAGCAGCAATGACGATAACCCCTGGATCCAGGTGAACCTGCTGCGG  
AGGATGTGGGTAAACAGGTGTGGTGACGCAGGGTGCCAGCCGCTTGGCCAGTCATGAG  
TACCTGAAGGCCTTCAAGGTGGCCTACAGCCTTAATGGACACGAATTCGATTTCATC  
CATGATGTTAATAAAAAACACAAGGAGTTTGTGGGTAACCTGGAACAAAAACGCGGTG  
CATGTCAACCTGTTTGAGACCCCTGTGGAGGCTCAGTACGTGAGATTGTACCCACG  
AGCTGCCACACGGCCTGCACTCTGCGCTTTGAGCTACTGGGCTGTGAGCTGAACGGA  
TGCGCCAATCCCCTGGGCCTGAAGAATAACAGCATCCCTGACAAGCAGATCACGGCC  
TCCAGCAGCTACAAGACCTGGGGCTTGCATCTCTTCAGCTGGAACCCCTCCTATGCA  
CGGCTGGACAAGCAGGGCAACTTCAACGCCTGGGTTGCGGGGAGCTACGGTAACGAT  
CAGTGGCTGCAGATCTTCCCTGGCAACTGGGACAACCACTCCCACAAGAAGAACTTG  
TTTGAGACGCCCATCCTGGCTCGCTATGTGCGCATCCTGCCTGTAGCCTGGCACAAAC  
CGCATCGCCCTGCGCCTGGAGCTGCTGGGCTGTTAG (SEQ ID NO: 287) >Artificial  
Sequence; **mPDL1-C1C2**, Amino Acid

**MRIFAGIIFTACCHLLRAFTITAPKDLYVVEYGSNVTMECRFPVERELDLLALVVYW  
EKEDEQVIQFVAGEEDLKPQHSNFSGRASLPKDQLLKGNALQITDVKLQDAGVYCC  
IISYGGADYKRITLKVNPYRKINQRISVDPATSEHELICQAEGYPEAEVIWTNSDH  
QPVSGKRSVTTSRTEGMLLNVTSSLRVNATANDVFYCTFWRSQPGQNHTAELIIPEL  
PATHPPQNRITIDVEPLGMENGNIANSQIAASSVRVTFLGLQHWVPELARLN RAGMVN  
AWTPSSNDNPWIQVNLLRRMWVTGVVTQGASRLASHEYLKAFKVAYSLNGHEFDI  
HDVNKKHKEFVGNWNKNVHVNLFETPVEAQYVRLYPTSCHTACTLRFELLGCELNG  
CANPLGLKNNSIPDKQITASSSYKTWGLHLFSWNPSYARLDKQGNFNAWVAGSYGND  
QWLQIFPGNWDNHSHKKNLFETPILARYVRILPVAWHNRIALRLELLGC (SEQ ID NO:  
288) mPDL1-Fc-GPI >Artificial Sequence; **mPDL1-Fc-GPI**, DNA**

ATGAGGATATTTGCTGGCATTATATTCACAGCCTGCTGTCACCTTGCTACGGGCGTTT  
ACTATCACGGCTCCAAAGGACTTGTACGTGGTGGAGTATGGCAGCAACGTCACGATG  
GAGTGCAGATTCCCTGTAGAACGGGAGCTGGACCTGCTTGCGTTAGTGGTGTACTGG  
GAAAAGGAAGATGAGCAAGTGATTTCAGTTTGTGGCAGGAGAGGAGGACCTTAAGCCT  
CAGCACAGCAACTTCAGGGGGGAGAGCCTCGCTGCCAAAGGACCAGCTTTTGAAGGGA  
AATGCTGCCCTTCAGATCACAGACGTCAAGCTGCAGGACGCAGGCGTTTACTGCTGC  
ATAATCAGCTACGGTGGTGCGGACTACAAGCGAATCACGCTGAAAGTCAATGCCCCA  
TACCGCAAAATCAACCAGAGAATTTCCGTGGATCCAGCCACTTCTGAGCATGAACTA  
ATATGTCAGGCCGAGGGTTATCCAGAAGCTGAGGTAATCTGGACAAACAGTGACCAC  
CAACCCGTGAGTGGGAAGAGAAGTGTCACCACTTCCCGGACAGAGGGGGATGCTTCTC  
AATGTGACCAGCAGTCTGAGGGTCAACGCCACAGCGAATGATGTTTTCTACTGTACG  
TTTTGGAGATCACAGCCAGGGCAAACACACAGCGGAGCTGATCATCCAGAACTG  
CCTGCAACACATCCTCCACAGAACAGGACTATCGATGACAAAACCTCACACATGCCCA  
CCGTGCCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCTCTTCCCCCAAAA  
CCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTGCTGGTGGAC  
GTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGAGGTG  
CATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCAGTACCGTGTGGTC  
AGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAG  
GTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAGCCAAAGGG  
CAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAG  
AACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAGCGACATCGCCGTG  
GAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCCGTGCTG  
GACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGG  
CAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCACGAGGCTCTGCACAACCACTAC  
ACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAAATCGATCCAAATAAAGGAAGTGGA  
ACCACTTCAGGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCAGTTGACAGGT



TTGCTTGGGACGCTAGTACCATGGGCTTGCTGACTTAG (SEQ ID NO: 289)

>Artificial Sequence; **mPDL1-Fc-GPI**, Amino Acid

**MRIFAGIIFTACCHLLRAFTITAPKDLYVVEYGSNVTMECRFPVERELDLLALVVYW  
EKEDEQVIQFVAGEEDLKPQHSNFRGRASLPKDQLLKGNAAALQITDVKLQDAGVYCC  
IISYGGADYKRITLKVNPYRKINQRISVDPATSEHELICQAEGYPEAEVIWTNSDH  
QPVSGKRSVTTSRTEGMLLNVTSSLRVNATANDVPYCTFWRSQPGQNHTAELIPEL  
PATHPPQNRRTIDDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVD  
VSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCK  
VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAV  
EWESNGQPENNYKTTTPVLDSDGSEFLYSLKLTVDKSRWQQGNVFSCSVMHEALHNHY  
TQKSLSLSPGKIDPNKGS GTTSGTTRLLSGHTCFTLTGLLGLVTMGLLT** (SEQ ID NO:

290) **mPDL1-GPI** >Artificial Sequence; **mPDL1-GPI**, DNA

**ATGAGGATATTTGCTGGCATTATATTCACAGCCTGCTGCTACTTGCTACGGGCGTTT  
ACTATCACGGCTCCAAAGGACTTGTACGTGGTGGAGTATGGCAGCAACGTCACGATG  
GAGTGCAGATTCCCTGTAGAACGGGAGCTGGACCTGCTTGCGTTAGTGGTGTACTGG  
GAAAAGGAAGATGAGCAAGTGATTCAAGTTTGTGGCAGGAGAGGAGGACCTTAAGCCT  
CAGCACAGCAACTTCAGGGGGGAGAGCCTCGCTGCCAAAGGACCAGCTTTTGAAGGGA  
AATGCTGCCCTTCAGATCACAGACGTCAAGCTGCAGGACGCAGGCGTTTACTGCTGC  
ATAATCAGCTACGGTGGTGCGGACTACAAGCGAATCACGCTGAAAGTCAATGCCCCA  
TACCGCAAAATCAACCAGAGAATTTCCGTGGATCCAGCCACTTCTGAGCATGAACTA  
ATATGTCAGGCCGAGGGTTATCCAGAAGCTGAGGTAATCTGGACAAACAGTGACCAC  
CAACCCGTGAGTGGGAAGAGAAGTGTCACCACTTCCCGGAZAGAGGGGGATGCTTCTC  
AATGTGACCAGCAGTCTGAGGGTCAACGCCACAGCGAATGATGTTTTCTACTGTACG  
TTTTGGAGATCACAGCCAGGGCAAACACACAGCGGAGCTGATCATCCCAGAACTG  
CCTGCAACACATCCTCCACAGAACAGGACTCCAAATAAAGGAAGTGGAACCACTTCA  
GGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACAGGTTTGCTTGGG  
ACGCTAGTAACCATGGGCTTGCTGACTTAG** (SEQ ID NO: 291)

>Artificial Sequence;  
**mPDL1-GPI**, Amino Acid

**MRIFAGIIFTACCHLLRAFTITAPKDLYVVEYGSNVTMECRFPVERELDLLALVVYW  
EKEDEQVIQFVAGEEDLKPQHSNFRGRASLPKDQLLKGNAAALQITDVKLQDAGVYCC  
IISYGGADYKRITLKVNPYRKINQRISVDPATSEHELICQAEGYPEAEVIWTNSDH  
QPVSGKRSVTTSRTEGMLLNVTSSLRVNATANDVFYCTFWRSQPGQNHTAELIPEL  
PATHPPQNRTPNKGS GTTSGTTRLLSGHTCFTLTGLLGLVTMGLLT** (SEQ ID NO: 292)

**mPDL2-C1C2** >Artificial Sequence; **mPDL2-C1C2**, DNA

**ATGCTGCTCCTGCTGCCGATACTGAACCTGAGCTTACAACCTTCATCCTGTAGCAGCT  
TTATTCACCGTGACAGCCCCTAAAGAAGTGACACCGTAGACGTCGGCAGCAGTG  
AGCCTGGAGTGCGATTTTGACCGCAGAGAATGCACTGAACTGGAAGGGATAAGAGCC  
AGTTTGCAGAAGGTAGAAAATGATACGTCTCTGCAAAGTGAAAGAGCCACCCTGCTG  
GAGGAGCAGCTGCCCCTGGGAAAGGCTTTGTTCCACATCCCTAGTGTCGAAGTGAGA  
GATTCCGGGCAGTACCGTTGCCTGGTCATCTGCGGGGCGCCTGGGACTACAAGTAC  
CTGACGGTGAAAGTCAAAGCTTCTTACATGAGGATAGACACTAGGATCCTGGAGGTT  
CCAGGTACAGGGGAGGTGCAGCTTACCTGCCAGGCTAGAGGTTATCCCCTAGCAGAA  
GTGTCCTGGCAAATGTCAGTGTTCTGCCAACACCAGCCACATCAGGACCCCCGAA  
GGCCTCTACCAGGTCACCAGTGTTCTGCGCCTCAAGCCTCAGCCTAGCAGAACTTC  
AGCTGCATGTTCTGGAATGCTCACATGAAGGAGCTGACTTCAGCCATCATTGACCCT  
CTGAGTCGGATGGAACCCAAAGTCCCCAGAACGATCGATGTCGAGCCACTGGGGCATG  
GAGAATGGGAACATTGCCAACTCACAGATCGCCGCCTCATCTGTGCGTGTGACCTTC  
TTGGGTTTGCAGCATTGGGTCCCGGAGCTGGCCCGCCTGAACCGCGCAGGCATGGTC  
AATGCCTGGACACCCAGCAGCAATGACGATAACCCCTGGATCCAGGTGAACCTGCTG  
CGGAGGATGTGGGTAACAGGTGTGGTGACGCAGGGTGCCAGCCGCTTGGCCAGTCAT**

GATTGACAGCTTCAAGGCTGGACACGAATTCGATTTC  
ATCCATGATGTTAATAAAAAACACAAGGAGTTTGTGGGTAACTGGAACAAAAACGCG  
GTGCATGTCAACCTGTTTGAGACCCCTGTGGAGGCTCAGTACGTGAGATTGTACCCC  
ACGAGCTGCCACACGGCCTGCACTCTGCGCTTTGAGCTACTGGGCTGTGAGCTGAAC  
GGATGCGCCAATCCCCTGGGCCTGAAGAATAACAGCATCCCTGACAAGCAGATCACG  
GCCTCCAGCAGCTACAAGACCTGGGGCTTGCATCTCTTCAGCTGGAACCCCTCCTAT  
GCACGGCTGGACAAGCAGGGCAACTTCAACGCCTGGGTTGCGGGGAGCTACGGTAAC  
GATCAGTGGCTGCAGATCTTCCCTGGCAACTGGGACAACCACTCCCACAAGAAGAAC  
TTGTTTGAGACGCCCATCCTGGCTCGCTATGTGCGCATCCTGCCTGTAGCCTGGCAC  
AACCGCATCGCCCTGCGCCTGGAGCTGCTGGGCTGTTAG (SEQ ID NO: 293)

>Artificial Sequence; **mPDL2-C1C2**, Amino Acid

**MLLLLPILNLSLQLHPVAALFTVTAPKEVYTVDVVGSSVSLECFDRRECTELEGIRA  
SLQKVENDTSLQSERATLLEEQLPLGKALFHIPSVQVRDSGQYRCLVICGAAWDYKY  
LTVKVKASYMRIDTRILEVPGTGEVQLTCQARGYPLAEVSWQNVSPANTSHIRTPE  
GLYQVTSVLRLKPQPSRNFSCMFWNAHMKELTSIIDPLSRMEPKVPRTIDVEPLGM  
ENGNANSQIAASSVRVTFGLGLQHWVPELARLNRAGMVNAWTPSSNDDNPWIQVNNL  
RRMWVTGVVTQGASRLASHEYLKAFKVAYSLNGHEFDFIHDVNKKHKEFVGNWNKNA  
VHVNLFETPVEAQYVRLYPTSCHTACTLRFELLGCELNGCANPLGLKNNSIPDKQIT  
ASSSYKTWGLHLFSWNPSYARLDKQGNFNAWVAGSYGNDQWLQIFPGNWDNHSHKKN  
LFETPILARYVRILPVAWHNRIALRLELLGC (SEQ ID NO: 294)** mPDL2-Fc-GPI

>Artificial Sequence; **mPDL2-Fc-GPI**, DNA

**ATGCTGCTCCTGCTGCCGATACTGAACCTGAGCTTACAACCTTCATCCTGTAGCAGCT  
TTATTCACCGTGACAGCCCCTAAAGAAGTGTACACCGTAGACGTCGGCAGCAGTGTG  
AGCCTGGAGTGCGATTTTGACCGCAGAGAATGCACTGAACTGGAAGGGATAAGAGCC  
AGTTTGCAGAAGGTAGAAAATGATACGTCTCTGCAAAGTGAAAGAGCCACCCTGCTG  
GAGGAGCAGCTGCCCCCTGGGAAAGGCTTTGTTCCACATCCCTAGTGTCCAAGTGAGA  
GATTCGCGGCAGTACCGTTGCCTGGTCATCTGCGGGGGCCGCTGGGACTACAAGTAC  
CTGACGGTGAAAGTCAAAGCTTCTTACATGAGGATAGACACTAGGATCCTGGAGGTT  
CCAGGTACAGGGGAGGTGCAGCTTACCTGCCAGGCTAGAGGTTATCCCCTAGCAGAA  
GTGTCCTGGCAAATGTCAGTGTTCTGCCAACACCAGCCACATCAGGACCCCCGAA  
GGCCTCTACCAGGTCACCAGTGTTCTGCGCCTCAAGCCTCAGCCTAGCAGAACTTC  
AGCTGCATGTTCTGGAATGCTCACATGAAGGAGCTGACTTCAGCCATCATTGACCCT  
CTGAGTCGGATGGAACCCAAAGTCCCCAGAACGATCGATGACAAAACCTCACACATGC  
CCACCGTGCCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCCTCTTCCCCCA  
AAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGTG  
GACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGAG  
GTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTG  
GTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGC  
AAGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAA  
GGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACC  
AAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCC  
GTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACCTACAAGACCACGCCTCCCGTG  
CTGGACTCCGACGGCTCCTTCTTCCTCTACAGCAAGCTCACCGTGGACAAGAGCAGG  
TGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCACGAGGCTCTGCACAACCAC  
TACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAAATCGATCCAAATAAAGGAAGT  
GGAACCACTTCAGGTACTACCCGTCTTCTATCTGGGCACACGTGTTTCACGTTGACA  
GGTTTGCTTGGGACGCTAGTAACCATGGGCTTGCTGACTTAG (SEQ ID NO: 295)**

>Artificial Sequence; **mPDL2-Fc-GPI**, Amino Acid

**MLLLLPILNLSLQLHPVAALFTVTAPKEVYTVDVVGSSVSLECFDRRECTELEGIRA  
SLQKVENDTSLQSERATLLEEQLPLGKALFHIPSVQVRDSGQYRCLVICGAAWDYKY**

LTVKVMRIDRTILEVPTGTEVQVLTQARGYPLAEVSWQNVSPANTSHIRTP  
GLYQVTSVLRLKPQPSRNFSCMFWNAHNKELTSAIIDPLSRMEPKVPRTIDDKTHTC  
PPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVE  
VHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAK  
GQPREPQVYITLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPV  
LDSDGSEFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGKIDPNKGS  
GTTSGTTRLLSGHTCFTLTGLLGLTVMGLLT (SEQ ID NO: 296) mPDL1-mFc-GPI  
>Artificial Sequence; **mPDL1-mFc-GPI**, DNA

ATGAGGATATTTGCTGGCATTATATTCACAGCCTGCTGTCACCTTGCTACGGGCGTTT  
ACTATCACGGCTCCAAAGGACTTGTACGTGGTGGAGTATGGCAGCAACGTCACGATG  
GAGTGCAGATTCCCTGTAGAACGGGAGCTGGACCTGCTTGCGTTAGTGGTGTACTGG  
GAAAAGGAAGATGAGCAAGTGATTCAAGTTTGTGGCAGGAGAGGAGGACCTTAAGCCT  
CAGCACAGCAACTTCAGGGGGGAGAGCCTCGCTGCCAAAGGACCAGCTTTTGAAGGGA  
AATGCTGCCCTTCAGATCACAGACGTCAAGCTGCAGGACGCAGGCGTTTACTGCTGC  
ATAATCAGCTACGGTGGTGCGGACTACAAGCGAATCACGCTGAAAGTCAATGCCCCA  
TACCGCAAAATCAACCAGAGAATTTCCGTGGATCCAGCCACTTCTGAGCATGAACTA  
ATATGTCAGGCCGAGGGTTATCCAGAAGCTGAGGTAATCTGGACAAACAGTGACCAC  
CAACCCGTGAGTGGGAAGAGAAGTGTCACCACTTCCCGGACAGAGGGGGATGCTTCTC  
AATGTGACCAGCAGTCTGAGGGTCAACGCCACAGCGAATGATGTTTTCTACTGTACG  
TTTTGGAGATCACAGCCAGGGCAAACACACAGCGGAGCTGATCATCCAGAACTG  
CCTGCAACACATCCTCCACAGAACAGGACTGGTTGTAAGCCTTGCATATGTACAGTC  
CCAGAAGTATCATCTGTCTTCATCTTCCCCCCAAAGCCCAAGGATGTGCTCACCATT  
ACTCTGACTCCTAAGGTCACGTGTGTTGTGGTAGACATCAGCAAGGATGATCCCGAG  
GTCCAGTTCAGCTGGTTTGTAGATGATGTGGAGGTGCACACAGCTCAGACGCAACCC  
CGGGAGGAGCAGTTCAACAGCACTTTCGGCTCAGTCAGTGAACCTCCCATCATGCAC  
CAGGACTGGCTCAATGGCAAGGAGTTCAAATGCAGGGTCAACAGTGCAGCTTTCCCT  
GCCCCCATCGAGAAAACCATCTCCAAAACCAAAGGCAGACCGAAGGCTCCACAGGTG  
TACACCATTCCACCTCCCAAGGAGCAGATGGCCAAGGATAAAGTCAGTCTGACCTGC  
ATGATAACAGACTTCTTCCCTGAAGACATTACTGTGGAGTGGCAGTGGAAATGGGCAG  
CCAGCGGAGAACTACAAGAACTCAGCCCATCATGGACACAGATGGCTCTTACTTC  
GTCTACAGCAAGCTCAATGTGCAGAAGAGCAACTGGGAGGCAGGAAATACTTTCACC  
TGCTCTGTGTTACATGAGGGCCTGCACAACCACCATACTGAGAAGAGCCTCTCCAC  
TCTCCTGGTAAACCAAATAAAGGAAGTGGAACCACTTCAGGTACTACCCGTCTTCTA  
TCTGGGCACACGTGTTTCACGTTGACAGGTTTGCTTGGGACGCTAGTAACCATGGGC  
TTGCTGACTTAG (SEQ ID NO: 297) >Artificial Sequence; **mPDL1-mFc-GPI**,  
Amino Acid

MRIFAGIIFTACCHLLRAFTITAPKDLYVVEYGSNVTMECRFPVERELDLLALVVYW  
EKEDEQVIQFVAGEEDLKPQHSNFRGRASLPKDQLLKGNAAALQITDVKLQDAGVYCC  
IISYGGADYKRITLKVNPYRKINQRISVDPATSEHELICQAEGYPEAEVIWTNSDH  
QPVSGKRSVTTSRTEGMLLNVTSSLRVNATANDVFYCTFWRSQPGQNHTAELIPEL  
PATHPPQNRTGCKPCICTVPEVSSVFIFPPKPKDVLTTITLPKVTCVVVDISKDDPE  
VQFSWFVDDVEVHTAQTPREEQFNSTFRSVSELPIMHQDWLNGKEFKCRVNSAAFP  
APIEKTISKTKGRPKAPQVYTIPPPKEQMAKDKVSLTCMITDFFPEDITVEWQWNGQ  
PAENYKNTQPIMDTDGSYFVYSKLVNQKSNWEAGNTFTCSVLHEGLHNHHTEKSLSH  
SPGKPNKGSGTTSGTTRLLSGHTCFTLTGLLGLTVMGLLT (SEQ ID NO: 298) mPDL2-  
GPI >Artificial Sequence; **mPDL2-GPI**, DNA

ATGCTGCTCCTGCTGCCGATACTGAACCTGAGCTTACAACCTTCATCCTGTAGCAGCT  
TTATTCACCGTGACAGCCCCTAAAGAAGTGACACCGTAGACGTCGGCAGCAGTGTG  
AGCCTGGAGTGCGATTTTGACCGCAGAGAATGCACTGAACTGGAAGGGATAAGAGCC  
AGTTTGCAGAAGGTAGAAAATGATACGTCTCTGCAAAGTGAAAGAGCCACCCTGCTG

**mPDL1-GPI-P2A-mHVEM-GPI, Amino Acid**  
**MRIFAGIIFTACCHLLRAFTITAPKDLVVEYGSNVTMECRFPVERELDLLALVYIW**  
**EKEDEQVIQFVAGEEDLKPQHSNFRGRASLPKDQLKGNALQITDVKLQDAGVYCC**  
**IISYGGADYKRITLKVNPYRKINQRISVDPATSEHELICQAEGYPEAEVIWTNSDH**

QPVSGKTRVNTSLRNVNATANADVIFYCTFWRSQPGQNHTAELIPEL  
PATHPPQNRTPNKGS GTTSGTTRLLSGHTCFTLTGLLGTLVTMGLLTGSGATNFSLL  
KQAGDVEENPGPMEPLPGWGSAPWSQAPTDNTFRLVPCVFLNLLQRISAQPSCRQE  
EFLVGDECCPMCNPGYHV KQVCSEHTGTVCAPCPPQTYTAHANGLSKCLPCGVCDPD  
MGLLTWQECSSWKDTVCR CIPGYFCENQD GSHCSTCLQH TTCPPGQ RVEKRGTHDQD  
TVCADCLTGT FSLGGTQEECLPWTNCSAFQQEVRRGTNSTDTTCS SQPNKGS GTTSG  
TTRLLSGHTCFTLTGLLGTLVTMGLLT (SEQ ID NO: 302) hPDL1-ADAM10 >Artificial  
Sequence; hPDL1-ADAM10, DNA

ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATT  
ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT  
GAATGCAAATTCCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG  
GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT  
CAGCATAGTAGCTACAGACAGAGGGGCCCGGCTGTTGAAGGACCAGCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGAGAGGAGAAGCTT  
TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGAA  
CTACCTCTGGCACATCCTCCAAATGAAAGGTGTGGAAATGGAATGGTAGAACAAGGT  
GAAGAATGTGATTGTGGCTATAGTGACCAGTGTAAGATGAATGCTGCTTCGATGCA  
AATCAACCAGAGGGAAGAAAATGCAAACCTGAAACCTGGGAAACAGTGCAAGTCCAAGT  
CAAGGTCCTTGTGTACAGCACAGTGTGCATTCAAGTCAAAGTCTGAGAAGTGTCCG  
GATGATT CAGACTGTGCAAGGGAAGGAATATGTAATGGCTTCACAGCTCTCTGCCCA  
GCATCTGACCCTAAACCAAACCTTCACAGACTGTAATAGGCATACACAAGTGTGCATT  
AATGGGCAATGTGCAGGTTCTATCTGTGAGAAATATGGCTTAGAGGAGTGTACGTGT  
GCCAGTTCTGATGGCAAAGATGATAAAGAATTATGCCATGTATGCTGTATGAAGAAA  
ATGGACCCATCAACTTGTGCCAGTACAGGGTCTGTGCAGTGGAGTAGGCACTTCAGT  
GGTCGAACCATCACCTTGCAACCTGGATCCCCCTTGCAACGATTTTAGAGGTTACTGT  
GATGTTTTTCATGCGGTGCAGATTAGTAGATGCTGATGGTCCTCTAGCTAGGCTTAAA  
AAAGCAATTTT TAGTCCAGAGCTCTATGAAAACATTGCTGAATGGATTGTGGCTCAT  
TGGTGGGCAGTATTACTTATGGGAATTGCTCTGATCATGCTAATGGCTGGATTATT  
AAGATATGCAGTGTTaATACTCCAAGTAGTAATCCAAAGTTGCCTCCTCCTAAACCA  
CTTCCAGGCACTTTAAAGAGGAGGAGACCTCCACAGCCCATT CAGCAACCCCAGCGT  
CAGCGGCCCCGAGAGAGTTATCAAATGGGACACATGAGACGCTAA (SEQ ID NO: 303)

>Artificial Sequence; hPDL1-ADAM10, Amino Acid

MRIFAVFIFMTYWHLLNAFTVTVPKDLYVVEYGSNM TIECKEPVEKQLDLAALIVYW  
EMEDKNIIQFVHGEEDLKVQHSSYRQRRARLLKDQLSLGNAALQITDVKLQDAGVYRC  
MISYGGADYKRITVKVNAPYNKINQRILVVD PVTSEHELTCQAEGYPKAEVIWTSSD  
HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPE  
LPLAHPPNERCGNGMVEQGEEDCGYSDQCKDECCFDANQPEGRKCKLKPGKQCSPS  
QGPCCTAQCAFKSKSEKCRDDSDCAREGICNGFTALCPASDPKPNFTDCNRHTQVCI  
NGQCAGSICEKYGLEECTASSDGKDDEELCHVCCMKKMDPSTCASTGSVQWSRHFS  
GRTITLQPGSPCNDFRGYCDVFMRCRLVDADGPLARLKKAI FSPELYENIAEWIVAH  
WWAVLLMGIALIMLMAGFIKICSVHTPSSNPKLPPP KPLPGTLKRRRPPQPIQQPQR  
QRPRESYQMGMHMR (SEQ ID NO: 304) hPDL1-4Fc- >Artificial Sequence; hPDL1-  
4Fc-CD9tm2, DNA CD9tm2

ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATT  
ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT

MRIFAVFIEMTYWHLLNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVYW  
EMEDKNIIQFVHGEEDLKVQHSSYRQRARLLKDQLSLGNAALQITDVKLQDAGVYRC  
MISYGGADYKRITVKVNAPYNKINQRILVVDPTSEHELTCQAEGYPKAEVIWTSSD  
HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPE  
LPLAHPPNERESKYGPPCPSCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVD  
VSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVRLTVLHQDWLNGKEYKCK  
VSNGKLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAV  
EWESNGQPEDNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHENHY  
TQKLSLSLSPGKFYTGVIILIGAGALMMLVGFLGCCGAVQESQCVIM (SEQ ID NO: 306)

ACTGTCACGGTTCCTCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT  
GAATGCAAATTCCTCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG  
GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT  
CAGCATAGTAGCTACAGACAGAGGGCCCGGCTGTTGAAGGACCAGCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCAATTCCAAGAGAGAGGAGAAGCTT  
TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGAA  
CTACCTCTGGCACATCCTCCAAATGAAAGGGAGTCCAAATATGGTCCCCCATGCCCA

TCATGCCCAGCACCTGATGTTCTCTGGGGGGACCTGATCTTCTTCCCCCCCCAAAA  
CCCAAGGACACTCTCATGATCTCCCCGGACCCCTGAGGTCACGTGCGTGGTGGTGGAC  
GTGAGCCAGGAAGACCCCGAGGTCCAGTTCAACTGGTACGTGGATGGCGTGGAGGTG  
CATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTTCAACAGCACGTACCGTGTGGTC  
AGGGTCCTCACCGTCCTGCACCAGGACTGGCTGAACGGTAAGGAGTACAAGTGCAAG  
GTCTCCAACAAAGGCCTCCCGTCCTCCATCGAGAAAACCATCTCCAAAGCCAAAGGG  
CAGCCCCGAGAGCCACAGGTGTACACCCTGCCCCCATCCCAGGAGGAGATGACCAAG  
AACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTACCCCAGCGACATCGCCGTG  
GAGTGGGAGAGCAATGGGCAGCCGGAGGACAACCTACAAGACCACGCCTCCCGTGCTG  
GACTCCGACGGCTCCTTCTTCTCTACAGCAGGCTAACCGTGGACAAGAGCAGGTGG  
CAGGAGGGGAATGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTAC  
ACACAGAAGAGCCTCTCCCTGTCTCCGGGTAAATTCTACACAGGAGTCTATATTCTG  
ATCGGAGCCGGCGCCCTCATGATGCTGGTGGGCTTCTTGGGCTGCTGCGGGGGCTGTG  
CAGGAGTCCCAGTGCAAAAAGAAGAAAAAGAAGAAGAAGACAAAGTGTGTAATTATG TAA  
(SEQ ID NO: 307) >Artificial Sequence; **hPDL1-4Fc-CD9tm2-KRAS**, Amino Acid  
**MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVYW**  
**EMEDKNIIQFVHGEEDLKVQHSSYRQRRARLLKDQLSLGNAALQITDVKLQDAGVYRC**  
**MISYGGADYKRITVKVNAPYNKINQRILVVD PVTSEHELTCQAEGYPKAEVIWTSSD**  
**HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPE**  
**LPLAHPPNERESKYGP**PCSCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVD  
VSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVRVLTVLHQDWLNGKEYKCK  
VSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQQEEMTKNQVSLTCLVKGFYPSDIAV  
EWESNGQPEDNYKTTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHY  
TQKSLSLSPGKFYTGVIYLIGAGALMMLVGFLGCCGAVQESQCKKKKKKKKTKCVIM (SEQ  
ID NO: 308) **hPDL1-Fc- >Artificial Sequence; hPDL1-Fc-CD9tm2**, DNA **CD9tm2**  
**ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATTT**  
**ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT**  
**GAATGCAAATTC**CCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG  
GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT  
CAGCATAGTAGCTACAGACAGAGGGGCCCGGCTGTTGAAGGACCAGCTCTCCCTGGGA  
AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC  
ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCA  
TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA  
CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTGGACAAGCAGTGAC  
CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGAGAGGAGAAGCTT  
TTCAATGTGACCAGCACACTGAGAATCAACACAACACTAATGAGATTTTCTACTGC  
ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGAA  
CTACCTCTGGCACATCCTCCAAATGAAAGGATCGATGACAAAACTCACACATGCCCA  
CCGTGCCCAGCACCTGAACTCCTGGGGGGGACCGTCAGTCTTCTCTTCCCCCCCCAAAA  
CCCAAGGACACCCTCATGATCTCCCCGGACCCCTGAGGTCACATGCGTGGTGGTGGAC  
GTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTG  
CATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTC  
AGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAG  
GTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGG  
CAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAG  
AACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTG  
GAGTGGGAGAGCAATGGGCAGCCGGAGAACAACTACAAGACCACGCCTCCCGTGCTG  
GACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGG  
CAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCACGAGGCTCTGCACAACCACTAC  
ACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAAATCGATTTCTACACAGGAGTCTAT

ATTCTGATCGGATCGGAGCCCTCATGCTGGGCTTCCTGGGCTGCTGCGGG  
GCTGTGCAGGAGTCCCAGTGCCTAATTATGTAA (SEQ ID NO: 309) >Artificial  
Sequence; **hPDL1-Fc-CD9tm2**, Amino Acid  
**MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVYW**  
**EMEDKNIIQFVHGEEDLKVQHSSYRQRRARLLKDQLSLGNAALQITDVKLQDAGVYRC**  
**MISYGGADYKRITVKVNAPYNKINQRILVVD PVTSEHELTCQAEGYPKAEVIWTSSD**  
**HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPE**  
**LPLAHPPNERIDDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVD**  
VSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCK  
VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAV  
EWESNGQOPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHY  
TQKSLSLSPGKIDFYTGVIYILIGAGALMMLVGFLGCCGAVQESQCVIM (SEQ ID NO: 310)

**hPDL1-Fc- >Artificial Sequence; hPDL1-Fc-CD9tm2-KRAS**, DNA CD9tm2-  
**ATGAGGATATTTGCTGTCTTTATATTCATGACCTACTGGCATTGCTGAACGCATTT**  
modified KRAS  
**ACTGTCACGGTTCCCAAGGACCTATATGTGGTAGAGTATGGTAGCAATATGACAATT**  
**GAATGCAAATTCCCAGTAGAAAAACAATTAGACCTGGCTGCACTAATTGTCTATTGG**  
**GAAATGGAGGATAAGAACATTATTCAATTTGTGCATGGAGAGGAAGACCTGAAGGTT**  
**CAGCATAGTAGCTACAGACAGAGGGGCCCGCTGTTGAAGGACCAGCTCTCCCTGGGA**  
**AATGCTGCACTTCAGATCACAGATGTGAAATTGCAGGATGCAGGGGTGTACCGCTGC**  
**ATGATCAGCTATGGTGGTGCCGACTACAAGCGAATTACTGTGAAAGTCAATGCCCCA**  
**TACAACAAAATCAACCAAAGAATTTTGGTTGTGGATCCAGTCACCTCTGAACATGAA**  
**CTGACATGTCAGGCTGAGGGCTACCCCAAGGCCGAAGTCATCTCTGGAAGCAGTGAC**  
**CATCAAGTCCTGAGTGGTAAGACCACCACCACCAATTCCAAGAGAGAGGAGAAGCTT**  
**TTCAATGTGACCAGCACACTGAGAATCAACACAACAATAATGAGATTTTCTACTGC**  
**ACTTTTAGGAGATTAGATCCTGAGGAAAACCATACAGCTGAATTGGTCATCCCAGAA**  
**CTACCTCTGGCACATCCTCCAAATGAAAGGATCGATGACAAAACCTCACACATGCCCCA**  
CCGTGCCCAGCACCTGAACTCCTGGGGGGGACCGTCAGTCTTCCTCTTCQCCCCAAAA  
CCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGGAC  
GTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTG  
CATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTC  
AGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAG  
GTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAGCCAAAGGG  
CAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAG  
AACCAGGTGAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTG  
GAGTGGGAGAGCAATGGGCAGCCGGAGAACAATAAGACCACGCCTCCCGTGCTG  
GA  
CTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGG  
CAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCACGAGGCTCTGCACAACCACTAC  
ACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAAATCGATTTCTACACAGGAGTCTAT  
ATTCTGATCGGAGCCGGCGCCCTCATGATGCTGGTGGGCTTCTGGGCTGCTGCGGG  
GCTGTGCAGGAGTCCCAGTGCAAAAAGAAGAAAAAGAAGAAGACAAAGTGTGTA  
ATTATGTAA (SEQ ID NO: 311) >Artificial Sequence; **hPDL1-Fc-CD9tm2-KRAS**,  
Amino Acid

**MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVYW**  
**EMEDKNIIQFVHGEEDLKVQHSSYRQRRARLLKDQLSLGNAALQITDVKLQDAGVYRC**  
**MISYGGADYKRITVKVNAPYNKINQRILVVD PVTSEHELTCQAEGYPKAEVIWTSSD**  
**HQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPE**  
**LPLAHPPNERIDDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVD**  
VSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCK  
VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAV



EWESNGQPENNYKIDFTPPVLDSDGSEFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHY  
TQKSLSLSPGKIDFYTGVIYILIGAGALMMLVGFLGCCGAVQESQCKKKKKKKKTKCV IM  
(SEQ ID NO: 312) mPDL1-mFc- >Artificial Sequence; **mPDL1-mFc-CD9tm2**, DNA  
CD9tm2

ATGAGGATATTTGCTGGCATTATATTCACAGCCTGCTGTCACTTGCTACGGGCGTTT  
ACTATCACGGCTCCAAAGGACTTGTACGTGGTGGAGTATGGCAGCAACGTCACGATG  
GAGTGCAGATTCCCTGTAGAACGGGAGCTGGACCTGCTTGCGTTAGTGGTGTACTGG  
GAAAAGGAAGATGAGCAAGTGATTCAGTTTGTGGCAGGAGAGGAGGACCTTAAGCCT  
CAGCACAGCAACTTCAGGGGGAGAGCCTCGCTGCCAAAGGACCAGCTTTTGAAGGGA  
AATGCTGCCCTTCAGATCACAGACGTCAAGCTGCAGGACGCAGGCGTTTACTGCTGC  
ATAATCAGCTACGGTGGTGC GGACTACAAGCGAATCACGCTGAAAGTCAATGCCCCA  
TACCGCAAAATCAACCAGAGAATTTCCGTGGATCCAGCCACTTCTGAGCATGAACTA  
ATATGTCAGGCCGAGGGTTATCCAGAAGCTGAGGTAATCTGGACAAACAGTGACCAC  
CAACCCGTGAGTGGGAAGAGAAGTGTCACCACTTCCCGGACAGAGGGGGATGCTTCTC  
AATGTGACCAGCAGTCTGAGGGTCAACGCCACAGCGAATGATGTTTTCTACTGTACG  
TTTTGGAGATCACAGCCAGGGCAAACACACAGCGGAGCTGATCATCCAGAAGCTG  
CCTGCAACACATCCTCCACAGAACAGGACTGGTTGTAAGCCTTGCATATGTACAGTC  
CCAGAAGTATCATCTGTCTTCATCTTCCCCCAAAGCCCAAGGATGTGCTCACCATT  
ACTCTGACTCCTAAGGTCACGTGTGTTGTGGTAGACATCAGCAAGGATGATCCCGAG  
GTCCAGTTCAGCTGGTTTGTAGATGATGTGGAGGTGCACACAGCTCAGACGCAACCC  
CGGGAGGAGCAGTTCAACAGCACTTTCCGCTCAGTCAGTGAACTTCCCATCATGCAC  
CAGGACTGGCTCAATGGCAAGGAGTTCAAATGCAGGGTCAACAGTGCAGCTTTCCCT  
GCCCCCATCGAGAAAACCATCTCCAAAACCAAAGGCAGACCGAAGGCTCCACAGGTG  
TACACCATTCCACCTCCCAAGGAGCAGATGGCCAAGGATAAAGTCAGTCTGACCTGC  
ATGATAACAGACTTCTTCCCTGAAGACATTACTGTGGAGTGGCAGTGGGAATGGGCAG  
CCAGCGGAGAACTACAAGAACTCAGCCCATCATGGACACAGATGGCTCTTACTTC  
GTCTACAGCAAGCTCAATGTGCAGAAGAGCAACTGGGAGGCAGGAAATACTTTCACC  
TGCTCTGTGTTACATGAGGGCCTGCACAACCACCATACTGAGAAGAGCCTCTCCAC  
TCTCCTGGTAAATTCTACACAGGAGTCTATATTCTGATCGGAGCCGGCGCCCTCATG  
ATGCTGGTGGGCTTCCTGGGCTGCTGCGGGGCTGTGCAGGAGTCCCAGTGCGTAATT  
ATGTAA (SEQ ID NO: 313) >Artificial Sequence; **mPDL1-mFc-CD9tm2**, Amino  
Acid

MRIFAGIIFTACCHLLRAFTITAPKDLYVVEYGSNVTMECRFPVERELDLLALVVYW  
EKEDEQVIQFVAGEEDLKPQHSNFRGRASLPKDQLLKGNAALQITDVKLQDAGVYCC  
IISYGGADYKRITLKVNPYRKINQRISVDPATSEHELICQAEGYPEAEVIWTNSDH  
QPVSGKRSVTTSRTEGMLLNVTSSLRVNATANDVFYCTFWRSQPGQNHTAELIPEL  
PATHPPQNRTGCKPCICTVPEVSSVFIFPPKPKDVLTTITLTPKVTCVVVDISKDDPE  
VQFSWFVDDVEVHTAQTQPREEQFNSTFRSVSELPIMHQDWLNGKEFKCRVNSAAFP  
APIEKTISKTKGRPKAPQVYTIPPPKEQMAKDKVSLTCMITDFFPEDITVEWQWNGQ  
PAENYKNTQPIMDTDGSYFVYSKLVNQKSNWEAGNTFTCSVLHEGLHNHHTTEKSLSH  
SPGKPFYTGVIYILIGAGALMMLVGFLGCCGAVQESQCVIM (SEQ ID NO: 314) mPDL1-  
mFc- >Artificial Sequence; **mPDL1-mFc-CD9tm2-KRAS**, DNA CD9tm2-

ATGAGGATATTTGCTGGCATTATATTCACAGCCTGCTGTCACTTGCTACGGGCGTTT  
modified KRAS  
ACTATCACGGCTCCAAAGGACTTGTACGTGGTGGAGTATGGCAGCAACGTCACGATG  
GAGTGCAGATTCCCTGTAGAACGGGAGCTGGACCTGCTTGCGTTAGTGGTGTACTGG  
GAAAAGGAAGATGAGCAAGTGATTCAGTTTGTGGCAGGAGAGGAGGACCTTAAGCCT  
CAGCACAGCAACTTCAGGGGGAGAGCCTCGCTGCCAAAGGACCAGCTTTTGAAGGGA  
AATGCTGCCCTTCAGATCACAGACGTCAAGCTGCAGGACGCAGGCGTTTACTGCTGC  
ATAATCAGCTACGGTGGTGC GGACTACAAGCGAATCACGCTGAAAGTCAATGCCCCA

TACCGCAACCAATGCAATTTCCGTGGATCCAGCCACTTCTGAGCATGAACTA  
 ATATGTCAGGCCGAGGGTTATCCAGAAGCTGAGGTAATCTGGACAAACAGTGACCAC  
 CAACCCGTGAGTGGGAAGAGAAGTGTCAACCACTTCCCGGACAGAGGGGATGCTTCTC  
 AATGTGACCAGCAGTCTGAGGGTCAACGCCACAGCGAATGATGTTTTCTACTGTACG  
 TTTTGGAGATCACAGCCAGGGCAAACACACAGCGGAGCTGATCATCCCAGAACTG  
 CCTGCAACACATCCTCCACAGAACAGGACTGGTTGTAAGCCTTGCATATGTACAGTC  
 CCAGAAGTATCATCTGTCTTTCATCTTCCCCCAAAGCCCAAGGATGTGCTCACCATT  
 ACTCTGACTCCTAAGGTCACGTGTGTTGTGGTAGACATCAGCAAGGATGATCCCGAG  
 GTCCAGTTCAGCTGGTTTGTAGATGATGTGGAGGTGCACACAGCTCAGACGCAACCC  
 CGGGAGGAGCAGTTCAACAGCACTTTCGCTCAGTCAGTGAAGTTCATCATGCAC  
 CAGGACTGGCTCAATGGCAAGGAGTTCAAATGCAGGGTCAACAGTGCAGCTTTCCT  
 GCCCCCATCGAGAAAACCATCTCCAAAACCAAAGGCAGACCGAAGGCTCCACAGGTG  
 TACACCATTCCACCTCCCAAGGAGCAGATGGCCAAGGATAAAGTCAGTCTGACCTGC  
 ATGATAACAGACTTCTTCCCTGAAGACATTACTGTGGAGTGGCAGTGGGAATGGGCAG  
 CCAGCGGAGAACTACAAGAACAATCAGCCCATCATGGACACAGATGGCTCTTACTTC  
 GTCTACAGCAAGCTCAATGTGCAGAAGAGCAACTGGGAGGCAGGAAATACTTTCACC  
 TGCTCTGTGTTACATGAGGGCCTGCACAACCACCATACTGAGAAGAGCCTCTCCCAC  
 TCTCCTGGTAAATTCTACACAGGAGTCTATATTCTGATCGGAGCCGGCGCCCTCATG  
 ATGCTGGTGGGCTTCCCTGGGCTGCTGCGGGGCTGTGCAGGAGTCCCAGTGCAAAAAG  
 AAGAAAAAGAAGAAGAAGACAAAGTGTGTAATTATGTAA (SEQ ID NO: 315) >Artificial  
 Sequence; **mPDL1-mFc-CD9tm2-KRAS**, Amino Acid

**MRIFAGIIFTACCHLLRAFTITAPEDLYVVEYGSNVTMECRFPVERELDLLALVVYW  
 EKEDEQVIQFVAGEEDLKQPQHSNFRGRASLPKDQLLKGNALQITDVKLQDAGVYCC  
 IISYGGADYKRITLKVNPYRKINQRISVDPATSEHELICQAEGYPEAEVIWTNSDH  
 QPVS GKR SVTTSRTEGMLLNVTSSLRVNATANDVFYCTFWRSQPGQNHTAELIPEL  
 PATHPPQNRTGCKPCICTVPEVSSVFIFPPKPKDVLTTITLTPKVTCVVVDISKDDPE  
 VQFSWFVDDVEVHTAQTQPREEQFNSTFRSVSELPIMHQDWLNGKEFKCRVNSAAFP  
 APIEKTISKTKGRPKAPQVYTIPPPKEQMAKDKVSLTCMITDFFPEDITVEWQWNGQ  
 PAENYKNTQPIMDTDGSYFVYSKLVQKSNWEAGNTFTCSVLHEGLHNHHTEKSLSH  
 SPGKFYTGVIYILIGAGALMMLVGFLGCCGAVQESQCKKKKKKKKTKCVIM** (SEQ ID NO:  
 316)

(98) In some embodiments of any of the aspects, the fusion polypeptides provided herein comprise two or more POI domains. The specific combinations of POI domains can be used to regulate inflammatory immune responses. Non-limiting examples of additive and synergistic combinations of POIs that can modulate inflammatory signaling pathways are provided in Table 5 (below).

(99) TABLE-US-00007 TABLE 5 Exemplary POI combinations and combined targets for modulating inflammation. COMBINED PUTATIVE ADDITIVE or POIs (LIGANDS) TARGETS SYNERGISTIC MOA PD-L1 or PD-L2 PD-1 Differential use of Shp HVEM BTLA phosphatases. BTLA inhibits both TCR and CD28 phosphorylation (via Shp1) while PD-1 inhibits CD28 phosphorylation (via Shp2). PD-L1 or PD-L2 PD-1 LAG-3 exerts differential FGL1 LAG-3 inhibitory impacts on various types of lymphocytes and shows synergy with PD-1 to inhibit immune responses. PD-L1 or PD-L2 PD-1 PD-1 and Tim-3 have non- CEACAM-1 or GAL9 TIM-3 redundant downstream signaling mechanisms. PD-L1 or PD-L2 PD-1 Differential use of Shp CD155 TIGIT phosphatases. Non-redundantly regulate T cell responses. PD-L1 or PD-L2 PD-1 PD-1 and VISTA non- VSIG3 VISTA redundantly regulate T cell responses. VISTA contains cytosolic SH3 binding domains for adapter proteins. CEACAM-1 or GAL9 TIM-3 TIGIT and TIM-3 have non- CD155 TIGIT redundant downstream signaling mechanisms. PD-L1 or PD-L2 PD-1 PD-1, LAG-3 and TIM-3 have FGL1 LAG-3 non-redundant downstream CEACAM-1 or GAL9 TIM-3 signaling mechanisms.

Methods of Preparing Extracellular Vesicle Compositions

(100) In another aspect, provided herein is a method of preparing an engineered extracellular vesicle provided herein. Generally, the method comprises providing a population of cells expressing a vector construct encoding one or more sticky binder (vesicle targeting domain) and one or more signaling domains (POI domain).

(101) The EVs provided herein can be isolated and purified from any biological source, e.g., cells. The cells that produce the engineered EVs provided herein can be from any viable non-human source or organism. Usually the organism is an animal, vertebrate, or mammal. In some embodiments, the cell described herein is from a human. The cells described herein can be from any tissue isolated from an organism by methods known in the art. The scientific literature provides guidance for one of ordinary skill in the art to isolate, prepare, and culture cells as necessary for use in the compositions and methods described herein. One of skill in the art can appreciate that the cell source of the EVs may alter the cellular protein expression and the native or endogenous cargo within the EV. It is contemplated herein that this can be leveraged for therapeutic effect depending on the disease or disorder being treated.

(102) In some embodiments, the population of cells has been altered by exposure to environmental conditions (e.g., hypoxia), small molecule addition, presence/absence of exogenous factors (e.g., growth factors, cytokines) at the time, or substantially contemporaneous with, isolating the plurality of artificial synapses in a manner altering the regulatory state of the cell. In various embodiments, the cells are HEK 293 cells, MSCs, PER.C, fibrosarcoma HT-1080 or HuH7 cell lines.

(103) The method comprises providing a population of cells and culturing the cells in serum-free or un-concentrated conditioned medium. This includes, for example, artificial synapses secreted into media as conditioned by a population of cells in culture, further including cell lines capable of serial passaging. In certain embodiments, the cells in culture are grown to 10, 20, 30, 40, 50, 60, 70, 80, 90, or 90% or more confluency when artificial synapses (engineered EVs) are isolated.

(104) The methods provided herein further comprise contacting the cells provided herein with a nucleic acid vector encoding the at least one fusion polypeptide provided herein. The vector can be added to the cell culture medium of the cells by methods known in the art and discussed further below.

(105) A vector is a nucleic acid construct designed for delivery to a host cell or for transfer of genetic material between different host cells. As used herein, a vector can be viral or non-viral. The term “vector” encompasses any genetic element that is capable of replication when associated with the proper control elements and that can transfer genetic material to cells. A vector can include, but is not limited to, a cloning vector, an expression vector, a plasmid, phage, transposon, cosmid, artificial chromosome, virus, virion, etc. In some embodiments of any of the aspects, the vector is selected from the group consisting of: a plasmid, a cosmid and a viral vector.

(106) “Expression” refers to the cellular processes involved in producing RNA and proteins and as appropriate, secreting proteins, including where applicable, but not limited to, for example, transcription, transcript processing, translation and protein folding, modification and processing. “Expression products” include RNA transcribed from a gene, and polypeptides obtained by translation of mRNA transcribed from a gene.

(107) In some embodiments, a vector is capable of driving expression of one or more sequences in a mammalian cell; i.e., the vector is a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, 1987. *Nature* 329:840) and pMT2PC (Kaufman, et al., 1987. *EMBO J.* 6:187-195). When used in mammalian cells, the expression vector's control functions are typically provided by one or more regulatory elements. For example, commonly used promoters are derived from polyoma, adenovirus 2, cytomegalovirus, simian virus 40, and others disclosed herein and known in the art. For other suitable expression systems for both prokaryotic and eukaryotic cells see, e.g., Chapters 16 and 17 of Sambrook, et al., *MOLECULAR CLONING: A LABORATORY MANUAL*. 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor

Laboratory Press, Cold Spring Harbor, N.Y., 1989.

(108) In some embodiments, the recombinant expression vector is capable of directing expression of the exogenous fusion polypeptide nucleic acid sequence preferentially in a particular cell type (e.g., via tissue-specific regulatory elements).

(109) Tissue-specific and inducible regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert, et al., 1987. *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton, 1988. *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore, 1989. *EMBO J.* 8:729-733) and immunoglobulins (Baneiji, et al., 1983. *Cell* 33:729-740; Queen and Baltimore, 1983. *Cell* 33:741-748), neuron-specific promoters (e.g., the neurofilament promoter; Byrne and Ruddle, 1989. *Proc. Natl. Acad. Sci. USA* 86:5473-5477), pancreas-specific promoters (Edlund, et al., 1985. *Science* 230:912-916), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Pat. No. 4,873,316 and European Application Publication No. 264, 166). Developmentally-regulated promoters are also encompassed, e.g., the murine hox promoters (Kessel and Gruss, 1990. *Science* 249:374-379) and the  $\alpha$ -fetoprotein promoter (Campes and Tilghman, 1989. *Genes Dev.* 3:537-546).

(110) In some embodiments, the at least one nucleic acid sequence described herein is delivered to the cell described herein via an integrating vector. Integrating vectors have their delivered genetic material (or a copy of it) permanently incorporated into a host cell chromosome. Non-integrating vectors remain episomal which means the nucleic acid contained therein is never integrated into a host cell chromosome. Examples of integrating vectors include retroviral vectors, lentiviral vectors, hybrid adenoviral vectors, and herpes simplex viral vectors.

(111) In some embodiments, the at least one nucleic acid sequence described herein is delivered to the cell described herein via a non-integrative vector. Non-integrative vectors include non-integrative viral vectors. Non-integrative viral vectors eliminate one of the primary risks posed by integrative retroviruses, as they do not incorporate their genome into the host DNA. One example is the Epstein Barr oriP/Nuclear Antigen-1 ("EBNA1") vector, which is capable of limited self-replication and known to function in mammalian cells. Containing two elements from Epstein-Barr virus, oriP and EBNA1, binding of the EBNA1 protein to the virus replicon region oriP maintains a relatively long-term episomal presence of plasmids in mammalian cells. This particular feature of the oriP/EBNA1 vector makes it ideal for generation of integration-free host cells. Other non-integrative viral vectors include adenoviral vectors and the adeno-associated viral (AAV) vectors.

(112) Another non-integrative viral vector is RNA Sendai viral vector, which can produce protein without entering the nucleus of an infected cell. The F-deficient Sendai virus vector remains in the cytoplasm of infected cells for a few passages, but is diluted out quickly and completely lost after several passages (e.g., 10 passages). This permits a self-limiting transient expression of a chosen heterologous gene or genes in a target cell. This aspect can be helpful, e.g., for the transient introduction of reprogramming factors, among other uses. As noted above, in some embodiments, the nucleic acid sequence described herein is expressed in the cells from a viral vector.

(113) A "viral vector" includes a nucleic acid vector construct that includes at least one element of viral origin and has the capacity to be packaged into a viral vector particle. The viral vector can contain a nucleic acid encoding a polypeptide described herein in place of non-essential viral genes. The vector and/or particle can be utilized for the purpose of transferring nucleic acids into cells either in vitro or in vivo.

(114) The nucleic acids described herein can be delivered using any transfection reagent or other physical means that facilitates entry of nucleic acids into a cell. Methods of non-viral delivery of nucleic acids include lipofection, nucleofection, microinjection, electroporation, biolistics, virosomes, liposomes, immunoliposomes, polycation or lipid: nucleic acid conjugates, naked DNA, artificial virions, and agent-enhanced uptake of DNA. Lipofection is described in e.g., U.S. Pat. Nos. 5,049,386, 4,946,787; and 4,897,355) and lipofection reagents are sold commercially (e.g.,

Transfectam™ and Lipofectin™). Cationic and neutral lipids that are suitable for efficient receptor-recognition lipofection of polynucleotides include those of Felgner, WO 91/17424; WO 91/16024. Delivery can be to cells (e.g., in vitro or ex vivo administration) or target tissues (e.g., in vivo administration).

(115) The preparation of lipid: nucleic acid complexes, including targeted liposomes such as immunolipid complexes, is well known to one of skill in the art (see, e.g., Crystal, Science 270:404-410 (1995); Blaese et al., Cancer Gene Ther. 2:291-297 (1995); Behr et al., Bioconjugate Chem. 5:382-389 (1994); Remy et al., Bioconjugate Chem. 5:647-654 (1994); Gao et al., Gene Therapy 2:710-722 (1995); Ahmad et al., Cancer Res. 52:4817-4820 (1992); U.S. Pat. Nos. 4,186,183, 4,217,344, 4,235,871, 4,261,975, 4,485,054, 4,501,728, 4,774,085, 4,837,028, and 4,946,787).

(116) An “agent that increases cellular uptake” is a molecule that facilitates transport of a molecule, e.g., nucleic acid, or peptide or polypeptide, or other molecule that does not otherwise efficiently transit the cell membrane across a lipid membrane. For example, a nucleic acid can be conjugated to a lipophilic compound (e.g., cholesterol, tocopherol, etc.), a cell penetrating peptide (CPP) (e.g., penetratin, TAT, Syn1B, etc.), or a polyamine (e.g., spermine). Further examples of agents that increase cellular uptake are disclosed, for example, in Winkler (2013). *Oligonucleotide conjugates for therapeutic applications. Ther. Deliv.* 4 (7); 791-809. The one or more nucleic acid sequences encoding the fusion polypeptides provided herein can be delivered to the cell by any method discussed above or known in the art.

(117) In some embodiments of any of the aspects, the vectors provided herein comprise a nucleic acid modification by methods known in the art. In some embodiments, the cell can be genetically manipulated to express one or more vectors, each encoding one or more vesicle targeting domains and/or one or more signaling domains. In certain embodiments, the population of cells has been genetically manipulated. This includes, for example, knockout (KO) or transgenic (TG) cell lines, wherein an endogenous gene has been removed and/or an exogenous introduced in a stable, persistent manner. In certain embodiments, this further includes transient knockdown of one or more genes and associated coding and non-coding transcripts within the population of cells, via any number of methods known in the art, such as introduction of dsRNA, siRNA, microRNA, etc. This further includes transient expression of one or more genes and associated coding and non-coding transcripts within the population of cells, via any number of methods known in the art, such as introduction of a vector, plasmid, artificial plasmid, replicative and/or non-replicative virus, etc.

(118) In certain embodiments the cell population has been manipulated to knockout the expression of one or more endogenous gene sequences that encode for metalloendopeptidases. In certain embodiments the cell population has been manipulated to knockout the expression of one or more endogenous gene sequences that code for metalloproteinases. In certain embodiments the cell population has been manipulated to knockout the expression of one or more endogenous gene sequences that encode for a disintegrin and metalloproteinase (ADAM). For example, the cell population can be manipulated to knock of the expression of one or more gene sequences that encode for ADAM1, ADAM2, ADAM7, ADAM8, ADAM9, ADAM10, ADAM11, ADAM12, ADAM15, ADAM17, ADAM18, ADAM19, ADAM20, ADAM21, ADAM22, ADAM23, ADAM28, ADAM29, ADAM30, ADAM33, etc.

(119) In certain embodiments the cell population has been manipulated to knockout the expression of one or more endogenous genes that encode for enzymes that hydrolyze the inositol phosphate linkage in proteins anchored by phosphatidylinositol glycans, thereby preventing the release of proteins attached to the plasma membrane via GPI anchors. For example, the cell population can be manipulated to knock of the expression of phosphatidylinositol-glycan-specific phospholipase D (GPLD1).

(120) In certain embodiments, the population of cells has been genetically manipulated. This includes, for example, knock-in of an exogenous genetic sequence, wherein the exogenous genetic

sequence is expressed in a stable, persistent manner. In certain embodiments, the cell population has been manipulated to knock-in recombinase recognition sequences (e.g., FRT), transgenic reporters such as antibiotic resistance genes, fluorescent or enzymatic reporter genes, etc. or the like.

(121) In some embodiments, the method comprises a step of isolating the engineered extracellular vesicles provided herein. Particulates within the medium are removed by a series of specific centrifugation steps and the media is filtered. The general method of isolating extracellular vesicles as provided herein is depicted in FIG. 21 of the working examples. Methods of isolating and purifying the extracellular vesicles and exosomes are known in the art and further described, e.g., in Whitford W, Guterstam P. Exosome manufacturing status. *Future Med Chem.* 2019 May; 11(10):1225-1236. doi: 10.4155/fmc-2018-0417. PMID: 31280675, Patel D B, Santoro M, Born L J, Fisher J P, Jay S M. Towards rationally designed biomanufacturing of therapeutic extracellular vesicles: impact of the bioproduction microenvironment. *Biotechnol Adv.* 2018 December; 36(8):2051-2059 . . . doi: 10.1016/j.biotechadv.2018.09.001. Epub 2018 Sep. 12. PMID: 30218694; PMCID: PMC6250573, Ng K S, Smith J A, McAteer M P, Mead B E, Ware J, Jackson F O, Carter A, Ferreira L, Bure K, Rowley J A, Reeve B, Brindley D A, Karp J M. Bioprocess decision support tool for scalable manufacture of extracellular vesicles. *Biotechnol Bioeng.* 2019 February; 116(2):307-319. doi: 10.1002/bit.26809. Epub 2018 Nov. 8. PMID: 30063243; PMCID: PMC6322973, Paganini C, Capasso Palmiero U, Pocsfalvi G, Touzet N, Bongiovanni A, Arosio P. Scalable Production and Isolation of Extracellular Vesicles: Available Sources and Lessons from Current Industrial Bioprocesses. *Biotechnol J.* 2019 October; 14(10):e1800528. doi: 10.1002/biot.201800528. Epub 2019 Jul. 8. PMID: 31140717, which are incorporated herein by reference in their entireties.

(122) In some embodiments, isolating the plurality of engineered EVs (artificial synapses) includes precipitation, centrifugation, filtration, immuno-separation, tangential flow, liquid chromatography, and/or flow fractionation. For example, differential ultracentrifugation has become a technique wherein secreted exosomes are isolated from the supernatants of cultured cells. This approach allows for separation of exosomes from non-membranous particles, by exploiting their relatively low buoyant density. Size exclusion allows for their separation from biochemically similar, but biophysically different microvesicles, which possess larger diameters of up to 1,000 nm. Differences in floatation velocity further allows for separation of differentially sized exosomes. In general, exosome sizes will possess a diameter ranging from 30-300 nm, including sizes of 30-150 nm. Further purification may rely on specific properties of the particular exosomes of interest. This includes, for example, use of immunoabsorption with a protein of interest to select specific vesicles with exoplasmic or outward orientations.

(123) Among current methods (differential centrifugation, discontinuous density gradients, immunoaffinity, ultrafiltration and liquid chromatography (e.g., fast protein liquid chromatography (FPLC)), differential ultracentrifugation is the most commonly used for exosome isolation. This technique utilizes increasing centrifugal force from 2000×g to 10,000×g to separate the medium- and larger-sized particles and cell debris from the exosome pellet at 100,000×g. Centrifugation alone allows for significant separation/collection of exosomes from a conditioned medium, although it is insufficient to remove various protein aggregates, genetic materials, particulates from media and cell debris that are common contaminants. Enhanced specificity of exosome purification may deploy sequential centrifugation in combination with ultrafiltration, or equilibrium density gradient centrifugation in a sucrose density gradient, to provide for the greater purity of the exosome preparation (floatation density 1.1-1.2 g/ml) or application of a discrete sugar cushion in preparation.

(124) Ultrafiltration can be used to purify exosomes without compromising their biological activity. Membranes with different pore sizes—such as 100 kDa molecular weight cut-off (MWCO) or 300 kDa MWCO and gel filtration to eliminate smaller particles—have been used to avoid the use of a

nonneutral pH or non-physiological salt concentration. Currently available tangential flow filtration (TFF) systems are scalable (to >10,000 L), allowing one to not only purify, but concentrate the exosome fractions, and such approaches are less time consuming than differential centrifugation. Liquid Chromatography can also be used to purify exosomes to homogeneously sized particles and preserve their biological activity as the preparation is maintained at a physiological pH and salt concentration.

(125) Other chemical methods have exploit differential solubility of exosomes for precipitation techniques, addition to volume-excluding polymers (e.g., polyethylene glycols (PEGs)), possibly combined additional rounds of centrifugation or filtration. For example, a precipitation reagent, ExoQuick®, can be added to conditioned cell media to quickly and rapidly precipitate a population of exosomes, although re-suspension of pellets prepared via this technique may be difficult. Flow field-flow fractionation (FIFFF) is an elution-based technique that is used to separate and characterize macromolecules (e.g., proteins) and nano- to micro-sized particles (e.g., organelles and cells) and which has been successfully applied to fractionate exosomes from culture media.

(126) Beyond these techniques relying on general biochemical and biophysical features, focused techniques may be applied to isolated specific exosomes of interest. This includes relying on antibody immunoaffinity to recognizing certain exosome-associated antigens. Conjugation to magnetic beads, chromatography matrices, plates or microfluidic devices allows isolating of specific exosome populations of interest as may be related to their production from a parent cell of interest or associated cellular regulatory state. Other affinity-capture methods use lectins which bind to specific saccharide residues on the exosome surface.

(127) In several embodiments, isolating a plurality of artificial synapses from the population of cells includes centrifugation of the cells and/or media conditioned by the cells. In several embodiments, ultracentrifugation is used. In several embodiments, isolating a plurality of artificial synapses from the population of cells is via size-exclusion filtration. In other embodiments, isolating a plurality of artificial synapses from the population of cells includes use of discontinuous density gradients, immunoaffinity, ultrafiltration, tangential flow and/or liquid chromatography.

(128) In certain embodiments, differential ultracentrifugation includes using centrifugal force from 1000-2000×g, 2000-3000×g, 3000-4000×g, 4000-5000×g, 5000×g-6000×g, 6000-7000×g, 7000-8000×g, 8000-9000×g, 9000-10,000×g, to 10,000×g or more to separate larger-sized particles from a plurality of artificial synapses derived from the cells.

(129) In other embodiments, isolating a plurality of artificial synapses from the population of cells includes use of filtration or ultrafiltration. In certain embodiments, a size exclusion membrane with different pore sizes is used. For example, a size exclusion membrane can include use of a filter with a pore size of 0.1-0.5 micron (μm), 0.5-1.0 μm, 1-2.5 μm, 2.5-5 μm, 5 or more μm. In certain embodiments, the pore size is about 0.2 μm. In certain embodiments, filtration or ultrafiltration includes size exclusion ranging from 100-500 daltons (Da), 500-1 kDa, 1-2 kDa, 2-5 kDa, 5-10 kDa, 10-25 kDa, 25-50 kDa, 50-100 kDa, 100-250 kDa, 250-500 kDa, 500 or more kDa. In certain embodiments, the size exclusion is for about 2-5 kDa. In certain embodiments, the size exclusion is for about 3 kDa. In other embodiments, filtration or ultrafiltration includes size exclusion includes use of hollow fiber membranes capable of isolating particles ranging from 100-500 daltons (Da), 500-1 kDa, 1-2 kDa, 2-5 kDa, 5-10 kDa, 10-25 kDa, 25-50 kDa, 50-100 kDa, 100-250 kDa, 250-500 kDa, 500 or more kDa. In certain embodiments, the size exclusion is for about 2-5 kDa. In certain embodiments, the size exclusion is for about 3 kDa. In other embodiments, a molecular weight cut-off (MWCO) gel filtration capable of isolating particles ranging from 100-500 daltons (Da), 500-1 kDa, 1-2 kDa, 2-5 kDa, 5-10 kDa, 10-25 kDa, 25-50 kDa, 50-100 kDa, 100-250 kDa, 250-500 kDa, 500 or more kDa. In certain embodiments, the size exclusion is for about 2-5 kDa. In certain embodiments, the size exclusion is for about 3 kDa. In various embodiments, such systems are used in combination with variable fluid flow systems. In certain embodiments, a size exclusion membrane with different pore sizes is used to purify extracellular vesicles from a solution

comprising undesirable proteins or nucleic acids.

(130) In other embodiments, isolating a plurality of artificial synapses from the population of cells includes use of tangential flow filtration (TFF) systems are used purify and/or concentrate the exosome fractions. In other embodiments, isolating a plurality of artificial synapses from the population of cells includes use of liquid chromatography can also be used to purify artificial synapses to homogeneously sized particles. In various embodiments, density gradients as used, such as centrifugation in a sucrose density gradient or application of a discrete sugar cushion in preparation.

(131) In other embodiments, isolating a plurality of artificial synapses from the population of cells includes use of a precipitation reagent. For example, a precipitation reagent, ExoQuick®, can be added to conditioned cell media to quickly and rapidly precipitate a population of artificial synapses. In other embodiments, isolating a plurality of artificial synapses from the population of cells includes use of volume-excluding polymers (e.g., polyethylene glycols (PEGs)) are used. In another embodiment, isolating a plurality of artificial synapses from the population of cells includes use of flow field-flow fractionation (FIFFF), an elution-based technique.

(132) In certain embodiments, isolating a plurality of artificial synapses from the population of cells includes use of one or more capture agents to isolate one or more artificial synapses possessing specific biomarkers or containing particular biological molecules. In one embodiment, one or more capture agents include at least one antibody. For example, antibody immunoaffinity recognizing exosome-associated antigens is used to capture specific artificial synapses. In other embodiments, the at least one antibody are conjugated to a fixed surface, such as magnetic beads, chromatography matrices, plates or microfluidic devices, thereby allowing isolation of the specific exosome populations of interest. In other embodiments, isolating a plurality of artificial synapses from the population of cells includes use of one or more capture agents that is not an antibody. This includes, for example, use of a “bait” molecule presenting an antigenic feature complementary to a corresponding molecule of interest on the exosome surface, such as a receptor or other coupling molecule. In one embodiment, the non-antibody capture agent is a lectin capable of binding to polysaccharide residues on the exosome surface.

(133) In other embodiments, isolating a plurality of artificial synapses from the population of cells includes use of ion exchange chromatography. In other embodiments, isolating a plurality of artificial synapses from the population of cells includes use of anion exchange chromatography. In other embodiments, isolating a plurality of artificial synapses from the population of cells includes use of cation exchange chromatography. In certain embodiments, ion exchange chromatography comprises a chromatography resin with a functional group selected from the group consisting of diethylaminoethyl (DEAE), quaternary aminoethyl (QAE), quaternary ammonium (Q), carboxymethyl (CM), sulfopropyl (SP), or methyl sulfate(S). In certain embodiments, ion exchange chromatography comprises a chromatography resin which may have properties of a weak acid, strong acid, weak base, or strong basic. In certain embodiments, ion exchange chromatography comprises a chromatography selected from the group consisting of DEAE cellulose, DEAE Sephadex, Mono Q, Mini Q, HiTrap Capto, Capto Core 700, HiPrep Q, QAE Sephadex, Q Sepharose, CM Cellulose, SP Sepharose, SOURCE S, EAH-Sepharose, sulfoxyethyl cellulose, CM Sephadex, or CM Sepharose. Isolating a plurality of artificial synapses can be prepared by any of a variety of ion exchange chromatography techniques that are known in the art.

(134) In other embodiments, isolating a plurality of artificial synapses from the population of cells includes use of a nuclease enzyme (e.g., a DNase or RNase). For example, a working concentration of Benzonase® nuclease may be added to an extracellular vesicle sample preparation in the presence of a divalent cation, for example 1-2 mM Mg.sup.2+, 2-5 mM Mg.sup.2+, 10-20 mM Mg.sup.2+, 20-50 mM Mg.sup.2+, 50-100 mM Mg.sup.2+, or more than 100 mM Mg.sup.2+.

(135) Following isolation and purification of the engineered EVs provided herein, EVs can be further evaluated for the desired structural and functional properties by methods known in the art.



For example, the engineered exosomes provided herein can be assayed for functional activity on a target cell using a cell-based bioassays (e.g., those commercially available, PROMEGA® DISCOVERX®), ligand-receptor binding assays, vesicle flow cytometric assays, enzyme-linked immunosorbent assays, tunable resistive pulse sensing (TRPS), nanoparticle tracking analysis (NTA), surface plasmon resonance (SPR), nucleotide sequencing, lipidomics, proteomics, colorimetric assays, fluorescence assays, luminescence assays, immunoblotting, radioimmunoassays, electron microscopy, or EV automated analysis (e.g., EXOVIEW®). Additional methods of characterizing EVs are found, e.g., in Zhang Y, Bi J, Huang J, Tang Y, Du S, Li P. Exosome: A Review of Its Classification, Isolation Techniques, Storage, Diagnostic and Targeted Therapy Applications. *Int J Nanomedicine*. 2020 Sep. 22; 15:6917-6934. doi: 10.2147/IJN.S264498. PMID: 33061359; PMCID: PMC7519827, Kluszczyńska K, Czernek L, Cypriak W, Pęczek Ł, Dürlichler M. Methods for the Determination of the Purity of Exosomes. *Curr Pharm Des*. 2019; 25(42):4464-4485. doi: 10.2174/1381612825666191206162712. PMID: 31808383, Nolan J P, Duggan E. Analysis of Individual Extracellular Vesicles by Flow Cytometry. *Methods Mol Biol*. 2018; 1678:79-92. doi: 10.1007/978-1-4939-7346-0\_5. PMID: 29071676; Doyle L M, Wang M Z. Overview of Extracellular Vesicles, Their Origin, Composition, Purpose, and Methods for Exosome Isolation and Analysis. *Cells*. 2019 Jul. 15; 8(7):727. doi: 10.3390/cells8070727. PMID: 31311206; PMCID: PMC6678302, Pugholm L H, Revenfeld A L, Søndergaard E K, Jørgensen M M. Antibody-Based Assays for Phenotyping of Extracellular Vesicles. *Biomed Res Int*. 2015; 2015:524817. doi: 10.1155/2015/524817. Epub 2015 Dec. 3. PMID: 26770974; PMCID: PMC4681819, Shao H, Im H, Castro C M, Breakefield X, Weissleder R, Lee H. New Technologies for Analysis of Extracellular Vesicles. *Chem Rev*. 2018 Feb. 28; 118(4):1917-1950. doi: 10.1021/acs.chemrev.7b00534. Epub 2018 Jan. 31. PMID: 29384376; PMCID: PMC6029891, which are incorporated herein by reference in their entireties.

#### Pharmaceutical Compositions

(136) Provided herein are compositions comprising the engineered extracellular vesicles (artificial synapses) provided herein.

(137) In one aspect, provided herein is a composition comprising: a plurality of the engineered extracellular vesicles provided herein. In some embodiments of any of the aspects, the compositions and engineered EVs provided herein further comprise a pharmaceutically acceptable carrier.

(138) For clinical use of the methods and compositions described herein, administration of the engineered EVs/artificial synapses provided herein can include formulation into pharmaceutical compositions or pharmaceutical formulations for parenteral administration, e.g., intravenous; mucosal, e.g., intranasal; ocular, or other mode of administration. In some embodiments, the engineered EVs described herein can be administered along with any pharmaceutically acceptable carrier compound, material, or composition which results in an effective treatment in the subject. Thus, a pharmaceutical formulation for use in the methods described herein can contain the engineered EVs described herein in combination with one or more pharmaceutically acceptable ingredients. The phrase “pharmaceutically acceptable” refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio. The phrase “pharmaceutically acceptable carrier” as used herein means a pharmaceutically acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent, media, encapsulating material, manufacturing aid (e.g., lubricant, talc magnesium, calcium or zinc stearate, or steric acid), or solvent encapsulating material, involved in maintaining the stability, solubility, or activity of, an engineered EV as described herein. Each carrier must be “acceptable” in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient. The terms “excipient,” “carrier,” “pharmaceutically

acceptable carrier” or the like are used interchangeably herein.

(139) The engineered EVs provided herein can be formulated for administration of the compound to a subject in solid, liquid, or gel form, including those adapted for the following: (1) parenteral administration, for example, by subcutaneous, intramuscular, intravenous or epidural injection as, for example, a sterile solution or suspension, or sustained-release formulation; (2) transdermally; (3) transmucosally; (4) via bronchoalveolar lavage.

(140) In some embodiments, the compositions described herein comprise a particle or polymer-based vehicle. Exemplary particle or polymer-based vehicles include, but are not limited to, nanoparticles, microparticles, polymer microspheres, or polymer-drug conjugates.

(141) In one embodiment of any of the aspects, the compositions described herein further comprise a lipid vehicle. Exemplary lipid vehicles include, but are not limited to, liposomes, phospholipids, micelles, lipid emulsions, and lipid-drug complexes.

(142) Formulations can be adapted for delivery to the airway, e.g., to address respiratory inflammation. Such formulations can be adapted for delivery as an aerosol, e.g., for inhalation. In some embodiments, the compositions described herein are formulated for aerosol administration, nebulizer administration, tracheal lavage administration, or for a pulmonary delivery device.

(143) As used herein, the term “pulmonary delivery device” refers to a device used to deliver a therapeutic dose of a composition of the present invention to the respiratory system including, but not limited to, a nebulizer, metered-dose inhaler, or dry powder inhaler.

(144) Examples of nebulizers include, but are not limited to, soft mist inhalers (for example Respimat® Boehringer Ingelheim) jet nebulizers (use compressed gas or air), ultrasonic nebulizers (produce aerosols using a piezoelectric crystal vibrating at high frequencies), and vibrating mesh nebulizers.

(145) As used herein, the term “jet nebulizer” refers to a device that flows compressed air or gas through a composition of the present invention for aerosolization. The aerosolized composition of the present invention may be inhaled by a patient. Jet nebulizer may include, but is not limited to, jet nebulizers with a corrugated tube, jet nebulizers with a collection bag, breath enhanced jet nebulizers, breath actuated jet nebulizers, and metered-dose inhalers. Examples of jet nebulizers include, but are not limited to, Circulaire (Westmed INC, Tucson, AZ), Pari Inhalierboy (PARI, Midlothian, VA), Pari LC Plus (PARI, Midlothian, VA), NebuTech (Salter Labs, Arvin, CA), AeroEclipse (Monaghan/Trudell Medical International, London, Ontario, Canada), and Maxin MA-2 (MA-2; Clinova Medical AB, Malmö, Sweden). Examples of ultrasonic nebulizers include, but are not limited to, De Vilbiss-Pulmosonic (Somerset, PA), Omron-Microair (Omron, Kyoto, Japan), Omron NE-U17 (Omron, Kyoto, Japan), Rhone Poulenc-Rorer-Fisoneb (Sanofi, Paris, France), and Beurer Nebulizer IH30 (Beurer GmbH, Neu-Ulm, Germany).

(146) As used herein, the term “mesh nebulizer” refers to forcing a liquid, gel, fluid, solution, tincture, or the like through apertures in a mesh or aperture plate to generate aerosol. Mesh nebulizer may include, but is not limited to, active mesh nebulizers and passive mesh nebulizers. Examples of active mesh nebulizers include, but is not limited to, Aeroneb® (Aerogen, Galway, Ireland) and eFlow® (PARI, Midlothian, VA). Examples of passive mesh nebulizers are, but not limited to, I-neb (Philips Respironics, Newark, USA), AKITA (Activaero, Gemunden/Wohra, Germany), and Microair NE-U22® (Omron, Kyoto, Japan).

(147) For use as aerosols, the compositions described herein can be prepared in a solution or suspension and may be packaged in a pressurized aerosol container together with suitable propellants, for example, hydrocarbon propellants like propane, butane, or isobutane with conventional excipients.

(148) The engineered EVs provided herein can also be administered in a non-pressurized form such as in a nebulizer or atomizer that reduces a liquid to a fine spray. Preferably, by such nebulization small liquid droplets of uniform size are produced from a larger body of liquid in a controlled manner. Nebulization can be achieved by any suitable means therefor, including by using many

nebulizers known and marketed today. For example, an AEROMIST™ pneumatic nebulizer available from Inhalation Plastic, Inc. of Niles, Ill.

(149) When the active ingredients are adapted to be administered, either together or individually, via nebulizer(s) they can be in the form of a nebulized aqueous suspension or solution, with or without a suitable pH or tonicity adjustment, either as a unit dose or multi-dose device.

(150) Furthermore, any suitable gas can be used to apply pressure during the nebulization, with preferred gases to date being those which are chemically inert. Exemplary gases including, but are not limited to, nitrogen, argon, or helium can be used to advantage.

(151) In some embodiments, the compositions described herein can also be administered directly to the airways in the form of a dry powder. Thus, the engineered EVs can be administered via an inhaler. Exemplary inhalers include metered dose inhalers and dry powdered inhalers.

(152) A metered dose inhaler or “MDI” is a pressure resistant canister or container filled with a product such as a pharmaceutical composition dissolved in a liquefied propellant or micronized particles suspended in a liquefied propellant. The propellants which can be used include chlorofluorocarbons, hydrocarbons or hydrofluoroalkanes. Commonly used propellants are P134a (tetrafluoroethane) and P227 (heptafluoropropane) each of which may be used alone or in combination. They are optionally used in combination with one or more other propellants and/or one or more surfactants and/or one or more other excipients, for example ethanol, a lubricant, an anti-oxidant and/or a stabilizing agent.

(153) As used herein, the term “dry powder inhaler” refers to a device that delivers a therapeutic dose of a composition of the present invention in a powdered form without propellants to the respiratory system. A dry powder inhaler (i.e., Turbuhaler™ (Astra AB)) is a system operable with a source of pressurized air to produce dry powder particles of a pharmaceutical composition that is compacted into a very small volume. Examples of dry powder inhalers include, but are not limited to, Spinhaler® (Fisons Pharmaceuticals, Rochester, NY), Rotahaler® (GlaxoSmithKline, NC), Turbuhaler® (AstraZeneca, UK), and Diskhaler® (GlaxoSmithKline, NC).

(154) Dry powder aerosols for inhalation therapy are generally produced with mean diameters primarily in the range of <5 μm. As the diameter of particles exceeds 3 μm, there is increasingly less phagocytosis by macrophages. However, increasing the particle size also has been found to minimize the probability of particles (possessing standard mass density) entering the airways and acini due to excessive deposition in the oropharyngeal or nasal regions.

(155) Suitable powder compositions include, by way of illustration, powdered preparations including the engineered EVs described herein. These can be intermixed with lactose, or other inert powders acceptable for intrabronchial administration. The powder compositions can be administered via an aerosol dispenser or encased in a breakable capsule which may be inserted by the patient or clinician into a device that punctures the capsule and blows the powder out in a steady stream suitable for inhalation. The compositions can include propellants, surfactants, and co-solvents and may be filled into conventional aerosol containers that are closed by a suitable metering valve.

(156) Aerosols for the delivery to the respiratory tract are described, for example, by Adjei, A. and Garren, J. *Pharm. Res.*, 1:565-569 (1990); Zanen, P. and Lamm, J.-W. *J. Int. J. Pharm.*, 114:111-115 (1995); Gonda, I. “Aerosols for delivery of therapeutic and diagnostic agents to the respiratory tract,” in *Critical Reviews in Therapeutic Drug Carrier Systems*, 6:273-313 (1990); Anderson et al., *Am. Rev. Respir. Dis.*, 140:1317-1324 (1989)) and have potential for the systemic delivery of peptides and proteins as well (Patton and Platz, *Advanced Drug Delivery Reviews*, 8:179-196 (1992)); Timsina et al., *Int. J. Pharm.*, 101:1-13 (1995); and Tansey, I. P., *Spray Technol. Market*, 4:26-29 (1994); French, D. L., Edwards, D. A. and Niven, R. W., *Aerosol Sci.*, 27:769-783 (1996); Visser, J., *Powder Technology* 58:1-10 (1989)); Rudt, S. and R. H. Muller, *J. Controlled Release*, 22:263-272 (1992); Tabata, Y. and Y. Ikada, *Biomed. Mater. Res.*, 22:837-858 (1988); Wall, D. A., *Drug Delivery*, 2:10 1-20 (1995); Patton, J. and Platz, R., *Adv. Drug Del. Rev.*, 8:179-196 (1992);

Bryon, P., Adv. Drug. Del. Rev., 5:107-132 (1990); Patton, J. S., et al., Controlled Release, 28:15 79-85 (1994); Damms, B. and Bains, W., Nature Biotechnology (1996); Niven, R. W., et al., Pharm. Res., 12 (9): 1343-1349 (1995); and Kobayashi, S., et al., Pharm. Res., 13 (1): 80-83 (1996), the contents of each of which are incorporated herein by reference in their entirety.

(157) Microemulsification technology can improve bioavailability of some lipophilic (water insoluble) pharmaceutical agents. Examples include Trimetrine (Dordunoo, S. K., et al., Drug Development and Industrial Pharmacy, 17 (12), 1685-1713, 1991 and REV 5901 (Sheen, P. C., et al., J Pharm Sci 80 (7), 712-714, 1991). Among other things, microemulsification provides enhanced bioavailability by preferentially directing absorption to the lymphatic system instead of the circulatory system, which thereby bypasses the liver, and prevents destruction of the cell-based compositions in the hepatobiliary circulation.

(158) The engineered EVs described herein can be formulated with an amphiphilic carrier. Amphiphilic carriers are saturated and monounsaturated polyethyleneglycolized fatty acid glycerides, such as those obtained from fully or partially hydrogenated various vegetable oils. Such oils may advantageously consist of tri-, di-, and mono-fatty acid glycerides and di- and mono-polyethyleneglycol esters of the corresponding fatty acids, with a particularly preferred fatty acid composition including capric acid 4-10, capric acid 3-9, lauric acid 40-50, myristic acid 14-24, palmitic acid 4-14 and stearic acid 5-15%. Another useful class of amphiphilic carriers includes partially esterified sorbitan and/or sorbitol, with saturated or mono-unsaturated fatty acids (SPAN-series) or corresponding ethoxylated analogs (TWEEN-series).

(159) Commercially available amphiphilic carriers are particularly contemplated, including Gelucire-series, Labrafil, Labrasol, or Lauroglycol (all manufactured and distributed by Gattefosse Corporation, Saint Priest, France), PEG-mono-oleate, PEG-di-oleate, PEG-mono-laurate and di-laurate, Lecithin, Polysorbate 80, etc. (produced and distributed by a number of companies in USA and worldwide).

(160) The engineered EV compositions provided herein can be formulated with hydrophilic polymers. Hydrophilic polymers are water-soluble, can be covalently attached to a vesicle-forming lipid, and which are tolerated in vivo without toxic effects (i.e., are biocompatible). Suitable polymers include polyethylene glycol (PEG), polylactic (also termed polylactide), polyglycolic acid (also termed polyglycolide), a polylactic-polyglycolic acid copolymer, and polyvinyl alcohol. Other hydrophilic polymers which may be suitable include polyvinylpyrrolidone, polymethoxazoline, polyethyloxazoline, polyhydroxypropyl methacrylamide, polymethacrylamide, polydimethylacrylamide, and derivatized celluloses such as hydroxymethylcellulose or hydroxyethylcellulose.

(161) In certain embodiments, a pharmaceutical composition as described herein comprises a biocompatible polymer selected from the group consisting of polyamides, polycarbonates, polyalkylenes, polymers of acrylic and methacrylic esters, polyvinyl polymers, polyglycolides, polysiloxanes, polyurethanes and co-polymers thereof, celluloses, polypropylene, polyethylenes, polystyrene, polymers of lactic acid and glycolic acid, polyanhydrides, poly(ortho) esters, poly(butic acid), poly(valeric acid), poly(lactide-co-caprolactone), polysaccharides, proteins, polyhyaluronic acids, polycyanoacrylates, and blends, mixtures, or copolymers thereof.

(162) In certain embodiments, a pharmaceutical composition described herein is formulated as a liposome. Liposomes can be prepared by any of a variety of techniques that are known in the art. See, e.g., U.S. Pat. No. 4,235,871; Published PCT applications WO 96/14057; New RRC, Liposomes: A practical approach, IRL Press, Oxford (1990), pages 33-104; Lasic DD, Liposomes from physics to applications, Elsevier Science Publishers BV, Amsterdam, 1993.

(163) Therapeutic formulations of the engineered EV compositions as described herein can be prepared for storage by with optional pharmaceutically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are

nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g., Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG).

(164) Vaccine or other pharmaceutical compositions comprising an engineered EV composition as described herein can contain a pharmaceutically acceptable salt, typically, e.g., sodium chloride, and preferably at about physiological concentrations. The formulations of the vaccine or other pharmaceutical compositions described herein can contain a pharmaceutically acceptable preservative. In some embodiments, the preservative concentration ranges from 0.1 to 2.0%, typically v/v. Suitable preservatives include those known in the pharmaceutical arts. Benzyl alcohol, phenol, m-cresol, methylparaben, and propylparaben are examples of preservatives. The formulations of the vaccine or other pharmaceutical compositions described herein can include a pharmaceutically acceptable surfactant at a concentration of 0.005 to 0.02%.

(165) Therapeutic pharmaceutical compositions described herein can also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other.

(166) In some embodiments in which the engineered EVs are formulated for use in or with a vaccine, the vaccine composition can be formulated with the engineered EVs as an adjuvant. In other embodiments the vaccine composition can be formulated with the engineered EVs and an additional adjuvant, e.g., as known in the art.

(167) As used herein in the context of immunization, immune response and vaccination, the term “adjuvant” refers to any substance than when used in combination with a specific antigen produces a more robust immune response than the antigen alone. When incorporated into a vaccine formulation, an adjuvant acts generally to accelerate, prolong, or enhance the quality of specific immune responses to the vaccine antigen(s). Adjuvants typically promote the accumulation and/or activation of accessory cells or factors to enhance antigen-specific immune responses and thereby enhance the efficacy of vaccines, i.e., antigen-containing or encoding compositions used to induce protective immunity against the antigen.

(168) Adjuvants, in general, include adjuvants that create a depot effect, immune-stimulating adjuvants, and adjuvants that create a depot effect and stimulate the immune system. An adjuvant that creates a depot effect is an adjuvant that causes the antigen to be slowly released in the body, thus prolonging the exposure of immune cells to the antigen. This class of adjuvants includes but is not limited to alum (e.g., aluminum hydroxide, aluminum phosphate); emulsion-based formulations including mineral oil, non-mineral oil, water-in-oil or oil-in-water-in oil emulsion, oil-in-water emulsions such as Seppic ISA series of Montanide adjuvants (e.g., Montanide ISA 720; AirLiquide, Paris, France); MF-59 (a squalene-in-water emulsion stabilized with Span 85 and Tween 80; Chiron Corporation, Emeryville, Calif.); and PROVAX™ (an oil-in-water emulsion containing a stabilizing detergent and a micelle-forming agent; IDEC Pharmaceuticals Corporation, San Diego, Calif.).

(169) An immune-stimulating adjuvant is an adjuvant that causes activation of a cell of the immune system. It may, for instance, cause an immune cell to produce and secrete cytokines and interferons. This class of adjuvants includes but is not limited to saponins purified from the bark of

the *Q. saponaria* tree, such as QS21 (a glycolipid that elutes in the 21st peak with HPLC fractionation; Aquila Biopharmaceuticals, Inc., Worcester, Mass.); poly[di(carboxylatophenoxy)phosphazene (PCPP polymer; Virus Research Institute, USA); derivatives of lipopolysaccharides such as monophosphoryl lipid A (MPL; Ribí ImmunoChem Research, Inc., Hamilton, Mont.), muramyl dipeptide (MDP; Ribí) and threonyl-muramyl dipeptide (t-MDP; Ribí); OM-174 (a glucosamine disaccharide related to lipid A; OM Pharma SA, Meyrin, Switzerland); and *Leishmania* elongation factor (a purified *Leishmania* protein; Corixa Corporation, Seattle, Wash.). This class of adjuvants also includes CpG DNA.

(170) Adjuvants that create a depot effect and stimulate the immune system are those compounds which have both of the above-identified functions. This class of adjuvants includes but is not limited to ISCOMS (immunostimulating complexes which contain mixed saponins, lipids and form virus-sized particles with pores that can hold antigen; CSL, Melbourne, Australia); SB-AS2 (SmithKline Beecham adjuvant system #2 which is an oil-in-water emulsion containing MPL and QS21; SmithKline Beecham Biologicals [SBB], Rixensart, Belgium); SB-AS4 (SmithKline Beecham adjuvant system #4 which contains alum and MPL; SBB, Belgium); non-ionic block copolymers that form micelles such as CRL 1005 (these contain a linear chain of hydrophobic polyoxypropylene flanked by chains of polyoxyethylene; Vaxcel, Inc., Norcross, Ga.); and Syntex Adjuvant Formulation (SAF, an oil-in-water emulsion containing Tween 80 and a nonionic block copolymer; Syntex Chemicals, Inc., Boulder, Colo.).

(171) The active ingredients of the pharmaceutical compositions described herein can also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980).

(172) In some embodiments, sustained-release preparations can be used. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing a composition described herein in which the matrices are in the form of shaped articles, e.g., films, or microcapsule. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and  $\gamma$  ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated, the composition can remain in the body for a long time (e.g., up to about 1 hour, between 1-12 hours, 12-24 hours, 24 hours to 2 days, 2-3 days, 3-4 days, 4-5 days, 5-6 days, 6-7 days, 1-2 weeks, 3-4 weeks, 4 weeks to 2 months, 2-3 months, 3-4 months, 4-5 months, 5-6 months, or more than 6 months, or a variation thereof), denature, or aggregate as a result of exposure to moisture at 37° C., resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S—S— bond formation through thio-disulfide interchange, stabilization can be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

Administration, Dosing, Efficacy

(173) The engineered EV compositions, pharmaceutical compositions, or vaccine compositions described herein can be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being

treated, the particular subject being treated, the clinical condition of the individual subject, the cause of the disorder, the site of delivery of the vaccine composition, the method of administration, the scheduling of administration, and other factors known to medical practitioners.

(174) Generally, application of artificial synapses as therapy will take into account similar parameters as other therapeutic strategies, including concentration, timing of delivery, and sustained bioavailability at injury/disease site. Extracellular vesicle can be delivered via a number of routes: intravenous, intracoronary, and intramyocardial. Extracellular vesicles (e.g., exosomes), also allow for new delivery routes that were previously infeasible for cell therapy, such as inhalation or injection. These various approaches are described below, including injection, topical application, enteral administration, and pulmonary delivery.

(175) The engineered EV compositions provided herein can be administered to a subject in need thereof by any appropriate route which results in an effective treatment in the subject. As used herein, the terms “administering,” and “introducing” are used interchangeably and refer to the placement of a composition provided herein into a subject by a method or route which results in at least partial localization of such compositions at a desired site, such as a site of inflammation or a tumor, such that a desired effect(s) is produced. The compositions can be administered to a subject by any mode of administration that delivers the composition systemically or to a desired surface or target, and can include, but is not limited to, injection, infusion, instillation, and inhalation administration. To the extent that the composition can be protected from inactivation in the gut, oral administration forms are also contemplated. “Injection” includes, without limitation, intravenous, intramuscular, intra-arterial, intrathecal, intraventricular, intracapsular, intraorbital, retro-orbital, intravitreal, intraocular, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, sub capsular, subarachnoid, intraspinal, intracerebral, intratarsal, and intrasternal, intratumoral injection, and infusion or the like as known in the art.

(176) A therapeutic dose of the present invention may be delivered to a patient by means of controlled release, for example but not limited to, implantable pump and implantable cannulas to provide continuous access to the venous or arterial system.

(177) Topical application refers to applying or spreading a composition of the present invention onto surfaces on or in the body, both internally and/or externally, in a therapeutically effective amount for local and/or systemic treatment. Topical application may be epicutaneous wherein a composition of the present invention may be directly applied onto a localized surface of the skin or mucous membranes. Topical application may include transdermal application wherein a composition of the present invention may be absorbed into the body to obtain systemic delivery and systemic distribution. Topical application formulations may include, but are not limited to, creams, foams, gels, lotions, solutions, ointments, dermal patch, transdermal patches, powder, solid, sponge, tape, vapor, paste, film, liposomes, balm, shampoo, spray, or tincture or the like or a combination thereof. A therapeutic dose of a composition of the present invention may be delivered vaginally (for example a vaginal suppository, vaginal ring, douche, intrauterine device, intravesical infusion, and the like) or urethra or the like or a combination thereof.

(178) Enteral administration refers to a composition of the present invention administered via the gastrointestinal tract in a therapeutically effective amount for local or systemic treatment. Enteral administration may include, but is not limited to, delivery of a composition of the present invention via the mouth, sublingual, esophagus, gastric (for example the stomach), small intestines, large intestines or rectum. Oral delivery of the present invention may include, but is not limited to, the use of a capsule, pastille, pill, tablet, solution, gel, suspension, emulsion, syrup, elixir, tincture, mouthwash, lozenges, chewing gum, lollipop, cream, foam, solution, powder, solid, vapor, liposomes, spray, or tincture osmotic-controlled release oral delivery system, or the like. Gastric delivery may involve the use of a tube or nasal passage that leads directly to the stomach, for example, a percutaneous endoscopic gastrostomy tube. Gastric delivery may involve direct injection made through the abdominal wall. Rectal delivery may involve, but is not limited to, the

use of a suppository, ointment, enema, Murphy drip, or the like. A therapeutic dose of the present invention may be delivered to a patient by means of controlled release, for example but not limited to, controlled release drug delivery pellet or pill.

(179) Inhalation (i.e., pulmonary delivery, pulmonary administration refers to delivery to the respiratory system through the respiratory route, including but not limited to, intranasal administration, oral administration, and oral inhalative administration (e.g., intratracheal instillation and intratracheal inhalation) of a therapeutically effective amount for local or systemic treatment. Pulmonary delivery of a therapeutically effective amount of a composition of the present invention may be achieved by dispersion, for example by using a syringe. Pulmonary delivery of a composition of the present invention may be achieved by aerosol administration, wherein aerosol administration may deposit a therapeutically effective amount of the present invention by gravitational sedimentation, inertial impaction, or diffusion.

(180) Intravenous delivery technique can occur through a peripheral or central venous catheter. As the simplest delivery mode, this technique avoids the risk of an invasive procedure. However, intravenous may be regarded as a comparatively inefficient and less localized delivery method, as a high percentage of infused cell exosomes may become sequestered in organs such as the lung, liver, or spleen. Such sequestration may result in few or no cellular exosomes reaching broader circulation or have unintended systemic effects following their distribution.

(181) In certain embodiments, administration can include delivery to a tissue or organ site that is the same as the site of diseased and/or dysfunctional tissue. In certain embodiments, administration can include delivery to a tissue or organ site that is different from the site of diseased and/or dysfunctional tissue. In certain embodiments, the delivery is via inhalation or oral administration. In various embodiments, administration of artificial synapses can include combinations of multiple delivery techniques.

(182) In some embodiments, the compositions described herein are administered by aerosol administration, nebulizer administration, or tracheal lavage administration.

(183) The term “effective amount” as used herein refers to the amount of an engineered EV composition needed to alleviate or prevent at least one or more symptom of a disease or disorder (e.g., autoimmune disease or cancer), and relates to a sufficient amount of pharmacological composition to provide the desired effect, e.g., reduce the pathology, or any symptom associated with or caused by a disease. The term “therapeutically effective amount” therefore refers to an amount of an engineered EV composition or vaccine composition described herein using the methods as disclosed herein, that is sufficient to affect a particular disease state when administered to a typical subject. An effective amount as used herein would also include an amount sufficient to delay the development of a symptom of the disease, alter the course of a symptom disease (for example, but not limited to, slow the progression of a symptom of the disease), or reverse a symptom of the disease. Thus, it is not possible to specify the exact “effective amount.” However, for any given case, an appropriate “effective amount” can be determined by one of ordinary skill in the art using only routine experimentation.

(184) Effective amounts, toxicity, and therapeutic efficacy can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dosage can vary depending upon the dosage form employed and the route of administration utilized. The dose ratio between toxic and therapeutic effects is the therapeutic index and can be expressed as the ratio LD50/ED50. Compositions and methods that exhibit large therapeutic indices are preferred. A therapeutically effective dose can be estimated initially from cell culture assays. Also, a dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (i.e., the concentration of the engineered EVs or fusion polypeptides provided herein), which achieves a half-maximal inhibition of symptoms) as determined in cell culture, or in an appropriate animal model. Levels of therapeutic engineered EVs



in plasma can be measured, for example, by high performance liquid chromatography, enzyme linked immunosorbent assay (ELISA), flow cytometry, FACS sorting, western blot, mass spectroscopy, tunable resistive pulse sensing, EXOVIEW®, qRT-PCR, next generation sequencing (NGS), or by any analysis technique known by one of ordinary skill in the art. The effects of any particular dosage can be monitored by a suitable bioassay. The dosage can be determined by a physician and adjusted, as necessary, to suit observed effects of the treatment.

(185) The engineered EV compositions, pharmaceutical compositions, or vaccine compositions described herein can be formulated, in some embodiments, with one or more additional therapeutic agents currently used to prevent or treat the infection, for example. The effective amount of such other agents depends on the amount of an engineered EV in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as used herein before or about from 1 to 99% of the heretofore employed dosages.

(186) The dosage ranges for the pharmaceutical compositions described herein depend upon the potency and encompass amounts large enough to produce the desired effect. The dosage should not be so large as to cause unacceptable adverse side effects. Generally, the dosage will vary with the age, condition, health, and sex of the patient and can be determined by one of skill in the art. The dosage can also be adjusted by the individual physician in the event of any complication. In some embodiments, the dosage ranges from 0.001 mg/kg body weight to 100 mg/kg body weight. In some embodiments, the dose range is from 5 µg/kg body weight to 100 µg/kg body weight. Alternatively, the dose range can be titrated to maintain serum levels between 0.1 µg/mL and 1000 µg/mL. For systemic administration, subjects can be administered a therapeutic amount, such as, e.g., 0.1 mg/kg, 0.5 mg/kg, 1.0 mg/kg, 2.0 mg/kg, 2.5 mg/kg, 5 mg/kg, 7.5 mg/kg, 10 mg/kg, 15 mg/kg, 20 mg/kg, 25 mg/kg, 30 mg/kg, 40 mg/kg, 50 mg/kg, or more. These doses can be administered by one or more separate administrations, or by continuous infusion. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until, for example, the infection is treated, as measured by the methods described above or known in the art. However, other dosage regimens can be useful.

(187) In various embodiments, the quantities of artificial synapses that are administered to achieve these effects range from  $1 \times 10^6$  to  $1 \times 10^7$ ,  $1 \times 10^7$  to  $1 \times 10^8$ ,  $1 \times 10^8$  to  $1 \times 10^9$ ,  $1 \times 10^9$  to  $1 \times 10^{10}$ ,  $1 \times 10^{10}$  to  $1 \times 10^{11}$ ,  $1 \times 10^{11}$  to  $1 \times 10^{12}$ ,  $1 \times 10^{12}$  to  $1 \times 10^{13}$ ,  $1 \times 10^{13}$  to  $1 \times 10^{14}$ ,  $1 \times 10^{14}$  to  $1 \times 10^{15}$ ,  $1 \times 10^{15}$  or more Evs/artificial synapses. In other embodiments, the numbers of artificial synapses are relative to the number of cells used in a clinically relevant dose for a cell-therapy method. For example, defining an effective dose range, dosing regimen and route of administration, may be guided by studies using fluorescently labeled artificial synapses, and measuring target tissue retention, which can be  $>10\times$ ,  $>50\times$ , or  $>100\times$  background, as measured 5, 10, 15, 30, or 30 or more min as a screening criterion. In certain embodiments,  $>100\times$  background measured at 30 mins is a baseline measurement for a low and high dose that is then assess for safety and bioactivity (e.g., using MRI endpoints: scar size, global and regional function of the target organ being treated). In various embodiments, single doses are compared to two, three, four, four or more sequentially-applied doses. In various embodiments, the repeated or sequentially-applied doses are provided for treatment of an acute disease and/or condition. In various embodiments, the repeated or sequentially-applied doses are provided for treatment of a chronic disease and/or condition. In other embodiments, administration of the plurality of artificial synapses is adjunctive to standard therapy.

(188) In other embodiments, administering a composition includes  $1 \times 10^{10}$  or more artificial synapses in a single dose. In various embodiments, exosome quantity may be defined by protein quantity, such as dosages including 1-10, 10-25, 25-50, 50-75, 75-100, or 100 or more mg exosome protein. In other embodiments, a single dose is administered multiple times to the subject. In other

embodiments, administering a composition consists of one or more of: injection, topical administration, enteral, intravenous, intra-arterial, or inhalation.

(189) In various embodiments, exosome quantity may be defined by protein quantity, such as dosages including 1-10, 10-25, 25-50, 50-75, 75-100, or 100 or more mg exosome protein. In various embodiments, administering a composition includes multiple dosages of the artificial synapses. In various embodiments, the repeated or sequentially-applied doses are provided for treatment of an acute disease and/or condition. In various embodiments, the repeated or sequentially-applied doses are provided for treatment of a chronic disease and/or condition.

(190) In other embodiments, administering a composition including a plurality of artificial synapses to the subject is adjunctive to standard therapy.

(191) The duration of a therapy using the methods described herein will continue for as long as medically indicated or until a desired therapeutic effect (e.g., those described herein) is achieved. In certain embodiments, the administration of the vaccine composition described herein is continued for 1 month, 2 months, 4 months, 6 months, 8 months, 10 months, 1 year, 2 years, 3 years, 4 years, 5 years, 10 years, 20 years, or for a period of years up to the lifetime of the subject.

(192) As will be appreciated by one of skill in the art, appropriate dosing regimens for a given composition can comprise a single administration/immunization or multiple ones. Subsequent doses may be given repeatedly at time periods, for example, about two weeks or greater up through the entirety of a subject's life, e.g., to provide a sustained preventative effect. Subsequent doses can be spaced, for example, about two weeks, about three weeks, about four weeks, about one month, about two months, about three months, about four months, about five months, about six months, about seven months, about eight months, about nine months, about ten months, about eleven months, or about one year after a primary immunization.

(193) The precise dose to be employed in the formulation will also depend on the route of administration and should be decided according to the judgment of the practitioner and each patient's circumstances. Ultimately, the practitioner or physician will decide the amount of the engineered EV or composition thereof to administer to particular subjects.

#### Methods of Modulating Inflammation and Treating Autoimmune Diseases

(194) The artificial synapses/engineered Evs and compositions thereof provided herein can be deployed in a therapeutic strategy against virtually any injury/disease, as providing a platform for altering biological signaling. This includes, for example, inflammation and immune signaling, which plays a role in virtually all injuries and diseases in living organisms.

(195) Thus, described herein is a method of modulating inflammation, including selecting a subject afflicted with an inflammatory related disease and/or condition; and administering to the subject a composition including a plurality of artificial synapses (engineered Evs) to the subject, wherein administration of the composition modulates inflammation.

(196) As used herein, the term “inflammation” or “inflamed” refers to activation or recruitment of the immune system or immune cells (e.g., T cells, B cells, macrophages). A tissue that has inflammation can become reddened, white, swollen, hot, painful, sensitivity, exhibit a loss of function, or have a film or mucus. Methods of identifying inflammation are well known in the art. Inflammation typically occurs following injury, infection by a microorganism, exposure to a substance (e.g., a toxin, chemical, or dust) or autoimmune dysfunction. Onset of inflammation may be rapid (e.g., immediately following injury) or slow (e.g., repeated exposure to an irritant such as a chemical over time) with a duration of minutes, hours, days, months, years, or an individual's life.

(197) Inflammation plays a vital role in alerting the immune system of potential danger and damage within a body. Inflammation is necessary to control and repair injury. For example, acute inflammation is a response to physical trauma, infection, and stress. Acute inflammation helps prevent further injury and triggers healing and recovery. Unfortunately, inflammation can become excessive and inappropriately active, lasting beyond the typical recovery time from an injury or infection. Wherein healthy inflammation helps a body respond to injury, chronic inflammation

perpetuates injury and may lead to negative consequences to one's health. In particular, autoimmune diseases are chronic diseases from a host's immune system attacking itself, often due to aberrant biological signaling in the host. Restoring normal homeostatic signaling via application of artificial synapses, particularly targeting immune checkpoints, represents a highly promising avenue. For example, surface bound immune-checkpoint proteins or fragments thereof may modulate immune cell stimulation and affect suppression of immune cell function when delivered via artificial synapses. Injection, inhalation, ingestion or topical application of artificial synapses with surface bound immune-checkpoint proteins or fragments thereof may be used to treat immune, auto-immune, inflammatory, and auto-inflammatory conditions. Examples include chronic obstructive pulmonary disease (COPD) which is an inflammatory, progressive, life-threatening lung disease, psoriasis, a common chronic noncommunicable inflammatory skin disease, arthritis, a debilitating and painful degeneration of joints, among others well-understood to one of skill in the art.

(198) In other embodiments, the inflammatory related disease and/or condition is acute, for example septicemia. In other embodiments, the inflammatory related disease and/or condition is chronic, for example chronic obstructive pulmonary disease. In other embodiments, the inflammatory condition is an autoimmune disease wherein the autoimmune disease and/or condition is one or more of: polymyositis, dermatomyositis, Graves' disease, Hashimoto's thyroiditis, myasthenia gravis, vasculitis, multiple sclerosis, psoriasis, rheumatoid arthritis, psoriatic arthritis, scleroderma, systemic lupus erythematosus, inflammatory bowel disease, Crohn's disease, hyperthyroidism, autoimmune adrenal insufficiency, Sjogren syndrome, type I diabetes mellitus, autoimmune hemolytic anemia, idiopathic thrombocytopenic purpura, myasthenia gravis, ulcerative colitis, uveitis, polyarteritis nodosa, relapsing polychondritis, Behcet's disease, reactive arthritis, ankylosing spondylitis, Guillain-Barre syndrome, or optic neuropathy. In other embodiments, the disease and/or condition is chronic obstructive pulmonary disease, rheumatoid arthritis, uveoretinitis, psoriasis, and eczema. In other embodiments, the disease and/or condition is irritable bowel disease, multiple sclerosis or lupus.

(199) In other embodiments, the inflammatory related disease and/or condition is an ocular disease. As used herein, the terms "ocular disease", "eye disorder" and "eye disease" are used interchangeably and refer to a disease or disorder that affects the health and/or vision of either one or both eyes or the general area of the eye(s), eye lid(s), or area surrounding or in near proximity to the eye(s). Eye disease may include, but are not limited to, macular degeneration (e.g., age-related macular degeneration), cataracts, diabetic retinopathy, diabetic macular edema, eye floaters, eye flashes, glaucoma, amblyopia, strabismus, retinitis (e.g., CMV retinitis), color blindness, keratoconus, retinal detachment, eyelid twitching, ocular hypertension, blepharitis, uveitis, Bietti's crystalline dystrophy, blepharospasm, cornea and corneal diseases, dry eye, histoplasmosis, macular hole, macular pucker, conjunctivitis, presbyopia, retinoblastoma, retinitis pigmentosa, retinopathy, Stargardt disease, Usher syndrome, uveal Coloma, and vitreous detachment, or the like.

(200) Described herein is a method for treatment including, selecting a subject in need of treatment, administering a composition including a plurality of artificial synapses to the individual, wherein administration of the composition treats the subject. In certain embodiments, the subject is in need to treatment for a disease and/or condition involving tissue damage or dysfunction.

(201) Described herein is a method of treating an autoimmune disease, inflammation, inflammatory disease or condition, or cancer in a subject, the method comprising: administering to a subject an engineered EV or composition thereof as provided herein to the subject.

(202) Measured or measurable parameters include clinically detectable markers of disease, for example, elevated or depressed levels of a clinical or biological marker, as well as parameters related to a clinically accepted scale of symptoms or markers for a disease or disorder. It will be understood, however, that the total usage of the compositions and formulations as disclosed herein

will be decided by the attending physician within the scope of sound medical judgment. The exact amount required will vary depending on factors such as the type of disease being treated.

(203) Non-limiting examples of clinical tests that can be used to assess autoimmune diseases, inflammatory conditions, or inflammation parameters include blood tests, skin biopsy, MRI, eye examination, ocular pressure tests, etc. Where necessary or desired, animal models of injury or disease can be used to gauge the effectiveness of a particular composition as described herein. For example, an EAU animal model, as demonstrated in the working examples can be used.

(204) In various embodiments, administration of the plurality of artificial synapses alters gene expression in the damaged or dysfunctional tissue, improves viability of the damaged tissue, and/or enhances regeneration or production of new tissue in the individual. In various embodiments, administration of the plurality of artificial synapses alters gene expression in the damaged or dysfunctional tissue, improves viability of the damaged tissue, and/or enhances regeneration or production of new tissue in the individual.

(205) In various embodiments, the damaged or dysfunctional tissue is in need of repair, regeneration, or improved function due to an acute event. Acute events include, but are not limited to, trauma such as laceration, crush or impact injury, shock, loss of blood or oxygen flow, infection, chemical or heat exposure, poison or venom exposure, drug overuse or overexposure, and the like. Other sources of damage also include, but are not limited to, injury, age-related degeneration, cancer, and infection. In several embodiments, the regenerative cells used to prepare the engineered EVs provided herein are from the same tissue type as is in need of repair or regeneration. In several other embodiments, the regenerative cells are from a tissue type other than the tissue in need of repair or regeneration. In some embodiments, the engineered EVs provided herein are derived from the subject being treated. In some embodiments, the engineered EVs are derived from a donor subject.

(206) In other embodiments, the damaged or dysfunctional tissue is in need of repair, regeneration, or improved function due to damage from chronic disease.

#### Some Selected Definitions

(207) All references cited herein are incorporated by reference in their entirety as though fully set forth. Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Allen et al., *Remington: The Science and Practice of Pharmacy* 22<sup>nd</sup> ed., Pharmaceutical Press (Sep. 15, 2012); Hornyak et al., *Introduction to Nanoscience and Nanotechnology*, CRC Press (2008); Singleton and Sainsbury, *Dictionary of Microbiology and Molecular Biology* 3<sup>rd</sup> ed., revised ed., J. Wiley & Sons (New York, NY 2006); Smith, *March's Advanced Organic Chemistry Reactions, Mechanisms and Structure* 7<sup>th</sup> ed., J. Wiley & Sons (New York, NY 2013); Singleton, *Dictionary of DNA and Genome Technology* 3<sup>rd</sup> ed., Wiley-Blackwell (Nov. 28, 2012); and Green and Sambrook, *Molecular Cloning: A Laboratory Manual* 4<sup>th</sup> ed., Cold Spring Harbor Laboratory Press (Cold Spring Harbor, NY 2012), provide one skilled in the art with a general guide to many of the terms used in the present application. For references on the preparation and structure of antibodies and fusion polypeptides, see, e.g., Greenfield, *Antibodies A Laboratory Manual* 2<sup>nd</sup> ed., Cold Spring Harbor Press (Cold Spring Harbor NY, 2013); Köhler and Milstein, *Derivation of specific antibody-producing tissue culture and tumor lines by cell fusion*, Eur. J. Immunol. 1976 Jul. 6 (7): 511-9; Queen and Selick, Humanized immunoglobulins, U.S. Pat. No. 5,585,089 (1996 December); and Riechmann et al., *Reshaping human antibodies for therapy*, Nature 1988 Mar. 24, 332 (6162): 323-7. See also, Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991), Lanzavecchia et al., Eur. J. Immunol. 17, 105 (1987), Huston et al., Proc. Natl. Acad. Sci. U.S.A., 85, 5879-5883 (1988), Bird et al., Science 242, 423-426 (1988), Brinkman et al. mAbs Vol 9, No. 2, 182-212 (2017), Chothia & Lesk, J. Mol. Biol, 196:901-917 (1987), Chothia et al., Nature 342:877-883 (1989), Holliger et al. (1993) Proc. Natl. Acad. Sci.

USA 90:6444-6448; Poljak (1994) Structure 2:1121-1123); Kontermann and Dubel eds., Antibody Engineering, Springer-Verlag, N.Y. (2001), p. 790 (ISBN 3-540-41354-5, Zapata et al. (1995) Protein Eng. 8 (10): 1057-1062; Morrison, et al., Proc. Natl. Acad. Sci. USA, 81:6851 (1984), U.S. Pat. Nos. 4,816,567, 5,693,780, which are incorporated herein by reference in their entireties.

(208) One skilled in the art will recognize many methods and materials similar or equivalent to those described herein, which could be used in the practice of the present invention. Indeed, the present invention is in no way limited to the methods and materials described. For purposes of the present invention, the following terms are defined below.

(209) As used herein, the term “extracellular vesicle” and “vesicle” are used interchangeably and refer to a particle, wherein the particle comprises a phospholipid bilayer that encloses an internal space and an exterior surface and may or may not be derived from a cell. The size of extracellular vesicles can range between 20 nm to 3  $\mu$ m in diameter but may be smaller than 20 nm or larger than 3  $\mu$ m. Examples of extracellular vesicles include, but is not limited to, exosomes (for example small exosomes and large exosomes), ectosomes, macrovesicles, microparticles, apoptotic bodies, vesicular organelles, oncosomes (for examples large oncosomes), exospheres, exomeres, cell derived nanovesicles (CDN) (e.g., by genesis via grating or shearing cells), liposomes or the like known by one of ordinary skill in the art. Extracellular vesicles may originate naturally via known or unknown biosynthetic pathways. Extracellular vesicles may be promoted to originate by using mechanical methods such as cell grating or cell shearing wherein a cell is grated or sheared causing portions or parts of the cell membrane to form vesicles. For example, CDNs may be formed by using mechanical methods such as cell grating or cell shearing wherein a cell is grated or sheared causing portions or parts of the cell membrane to form vesicles. Additional non-limiting examples of mechanical methods that can be used to form cell derived nanovesicles are further described in detail, e.g., Goh, W. J., Zou, S., Ong, W. Y. et al. Bioinspired Cell-Derived Nanovesicles versus Exosomes as Drug Delivery Systems: a Cost-Effective Alternative. *Sci Rep* 7, 14322 (2017). doi: 10.1038/s41598-017-14725-x, the contents of which are incorporated herein by reference in their entireties.

(210) Extracellular vesicles comprise cargo, wherein the term “cargo” refers to peptides, proteins, nucleic acids, lipids, metabolites, carbohydrates, biomolecules, small molecules, large molecules, vesicles, organelles, or fragments thereof. In some embodiments, cargo may refer to existing drugs or therapeutics known in the art. Extracellular vesicle cargo may be located within the internal space of the extracellular vesicle. Extracellular vesicle cargo may be membrane bound and span one or both layers of the extracellular vesicle phospholipid bilayer (for example a transmembrane protein). Extracellular vesicle cargo may be in contact with the external or internal surface of the extracellular vesicle, for example through a covalent bond or a non-covalent bond. The phospholipid bilayer of the extracellular vesicle may comprise one or more transmembrane proteins, wherein a portion of the one or more transmembrane membrane proteins is located within the internal space of the extracellular vesicle. The phospholipid bilayer of the extracellular vesicle may comprise one or more transmembrane proteins, wherein the one or more transmembrane membrane proteins comprises a domain on the exterior of the extracellular vesicle. The phospholipid bilayer of the extracellular vesicle may comprise one or more transmembrane proteins, wherein the one or more transmembrane membrane proteins comprises a domain on the interior of the extracellular vesicle. Cargo may refer to a protein on the luminal side (e.g., in the internal space) of the extracellular vesicle wherein said protein encodes a vesicle targeting domain that may be in contact with the interior phospholipid layer of the extracellular vesicle. Cargo may refer to a protein on the luminal side (e.g., in the internal space) of the extracellular vesicle wherein said protein encodes a vesicle targeting domain that may be in contact with the interior phospholipid layer of the extracellular vesicle and wherein said protein may be presented into the internal space of the extracellular vesicle.

(211) As used herein, the terms “sticky binder” and “vesicle targeting domain” and “anchor

protein” are used interchangeably and refer to a protein that is covalently or non-covalently attached to at least one lipid wherein the one or more lipid is embedded within a membrane (e.g., a cell membrane), and the lipid serves to anchor the protein to the membrane. The terms “sticky binder” and “vesicle targeting domain” and “anchor protein” can also mean a protein sequence that encodes for one or more transmembrane domains wherein the one or more transmembrane domains spans at least partly through a phospholipid bilayer, for example the phospholipid bilayer of an extracellular vesicle. The transmembrane domain can be of a Type I or Type II membrane protein. Transmembrane domains can be structurally identified using methods known to those of skill in the art, such as sequence analysis programs that identify hydrophobic and hydrophilic domains (for example TMHMM Server, v. 2.0-DTU, Erik L. L. Sonnhammer, Gunnar von Heijne, and Anders Krogh: *A hidden Markov model for predicting transmembrane helices in protein sequences. In Proc. of Sixth Int. Conf. on Intelligent Systems for Molecular Biology*, p 175-182 Ed J. Glasgow, T. Littlejohn, F. Major, R. Lathrop, D. Sankoff, and C. Sensen Menlo Park, CA: AAAI Press, 1998, which is incorporated herein by reference in its entirety.)

(212) A vesicle targeting domain may include, but is not limited to, one or more prenylation site, fatty acylation site, and/or glycosylphosphatidylinositol (GPI) linked protein. One preferred embodiment of a vesicle targeting domain is the GPI sequence from CD55. Another preferred embodiment of a vesicle targeting domain is the GPI sequence from CD59. Another embodiment of a vesicle targeting domain is the C1C2 domain from MFGE8. Other embodiments of sequences for vesicle targeting domains include transmembrane regions of CD9 (for example transmembrane 2 or 3 of CD9, CD9tm2 or CD9tm3, respectively), K-Ras (for example K-Ras4A and K-Ras4B), transmembrane domain from A Disintegrin and Metalloproteinase Domain-containing protein 10 (ADAM10, also known as CDw156 or CD156c) or other ADAM proteins. Vesicle targeting domains may include one or more sequences from 4F2 (for example 4F2 encoded by the solute carrier family 3 member 2 (SLC3A2) gene which makes up the heavy subunit of CD98). Vesicle targeting domains can include a sequence for one or more myristoylation sites. For example, the protein sequence for a myristoylation site from myristoylated alanine-rich C-kinase substrate (MARCKS) protein. Vesicle targeting domains can include a sequence for one or more palmitoylation sites. For example, the myristoylation sequence from the MARCKS protein may be modified to encode for a palmitoylation site. All variants, isoforms, or fragments or the like known by one of ordinary skill in the art are encompassed by the present invention.

(213) Vesicle targeting domains may include transmembrane sequences from *Homo sapiens* transferrin receptor 2 (TFR2), transcript variant 1 (transferrin receptor protein 2 isoform 1) or versions therefore. In a preferred embodiment, the vesicle targeting domain may be a transmembrane domain from CD298.

(214) As used herein, the terms “proteins” and “peptides” and “polypeptides” are used interchangeably herein to designate a series of amino acid residues connected to the other by peptide bonds between the alpha-amino and carboxy groups of adjacent residues. Although “protein” is often used in reference to relatively large polypeptides, and “peptide” is often used in reference to small polypeptides, usage of these terms in the art overlaps and varies. The term “peptide” as used herein refers to peptides, polypeptides, proteins and fragments of proteins, unless otherwise noted. The terms “protein” and “peptide” are used interchangeably herein when referring to a gene product and fragments thereof. Thus, exemplary peptides or proteins include gene products, naturally occurring proteins, homologs, orthologs, paralogs, fragments and other equivalents, variants, fragments, and analogs of the foregoing.

(215) As used herein, the term “linker” refers to a synthetic protein sequence of amino acids that is used to connect two polypeptide domains via peptide bonds.

(216) As used herein, the term “fusion protein” refers to a single chimeric protein comprising a protein of interest (e.g., checkpoint protein) joined to an exogenous protein or protein fragment (e.g., an anchor protein), wherein the components of the fusion protein are linked to each other by

peptide-bonds, either directly or through a peptide linker. The anchor protein of the fusion protein may enhance incorporation of the fusion protein onto and/or into the membrane of a vesicle, for example the internal and/or external leaflet of the phospholipid bilayer of an exosome membrane. The fusion protein may have at least a part of an amino acid sequence of an immune checkpoint protein or proteins involved in immune synapses. The fusion protein may have at least a part of an amino acid sequence of A2AR, VTCN1, Galectin 9, FGL-1, PECAM-1, TSG-6, STAB-1, NRP1, NRP2, SEMA3A, SEMA3F, RGMB, TIM-3, TIGIT, HLA class I, HLA class II, VISTA, HMGB1, phosphatidylserine, T-cell receptor (TCR), SHP-1, SHP-2, FBXO38, SH2D1A, B7RP1, IDO, NOX2, TNFRSF18, B7-H4, B7-H5, SISP1, B7-H6, B7-H7, APLNR, IFN  $\gamma$ , PD-1, WNT5A, IL-6, IL-10, NKG2 family of C-type lectin receptors, ligands of NKG2 family, killer cell immunoglobulin-like receptors, CD2, CD4, CD8, CD27, CD27 ligand (CD70), CD28, CD28H, CD39, CD40, CD44, CD47, CD63, CD66a, CD80, B7-2, CD86, CD73, CD94, CD96, CD101, CD112, CD112R, CD122, CD134, CD137 (4-1BB), CD137 ligand (4-1BBL), CD152, CD154, CD155, CD158, CD158a, CD158g, CD158h, KIR2DL1, KIR2DS1, KIRDS3, KIR2DS5, CD160, CD172a, CD200, CD200R, CD223, CD226, CD252, CD270, CD272, CD273, CD274, CD275, CD276, CD278, CD279 (PD-1), CD279 ligand (PD-L1/PDL-2), CD328, CD329, and/or CD337. The fusion protein may have a polypeptide linker sequence (e.g., an Fc domain and/or a GSSG linker (SEQ ID NO: 319)), followed by an amino acid sequence coding for an anchor protein sequence (e.g., a prenylation site, fatty acylation site, or a GPI sequence) or any isoform, fragment, variation thereof, or a ligand to the aforementioned proteins thereof, or the like known by one of ordinary skill in the art. All variants are encompassed by the present invention.

(217) As used herein, the term “immune synapse” and “cell synapse” are used interchangeably and refer to cell-to-cell interaction wherein said interaction results in activation, suppression, and/or adhesion of either one or more cells. Immune synapse or cell synapse are mediated by proteins that may be cytoplasmic, membrane bound, membrane associated, and/or secreted. Immune or cell synapses may be mediated by one or more “immune checkpoint proteins” which herein refers to any protein that is involved in maintaining immune homeostasis or plays a role in regulating immune activation or suppression. Immune checkpoint proteins may be cytoplasmic, membrane bound, membrane associated, and/or secreted.

(218) As used herein, the term “fragment” or “active fragment” refers to a portion of a nucleic acid or polypeptide provided herein that retains the ability to be expressed by the engineered EVs provided herein. In some embodiments, the active fragment retains the ability to activate a target polypeptide, thereby increasing the activity of said target polypeptide (e.g., suppressing an immune response).

(219) As used herein, the terms “specifically bind” and/or “specifically recognize” or “substantially binds” refers to the affinity of a binding molecule for a target molecule compared to the binding molecule's affinity for non-target molecules. A binding molecule (e.g., a POI domain) that specifically binds a target molecule (e.g., a target polypeptide provided herein) does not substantially recognize or bind non-target molecules. e.g., an antibody “specifically binds” and/or “specifically recognize” another molecule, meaning that this interaction is dependent on the presence of the binding specificity of the molecule structure, e.g., an antigenic epitope. As used herein, “non-specific binding” and “background binding” refers to the interaction that does not depend on the presence of specific structure (e.g., a specific antigenic epitopes). Methods of measuring binding of a polypeptide to a target are known in the art (e.g., differential scanning calorimetry, isothermal titration calorimetry, spectroscopy, crystallography, surface plasmon resonance, co-immunoprecipitation, pulldown assays, crosslinking, yeast two-hybrid system, tandem affinity purification-mass spectroscopy, protein microarrays, bio-layer interferometry, far-Western blots, computational prediction, analytical ultracentrifugation, light scattering, fluorescence spectroscopy, resonance energy transfer, ELISA or ELISPOT assays, or any other assays known in the art).

(220) As used herein, the terms “treat,” “treatment,” “treating,” or “amelioration” refer to therapeutic treatments, wherein the object is to reverse, alleviate, ameliorate, inhibit, slow down or stop the progression or severity of a condition associated with, a disease or disorder. The term “treating” includes reducing or alleviating at least one adverse effect or symptom of a condition, disease or disorder associated with an infection or a cancer. Treatment is generally “effective” if one or more symptoms or clinical markers are reduced. Alternatively, treatment is “effective” if the progression of a disease is reduced or halted. That is, “treatment” includes not just the improvement of symptoms or markers, but also a cessation or at least slowing of progress or worsening of symptoms that would be expected in absence of treatment. Beneficial or desired clinical results include, but are not limited to, alleviation of one or more symptom(s), diminishment of extent of disease, stabilized (i.e., not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and remission (whether partial or total), whether detectable or undetectable. The term “treatment” of a disease also includes providing relief from the symptoms or side-effects of the disease (including palliative treatment).

(221) As used herein “preventing” or “prevention” refers to any methodology where the disease state does not occur due to the actions of the methodology (such as, for example, administration of a composition or construct as described herein). In one aspect, it is understood that prevention can also mean that the disease is not established to the extent that occurs in untreated controls. Accordingly, prevention of a disease encompasses a reduction in the likelihood that a subject can develop the disease, relative to an untreated subject (e.g., a subject who is not treated with the methods or compositions described herein).

(222) As used herein, the terms “autoimmune condition” and “autoimmune disease” are used interchangeably and refer to any disease characterized by abnormal functioning of the immune system and may include, but is not limited to, achalasia, Addison's disease, adult Still's disease, agammaglobulinemia, alopecia areata, amyloidosis, ankylosing spondylitis, anti-GBM/Anti-TBM nephritis, antiphospholipid syndrome, autoimmune angioedema, autoimmune dysautonomia, autoimmune encephalomyelitis, autoimmune hepatitis, autoimmune inner ear disease (AIED), autoimmune myocarditis, autoimmune oophoritis, autoimmune orchitis, autoimmune pancreatitis, autoimmune retinopathy, autoimmune urticaria, axonal & neuronal neuropathy (AMAN), Baló disease, Behcet's disease, benign mucosal pemphigoid, bullous pemphigoid, Castleman disease, celiac disease, Chagas disease, chronic inflammatory demyelinating polyneuropathy (CIDP), chronic recurrent multifocal osteomyelitis (CRMO), Churg-Strauss syndrome (CSS), eosinophilic granulomatosis (EGPA), cicatricial pemphigoid, Cogan's syndrome, cold agglutinin disease, congenital heart block, coxsackie myocarditis, CREST syndrome, Crohn's disease, dermatitis herpetiformis, dermatomyositis, Devic's disease (neuromyelitis optica), discoid lupus, Dressler's syndrome, endometriosis, eosinophilic esophagitis (EoE), eosinophilic fasciitis, erythema nodosum, essential mixed cryoglobulinemia, Evans syndrome, fibromyalgia, fibrosing alveolitis, giant cell arteritis (temporal arteritis), giant cell myocarditis, glomerulonephritis, Goodpasture's syndrome, granulomatosis with polyangiitis, Graves' disease, Guillain-Barre syndrome, Hashimoto's thyroiditis, hemolytic anemia, Henoch-Schonlein purpura (HSP), Herpes gestationis or pemphigoid gestationis (PG), hidradenitis suppurativa (HS) (acne inversa), hypogammaglobulinemia, IgA nephropathy, IgG4-related sclerosing disease, immune thrombocytopenia purpura (ITP), inclusion body myositis (IBM), interstitial cystitis (IC), juvenile arthritis, type 1 diabetes, juvenile myositis (JM), Kawasaki disease, Lambert-Eaton syndrome, leukocytoclastic vasculitis, lichen planus, lichen sclerosus, ligneous conjunctivitis, linear IgA disease (LAD), lupus, Lyme disease chronic, Meniere's disease, microscopic polyangiitis (MPA), mixed connective tissue disease (MCTD), Mooren's ulcer, Mucha-Habermann disease, multifocal motor neuropathy (MMN) or MMNCB, multiple sclerosis, myasthenia gravis, myositis, narcolepsy, neonatal Lupus, neuromyelitis optica, neutropenia, ocular cicatricial pemphigoid, optic neuritis, palindromic rheumatism (PR), PANDA, paraneoplastic cerebellar degeneration (PCD), paroxysmal



nocturnal hemoglobinuria (PNH), Parry Romberg syndrome, pars planitis (peripheral uveitis), Parsonage-Turner syndrome, pemphigus, peripheral neuropathy, perivenous encephalomyelitis, pernicious anemia (PA), POEMS syndrome, polyarteritis nodosa, polyglandular syndromes type I, II, III, polymyalgia rheumatica, polymyositis, postmyocardial infarction syndrome, postpericardiotomy syndrome, primary biliary cirrhosis, primary sclerosing cholangitis, progesterone dermatitis, psoriasis, psoriatic arthritis, pure red cell aplasia (PRCA), pyoderma gangrenosum, Raynaud's phenomenon, reactive arthritis, reflex sympathetic dystrophy, relapsing polychondritis, restless legs syndrome (RLS), retroperitoneal fibrosis, rheumatic fever, rheumatoid arthritis, sarcoidosis, Schmidt syndrome, scleritis, scleroderma, Sjögren's syndrome, sperm & testicular autoimmunity, stiff person syndrome (SPS), subacute bacterial endocarditis (SBE), Susac's syndrome, sympathetic ophthalmia (SO), takayasu's arteritis, temporal arteritis/Giant cell arteritis, thrombocytopenia purpura (TTP), Tolosa-Hunt syndrome (THS), transverse myelitis, type 1 diabetes, ulcerative colitis (UC), undifferentiated connective tissue disease (UCTD), uveitis, vasculitis, vitiligo, Vogt-Koyanagi-Harada disease. An autoimmune condition or autoimmune diseases may be caused by, but not limited to, a natural predisposition, an infection (e.g., bacteria or virus), drugs, vaccination, environmental triggers (e.g., toxins or chemicals such as dust, silica, oil, benzene, tri- or per-chloroethylene etc.), stress, cancer, blood or tissue or organ transplantation, or unknown etiology. Autoimmune disorders may result in but not limited to the destruction of body tissue, abnormal growth of an organ or tissue, changes in organ or tissue function (e.g., changes in blood vessels, connective tissue, function of endocrine glands, joints, muscles, blood cells, skin, etc.).

(223) As used herein, the term “cancer” refers to a hyperproliferation of cells that exhibit a loss of normal cellular control that results in unregulated growth, lack of differentiation, local tissue invasion, and metastasis. The methods and compositions described herein can be used for the treatment of solid tumors (e.g., cancer) or non-solid tumors, such as leukemia, blood cell cancers, and the like. Solid tumors can be found in bones, muscles, the brain, or organs, and can be sarcomas or carcinomas. Where the methods and compositions described herein can overcome barriers of tumor treatment, including, but not limited to barriers to treatment or inhibition of metastases, it is contemplated that aspects of the technology described herein can be used to treat all types of solid and non-solid tumor cancers, including cancers not listed in the instant specification. The compositions and methods described herein, without limitation, include methods of treating cancer, methods of inhibiting metastases, and methods of inducing an anti-tumor immune response.

(224) As used herein, the terms “subject”, “individual”, “host”, and “patient” are used interchangeably and may refer to any animal, mammal, bird, fish, reptile, and amphibian, for example, human, monkey, dog, cat, horse, pig, cattle, ox, donkey, rabbit, sheep, goat, mouse, rat, guinea pig, llama, chicken, goose, duck, turkey, or the like receiving or registered to receive a therapeutic amount of a composition of the present invention for medical care or treatment.

(225) As used herein, the term “injection” refers to any process or method which allows the person skilled in the art to administer any therapeutic to a target site by penetration. Examples of injection are, but not limited to, subcutaneous, subcuticular, subcapsular, subarachnoid, intradermal, intramuscular, intravenous, intra-arterial, intraventricular, intracapsular, intraorbital, intraocular, intrathoracic, intraperitoneal, intravitreal, retro-orbital, intranasal, intracerebral, intrathymic, intraspinal, intrasternal, intra-articular, intracavernous, intracardiac, intraosseous, intrathecal, transtracheal, epidural, or the like as known in the art. A therapeutic of the present invention may be delivered to a patient by means of controlled release, for example but not limited to, implantable pump and implantable cannulas to provide continuous access to the venous or arterial system.

(226) As used herein, the term “topical application” refers to applying or spreading a composition of the present invention onto surfaces on or in the body, both internally and/or externally, in a

therapeutically effective amount for local and/or systemic treatment. Topical application may be epicutaneous wherein a composition of the present invention may be directly applied onto a localized surface of the skin or mucous membranes. Topical application may include transdermal application wherein a composition of the present invention may be absorbed into the body to obtain systemic delivery and systemic distribution. For example, a transdermal patch may be applied onto the body to deliver a therapeutic dose of a composition of the invention presented herein. Topical application formulations may include, but are not limited to, creams, foams, gels, lotions, solutions, ointments, dermal patch, transdermal patches, powder, solid, sponge, tape, vapor, paste, film, liposomes, balm, shampoo, spray, or tincture. A therapeutic dose of a composition of the present invention may be delivered vaginally (for example a vaginal suppository, vaginal ring, douche, intrauterine device, intravesical infusion, and the like) or urethra.

(227) As used herein, the term “enteral administration” refers to a composition of the present invention administered via the gastrointestinal tract in a therapeutically effective amount for local or systemic treatment. Enteral administration may include, but is not limited to, delivery of a composition of the present invention via the mouth, sublingual, esophagus, gastric (for example the stomach), small intestines, large intestines or rectum. Oral delivery of the present invention may include, but is not limited to, the use of a capsule, pastille, pill, tablet, solution, gel, suspension, emulsion, syrup, elixir, tincture, mouthwash, lozenges, chewing gum, lollipop, osmotic-controlled release oral delivery system, or the like. Gastric delivery may involve the use of a tube or nasal passage that leads directly to the stomach, for example, a percutaneous endoscopic gastrostomy tube. Gastric delivery may involve direct injection made through the abdominal wall. Rectal delivery may involve, but is not limited to, the use of a suppository, ointment, enema, murray drip, or the like. A therapeutic dose of the present invention may be delivered to a patient by means of controlled release, for example but not limited to, controlled release drug delivery pellet or pill.

(228) As used herein, the terms “pulmonary system” or “respiratory system” are used interchangeably and refer, but are not limited, to the respiratory region, conducting airways, nasal cavity, sinuses, nasopharynx, oropharynx, larynx, trachea, bronchi, bronchioles, respiratory bronchioles, alveolar ducts, alveolar sacs, respiratory epithelium (e.g., alveolar epithelial cells), endothelial cells, or the like.

(229) As used herein, the terms “pulmonary delivery” and “pulmonary administration” are used interchangeably and refer to delivering a composition of the present invention to the respiratory system through the respiratory route, including but not limited to, intranasal administration, oral administration, and oral inhalative administration (e.g., intratracheal instillation and intratracheal inhalation) of a therapeutically effective amount for local or systemic treatment. Pulmonary delivery of a therapeutically effective amount of a composition of the present invention may be achieved by dispersion, for example by using a syringe. Pulmonary delivery of a composition of the present invention may be achieved by aerosol administration, wherein aerosol administration may deposit a therapeutically effective amount of the present invention by gravitational sedimentation, inertial impaction, or diffusion.

(230) Pulmonary delivery of a therapeutically effective amount of a composition of the present invention may be deposited on any mucus layer of the respiratory system, for example, but not limited to, the mucus layer which coats the walls of conducting airways, the smaller airway, and/or alveolar space.

(231) As used herein, an “appropriate control” refers to an untreated, otherwise identical cell or population (e.g., a subject who was not administered the composition described herein, or was administered by only a subset of agents provided herein, as compared to a non-control cell).

(232) As used herein, a “reference level” can refer to one or more parameters or markers as measured for a normal, otherwise unaffected cell population or tissue (e.g., a biological sample obtained from a healthy subject, or a biological sample obtained from the subject at a prior time point, or a biological sample that has not yet been contacted with a pathogen as described herein).

For measuring or monitoring therapeutic efficacy, a level determined prior to treatment or earlier in treatment can also provide a reference level for a given parameter or value.

(233) As used herein, the term “modulates” refers to an effect including increasing or decreasing a given parameter as those terms are defined herein.

(234) The terms “increased,” “increase,” “increases,” or “enhance” or “activate” are all used herein to generally mean an increase of a property, level, or other parameter by a statistically significant amount; for the avoidance of any doubt, the terms “increased”, “increase” or “enhance” or “activate” means an increase of at least 10% as compared to a reference level, for example an increase of at least about 20%, or at least about 30%, or at least about 40%, or at least about 50%, or at least about 60%, or at least about 70%, or at least about 80%, or at least about 90% or up to and including a 100% increase or any increase between 10-100% as compared to a reference level, or at least about a 2-fold, or at least about a 3-fold, or at least about a 4-fold, or at least about a 5-fold or at least about a 10-fold increase, at least about a 20-fold increase, at least about a 50-fold increase, at least about a 100-fold increase, at least about a 1000-fold increase or more as compared to a reference level. For example, increasing activity can refer to activating a receptor or a signaling pathway (e.g., antibody production or inflammation).

(235) The terms “decrease”, “reduced”, “reduction”, or “inhibit” are all used herein to mean a decrease or lessening of a property, level, or other parameter by a statistically significant amount. In some embodiments of any of the aspects, “reduce,” “reduction” or “decrease” or “inhibit” typically means a decrease by at least 10% as compared to a reference level (e.g., the absence of a given treatment) and can include, for example, a decrease by at least about 10%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or more. As used herein, “reduction” or “inhibition” does not encompass a complete inhibition or reduction as compared to a reference level. “Complete inhibition” is a 100% inhibition as compared to a reference level. A decrease can be preferably down to a level accepted as within the range of normal for an individual without a given disorder.

(236) As used herein the term “comprising” or “comprises” is used in reference to compositions, methods, and respective component(s) thereof, that are essential to the method or composition, yet open to the inclusion of unspecified elements, whether essential or not.

(237) As used herein the term “consisting essentially of” refers to those elements required for a given embodiment. The term permits the presence of elements that do not materially affect the basic and novel or functional characteristic(s) of that embodiment.

(238) The term “consisting of” refers to compositions, methods, and respective components thereof as described herein, which are exclusive of any element not recited in that description of the embodiment.

(239) As used in this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural references unless the context clearly dictates otherwise. Thus for example, references to “the method” includes one or more methods, and/or steps of the type described herein and/or which will become apparent to those persons skilled in the art upon reading this disclosure and so forth. Similarly, the word “or” is intended to include “and” unless the context clearly indicates otherwise. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of this disclosure, suitable methods and materials are described below.

(240) The abbreviation, “e.g.” is derived from the Latin *exempli gratia*, and is used herein to indicate a non-limiting example. Thus, the abbreviation “e.g.” is synonymous with the term “for example.”

(241) The abbreviation, “etc.” is derived from the Latin *et cetera*, and is used herein to indicate a non-limiting list. Thus, the abbreviation “etc.” is synonymous with the term “and other similar

things”, or “and so forth”.

(242) Other than in the operating examples, or where otherwise indicated, all numbers expressing quantities of ingredients or reaction conditions used herein should be understood as modified in all instances by the term “about.” The term “about” when used in connection with percentages can mean  $\pm 1\%$ .

(243) The term “statistically significant” or “significantly” refers to statistical significance and generally means a two-standard deviation (2SD) difference, above or below a reference value. Additional definitions are provided in the text of individual sections below.

(244) It should be understood that this invention is not limited to the particular methodology, protocols, and reagents, etc., described herein and as such may vary. The terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention, which is defined solely by the claims.

(245) As used herein and in the claims, the singular forms include the plural reference and vice versa unless the context clearly indicates otherwise. The term “or” is inclusive unless modified, for example, by “either.” Other than in the operating examples, or where otherwise indicated, all numbers expressing quantities of ingredients or reaction conditions used herein should be understood as modified in all instances by the term “about.”

(246) Unless otherwise explained, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs.

(247) It is to be understood that the foregoing description and the following examples are illustrative only and are not to be taken as limitations upon the scope of the invention. Various changes and modifications to the disclosed embodiments, which will be apparent to those of skill in the art, may be made without departing from the spirit and scope of the present invention. Further, all patents, patent applications, and publications identified are expressly incorporated herein by reference for the purpose of describing and disclosing, for example, the methodologies described in such publications that might be used in connection with the present invention. These publications are provided solely for their disclosure prior to the filing date of the present application. Nothing in this regard should be construed as an admission that the inventors are not entitled to antedate such disclosure by virtue of prior invention or for any other reason. All statements as to the date or representation as to the contents of these documents are based on the information available to the applicants and do not constitute any admission as to the correctness of the dates or contents of these documents.

(248) All patents and other publications identified are expressly incorporated herein by reference for the purpose of describing and disclosing, for example, the methodologies described in such publications that could be used in connection with the present invention. These publications are provided solely for their disclosure prior to the filing date of the present application. Nothing in this regard should be construed as an admission that the inventors are not entitled to antedate such disclosure by virtue of prior invention or for any other reason. All statements as to the date or representation as to the contents of these documents is based on the information available to the applicants and does not constitute any admission as to the correctness of the dates or contents of these documents.

(249) Some embodiments of the technology described herein can be defined according to any of the following numbered paragraphs: 1. An engineered extracellular vesicle comprising: at least one fusion polypeptide comprising: (i) at least one protein of interest (POI) domain or a fragment thereof; and (ii) at least one vesicle targeting domain, wherein the POI domain is in an extracellular position relative to a lipid membrane of the extracellular vesicle. 2. The engineered extracellular vesicle of paragraph 1, wherein the extracellular vesicle is an exosome. 3. The engineered extracellular vesicle of paragraph 1 or paragraph 2, wherein the protein of interest (POI) domain or a fragment thereof is a N-terminal domain of the fusion polypeptide. 4. The engineered

extracellular vesicle of any one of paragraphs 1-3, wherein the vesicle targeting domain is a C-terminal domain of the fusion polypeptide. 5. The engineered extracellular vesicle of any one of paragraphs 1-4, wherein the fusion polypeptide comprises at least two POI domains and/or at least two exosome targeting domains. 6. The engineered extracellular vesicle of any one of paragraphs 1-5, wherein the fusion polypeptide further comprises a peptide linker. 7. The engineered extracellular vesicle of any one of paragraphs 1-6, wherein the fusion polypeptide further comprises a fragment crystallizable region (Fc) domain. 8. The engineered extracellular vesicle of any one of paragraphs 1-7, wherein the vesicle targeting domain is in a luminal position relative to the lipid membrane of the extracellular vesicle. 9. The engineered extracellular vesicle of any one of paragraphs 1-7, wherein the vesicle targeting domain is in an exterior position relative to the lipid membrane of the extracellular vesicle. 10. The engineered extracellular vesicle of any one of paragraphs 1-9, wherein the POI domain is selected from the group consisting of: Table 1. 11. The engineered extracellular vesicle of any one of paragraphs 1-10, wherein the POI domain is PD-L1 or a fragment thereof. 12. The engineered extracellular vesicle of any one of paragraphs 1-11, wherein the POI domain is PD-L2 or a fragment thereof. 13. The engineered extracellular vesicle of any one of paragraphs 1-12, wherein the POI domain is FGL1 or a fragment thereof. 14. The engineered extracellular vesicle of any one of paragraphs 1-13, wherein the POI domain is 4-1BBL or a fragment thereof. 15. The engineered extracellular vesicle of any one of paragraphs 1-14, wherein the POI domain is CTLA-4 or a fragment thereof. 16. The engineered extracellular vesicle of any one of paragraphs 1-15, wherein the POI domain substantially binds to one or more of a target polypeptide. 17. The engineered extracellular vesicle of paragraph 16, wherein the target polypeptide is selected from the group consisting of: Table 2. 18. The engineered extracellular vesicle of any one of paragraphs 1-17, wherein the vesicle targeting domain is selected from the group consisting of: Table 3. 19. The engineered extracellular vesicle of any one of paragraphs 1-18, wherein the linker is in an exterior position relative to the lipid membrane of the extracellular vesicle. 20. The engineered extracellular vesicle of any one of paragraphs 1-18, wherein the linker is a transmembrane linker. 21. The engineered extracellular vesicle of any one of paragraphs 1-18, wherein the linker is in a luminal position relative to the lipid membrane of the extracellular vesicle. 22. The engineered extracellular vesicle of any one of paragraphs 1-21, wherein the extracellular vesicle does not comprise an endogenous POI polypeptide. 23. A composition comprising a plurality of the engineered extracellular vesicles of any one of paragraphs 1-22. 24. The composition of paragraph 23, further comprising a pharmaceutically acceptable carrier. 25. An engineered extracellular vesicle comprising: (a) a first fusion polypeptide comprising: (i) at least one protein of interest (POI) domain or a fragment thereof; and (ii) at least one vesicle targeting domain, wherein the at least one POI domain is in an extracellular position relative to a lipid membrane of the extracellular vesicle, (b) a second fusion polypeptide comprising: (i) at least one protein of interest (POI) domain or a fragment thereof; and (ii) at least one vesicle targeting domain, wherein the POI domain is in an extracellular position relative to a lipid membrane of the extracellular vesicle, and wherein the at least one vesicle targeting domain is within a lipid membrane of the extracellular vesicle. 26. A composition comprising two or more of the engineered extracellular vesicles selected from any one of paragraphs 1-25. 27. An extracellular vesicle composition comprising: a plurality of artificial synapses, wherein each artificial synapse comprises (i) an extracellular vesicle; (ii) one or more sticky binders; and (iii) one or more signaling domains. The composition of paragraph 27, wherein the extracellular vesicle comprises an exosome. 28. The composition of paragraph 27, wherein the one or more sticky binders is selected from the group consisting of: a GPI anchor, a fatty acylation site, and a prenylation site. 30. The composition of paragraph 27, wherein the signaling domain comprises one or more of: PD-L1, PD-L2, CTLA-4 (CD152), 4-1BBL (CD137L), HVEM (CD270), FGL1, OX-2 (CD200), Galectin-9, PVR (CD155), Nectin-2 (CD112) isoform alpha, Nectin-2 (CD112) isoform beta, Nectin-2 (CD112) isoform delta, IL-10, TSG-6, B7-H3 (CD276), B7-H4 (VTCN1), B7-H5

(VISTA), B7-H7 (HHLA2), BTNL1, VSIG8, VSIG3 (IGSF11), VSIG4, TIM-3 (HAVCR2), TIM-4 (TIMD4), CEACAM1, BTN3A1, BTN3A2, BTN2A1, BTNL8, BTN2A2, BTN1A1, TIGIT, CD27L (CD70), CD30L (CD153), GITRL, CD40L (CD154), LIGHT (CD258), TL1, CD80, CD86, LFA-3 (CD58), SLAM (CD150), CD40, CD28, CD28H, CD2, LFA-3 (CD58), CD48, CD226, DR3, DcR3, FasL, TIM-1 (CD365), PD-1, or active fragment thereof. 29. A method of producing the engineered extracellular vesicle or the composition of any one of paragraphs 1-30, comprising: (a) providing a population of cells expressing a vector construct encoding one or more sticky binder and one or more signaling domains; and (b) isolating a plurality of artificial synapses from the population of cells. 30. A method of producing the engineered extracellular vesicle or the composition of any one of paragraphs 1-30, comprising: (a) providing a population of cells expressing a vector construct encoding one or more sticky binder and one or more signaling domains; and (b) isolating a plurality of artificial synapses from the population of cells; and (c) purifying the plurality of artificial synapses from the population of cells. 33. The method of paragraph 31 or paragraph 32, the isolating is via size exclusion chromatography. 34. The method of paragraph 32, wherein the purifying is via multimodal chromatography. 35. The method of any of paragraphs 31-34, further comprising performing an assay for POI binding to a target polypeptide. 36. The method of paragraph 35, wherein the vector construct further encodes a promoter. 37. The method of paragraph 36, wherein the promoter is a tissue-specific promoter or an inducible promoter. 38. A method of modulating inflammation in a subject, the method comprising: administering a composition comprising a plurality of engineered extracellular vesicles to a subject in need thereof, wherein the engineered extracellular vesicles comprise at least one fusion polypeptide comprising: (i) at least one protein of interest (POI) domain or a fragment thereof; and (ii) at least one vesicle targeting domain. 39. The method of paragraph 38, wherein the extracellular vesicle comprises an exosome. 40. The method of any one of paragraphs 38-39, further comprising selecting a subject that has or is suspected of having an autoimmune disease or an inflammatory disease or condition. 41. The method of any one of paragraphs 38-40, wherein the vesicle targeting domain is selected from the group consisting of: a Glycosylphosphatidylinositol (GPI) anchor, a fatty acylation site, and a prenylation site. 42. The method of any one of paragraphs 38-41, wherein the vesicle targeting domain is a GPI anchor. 43. The method of any one of paragraphs 38-41, wherein the vesicle targeting domain is C1C2. 44. The method of any one of paragraphs 38-43, wherein the protein of interest (POI) domain comprises one or more of: PD-L1, PD-L2, CTLA-4 (CD152), 4-1BBL (CD137L), HVEM (CD270), FGL1, OX-2 (CD200), Galectin-9, PVR (CD155), Nectin-2 (CD112) isoform alpha, Nectin-2 (CD112) isoform beta, Nectin-2 (CD112) isoform delta, IL-10, TSG-6, B7-H3 (CD276), B7-H4 (VTCN1), B7-H5 (VISTA), B7-H7 (HHLA2), BTNL1, VSIG8, VSIG3 (IGSF11), VSIG4, TIM-3 (HAVCR2), TIM-4 (TIMD4), CEACAM1, BTN3A1, BTN3A2, BTN2A1, BTNL8, BTN2A2, BTN1A1, TIGIT, CD27L (CD70), CD30L (CD153), GITRL, CD40L (CD154), LIGHT (CD258), TL1, CD80, CD86, LFA-3 (CD58), SLAM (CD150), CD40, CD28, CD28H, CD2, LFA-3 (CD58), CD48, CD226, DR3, DcR3, FasL, TIM-1 (CD365), PD-1, or active fragment thereof. 45. The method of any one of paragraphs 38-44, wherein the protein of interest (POI) domain is PD-L1 or a fragment thereof. 46. The method of any one of paragraphs 38-44, wherein the protein of interest (POI) domain is PD-L2 or a fragment thereof. 47. The method of any one of paragraphs 38-44, wherein the protein of interest (POI) domain is CTLA-4 or a fragment thereof. 48. The method of any one of paragraphs 38-44, wherein the protein of interest (POI) domain is HVEM or a fragment thereof. 49. The method of paragraph 40, wherein the inflammatory disease and/or condition is acute. 50. The method of paragraph 40, wherein the inflammatory related disease and/or condition is chronic. 51. The method of paragraph 38, wherein administering the composition comprises injection, topical administration, or inhalation. 52. Use of a composition comprising a plurality of engineered extracellular vesicles, the engineered extracellular vesicles each comprising: at least one fusion polypeptide comprising: (i) at least one protein of interest (POI) domain or a fragment thereof; and (ii) at least one vesicle

targeting domain for the treatment of an inflammatory disease or condition. 53. Use of a composition comprising a plurality of engineered extracellular vesicles, the engineered extracellular vesicles each comprising: at least one fusion polypeptide comprising: (i) at least one protein of interest (POI) domain or a fragment thereof; and (ii) at least one vesicle targeting domain for the treatment of an autoimmune disease or condition. 54. Use of a composition comprising a plurality of engineered extracellular vesicles, the engineered extracellular vesicles each comprising: at least one fusion polypeptide comprising: (i) at least one protein of interest (POI) domain or a fragment thereof; and (ii) at least one vesicle targeting domain for the treatment of cancer.

## EXAMPLES

(250) The following examples are provided by way of illustration, not limitation.

### Example 1

#### Design of Artificial Synapse

(251) As described, artificial synapses are engineered to induce and propagate biological signaling, including for example, antagonist and agonist signaling. Artificial synapses are designed to include hallmark biophysical and biochemical features of extracellular vesicles, further including vesicle targeting domains and signaling domains. Vesicle targeting domains capable of attaching to extracellular vesicles such as exosomes, signaling domains, optionally including a linker (e.g., Fc linker), can be organized in genetic vector constructs. Designs are shown in FIG. 1.

(252) Sticky binders are extracellular vesicle targeting sequences. Preliminary extracellular vesicle targeting sequences of interest are from, but not limited to, 4F2 (CD98), ADAM10, CD298, TFR2, transmembrane domains of CD9, MARCKS, KRAS, etc. or the like as appreciate by one of ordinary skill in the art. The Inventors discovered high efficiency when proteins are engineered with a GPI domain. Optionally, linker regions such as an Fc linker between the vesicle targeting domains and signaling domains can be added.

(253) A variety of signaling domains are of interest with proof-of-concept examples including PD-L1, PD-L2 and CTLA-4 (CD152). Artificial synapses including these three signaling domains are shown in FIGS. 2-5.

(254) Each of these elements are described in the following non-limiting examples.

### Example 2

#### Genetic Constructs

(255) Examples of constructs including these variable elements (e.g., sticky binders GPI or C1C2, or signaling domains including PD-L1, PD-L2 and CTLA-4 (CD152) were engineered into vectors shown in FIGS. 2-5.

### Example 3

#### Purification of hPD-L1 Tagged Artificial Synapses by a Multimodal Resin Marketed for Exosome Purification

(256) Upon expression of hPD-L1-Fc-GPI in mammalian cells, artificial synapses were further purified using a size exclusion resin marketed for exosome purification. Large MW artificial synapses elute in the first fraction as shown by the high hPD-L1 concentration and exosome quantity (2.26E9 artificial synapses/ml) in elution 1. Clean in place (CIP) fractions show bound and eliminated proteins from the Inventors' exosome elution. Results are shown in FIG. 6.

### Example 4

#### hPDL1-Fc-GPI Exosome Purification-Size Exclusion Chromatography Column

(257) Artificial synapses engineered from exosomes such as hPDL1-Fc-GPI after elution from size exclusion resin marketed for exosome purification can be further purified via a size exclusion column as shown here. Using a size exclusion chromatography (SEC), artificial synapses elute in fractions 7-9. Total protein (determined by Qubit™) and hPD-L1 ng/ml (determined by ELISA) of each fraction is shown in the graph. Bars show exosome number per ml (i.e., 1E10 artificial synapses/ml etc.). Fractions 7-9 contain >99% purified artificial synapses. Fractions 7-9 are pooled and may be concentrated using a filtration device, for example a 10K MWCO Amicon®

Centrifugal Filter. Final purified product is filtered through a low protein binding 0.2  $\mu\text{m}$  or 0.45  $\mu\text{m}$  filter, for example a PES filter. Results are shown in FIG. 7.

#### Example 5

##### hPD-L1 Expression on Artificial Synapses

(258) Exosome quantity and hPD-L1 concentration was determined in SEC fractions 7-9. Knowing the molecular weight of engineered hPD-L1, the Inventors can determine the number of hPD-L1 molecules per exosome to be approximately between 12 to 40 PD-L1/exosome. This value is consistent between different purification runs and constructs. Results are shown in FIG. 8.

#### Example 6

##### Purification of hPD-L2-Fc-GPI Artificial Synapses Via Multimodal Resin Chromatography Marketed for Exosome Purification

(259) This graph shows Abs 280 of multimodal resin chromatography fractions and quantity of hPDL2 in indicated fractions. Artificial synapses eluted in Elution 1.

(260) Clean in place (CIP) fractions show bound and eliminated proteins from the Inventors' exosome elution. Results are shown in FIG. 9.

#### Example 7

##### PD-L2 Purification Via Size Exclusion Chromatography

(261) Artificial synapses engineered from artificial synapses such as hPDL2-GPI after elution from size exclusion resin size exclusion resin marketed for exosome purification are further purified via size exclusion chromatography as shown. Results are shown in FIG. 10.

#### Example 8

##### hCTLA4-Fc-GPI Exosome Purification Via Size Exclusion Chromatography

(262) Using size exclusion chromatography marketed for exosome purification, artificial synapses elute in fractions 7-9. Total protein (determined by Qubit™) and hPD-L1 ng/ml (determined by ELISA) of each fraction is shown in the graph. Fractions 7-9 are pooled and contain >99% purified artificial synapses. Pooled artificial synapses engineered from artificial synapses fractions may then be concentrated using a filtration device, for example a 10K MWCO Amicon®. Final purified product is filtered through a low protein binding filter, for example a 0.2  $\mu\text{m}$  or 0.45  $\mu\text{m}$  PES filter. Results are shown in FIG. 11.

#### Example 9

##### PD-L1 and PD-L2 In Vitro Assay from DISCOVERX®

(263) To perform this validation method, the Inventors modified the PathHunter® PD-1 Signaling Bioassay from DISCOVERX®. Briefly, the PathHunter® PD-1 Signaling Bioassay relies on the well-established PathHunter® Enzyme Fragment Complementation (EFC) technology to interrogate receptor activity. EFC consists of a split  $\beta$ -galactosidase ( $\beta$ -gal) enzyme: the Enzyme Donor (ED) and Enzyme Acceptor (EA) fragments which independently have no  $\beta$ -gal activity. However, when forced to complement they form an active  $\beta$ -gal enzyme that will hydrolyze substrate to produce a chemiluminescent signal. The PathHunter® PD-1 Signaling Bioassay consists of human cells engineered to stably express an ED-tagged PD-1 receptor, while EA is fused to the phosphotyrosine-binding SH2 domain of the intracellular signaling protein, SHP1. Ligand or antibody-induced activation of the receptor results in phosphorylation of the receptor's cytosolic tail. The SH2-domain fused to EA binds the phosphorylated receptor, forcing complementation of ED and EA, resulting in formation of an active  $\beta$ -gal enzyme, which hydrolyzes the substrate to produce a chemiluminescent signal. Full-length PD-1 receptor was engineered with a small  $\beta$ -gal fragment (ED in red) fused to its C-terminus, and the SH2-domain of SHP1 was engineered with the complementing  $\beta$ -gal fragment (EA). These constructs were stably expressed in Jurkat cells (produced by DISCOVERX®), while PD-L1 and PD-L2 was stably expressed on artificial synapses produced by Diadem Biotherapeutics. Artificial synapses were engineered to have surface expressed human PD-L1 or PD-L2. Briefly, the gene sequence coding for the extracellular domain of human PD-L1 or PD-L2 was linked to the exosome via a



glycosylphosphatidylinositol (GPI) linker with an Fc domain between the linker and PD-L1 or PD-L2 (PD-L1-Fc-GPI and PD-L2-Fc-GPI). Additional variations of the Inventors' PD-L1 and PD-L2 artificial synapses include cloning a C1C2 linker (from MFGE8) in place of the GPI linker, and with or without the Fc domain. The Inventors also cloned murine versions of PD-L1 and PD-L2 extracellular domains in place of the human PD-L1 and PD-L2 all variations. Ligand engagement, through addition of ligand-presenting artificial synapses, results in phosphorylation of PD-1, leading to the recruitment of SHP1-EA

(264) The Inventors obtained approximately 1000× higher increase in Relative Light Units (RLU) in Jurkat signaling cells treated with PD-L1 or PD-L2 labeled artificial synapses when compared to soluble PD-L1-Fc or PD-L2-Fc ligand, respectively. Meaning, it took 1000× less µg/ml of PD-L1 or PD-L2 on artificial synapses than solubilized PD-L1-Fc or PD-L2 ligand to achieve the same RLU signaling. Results are shown in FIG. 12.

#### Example 10

PD-L1 In Vivo Assay-Experimental Autoimmune Uveoretinitis (EAU) in Lewis Rats Bioassay

(265) Experimental autoimmune uveoretinitis (EAU) is an organ-specific, T lymphocyte-mediated autoimmune disease, which serves as a model for several human ocular inflammations of an apparently autoimmune nature. There is a statistically significant initial reduction in EAU in mPDL1 artificial synapse treated rats via either the intravitreal and intravenous delivery modes. 2nd intravitreal and 3rd intravenous injections are performed on Day 12. There appears to be a more rapid rate of resolution in the 1× intravitreal and intravenous groups. (C) Simplified view of aforementioned results. (D) Weight of rats was monitored throughout the study. 3rd intravitreal and 4th intravenous injections are performed on Day 16. There does not appear to be any significant change in EAU in any of the test groups. The aforementioned results provide proof of principle of successfully immunizing the rats with human cell derived artificial synapses with mouse PDL1 injected into rats. Results are shown in FIG. 13.

#### Example 11

Engineered Exosome Multivalent Display

(266) The inventors have developed the following 3 types of protein display on or within exosomes: Type I membrane proteins wherein the N-Terminus is on the luminal (interior) side of the exosome membrane and the C-Terminus is on the exterior of the exosome. Type II membrane proteins wherein the N-Terminus is on the exterior while the C-Terminus is on the interior. Luminal internally loaded proteins which are linked to the exosome by a Myristoylation/Palmitoylation site which attaches proteins to the interior of the exosome membrane.

(267) FIGS. 14-21 demonstrate the various embodiments of the engineered extracellular vesicles.

(268) Additional embodiments or ligands displayed on the exosome surface (Type I and Type II membrane proteins) and internal luminal display can include the following: Type I: PD-L1, PD-L2, FGL1, OX40L. Type II: 4-1BBL, GITRL, CD27L, CD30L Luminal: NanoLuc® luciferase; Green fluorescent protein (GFP) (e.g., eGFP, etc.); Red fluorescent protein (RFP) (e.g., mScarlet, mCherry, mRuby, tdTomato, etc.); Cyan fluorescent protein (CFP); Yellow fluorescent protein (YFP); A therapeutic protein; and CRISPR/CAS-9

(269) FIG. 20 shows an exemplary multiple protein display construct. Sequences such as P2A, E2A, F2A, and T2A induce ribosomal slippage which prevent peptide bond formation, meaning that a single mRNA transcript with a 2A sequence will result in two separate peptides after translation. This allows the expression of two separate proteins from one promoter region and thus loading of two proteins on an exosome. Any combination of the proteins of interest domains provided herein can be engineered. Furthermore, a cell line with multiple transgene inserts under separate promoter control. Either method can be used to label Type I, Type II, and luminal display proteins.

#### Example 12a

Designed and Engineered Human Fusion Polypeptide Constructs

(270) The inventors have designed, engineered, and purified the following human fusion polypeptide constructs for therapeutic use (FIG. 5A-FIG. 5WW): pEF5-FRT-hPDL1-C1C2 (FIG. 5I) pEF5-FRT-hPDL2-C1C2 (FIG. 5J) pEF5-FRT-hPDL1-GPI-P2A-hFGL1-GPI (FIG. 5E) pEF5-FRT-hCTLA4-Fc-GPI (FIG. 5C) pEF5-FRT-hPDL2-Fc-GPI (FIG. 5H) pEF5-FRT-hPD-L1-GPI-P2A-hHVEM-GPI (FIG. 5D) pEF5-FRT-hPDL1-GPI (FIG. 5F) pcDNA5-FRT-hSecPDL1-GPI (FIG. 5O) pcDNA5-FRT-hPDL1-GPI (FIG. 5F) pcDNA5-FRT-hPDL1-Link-GPI (FIG. 5T) pcDNA5-FRT-4F2-h41BBL (FIG. 5K) pcDNA5-FRT-Tfr2-h41BBL (FIG. 5P) pEF5-FRT-hPDL1-Fc-GPI (FIG. 5G) pcDNA5-FRT-CD9tm3-h41BBL (FIG. 5Q) pcDNA5-FRT-hPDL1-Fc-GPI (FIG. 5G) pcDNA5-FRT-hPDL1-4Fc-CD9tm2 (FIG. 5RR) pcDNA5-FRT-hPDL1-Fc-CD9tm2KRAS (FIG. 5UU) pcDNA5-FRT-hPDL1-4Fc-CD9tm2KRAS (FIG. 5SS) pcDNA5-FRT-hPDL1-4Fc-GPI (FIG. 5L) pcDNA5-FRT-hPDL1-ADAM10 (FIG. 5QQ) pcDNA5-FRT-MyrPalm-4F2-h41BBL (FIG. 5R) pcDNA5-FRT-MyrPalm-h41BBL (FIG. 5S) pcDNA5-FRT-hPDL1-Fc-CD9tm2 (FIG. 5TT) pcDNA5-FRT-hSecPDL1-CD9tm4 (FIG. 5W) pcDNA5-FRT-hSecPDL1-CD9tm2KRas (FIG. 5V) pcDNA5-FRT-hSecPDL1-CD9tm2 (FIG. 5U) pcDNA5-FRT-hSecPDL1-CD81 (FIG. 5X) pEF5-FRT-hCD200-Fc-GPI (FIG. 5Y) pEF5-FRT-hCD200-GPI (FIG. 5BB) pEF5-FRT-hTSG6-GPI (FIG. 5FF) pEF5-FRT-hPDL2-GPI (FIG. 5EE) pEF5-FRT-hFGL-1-GPI (FIG. 5Z) pEF5-FRT-hHVEM-GPI (FIG. 5DD) pEF5-FRT-hGal9-GPI (FIG. 5CC) pEF5-FRT-hHVEM-Fc-GPI (FIG. 5GG); and pEF5-FRT-hGal9-Fc-GPI (FIG. 5AA)

#### Example 12b

##### Designed and Engineered Fusion Polypeptide Constructs

(271) The inventors have designed, engineered, and purified the following mouse fusion polypeptide constructs for therapeutic use (FIG. 5A-FIG. 5WW): pcDNA5-FRT-mPDL1-mFc-CD9tm2KRAS (FIG. 5WW) pcDNA5-FRT-mPDL1-mFc-CD9tm2 (FIG. 5VV) pcDNA5-FRT-mPDL1-mFc-GPI (FIG. 5NN) pcDNA5-FRT-mPDL1-GPI (FIG. 5KK) pEF5-FRT-mPDL2-GPI (FIG. 5OO) pEF5-FRT-mPDL1-GPI-P2A-mHVEM-GPI (FIG. 5PP) pEF5-FRT-mPDL1-GPI (FIG. 5KK) pEF5-FRT-mPDL2-Fc-GPI (FIG. 5MM) pEF5-FRT-mPDL1-Fc-GPI (FIG. 5JJ) pEF5-FRT-mCTLA4-Fc-GPI (FIG. 5HH) pEF5-FRT-mPDL1-C1C2 (FIG. 5II); and pEF5-FRT-mPDL2-C1C2 (FIG. 5LL).

#### Example 12c

##### Designed and Engineered Luminal Loaded Fusion Polypeptide Constructs

(272) The inventors have designed, engineered, and purified the following fusion polypeptide constructs for internal luminal loading of the fusion polypeptide: pcDNA5-FRT-Myr-NanoLuc (FIG. 5M) pcDNA5-FRT-Myr-mScarlet (FIG. 5N)

#### Example 13

##### Purification of Exosomes Labeled with Type I Membrane Fusion Polypeptides

(273) The inventors have purified engineered EVs, including hPD-L1-GPI; hPDL1-Fc-GPI; hPDL2-Fc-GPI; hCTLA4-Fc-GPI; mPDL1-GPI; and mPD-L1-Fc-GPI. The process for purification and analytical processing of the engineered EVs are shown in the flow chart provided in FIG. 21.

(274) Size exclusion chromatography was performed to purify hPD-L1-GPI (no Fc) exosomes (FIG. 24). Protein, RNA and DNA measurements in SEC fractions. Invitrogen Qubit™ fluorometric assays were used to measure biomolecules from unmodified concentrated cell media SEC fractions or hPD-L1-Exo-Tag concentrated cell media SEC fractions. PD-L1 was measured using an R&D systems PD-L1 ELISA kit. Dot-blot immunoblot analysis of SEC fractions. A 96-well dot blot apparatus was used to immobilize 50 µl of each SEC fraction onto PVDF. Exosome size and concentration was measured in fraction 7 by tunable resistive pulse sensing (TRPS). It was confirmed that GPI anchors the hPD-L1 fusion protein onto the exosomes (FIG. 25).

(275) Furthermore, a commercially available multimodal exosome purification resin was also used to purify and isolate PD-L1-GPI exosomes and PD-L1-Fc-GPI exosomes. Fraction 7 was further analyzed by dot blots (FIG. 28A-28B). In particular, FIG. 28B shows SEC purification results of various embodiments of human PD-L1 displayed on the surface of extracellular vesicles. One

embodiment is the hPD-L1-4Fc-GPI (CMV) construct as seen in the top dot blot (stained with rabbit monoclonal anti-PD-L1 antibody). Another embodiment is the hPD-L1-4Fc-GPI (EF1a) as seen in the top dot blot (stained with rabbit monoclonal anti-PD-L1 antibody).

(276) Large MW exosomes elute in the first fraction as shown by the high hPD-L1 concentration and exosome quantity ( $2.26 \times 10^9$  exosomes/ml) in elution 1. Clean in place (CIP) fractions show bound and eliminated proteins from our exosome elution. Exosome quantity and hPD-L1 concentration was determined in SEC fractions 7-9. Knowing the molecular weight of engineered hPD-L1, we can determine the number of hPD-L1 molecules per exosome to be approximately 12 PD-L1/exosome. This value is consistent between different purification runs and constructs (FIG. 8).

(277) Human hPD-L2 and hCTLA-4-Fc-GPI SEC fractions were purified. In addition, purification of the mouse PD-L1-FcGPI exosomes was performed (FIG. 29). The mouse Fc-PD-L1 expressing exosomes have a higher valency than those that do not comprise the Fc linker.

#### Example 14

##### Comparative Proteomics Analysis of the Engineered EVs

(278) Fc-GPI enables high density display and has a higher abundance than endogenous PTGFRN or CD81. Therefore, comparison proteomics of transprotein expression and surface labeling on the engineered exosomes, hPD-L1-Fc-GPI; hPD-L2-Fc-GPI; and hCTLA-Fc-GPI, was performed to determine the effects on endogenous protein expression in engineered exosomes. It was confirmed that the fusion polypeptide expression does not affect the relative expression of native and associated exosome proteins. However, the trans protein may crowd out abundant proteins like CD81 (data not shown).

#### Example 15

##### Scale-Up Production and Purification of mPD-L1-Fc-GPI Exosomes Using Microcarriers in a Stirred Tank Single-Use Bioreactor (STR)

(279)  $1 \times 10^7$  HEK 293 cells were utilized for the production of mPDL1-Fc-GPI exosomes. Cells were passaged on SoloHill® Microcarriers up to Passage 4, at which point cells were expanded in a 2.5 L Stirred Tank Single-Use Bioreactor. Passage 4 cells were cultured for an additional 5 days and media was harvested on Day 5 and used for exosome purification. The general aim and process is provided below

(280) AIM: Utilize SoloHill®'s Xeno-free microcarrier technology to scale up cells for engineering EVs and evaluate Microcarrier-stir tank bioreactor technology for production of therapeutic exosomes in the Xeno-free medium conditions.

(281) Passage 1:

(282) Thaw vial ( $1.00 \times 10^7$ ) of cells and seed Corning® T-150 & CellSTACK®2 tissue culture treated flask at  $1.00 \times 10^4$  cells per cm.<sup>sup.2</sup> seed density.

(283) Perform 100% medium exchange from both flasks on day 3.

(284) Harvest Corning® T-150 & CellSTACK®2 flasks on day 4 post seeding and seed spinner microcarrier culture.

(285) Passage 2:

(286) Expand cells in  $2 \times 200$  mL spinner flasks at 10 cm.<sup>sup.2</sup>/mL microcarrier density using SoloHill®'s Xeno-free prototype microcarrier.

(287) Seed microcarrier cultures at  $1.00 \times 10^4$  cells per cm.<sup>sup.2</sup> seed density and T-25 as flatware control flask.

(288) Perform 80% batch volume medium exchange from spinners and T-25 flasks on day 3.

(289) Harvest both microcarrier and T-25 flasks on day 4 post seeding and seed spinner microcarrier culture.

(290) Passage 3:

(291) Expand cells in  $3 \times 300$  mL spinner flasks at 10 cm.<sup>sup.2</sup>/mL microcarrier density using SoloHill®'s Xeno-free prototype microcarrier.

- (292) Seed microcarrier cultures at 1.00E+04 cells per cm.<sup>sup.2</sup> seed density and T-25 as flatware control flask.
- (293) Perform 80% batch volume medium exchange from spinners and T-25 flasks on day 3.
- (294) Harvest both microcarrier and T-25 flasks on day 4 post seeding.
- (295) Seed microcarrier-stir tank bioreactor for exosome production.
- (296) Passage 4:
- (297) Expand cells into a 2.5 L microcarrier-stir tank at 10 cm.<sup>sup.2</sup>/mL surface area to medium ratio.
- (298) Seed cultures at 1.00E+04 cells per cm.<sup>sup.2</sup> seed density and T-25 as flatware control flask.
- (299) Perform 80% batch volume medium exchange on day 2.
- (300) On day 3 rinse all cultures with 2×cell culture volumes of DPBS containing Ca and Mg.
- (301) Add exosome production medium (DMEM-1% GlutaMAX™) to all cultures at 10 cm.<sup>sup.2</sup>/mL surface area to medium volume ratio.
- (302) On day 5 collect harvest spent medium from all cultures, filter using 0.45 µm Nalgene rapid flow system and freeze at -20° C.
- (303) Procedures:
- (304) Medium Composition DMEM 1× (Corning ref #10-013-CV) 1% GlutaMAX™ (Thermo ref #35050061) 3% Human platelet lysate (Stemulate from Cook Reagentec PG-NH-500)
- Cell Harvest Protocol for Planar Culture
- (305) Settle microcarriers and remove maximum volume of spent medium without removing microcarriers.
- (306) Wash microcarrier culture with DPBS 2× time at 0.1 mL/cm.<sup>sup.2</sup> volume to surface area ratio.
- (307) Add 37° C. warmed TrypLE™ 5× enzyme at 0.012 mL/cm.<sup>sup.2</sup> and incubate flask at room temperature for ~15 minutes.
- (308) Add complete medium at 0.024 mL/cm.<sup>sup.2</sup> to quench TrypLE™ 5× activity.
- (309) Perform viable cell count using NC200 cell count instrument.
- (310) Nuclei Count Protocol for Microcarrier Culture
- (311) Obtain 4-5 mL of microcarrier culture from bioreactor or spinner flask
- (312) Settle microcarriers and remove maximum volume of spent medium without removing microcarriers.
- (313) Add 1.5 mL NucleoCounter® Reagent A to macrocarrier sample tube and vortex at high speed for a minute.
- (314) Add 1.5 mL NucleoCounter® Reagent B to macrocarrier sample tube and vortex at high speed for a minute.
- (315) Perform nuclei count using NC200 nuclei count instrument.
- (316) Medium Collection from STR Bioreactor
- (317) Stop all controls and settle microcarriers in the bioreactor vessel.
- (318) Pump out medium through screen bag into collection bottle at 200 mL/minute flowrate using peristaltic pump.
- (319) Inside BSC pour medium into 0.45 µm Nalgene™ Rapid-Flow™ filter system and remove free floating cells.
- (320) Freeze medium bottles in minus 20° C. freezer.
- (321) Medium Collection from Spinner Flasks
- (322) Inside BSC pour microcarrier culture into 0.45 µm Nalgene™ Rapid-Flow™ system and remove free floating cells as well as microcarriers.
- (323) Freeze medium bottles in minus 20° C. freezer.
- (324) TABLE-US-00008 Cell culture set points Tem- Dis- Incubator perature Agitation solved Oxygen CO.sub.2 ° C. rpm (DO) % pH setting % T-Flask 37 n/a n/a n/a 5 ± 1 CellSTACK 2 37 n/a n/a n/a 5 ± 1 Spinner flask 37 35 n/a n/a 5 ± 1 STR bioreactor 37 35 50 7.35 n/a

(325) FIG. 31 shows mPDL1-Fc-GPI production, growth parameters, and analyte concentrations from a 2.6-L culture in a Stirred Tank Single-Use (STR) bioreactor. Day 2: 80% batch volume medium was exchanged (1st increase in glucose and decreased in lactate) Day 3: rinse culture with 2× cell culture volumes of DPBS containing Ca and Mg. (2nd increase in glucose and decreased in lactate). Add exosome production medium (DMEM-1% GlutaMAX™) to culture at 10 cm.sup.2/mL surface area to medium volume ratio.

(326) mPDL1 was purified using the purification process outlined above (FIGS. 32-33).

#### Example 16

##### PD-L1-Fc-GPI and PDL2-Fc-GPI Exosomes Increase PD-1 Signaling

(327) The purified exosomes were tested using the modified DISCOVERX® Assay in FIG. 12A. Approximately a 1000× increase in Relative Light Units (RLU) was achieved for Jurkat signaling cells treated with PD-L1 or PD-L2 labeled exosomes when compared to soluble PD-L1-Fc or PD-L2-Fc ligands alone, respectively. Therefore, it takes 1000× less µg/ml of PD-L1 or PD-L2 on the engineered exosomes to activate PD-1 over solubilized ligands, PD-L1-Fc or PD-L2, achieve the same RLU signaling. FIG. 12B show a dose-response curves for the PD-L1 and PD-L2 exosomes vs soluble PD-L1 and PD-L2 signaling bioassay. FIG. 12B shows dose-response curves for the PD-L1 and PD-L2 exosomes comprising an Fc linker and GPI sticky binder vs. soluble ligands with an Fc domain linker. These results show that the PD-L1 and PD-L2 polypeptides fused with the Fc and GPI domains on EVs have a more potent effect on PD-1 signaling than the soluble ligands alone.

#### Example 17

##### In Vivo Assay—Therapeutic Effect of mPD-L1 Exosomes in an Experimental Autoimmune Uveoretinitis (EAU) Model in Lewis Rats

(328) Lewis rats were challenged with retinal antigen interphotoreceptor retinoid-binding protein (IRBP) peptide. This model can be used to study anterior and posterior chamber dependent EAU. Rats were immunized on Day 1 with EAU presenting typically at Day 6. Clinical scores in the rat were determined. The EAU dosing schedule is shown in FIG. 13A. EAU dosing test article are shown in the following table.

(329) TABLE-US-00009 EAU dosing test articles

Unmodified mPD-L1-Fc-GPI	mPD-L1-Fc-GPI	mPD-L1 Exosomes (IVT)	Exosomes 1X (IVT)	Exosomes 10X (IVT)	Exosomes (IV)	Dose
2 ul	2 ul	5 ml/kg	Total protein 40 ug/ml	40 ug/ml	400 ug/ml	40 ug/ml
concentration	Total protein 80 ng/eye	80 ng/eye	800 ng/eye	50 ug/animal	administered	Exosome 5.7 × 10 <sup>sup.10</sup> /ml
2.34 × 10 <sup>sup.10</sup> /ml	2.34 × 10 <sup>sup.11</sup> /ml	2.34 × 10 <sup>sup.10</sup> /ml	concentration	Total exosomes 4.7 × 10 <sup>sup.7</sup>	4.7 × 10 <sup>sup.7</sup>	4.7 × 10 <sup>sup.8</sup>
2.93 × 10 <sup>sup.10</sup>	administered	*IVT-intravitreal,	IV-intravenous			

(330) The study design is outlined below:

##### (331) TABLE-US-00010 Group Test Article N Route Concentration Dosage Regimen 1

Cyclosporine 8 p.o. 1 mg/mL 10 mg/kg BID from day 0 to Day 20 2 Negative control (PBS 8 Intravitreal both 1× 2-3 µL Day 6, Day 12, vehicle) eyes and Day 16 3 Unmodified exosomes 8 Intravitreal both 1× (~40 ug/ml) 2-3 µL Day 6 and Day (Control exosomes) eyes 12 4 mPD-L1-Fc-GPI (40 8 Intravitreal both 1× (~40 ug/ml) 2-3 µL Day 6, Day 12, ug/ml) eyes and Day 16 5 mPD-L1-Fc-GPI (40 4 Intravenous 1× (~40 ug/ml) 5 mL/kg Day 1, Day 6, ug/ml) Injection Day 12, and Day 16 6 No IRBP peptide but 4 Intravitreal both 1× (~40 ug/ml) 2-3 µL Day 6, Day 12, treated with Test Agent eyes and Day 16 B (for tolerability) 7 mPD-L1-Fc-GPI (400 8 Intravitreal both 1× (400 ug/ml) 2-3 µL Day 6, Day 12, ug/ml) eyes and Day 16

(332) Clinical Scores were determined as follows:

(333) TABLE-US-00011 EAU Clinical Scores in Rats Score Clinical Criteria 0 No disease; eye is translucent and reflects light (red reflex) 0.5 Dilated blood vessels in the iris (trace) 1 Engorged blood vessels in the iris; abnormal pupil contraction 2 Hazy anterior chamber; decreased red reflex 3 Moderately opaque anterior chamber, but pupil still visible; dull red reflex 4 Opaque anterior

chamber and obscured pupil; red reflex absent; proptosis

Each higher grade includes the criteria of the preceding one.

(334) It was discovered that there is a statistically significant initial reduction in EAU in mPDL1 exosome treated rats via either the intravitreal and intravenous delivery modes as compared with untreated animals (FIG. 13A). Rat weight did not change post immunization (FIG. 13C).

#### Example 18

##### Purification of Exosomes Labeled with Type II Membrane Proteins

(335) The inventors designed, engineered, and purified pcDNA5-FRT-4F2-4-1BBL exosomes by the methods provided herein (FIG. 34). Several embodiments of the 4-1BBL labeled exosomes are shown in FIG. 35. Cell expression of the 4F2-4-1BBL was confirmed (data not shown). FIGS.

92A-92B shows the purification of 4F2-4-1BBL exosomes.

#### Example 19

##### Purification of Luminal Labeled Exosomes (Internal Loading)

(336) In addition to Type I and Type II display fusion proteins on the surface of an EV, exosomes can be loaded with fusion proteins that are localized to the lumen of the phospholipid bilayer of the exosome (FIG. 37). The Myr/Palm sequence used herein when fused to mScarlet the fusion protein into the luminal interior of extracellular vesicles. Fluorescence at an excitation wavelength 470 nm and emission wavelength of 665-720 nm peaks in SEC fractions 7, 8, and 9. SEC fractions 7, 8, and 9 contain exosomes as demonstrated by the dot blot. Fraction 8 was further analyzed for exosome quantification using an EXOVIEW® system (FIG. 38). Unmodified exosomes do not show fluorescence. Exosomes show near 80% loading with Myr/Palm-mScarlet. The remaining 20% were out of the detection limit. Thus, nearly 100% internal loading was achieved using the specific Myr/Palm sequence.

(337) NanoLuc® luciferase expressing exosomes were also purified with the Myr/Palm sequence incorporated into the vector encoding the fusion polypeptide. A Qubit™ fluorometer was used to measure total protein and PROMEGA® Nano-Glo® substrate and plate luminometer to measure luminescence (FIG. 39A). Tetraspanin characterization of exosomes was performed and determined that the NanoLuc® luciferase exosomes were internally loaded and purified in fraction 8 (FIG. 39B).

## Claims

1. An engineered extracellular vesicle comprising a fusion protein, the fusion protein comprising: at least two signaling domains, each of the at least two signaling domains being a same signaling domain and selected from the group consisting of either full length or active fragments of 4-1BBL (CD137L), CD27L (CD70), CD30L (CD153), mCD30L, GITRL, CD40L (CD154), mCD140L, LIGHT (CD258), TL1, mTL1, FasL, NKG2 family, CD94, CD98, OX40L (CD252), and 4-1BB/TNFRSF4/CD137; at least one linker, and at least one vesicle targeting domain linked to the at least two signaling domains; wherein the at least one linker comprises a fragment crystallizable region (Fc) domain, wherein the at least one vesicle targeting domain comprises a Type II transmembrane domain, and wherein the at least two signaling domains are in an extracellular position relative to a lipid membrane of the extracellular vesicle.
2. The engineered extracellular vesicle of claim 1, wherein the at least one vesicle targeting domain comprises a fatty acylation site or a prenylation site, whereby the at least one vesicle targeting domain is embedded in a phospholipid bilayer of the engineered extracellular vesicle through covalent lipid attachment to the fatty acylation site or the prenylation site.
3. The engineered extracellular vesicle of claim 1, wherein the at least one vesicle targeting domain is selected from the group consisting of sequences encoding peptides or protein domains from ADAM10, TFR2, and MARCKS comprising modified myristoylation and palmitoylation tags; sequences encoding prenylation sites and fatty acylation sites; and sequences encoding lipid

affinity tags derived from KRAS, CD81, CD63, ALIX, TSG101, CD98, CD298, GPI, CD9, and CD105.

4. The engineered extracellular vesicle of claim 1, wherein the at least one vesicle targeting domain comprises at least two exosome targeting domains.

5. The engineered extracellular vesicle of claim 1, further comprising a tetraspanin selected from the group consisting of CD9, CD63, CD81, CD82, CD53, and CD37.

6. The engineered extracellular vesicle of claim 5, wherein the at least two signaling domains comprise three identical signaling domains.

7. The engineered extracellular vesicle of claim 1, comprising a plurality of the fusion protein, and wherein the density of the plurality of fusion protein is configured to support receptor clustering on a target cell.

8. The engineered extracellular vesicle of claim 1, wherein the at least two signaling domains substantially bind to one or more of a target polypeptide.

9. The engineered extracellular vesicle of claim 1, further comprising one or more secondary fusion proteins, each comprising a secondary signaling domain different from the at least two signaling domains of the fusion protein.

10. The engineered extracellular vesicle of claim 9, wherein the secondary signaling domains are each independently selected from the group consisting of either full-length or active fragments of PD-L1 (CD274), PD-L2 (CD273), CTLA-4 (CD152), 4-1BBL (CD137L), HVEM (CD270), FGL1, OX-2 (CD200), Galectin-9, PVR (CD155), Nectin-2 (CD112) isophorm alpha, Nectin-2 (CD112) isophorm delta, mNectin-2 beta, IL-10, TSG-6, B7-H3 (CD276), B7-H4 (VTCN1), B7-H5 (VISTA), B7-H7 (HLA2), VSIG8, VSIG3 (IGSF11), VSIG4, Tim-3 (HAVCR2), Tim-4, CEACAM1 (CD66a), BTN3A1, BTN3A2, BTN2A1, BTNLR, BTN2A2, mBTN2A2, BTN1A1, TIGIT, CD27L (CD70), CD30L (CD153), mCD30L, GITRL, CD40L (CD154), mCD140L, LIGHT (CD258), TL1, mTL1, B7-1 (CD80), B7-2 (CD86), LFA-3 (CD58), SLAM (CD150), mSLAM, CD40, CD28, mCD28, CD28H/TMIGD2/IGPR1, CD2, CD48, CD226, DR3, DcR3, FasL, Tim-1 (CD365), PD-1 (CD279), mScarlet, Nanoluciferase, A2AR, PECAM-1, STAB-1, Clever-I, NRP1, NRP2, SEMA3A, SEMA3F, RGMB/DRG11, HLA DII, HMGB1, TCR, SHP-1, SHP-2, FBOX38, SH2DIA, B7RP1, IDO, NOX2, TNFRSF18/GITR/CD357, SISP1, B7-H6/NCR3LG1, APLNR, IFNg receptor, WNTSA, PAK4, IL-6, NKG2 family, NKG2 family ligands, Killer cell Ig-like receptors, CD4, CD5, CD27, CD39, CD44, CD47, CD73, CD94, CD96, CD98, 1GSF2/CD101, PVIRG/CD112R, IL5RB/CD122, OX40L (CD252), 4-1BB/TNFRSF4/CD137, KIRs/CD158 family, CD160, SIRP alpha/CD172a, CD200R, LAG-3/CD223, CD244, BTLA/CD272, B7H2/ICOSLG/B7RP1/CD275, ICOS/CD278, LIAR-1/CD305, Collagen family members, SIGLEC7/CD328, SIGLEC9/CD329, NKp30/CD337, TNFR superfamily, Nectin-like binding receptors, Nectin, IL10RA, IL10RB, TNFRSF25, TNFRSF6B, CD113, CD30, TRAF family members, and TIM family members.

11. The engineered extracellular vesicle of claim 9, wherein the one or more secondary fusion proteins each comprises a secondary vesicle targeting domain linked to its secondary signaling domain.

12. The engineered extracellular vesicle of claim 11, wherein the one or more secondary vesicle targeting domains are each independently selected from the group consisting of sequence prides or protein domains from ADAM10, TFR2, and MARCKS comprising modified myristoylation and palmitoylation tags; sequences encoding prenylation sites and fatty acylation sites; and sequences encoding lipid affinity tags derived from KRAS, CD81, CD63, ALIX, TSG101, CD98, CD298, GPL CD9, and CD105.

13. An engineered extracellular vesicle comprising a fusion protein, the fusion protein comprising: at least two signaling domains, each of the at least two signaling domains being a same signaling domain and selected from the group consisting of either full-length or active fragments of 4-1BBL (CD137L), CD27L (CD70), CD30L (CD153), mCD30L, GITRL, CD40L (CD154), mCD140L,

LIGHT (CD258), TL1, mTL1 FasL, NKG2 family, CD94, CD98, OX40L (CD252), and 4-1BB/TNFRSF4/CD137; at least one linker, and at least one vesicle targeting domain linked to the at least two signaling domains; wherein the at least one linker comprises a fragment crystallizable region (Fc) domain and is between the at least two signaling domains and the at least one vesicle targeting domain, wherein the at least one vesicle targeting domain comprises a Type II transmembrane domain, and wherein the at least two signaling domains are in an extracellular position relative to a lipid membrane of the extracellular vesicle.

14. The engineered extracellular vesicle of claim 13, wherein the linker is an Fc from IgG1, Fc from IgG2, Fc from IgG3, or Fc from IgG4, or wherein the linker further comprises Gly-Ser-Ser-Gly (SEQ ID NO: 319), a cleavable 2A sequence, P2A, E2A, F2A, T2A, or (GGGGS (SEQ ID NO: 320)) n.

15. The engineered extracellular vesicle of claim 13, wherein the at least one vesicle targeting domain comprises a sequence encoding a fatty acylation site or a prenylation site, whereby the at least one vesicle targeting domain is embedded in a phospholipid bilayer of the engineered extracellular vesicle through covalent lipid attachment to the fatty acylation site or the prenylation site.

16. The engineered extracellular vesicle of claim 13, wherein the at least one vesicle targeting domain is selected from the group consisting of sequences encoding peptides or protein domains from ADAM10, TFR2, and MARCKS comprising modified myristoylation and palmitoylation tags; sequences encoding prenylation sites and fatty acylation sites; and sequences encoding lipid affinity tags derived from KRAS, CD81, CD63, ALIX, TSG101, CD98, CD298, GPI, CD9, and CD105.

17. The engineered extracellular vesicle of claim 13, further comprising a tetraspanin selected from the group consisting of CD9, CD63, CD81, CD82, CD53, and CD37.

18. The engineered extracellular vesicle of claim 17, wherein the at least two signaling domains comprise three identical signaling domains.

19. The engineered extracellular vesicle of claim 13, comprising a plurality of the fusion protein, and wherein the density of the plurality of fusion protein is configured to support receptor clustering on a target cell.

20. The engineered extracellular vesicle of claim 13, further comprising one or more secondary fusion proteins, each comprising a secondary signaling domain different from the signaling domain of the fusion protein.

21. The engineered extracellular vesicle of claim 20, wherein the secondary signaling domains are each independently selected from the group consisting of either full-length or active fragments of PD-L1, PD-L2, CTLA-4 (CD152), 4-1BBL (CD137L), HVEM (CD270), FOL1, OX-2 (CD200), Galectin-9, PVR (CD155), Nectin-2 (CD112) isophorm alpha, Nectin-2 (CD112) isophorm delta, mNectin-2 beta, IL-10, TSG-6, B7-H3 (CD276), B7-H4 (VTCN1), B7-H5 (VISTA), B7-H7 (HHLA2), VSIG8, VSIG3 (IGSF11), VSIG4, Tim-3 (HAVCR2), Tim-4, CEACAM1 (CD66a), BTN3A1, BTN3A2, BTN2A1, BTNL8, BTN2A2, mBTN2A2, BTN1A1, TIGIT, CD27L (CD70), CD30L (CD153), mCD30L, GITRL, CD40L (CD154), mCD140L, LIGHT (CD258), TL1, mTL1, B7-1 (CD80), B7-2 (CD86), LFA-3 (CD58), SLAM (CD150), mSLAM, CD40, CD28, mCD28, CD28H/TMIGD2/IGPR1, CD2, CD48, CD226, DR3, DcR3, FasL, Tim-1 (CD365), PD-1 (CD279), mScarlet, Nanoluciferase, AZAR, PECAM-I, STAB-1, Clever-1, NRP1, NRP2, SEMA3A, SEMA3F, RGMB/DRG11, HLA I/II, HMGB1, TCR, SHP-1, SHP-2, FBOX38, SH2D1A, B7RP1, IDO, NOX2, TNFRSF18/GITR/CD357, SISP1, B7-H6/NCR3LG1, APLNR, IFNg receptor, WNTSA, PAK4, IL-6, NKG2 family, NKG2 family ligands, Killer cell Ig-like receptors, CD4, CD8, CD27, CD39, CD44, CD47, CD73, CD94, CD96, CD98, IGSF2/CD101, NECTIN2/CD112, PVIRO/CD112R, ILSRB/CD122, OX40L (CD252), 4-1BB/TNFRSF4/CD137, CTLA-4/CD152, KIRs/CD158 family, CD160, SIRP alpha/CD172a, CD200R, LAG-3/CD223, CD244, BTLA/CD272, B7H2/ICOSLG/B7RP1/CD275, ICOS/CD278, LIAR-1/CD305, Collagen



family members, SIGLEC7/CD328, SIGLEC9/CD329, NKp30/CD337, TNER superfamily, Nectin-like binding receptors, Nectin, IL10RA, IL10RB, TNFRSF25, TNFRSF6B, CD113, CD30, TRAF family members, and TIM family members.

22. The engineered extracellular vesicle of claim 20, wherein the one or more secondary fusion proteins each comprises a secondary vesicle targeting domain linked to its secondary signaling domain.

23. The engineered extracellular vesicle of claim 22, wherein the one or more secondary vesicle targeting domains are each independently selected from the group consisting of sequences encoding peptides or protein domains from ADAM10, TFR2, and MARCKS comprising modified myristoylation and palmitoylation tags; sequences encoding prenylation sites and fatty acylation sites; and sequences encoding lipid affinity s derived from KRAS, CD81, CD63, ALIX, TSG101, CD98, CD298, GPI, CD9, and CD105.

---