



US 20250249107A1

(19) **United States**

(12) **Patent Application Publication**
DEMUTH et al.

(10) **Pub. No.: US 2025/0249107 A1**

(43) **Pub. Date: Aug. 7, 2025**

(54) **COMPOUNDS INCLUDING A MUTANT KRAS SEQUENCE AND A LIPID AND USES THEREOF**

(71) Applicant: **Elicio Therapeutics, Inc.**, Boston, MA (US)

(72) Inventors: **Peter C. DEMUTH**, Boston, MA (US);
Julian ADAMS, Boston, MA (US);
Martin P. STEINBUCK, Boston, MA (US)

(21) Appl. No.: **19/186,849**

(22) Filed: **Apr. 23, 2025**

Related U.S. Application Data

(62) Division of application No. 18/882,166, filed on Sep. 11, 2024, now abandoned, which is a division of application No. 16/977,155, filed on Sep. 1, 2020, now Pat. No. 12,138,301, filed as application No. PCT/US2019/020404 on Mar. 1, 2019.

(60) Provisional application No. 62/637,879, filed on Mar. 2, 2018.

Publication Classification

(51) **Int. Cl.**
A61K 47/54 (2017.01)
A61K 39/00 (2006.01)
A61K 39/39 (2006.01)
A61K 47/60 (2017.01)
A61P 35/00 (2006.01)
C12N 9/14 (2006.01)
C12N 9/96 (2006.01)

(52) **U.S. Cl.**
CPC *A61K 47/543* (2017.08); *A61K 39/001164* (2018.08); *A61K 39/39* (2013.01); *A61K 47/60* (2017.08); *A61P 35/00* (2018.01); *C12N 9/14* (2013.01); *C12N 9/96* (2013.01); *C12Y 306/05002* (2013.01); *A61K 2039/545* (2013.01); *A61K 2039/55561* (2013.01)

(57) **ABSTRACT**

The invention provides compounds including a KRAS sequence that is linked to a lipid by a linker, and related compositions and methods.

Specification includes a Sequence Listing.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6





FIG. 7

COMPOUNDS INCLUDING A MUTANT KRAS SEQUENCE AND A LIPID AND USES THEREOF

SEQUENCE LISTING

[0001] The instant application contains a Sequence Listing which has been submitted electronically in XML format and is hereby incorporated by reference in its entirety. Said XML copy, created on Feb. 7, 2024 is named 51026-026WO2_Sequence_Listing_2_7_24 and is 28,139 bytes in size.



BACKGROUND OF THE INVENTION

[0002] RAS proteins are essential components of cell signaling pathways. Oncogenic activation of RAS proteins by mutation is frequently detected in several types of cancer, including pancreatic cancer. There continues to be a need for further and more effective cancer treatments.

SUMMARY OF THE INVENTION

[0003] The invention provides compounds that can be used in therapeutic methods.

[0004] Accordingly, in the first aspect, the invention features a compound including a mutant KRAS sequence and a lipid, where the mutant KRAS sequence is conjugated to the lipid by a linker, and (i) the linker includes one or more polyethylene glycol blocks, (ii) the lipid is 1,2-distearoyl-sn-glycero-3-phosphoethanolamine (DSPE), and (iii) the mutant KRAS sequence comprises or consists of an amino acid sequence selected from the group consisting of YKLVVVGADGVGKSALTI (SEQ ID NO:23), YKLVVVGAVGVGKSALTI (SEQ ID NO:24), YKLVVVGARGVGKSALTI (SEQ ID NO:25), YKLVVVGAAGVGKSALTI (SEQ ID NO:26), YKLVVVGASGVGKSALTI (SEQ ID NO:27), YKLVVVGAGGVGKSALTI (SEQ ID NO:28), YKLVVVGATGVGKSALTI (SEQ ID NO:29), and YKLVVVGAGDVGKSALTI (SEQ ID NO:30).

[0005] In the second aspect, the invention features a compound including a mutant KRAS sequence and a lipid, where the mutant KRAS sequence is conjugated to the lipid by a linker, and (i) the linker includes one or more polyethylene glycol blocks, (ii) the lipid is 1,2-distearoyl-sn-glycero-3-phosphoethanolamine (DSPE), and (iii) the mutant KRAS sequence comprises or consists of an amino acid sequence selected from the group consisting of GYKLVVVGADGVGKSALTI (SEQ ID NO:1), CYKLVVVGAVGVGKSALTI (SEQ ID NO:2), CYKLVVVGARGVGKSALTI (SEQ ID NO:3), CYKLVVVGAAGVGKSALTI (SEQ ID NO:4), CYKLVVVGASGVGKSALTI (SEQ ID NO:5),

CYKLVVVGACGVGKSALTI (SEQ ID NO:6), CYKLVVVGATGVGKSALTI (SEQ ID NO:22), and CYKLVVVGAGDVGKSALTI (SEQ ID NO:7).

[0006] In one embodiment of the first and the second aspects of the invention, the mutant KRAS sequence, at its N-terminus, is conjugated to the linker through a cysteine-maleimide linkage.

[0007] In another embodiment of the first and the second aspects of the invention the linker includes 48 repeat units of polyethylene glycol.

[0008] In an additional embodiment of the first and the second aspects of the invention, the mutant KRAS sequence, at its N-terminus, is conjugated to the following structure:

[0009] In the third aspect, the invention features a composition including one or more compounds of the first and the second aspects of the invention and a pharmaceutically acceptable carrier.

[0010] In one embodiment of the third aspect of the invention, the composition includes (1) a compound including the amino acid sequence YKLVVVGADGVGKSALTI (SEQ ID NO:23), (2) a compound including the amino acid sequence YKLVVVGAVGVGKSALTI (SEQ ID NO:24), (3) a compound including the amino acid sequence YKLVVVGARGVGKSALTI (SEQ ID NO:25), (4) a compound including the amino acid sequence YKLVVVGAAGVGKSALTI (SEQ ID NO:26), (5) a compound including the amino acid sequence YKLVVVGASGVGKSALTI (SEQ ID NO:27), (6) a compound including the amino acid sequence YKLVVVGAGGVGKSALTI (SEQ ID NO:28) or a compound including the amino acid sequence YKLVVVGATGVGKSALTI (SEQ ID NO:29), and (7) a compound including the amino acid sequence YKLVVVGAGDVGKSALTI (SEQ ID NO:30).


[0011] In one embodiment of the third aspect of the invention, the composition includes (1) a compound including the amino acid sequence YKLVVVGADGVGKSALTI (SEQ ID NO:23), (2) a compound including the amino acid sequence YKLVVVGAVGVGKSALTI (SEQ ID NO:24), (3) a compound including the amino acid sequence YKLVVVGARGVGKSALTI (SEQ ID NO:25), (4) a compound including the amino acid sequence YKLVVVGAAGVGKSALTI (SEQ ID NO:26), (5) a compound including the amino acid sequence YKLVVVGASGVGKSALTI (SEQ ID NO:27), (6) a compound including the amino acid sequence YKLVVVGAGGVGKSALTI (SEQ ID NO:28), and (7) a compound including the amino acid sequence YKLVVVGAGDVGKSALTI (SEQ ID NO:30).

[0012] In another embodiment of the third aspect of the invention, the composition includes (1) a compound including the amino acid sequence CYKLVVVGADGVGKSALTI (SEQ ID NO:1), (2) a compound including the amino acid sequence CYKLVVVGAVGVGKSALTI (SEQ ID NO:2), (3) a compound including the amino acid sequence CYKLVVVGARGVGKSALTI (SEQ ID NO:3), (4) a compound including the amino acid sequence CYKLVVVGAAGVGKSALTI (SEQ ID NO:4), (5) a compound including the amino acid sequence CYKLVVVGASGVGKSALTI (SEQ ID NO:5), (6) a compound including the amino acid sequence CYKLVVVGACGVGKSALTI (SEQ ID NO:6) or a compound including the amino acid sequence CYKLVVVGATGVGKSALTI (SEQ ID NO:22), and (7) a compound including the amino acid sequence CYKLVVVGAGDVGKSALTI (SEQ ID NO:7).

[0013] In another embodiment of the third aspect of the invention, the composition includes (1) a compound including the amino acid sequence CYKLVVVGADGVGKSALTI (SEQ ID NO:1), (2) a compound including the amino acid sequence CYKLVVVGAVGVGKSALTI (SEQ ID NO:2), (3) a compound including the amino acid sequence CYKLVVVGARGVGKSALTI (SEQ ID NO:3), (4) a compound including the amino acid sequence CYKLVVVGAAGVGKSALTI (SEQ ID NO:4), (5) a compound including the amino acid sequence CYKLVVVGASGVGKSALTI (SEQ ID NO:5), (6) a compound including the amino acid sequence CYKLVVVGACGVGKSALTI (SEQ ID NO:6), and (7) a compound including the amino acid sequence CYKLVVVGAGDVGKSALTI (SEQ ID NO:7).

[0014] In some embodiments, 700 μ g of each compound is included in the composition.

[0015] In an additional embodiment of the third aspect of the invention, the composition further includes a compound consisting of the nucleotide sequence 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3' (SEQ ID NO:8), which, at its 5' end, is bonded or linked by a linker to the following lipid:



or a salt thereof, where X is O or S. In one embodiment, the nucleotide sequence is bonded to the lipid.


[0016] In another embodiment, all internucleoside groups connecting the nucleosides in 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3' (SEQ ID NO:8) are phosphorothioates.

[0017] In the fourth aspect, the invention features a method of treating a cancer in a human patient, the method including administering the composition of the third aspect of the invention to the patient.

[0018] In one embodiment of the fourth aspect of the invention, the method further includes administering an adjuvant, such as an adjuvant including a CpG nucleotide sequence (e.g., 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3'; SEQ ID NO:8).

[0019] In some embodiments, 0.1 mg, 0.5 mg, or 2.5 mg of the adjuvant is administered.

[0020] In one embodiment of the fourth aspect of the invention, the method further includes administering to the patient a compound consisting of the nucleotide sequence 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3' (SEQ ID NO:8), which, at its 5' end, is bonded or linked by a linker to the following lipid:




or a salt thereof, where X is O or S. In one embodiment, the nucleotide sequence is bonded to the lipid.

[0021] In one embodiment, 0.1 mg, 0.5 mg, or 2.5 mg of the compound is administered.

[0022] In another embodiment, all internucleoside groups connecting the nucleosides in 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3' (SEQ ID NO:8) are phosphorothioates.

[0023] In one embodiment of the fourth aspect of the invention, the cancer is a pancreatic cancer, a lung cancer, or a colorectal cancer.

[0024] In the fifth aspect, the invention features a kit including (i) a compound of any one of claims 1 to 5, or a composition of any one of claims 6-8, and (ii) a compound consisting of the nucleotide sequence 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3' (SEQ ID NO:8), which, at its 5' end, is bonded or linked by a linker to the following lipid:



or a salt thereof, where X is O or S. In one embodiment, the nucleotide sequence is bonded to the lipid.

[0025] In another embodiment, all internucleoside groups connecting the nucleosides in 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3' (SEQ ID NO:8) are phosphorothioates.

Definitions

[0026] A “carbonyl,” as used herein, refers to a group —C(O)— .

[0027] A “carboxylate,” as used herein, refers to a group —C(O)—OH or a salt thereof.

[0028] As used herein, “conjugated” refers to a covalent bonding of a mutant KRAS sequence to a linker and, optionally, further covalent bonding of the linker to a lipid. In various examples, the covalent bonds connecting the linker to the mutant KRAS sequence or to the lipid may be a covalent bond between a sulfur atom and a sp^3 -hybridized carbon atom in succinimide, between a nitrogen atom and a carbon atom of a carbonyl, or between oxygen atom and the phosphorus atom of a thiophosphoryl or phosphoryl. In various examples, each of the KRAS sequence and the lipid may contain a group for bonding to a complementary group in the linker. For example, the KRAS polypeptide may include a sulfur atom (e.g., the sulfur atom in a cysteine residue) bonded to a complementary group, e.g., succinimide, in the linker. The bond from the sulfur atom in the KRAS polypeptide to the succinimide in the linker may be formed by a reaction of a thiol (e.g., a thiol in a cysteine residue) in the KRAS polypeptide and a maleimide in the linker. Alternatively, the KRAS polypeptide may include a nitrogen atom or a carbonyl group bonded to a carbonyl group or a nitrogen atom, respectively, in the linker. The bond between the carbonyl group and the nitrogen atom may be formed by a reaction between an amine or carboxylate in the KRAS polypeptide and a carboxylate or an amine, respectively, in the linker.

[0029] A “lipid” is any of a group of organic compounds that are greasy to the touch, insoluble in water, and soluble in organic solvents, and in general includes fatty acids and their derivatives. One example of a lipid used in the invention is 1,2-distearoyl-sn-glycero-3-phosphoethanolamine (DSPE).

[0030] A “linker,” as used herein, refers to a monovalent or divalent group, in which one valency is covalently bonded to one biologically functional group, and the other valency

is covalently bonded to another biologically functional group. In one example, a linker connects a KRAS sequence to a lipid, as noted above. Optionally, such a linker can include one or more polyethylene glycol blocks. In another example, a linker connects a nucleotide sequence of, e.g., a CpG oligonucleotide, to a lipid (e.g., $\text{—P(X)(OH)—O—CH(CH}_2\text{NHCO—(CH}_2\text{)}_{16}\text{—CH}_3\text{)}_2$, or a salt thereof, where X is O or S, as described herein). Such linkers can optionally include one or more nucleotides, for example, a dinucleotide (e.g., GG).

[0031] A “pharmaceutically acceptable carrier,” as used herein, refers to a vehicle capable of suspending or dissolving an active compound, and having the properties of being nontoxic and non-inflammatory in a subject (e.g., a human patient) to whom it is administered. Moreover, a pharmaceutically acceptable carrier may include a pharmaceutically acceptable additive, such as a preservative, antioxidant, fragrance, emulsifier, dye, or excipient known or used in the field of drug formulation and that does not significantly interfere with the therapeutic effectiveness of the biological activity of the active agent, and that is non-toxic to the subject.

[0032] A “phosphoryl,” as used herein, refers to a group $\text{—P(O)(OR}^A\text{)(OR}^B\text{)}$, where R^A is H, and R^B is a valency.

[0033] A “polyethylene glycol,” as used herein, refers to a block $\text{—(OCH}_2\text{CH}_2\text{)}_n\text{—}$, where n is a number of repeat units and is an integer from 2 to 50 (e.g., 24 or 48).

[0034] A “thiol,” as used herein, refers to a group —SH .

[0035] A “thiophosphoryl,” as used herein, refers to a group $\text{—P(S)(OR}^A\text{)(OR}^B\text{)}$, where R^A is H, and R^B is a valency.

[0036] The terms “treat,” “treatment,” and “treating” refer to therapeutic approaches in which the goal is to reverse, alleviate, ameliorate, inhibit, slow down, or stop the progression or severity of a condition associated with a disease or disorder, e.g., cancer. These terms include reducing or alleviating at least one adverse effect or symptom of a condition, disease, or disorder. Treatment is generally “effective” if one or more symptoms or clinical markers are reduced, or if a desired response (e.g., a specific immune response) is induced. Alternatively, treatment is “effective” if the progression of a disease is reduced or halted.

[0037] The invention provides several advantages. For example, in including DSPE moieties, certain compounds of the invention bind to endogenous albumin in subjects to whom they are administered, which enhances delivery of the compounds to the lymph nodes of the subjects. This facilitates the induction of a therapeutic immune response against the KRAS sequences of the compounds, leading to effective cancer treatment.

[0038] Other features and advantages of the invention will be apparent from the following detailed description, the drawings, and the claims.

DESCRIPTION OF THE DRAWINGS

[0039] FIG. 1 shows that amphiphile-KRas (aKRas), in combination with amphiphile-CpG (aCpG), activated splenocytes, while soluble KRas, in combination with aCpG, and the untreated and soluble KRas and polyIC (pIC) control groups, did not.


[0040] FIG. 2 shows that both aKRas G12R and G12V, in combination with aCpG, activated splenocytes, while neither soluble KRas G12R nor G12V in combination with aCpG, and the untreated and soluble KRas and pIC control groups, did not.

[0041] FIG. 3 shows a KRas-specific CD8+ T-cell response for mice immunized with the combination of aKRas G12D and aCpG, but not for soluble KRas G12D in combination with CpG or soluble wild-type KRas in combination with CpG.

ages are phosphorothioate. The structure is presented as the sodium salt.


DETAILED DESCRIPTION OF THE INVENTION

[0046] The invention provides compounds that can be used in therapeutic methods. The compounds of the invention each include a mutant KRAS sequence (e.g., any one of SEQ ID NOS:1-7 and 22; see Table 1, below, or any one of SEQ ID NOS:23-30, see Table 2, below) and a lipid (e.g., 1,2-distearoyl-sn-glycero-3-phosphoethanolamine (DSPE)). These moieties (the mutant KRAS sequence and the lipid) are linked to one another by a linker, for example, a linker including one or more polyethylene glycol blocks. In one example, the linker is:



[0042] FIG. 4 shows a KRas-specific CD8+ T-cell response (9mers) and CD8+ or CD4+ T-cell response (18mer) for mice immunized with the combination of aKRas G12D and aCpG, but not for soluble KRas G12D in combination with CpG or the untreated control.

In a specific example, a mutant KRAS sequence (e.g., any one of SEQ ID NOS:1-7 or 22, or any one of SEQ ID NOS:23-30) is bonded at its N-terminus through a Cys residue to a linker, and the linker is bonded to a lipid, where the linker is:




[0043] FIG. 5 shows amphiphile-KRas (Amph-KRAS) and amphiphile-CpG (Amph-CpG) vaccines elicit superior endogenous T-cell responses compared to soluble KRas vaccines, as indicated by increased cytokine production by splenocytes in a cytokine bead array. Soluble KRAS G12D=YKLVVVGADGVGKSALTI (SEQ ID NO: 23); Amph-KRAS G12D=DSPE-amido-dPEG24-amido-dPEG24-CYKLVVVGADGVGKSALTI (SEQ ID NO: 1); KRAS G12D 18-mer=YKLVVVGADGVGKSALTI (SEQ ID NO:23); CBA=Cytokine bead array; IFN=interferon; IL=interleukin; KRAS=Kirsten Rat Sarcoma.

[0044] FIG. 6 shows the structure of the amphiphile-peptide G12D 4-21 (SEQ ID NO:1).

[0045] FIG. 7 shows the structure of amphiphile-CpG-7909. All sugars are deoxyribose. All internucleoside link-

and the lipid is:




or a salt thereof.

[0047] The invention also provides compositions that include one or more compounds of the invention, together with a pharmaceutically acceptable carrier or diluent. Optionally, the compositions can include seven different

compounds as described above, where each of the seven different compounds includes a different sequence selected from SEQ ID NOS:1-7 and 23 or SEQ ID NOS:23-30. Compositions including subsets (e.g., 2, 3, 4, 5, or 6) of these compounds are also included in the invention. Different compounds included in the compositions of the invention can optionally each include the same or different KRAS sequences, linkers, and/or lipids.

[0048] The compositions of the invention can be used in methods to induce immune responses to the KRAS sequences of the compounds. Accordingly, the compositions can be referred to as immunogenic or vaccine compositions. The compositions can thus optionally include or be administered with one or more adjuvants. In one example, the adjuvant used can include a CpG oligonucleotide (e.g., 5'-TCGTCGTTTGTGCGTTTGTGCGTT-3'; SEQ ID NO:8; CpG-7909) that, at its 5' end, is bonded or linked by a linker to a lipid, such as the following:



or a salt thereof, where X is O or S. Preferably, X is S.
[0049] The CpG oligonucleotide may be directly bonded to the lipid. Alternatively, the linker bonded to the CpG oligonucleotide and to the lipid may be GG. In the CpG oligonucleotide, all internucleoside groups are phosphorothioates (e.g., all internucleoside groups in the compound may be phosphorothioates).
[0050] The compounds and compositions of the invention can be used in therapeutic methods. In particular, the KRAS

sequences of the compounds can induce an immune response to KRAS, which is expressed in certain cancer cells. Accordingly, the invention provides methods of treating cancer in a subject (e.g., a human patient) by administering one or more compounds or compositions of the invention to the subject. The invention also includes methods of inducing an immune response against KRAS in a subject (e.g., a human patient) by administering one or more compounds or compositions of the invention to the subject. In various examples, the cancer is selected from the group consisting of pancreatic cancer, lung cancer, and colorectal cancer. Optionally, the methods of the invention can further include administering a compound or composition of the invention in combination with a second (or further) different approach to treatment.

[0051] The invention also provides kits that each contain, for example, a first vessel that includes one or more compounds of the invention, optionally together with a second vessel that includes an adjuvant, such as an adjuvant as described herein.

TABLE 1

(SEQ ID NOS: 1-7 and 22)						
Pep- tide	Name	Sequence	MW (Da)	Res- idues	Amph- Peptide MW (Da)	
1	G12D 4-21	CYKLVVVGAD <u>D</u> GVGKSALTI	1893	19	5049	
2	G12V 4-21	CYKLVVVGAV <u>G</u> GVGKSALTI	1877	19	5033	
3	G12R 4-21	CYKLVVVGAR <u>G</u> GVGKSALTI	1934	19	5090	
4	G12A 4-21	CYKLVVVGAA <u>G</u> GVGKSALTI	1849	19	5005	
5	G12S 4-21	CYKLVVVGAS <u>G</u> GVGKSALTI	1865	19	5021	
6	G12C 4-21 or G12T 4-21	YKLVVVGAC <u>G</u> GVGKSALTI or CYKLVVVGAT <u>G</u> GVGKSALTI	1778	19	4934	
7	G13D 4-21	CYKLVVVGAG <u>D</u> GVGKSALTI	1790	19	5049	

For each of the “Amph-Peptide” referred to in Table 1, the lipid poly(ethylene glycol) moiety was conjugated via a maleimide (MAL) -cysteine coupling to the peptide using the following synthon: DSPE-amido-dPEG24-amido-dPEG24-MAL.

TABLE 2

(SEQ ID NOS: 23-30)				
Pep- tide	Name	Sequence	MW (Da)	Resi- dues
1	G12D 4-21	YKLVVVGAD <u>D</u> GVGKSALTI	1790	18
2	G12V 4-21	YKLVVVGAV <u>G</u> GVGKSALTI	1774	18


TABLE 2-continued

(SEQ ID NOS: 23-30)			
Pep- tide Name	Sequence	MW (Da)	Resi- dues
3	G12R 4-21 YKLVVVGARGVGKSALTI	1831	18
4	G12A 4-21 YKLVVVGAGGVGKSALTI	1746	18
5	G12S 4-21 YKLVVVGASGVGKSALTI	1762	18


TABLE 2-continued

(SEQ ID NOS: 23-30)			
Pep- tide Name	Sequence	MW (Da)	Resi- dues
6	G12C 4-21 YKLVVVGACGVGKSALTI or G12T 4-21 YKLVVVGATGVGKSALTI	1778	18
7	G13D 4-21 YKLVVVGAGDVGKSALTI	1790	18


[0052] A compound of the invention may be prepared from a mutant KRAS polypeptide, a linker precursor, and a lipid using methods known in the art. In some variations, a linker precursor is first linked to a lipid, e.g., as follows:



Amidation reaction conditions are known in the art, for example, typical amidation conditions include the use of reagents, such as EDC/DMAP, HATU/HOAt, or HBTU/HOAt. Alternatively, the carboxylate may be replaced with an O-succinimide ester, pentafluorophenyl ester, or tetrafluorophenyl ester. The product of this reaction may then be reacted with a mutant KRAS polypeptide, e.g., as follows:




-continued





The 1,4-addition reaction may be carried out in an appropriate solvent, e.g., a polar organic solvent or water.

[0053] Alternatively, a compound of the invention may be prepared as follows. A linker precursor may be first linked to a lipid, e.g., as follows:




The product may then be reacted with an amine group (e.g., N-terminal amine) in the mutant KRAS polypeptide to produce a compound of the invention:






[0054] A CpG oligonucleotide may be bonded directly or linked by a linker to the lipid. These compounds may be produced using the ordinary phosphoramidite chemistry known in the art. In some examples, the CpG oligonucleotide or CpG oligonucleotide that, at its 5' end, is bonded to GG may be reacted with the following compound:



to produce an intermediate, which upon oxidation with (e.g., phosphite oxidation methods known in the art, e.g., a sulfurizing agent, such as 3-((N,N-dimethylaminomethylidene)amino)-3H-1,2,4-dithiazole-5-thione) and hydrolysis of the cyanoethyl group may produce a compound consisting of CpG oligonucleotide that, at its 5' end, is bonded or linked by a linker to the lipid:



or a salt thereof,

[0055] wherein X is O or S.

Dosing

[0056] The dose of each KRAS peptide administered can be in the range of 100 to 5000 μg per peptide (e.g., 700 μg /peptide, which would yield a peptide dose of 4900 μg for a group of 7 peptides).

[0057] If an adjuvant is administered with the KRAS peptide, the dose of the adjuvant can be, on a body weight basis, as high as 0.48 mg/kg. Exemplary dosing regimens are shown below in Table 3.

TABLE 3

Dosing	Adjuvant (e.g. Amph-CpG-7909; FIG. 6)	Peptide
Dose 1 (low)	0.1 mg	700 μg for each of 7 peptides
Dose 2 (medium)	0.5 mg	700 μg for each of 7 peptides
Dose 3 (high)	2.5 mg	700 μg for each of 7 peptides

[0058] In order that this invention be more fully understood, the following examples are set forth. These examples are for the purpose of illustration only and are not to be construed as limiting the scope of the invention in any way.

Examples

Example 1: Amphiphile KRas G12D Elicits an Immune Response

[0059] The efficacy of soluble-KRas (KRas) or amphiphile-KRas (aKRas) (FIG. 6) mutant sequences and an amphiphile-CpG adjuvant (aCpG) to elicit an immune response was determined. In these experiments five-times as much aCpG as amphiphile-KRas was used because CpG escalation experiments suggested that this dose may be more efficacious.

[0060] The experimental design included the following 4 groups of C57BL/6 mice immunized with the listed compounds ($n=5$ for each group):

[0061] 1. KRas G12D+ aCpG

[0062] 2. aKRasG12D+ aCpG

[0063] 3. KRas G12D+ pIC

[0064] 4. No immunization

[0065] PolyIC (pIC) was used as a benchmark adjuvant control.

[0066] The adjuvant, as well as peptide stock, solutions were dissolved in H_2O . Final injections were diluted with 1xPBS (phosphate buffered saline).

[0067] The aKRas peptides used were 18mer sequences of mutant sequences with a G12D substitution (amino acids 4-21 of wild-type+YKLVVVGAGGVGKSALTI (SEQ ID NO:9)). 20 μg in 100 μl were used for each injection.

[0068] The aCpG sequence used was the CpG1826 sequence (5'-tccatgacgttcctgacgtt-3'; SEQ ID NO:10) with two guanines added at the 5' end (5'-gg tccatgacgttcctgacgtt-3'; SEQ ID NO:11) at a concentration of 5 nmol for each 100 μl injection. CpG1826 is an optimal mouse sequence while CpG7909 is optimal for humans and poorly active in mice. CpG1826 and CpG7909 are in the same CpG class (class B) and generally have similar activity profiles in their respective species.

[0069] 50 µg of pIC were used for each 100 µl injection.
 [0070] Primer immunizations were given subcutaneously (s.c.) into the tail base (day 0) with one booster dose after 2 weeks (day 14). ELISpot analysis for IFN γ on splenocytes was performed, using standard protocols, 8 days after booster dose administration (day 21). Splenocytes (10⁶ cells/well) were activated with 5 µg/well of G12D sequence 18mer (aa4-21)→YKLVVVGADGVGKSALTI (SEQ ID NO:1). As shown in FIG. 1, aKRas in combination with aCpG activated splenocytes while soluble KRas in combination with aCpG, and the untreated and soluble KRas and pIC control groups, did not.

Example 2: Amphiphile KRas G12R and G12V
 Elicit an Immune Response

[0071] The efficacy of additional soluble-KRas (KRas) or amphiphile-KRas (aKRas) mutant sequences and an amphiphile-CpG adjuvant (aCpG) or soluble CpG (CpG) to elicit an immune response was determined.

[0072] The experimental design included the following 7 groups of C57BL/6 mice immunized with the listed compounds (n=5 for each group):

- [0073] 1. KRas G12R+CpG
- [0074] 2. KRas G12V+CpG
- [0075] 3. KRas G12R+ pIC
- [0076] 4. KRas G12V+ pIC
- [0077] 5. aKRas G12R+ aCpG
- [0078] 6. aKRas G12V+ aCpG
- [0079] 7. No immunization

[0080] The adjuvant as well as peptide stock solutions are dissolved in H₂O. Final injections are diluted with 1×PBS.

[0081] The aKRas peptides used were 18mer sequences of mutant sequences with G12R/V substitutions (SEQ amino acids 4-21 of wild-type+YKLVVVGAGGVGKSALTI (SEQ ID NO:9)). 20 µg in 100 µl were used for each injection.

[0082] The aCpG sequence used was the CpG1826 sequence (5'-tccatgacgttctgacgtt-3'; SEQ ID NO:10) with two guanines added at the 5' end (5'-gg tccatgacgttctgacgtt-3'; SEQ ID NO:11) at a concentration of 5 nmol for each 100 µl injection.

[0083] 50 µg of pIC were used for each 100 µl injection.

[0084] Primer immunizations were given subcutaneously (s.c.) into the tail base with one booster dose after 2 weeks (d14).

[0085] ELISpot analysis for IFN γ on splenocytes was performed, using standard protocols, 7 days after booster dose administration (d21). Splenocytes (10⁶ cells/well) were

activated with 5 µg/well of either: the G12V sequence 18mer (aa4-21)→YKLVVVGAVGVGKSALTI (SEQ ID NO:2) or the G12R sequence 18mer (aa4-21)→YKLVVVGARGVGKSALTI (SEQ ID NO:3).

[0086] As shown in FIG. 2, both aKRas G12R and G12V in combination with aCpG activated splenocytes while neither soluble KRasG12R nor G12V in combination with aCpG, and the untreated and soluble KRas and pIC control groups, did not.

Example 3: Amphiphile KRas G12D Elicits a
 CD8+ T-cell Immune Response

[0087] In wild-type B6 mice it is difficult to elicit a Kras-specific CD8+ T-cell response. B6 mice transgenic for the human HLA gene, A2.1, were immunized either with the soluble or amphiphile-conjugated forms of the KRas G12D antigen, as well as adjuvant (CpG or aCpG). The A2.1 allele is known to mount responses to the KRas peptides in humans.

[0088] The experimental design included the following 4 groups of B6 HLA-A2.1 Tg mice immunized with the listed compounds (n=5 for each group):

- [0089] 1. KRas wild-type+CpG
- [0090] 2. KRas12D+CpG
- [0091] 3. aKRas 12D+ aCpG
- [0092] 4. No immunization

[0093] The adjuvant as well as peptide stock solutions were dissolved in H₂O. Final injections were diluted with 1×PBS.

[0094] The aKRas peptides used were 18mer sequences of mutant sequences with a G12D substitution (amino acids 4-21 of wild-type+YKLVVVGAGGVGKSALTI (SEQ ID NO:9)). 20 µg in 100 µl were used for each injection.

[0095] The aCpG sequence used was the CpG1826 sequence (5'-tccatgacgttctgacgtt-3'; SEQ ID NO:10) with two guanines added at the 5' end (5'-gg tccatgacgttctgacgtt-3'; SEQ ID NO:11) at a concentration of 5 nmol for each 100 µl injection.

[0096] The vaccine was administered subcutaneously to female mice bilaterally at the tail base at 50 µl per side. Two booster doses were given in two-week intervals.

[0097] Intracellular cytokine staining (ICS) analysis for IFN γ on peripheral blood mononuclear cells (PBMCs) and ELISpot analysis for IFN γ on splenocytes was performed, using standard protocols, 7 days after the second and third booster dose, respectively.

[0098] PBMCs/Splenocytes (10⁶ cells/well and 2×10⁶ cells/well) were activated with 5 µg/well of either SEQ ID NO:1 or one of SEQ ID NOS:12-20 as shown below:

1. 12D sequence 18 mer (aa4-21)	→YKLVVVGADGVGKSALTI	(MHCII/CD4 epitope)
2. 12D sequence 17 mer (aa5-21)	→KLVVVGADGVGKSALTI	
3. 12D sequence 10 mer (aa5-14)	→KLVVVGADGV	(MHCI/CD8 epitope)
4. 12D sequence 9 mer (aa6-14)	→LVVVGADGV	
5. 12D sequence 9 mer (aa7-15)	→VVVGADGVG	
6. 12D sequence 9 mer (aa8-16)	→VVGADGVGK	
7. 12D sequence 9 mer (aa9-17)	→VGADGVGKS	
8. 12D sequence 9 mer (aa10-18)	→GADGVGKSA	
9. 12D sequence 9 mer (aa11-19)	→ADGVGKSAL	
10. 12D sequence 9 mer (aa12-20)	→DGVGKSALT	

[0099] Specific mutant peptides are given only to the mice that were immunized with the corresponding 18mer.

[0100] FIG. 3 shows a KRas-specific CD8+ T-cell response for mice immunized with the combination of aKRas G12D and aCpG, but not for soluble KRas G12D in combination with CpG or soluble wild-type KRas in combination with CpG.

Example 4: Analysis for Higher Antigen Concentration and Different Dose Intervals

[0101] The concentration of antigen was increased from 20 µg to 50 µg and bi-weekly (bw) or weekly (w) dose intervals were tested.

[0102] The experimental design included the following 5 groups of C57BL/6 mice immunized with the listed compounds (n=5 for each group):

[0103] 1. KRas 12D+CpG1826 (bw)

[0104] 2. KRas 12D+CpG1826 (w)

[0105] 3. aKRas 12D+ aCpG1826 (bw)

[0106] 4. aKRas 12D+ aCpG1826 (w)

[0107] 5. No immunization

[0108] The adjuvant as well as peptide stock solutions were dissolved in H₂O. Final injections were diluted with 1×PBS.

[0109] The aKRas peptides used were 18mer sequences of mutant sequences with a G12D substitution (amino acids 4-21 of wild-type+YKLVVVGAGGVGKSALTI (SEQ ID NO:9)). 50 µg in 100 µl were used for each injection.

[0110] The aCpG sequence used was the CpG1826 sequence (5'-tccatgacgttctctgacgtt-3'; SEQ ID NO:10) with two guanines added at the 5' end (5'-gg tccatgacgttctctgacgtt-3'; SEQ ID NO:11) at a concentration of 5 nmol for each 100 µl injection.

[0111] Primer immunizations were given subcutaneously (s.c.) into the tail base with one booster dose after 2 weeks.

[0112] ELISpot analysis for IFN γ , and CBA (cytometric bead array) analysis for IL6, IL10, IL12, TNF α , IFN γ , and MCP1 was performed on splenocytes, using standard protocols, 7 days after booster dose administration. Splenocytes (10⁶ cells/well) were activated with 5 µg/well of either SEQ ID NO:1 or one of SEQ ID NOS:12-21 as shown below:

[0113] Specific mutant peptides were given only to the mice that were immunized with the corresponding 18mer.

[0114] Separate plates were prepared for either short (9mer and 10mer) or long (17mer and 18mer) peptide stimulations. Cells from "No immunization" control mice were stimulated with all stimuli.

[0115] FIG. 4 shows a KRas-specific CD8+ T-cell response (9mers) and CD8+ or CD4+ T-cell response (18mer) for mice immunized with the combination of aKRas G12D and aCpG, but not for soluble KRas G12D in combination with CpG or the untreated control.

Example 5: Further Analysis of KRAS Amphiphiles

[0116] As described in the above Examples, the potential immunogenicity of a mutant KRAS peptide -amphiphile vaccine with aCpG adjuvant has been demonstrated in a C57BL/6 mouse model. These studies, and further studies, are summarized in the Table 4 and demonstrate that although mice historically mount poorer immunologic responses than humans to mutant KRAS peptides, due to species differences in MHC binding,

[0117] subcutaneous administration of mutated KRAS Amph-Peptides (G12D (FIG. 6), G12R, and G12V) is effective at inducing KRAS-specific splenic T-cells in mice as identified by ELISPOT and cytokine bead array (FIG. 1, FIG. 2, FIG. 4, and FIG. 5)

[0118] mutated KRAS Amph-Peptides (G12D, G12R, and G12V) combined with aCpG as the adjuvant are more effective at inducing KRAS-specific splenic T-cells than mutated KRAS peptides combined with either soluble CpG or polyinosinic:polycytidylic acid (poly IC) as the adjuvant (FIG. 1, FIG. 2, and FIG. 5) and

[0119] mutated KRAS Amph--Peptides (G12D) combined with aCpG are more effective than soluble KRAS peptides combined with soluble CpG at eliciting both CD4 and CD8 T-cell responses as demonstrated through effector cytokine response upon restimulation with either long (18-mer) or minimal (9-mer) peptide epitopes (FIG. 4).

1. 12D sequence 18 mer (aa4-21)	→YKLVVVGADGVGKSALTI	(MHCII/CD4 epitope)
2. 12D sequence 17 mer (aa5-21)	→KLVVVGADGVGKSALTI	
3. 12D sequence 10 mer (aa5-14)	→KLVVVGADGV	(MHCI/CD8 epitope)
4. 12D sequence 9 mer (aa6-14)	→LVVVGADGV	
5. 12D sequence 9 mer (aa7-15)	→VVVGADGVG	
6. 12D sequence 9 mer (aa8-16)	→VVGADGVGK	
7. 12D sequence 9 mer (aa9-17)	→VGADGVGKS	
8. 12D sequence 9 mer (aa10-18)	→GADGVGKSA	
9. 12D sequence 9 mer (aa11-19)	→ADGVGKSAL	
10. 12D sequence 9 mer (aa12-20)	→DGVGKSALT	

TABLE 4

Test Article Doses/Dose Volume	ROA/Treatment Regimen	Results
4 groups: 1) KRAS G12D + CpG; 2) Amph-KRAS G12D + aCpG; 3) KRAS G12D + Poly IC; 4) No Treatment KRAS peptides (amphiphilic or soluble) were dosed at 20 µg; aCpG-1826 at 5 nmoL; and Poly IC at 50 µg; stock solutions were dissolved in water and final dose solutions in phosphate- buffered saline (PBS). Dose volume: 100 µL administered divided as a SC injection on either side of the tail head (50 µL per injection site)	ROA: SC Prime: Day 0 Boost: Day 14	ELISPOT (IFN γ) Results: Amph-Peptide G12D + aCpG produced superior splenic T-cell responses when compared to either soluble G12D + soluble CpG or Poly IC
7 groups: 1) KRAS G12R + CpG; 2) KRAS G12V + CpG 3) KRAS G12R + Poly IC; 4) KRAS G12V + Poly IC; 5) Amph-KRAS G12R + aCpG; 6) Amph-KRAS G12V + aCpG; 7) No Treatment KRAS peptides (amphiphilic or soluble) were dosed at 20 µg; aCpG-1826 at 5 nmoL; and Poly IC at 50 µg; stock solutions were dissolved in water and final dose solutions in PBS. Dose volume: 100 µL administered divided as a SC injection on either side of the tail head (50 µL per injection site)	ROA: SC Prime: Day 0 Boost: Day 14	ELISPOT (IFN γ) Results: KRAS peptide sequences are more immunogenic when conjugated to a lipid moiety than are soluble peptides
5 groups: 1) KRAS G12D + CpG (bw); 2) KRAS G12D + CpG (w); 3) Amph-KRAS G12D + aCpG (bw); 4) Amph-KRAS G12D + aCpG (w); 5) No treatment KRAS peptides (amphiphilic or soluble) were dosed at 50 µg; aCpG-1826 at 5 nmol; stock solutions were dissolved in water and final dose solutions in PBS. Dose volume: 100 µL administered divided as a SC injection on either side of the tail head (50 µL per injection site)	ROA: SC 2 doses (prime + boost) were administered for the animals treated biweekly (bw) and 3 doses (prime + 2 boosts) were administered to the animals treated weekly (w): Prime: Day 0 (w and bw) Boost: Day 7 (w) Boost: Day 14 (w and bw)	ELISPOT (IFN γ) and CBA Results: Amphiphiles elicited endogenous CD8 and CD4 T-cell responses against mutant KRAS in C57BL/6 mice as assessed by both ELISPOT (IFN γ) and/or cytokine bead array (IFN γ , TNF α and IL-6)

CBA = cytometric bead array;

ELISPOT = enzyme-linked immunospot;

IFN γ = interferon gamma;

IL-6 = interleukin 6;

KRAS = Kirsten Rat Sarcoma;

poly IC = polyinosinic:polycytidylic acid;


ROA = route of administration;

SC = subcutaneous;

TNF α = tumor necrosis factor alpha

OTHER EMBODIMENTS

[0120] While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the



invention and including such departures from the present disclosure come within known or customary practice within the art to which the invention pertains and may be applied to the essential features hereinbefore set forth.

[0121] All publications, patents, and patent applications are herein incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety.

[0122] Some embodiments of the invention are within the following numbered paragraphs. 1. A compound comprising a mutant KRAS sequence and a lipid, wherein the mutant KRAS sequence is conjugated to the lipid by a linker, and (i) the linker comprises one or more polyethylene glycol blocks, (ii) the lipid is 1,2-distearoyl-sn-glycero-3-phosphoethanolamine (DSPE), and (iii) the mutant KRAS sequence comprises or consists of an amino acid sequence selected from the group 15 consisting of YKL V V V G A D G V G K S A L T I (SEQ ID NO:23), YKL V V V G A V G V G K S A L T I (SEQ ID NO:24), YKL V V V G A R G V G K S A L T I (SEQ ID NO:25), YKL V V V G A A G V G K S A L T I (SEQ ID NO:26), YKL V V V G A S G V G K S A L T I (SEQ ID NO:27), YKL V V V G A C G V G K S A L T I (SEQ ID NO:28), YKL V V V G A T G V G K S A L T I (SEQ ID NO:29), and YKL V V V G A G D V G K S A L T I (SEQ ID NO:30).

[0123] 2. A compound comprising a mutant KRAS sequence and a lipid, wherein the mutant KRAS sequence is conjugated to the lipid by a linker, and (i) the linker comprises one or more polyethylene glycol blocks, (ii) the lipid is 1,2-distearoyl-sn-glycero-3-phosphoethanolamine (DSPE), and (iii) the mutant KRAS sequence comprises or consists of an amino acid sequence selected from the group consisting of CYKL V V V G A D G V G K S A L T I (SEQ ID NO:1), CYKL V V V G A V G V G K S A L T I (SEQ ID NO:2), CYKL V V V G A R G V G K S A L T I (SEQ ID NO:3), CYKL V V V G A A G V G K S A L T I (SEQ ID NO:4), CYKL V V V G A S G V G K S A L T I (SEQ ID NO:5), CYKL V V V G A C G V G K S A L T I (SEQ ID NO:6), CYKL V V V G A T G V G K S A L T I (SEQ ID NO:22), and CYKL V V V G A G D V G K S A L T I (SEQ ID NO:7).

[0124] 3. The compound of paragraph 1 or 2, wherein the mutant KRAS sequence, at its N-terminus, is conjugated to the linker through a cysteine-maleimide linkage.


[0125] 4. The compound of any one of paragraphs 1-3, wherein linker comprises 48 repeat units of polyethylene glycol.

[0126] 5. The compound of paragraph 1 or 2, wherein the mutant KRAS sequence, at its N-terminus, is conjugated to the following structure:

[0127] 6. A composition comprising one or more compounds of any one of paragraphs 1-5 and a pharmaceutically acceptable carrier.

[0128] 7. The composition of paragraph 6, wherein the composition comprises (1) a compound comprising the amino acid sequence YKL V V V G A D G V G K S A L T I (SEQ ID NO:23), (2) a compound comprising the amino acid sequence YKL V V V G A V G V G K S A L T I (SEQ ID NO:24), (3) a compound comprising the amino acid sequence YKL V V V G A R G V G K S A L T I (SEQ ID NO:25), (4) a compound comprising the amino acid sequence YKL V V V G A A G V G K S A L T I (SEQ ID NO:26), (5) a compound comprising the amino acid sequence YKL V V V G A S G V G K S A L T I (SEQ ID NO:27), (6) a compound comprising the amino acid sequence YKL V V V G A C G V G K S A L T I (SEQ ID NO:28) or a compound comprising the amino acid sequence YKL V V V G A T G V G K S A L T I (SEQ ID NO:29), and (7) a compound comprising the amino acid sequence YKL V V V G A G D V G K S A L T I (SEQ ID NO:30).

[0129] 8. The composition of paragraph 6, wherein the composition comprises (1) a compound comprising the amino acid sequence CYKL V V V G A D G V G K S A L T I (SEQ ID NO:1), (2) a compound comprising the amino acid sequence CYKL V V V G A V G V G K S A L T I (SEQ ID NO:2), (3) a compound comprising the amino acid sequence CYKL V V V G A R G V G K S A L T I (SEQ ID NO:3), (4) a compound comprising the amino acid sequence CYKL V V V G A A G V G K S A L T I (SEQ ID NO:4), (5) a compound comprising the amino acid sequence CYKL V V V G A S G V G K S A L T I (SEQ ID NO:5), (6) a compound comprising the amino acid sequence CYKL V V V G A C G V G K S A L T I (SEQ ID NO:6) or a compound comprising the amino acid sequence CYKL V V V G A T G V G K S A L T I (SEQ ID NO:22), and (7) a compound comprising the amino acid sequence CYKL V V V G A G D V G K S A L T I (SEQ ID NO:7). 9. The composition of any one of paragraphs 6-8, wherein the composition further comprises a compound consisting of the nucleotide sequence 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3' (SEQ ID NO:8), which, at its 5' end, is bonded or linked by a linker to the following lipid:



or a salt thereof,
wherein X is O or S.

[0130] 10. The composition of paragraph 6, wherein the composition comprises 700 µg of each compound.

[0131] 11. A method of treating a cancer in a human patient, the method comprising administering the composition of any one of paragraphs 6-10 to the patient.


[0132] 12. The method of paragraph 11, wherein the method further comprises administering an adjuvant.

[0133] 13. The method of paragraph 12, wherein the adjuvant comprises a CpG nucleotide sequence.

[0134] 14. The method of paragraph 13, wherein the CpG nucleotide sequence comprises 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3' (SEQ ID NO: 8).

[0135] 15. The method of any one of paragraphs 12-14, wherein 0.1 mg, 0.5 mg, or 2.5 mg of the adjuvant is administered.


[0136] 16. The method of paragraph 10, wherein the method further comprises administering to the patient a compound consisting of the nucleotide sequence 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3' (SEQ ID NO:8), which, at its 5' end, is bonded or linked by a linker to the following lipid:



or a salt thereof,
wherein X is O or S.

[0137] 17. The composition of paragraph 9 or the method of paragraph 16, wherein the nucleotide sequence is bonded to the lipid.

[0138] 18. The method of paragraph 16 or 17, wherein 0.1 mg, 0.5 mg, or 2.5 mg of the compound consisting of the nucleotide sequence 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3' (SEQ ID NO:8), which, at its 5' end, is bonded or linked by a linker to the following lipid:




or a salt thereof,
wherein X is O or S,
is administered.

[0139] 19. The method of any one of paragraphs 11-16 or 18, wherein the cancer is a pancreatic cancer, a lung cancer, or a colorectal cancer.

[0140] 20. The method of any one of paragraphs 11-16 or 18, wherein all internucleoside groups connecting the nucleosides in 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3' (SEQ ID NO:8) are phosphorothioates.

[0141] 21. A kit comprising (i) a compound of any one of paragraphs 1-5, or a composition of any one of paragraphs 6-8, and (ii) a compound consisting of the nucleotide sequence 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3' (SEQ ID NO:8), which, at its 5' end, is bonded or linked by a linker to the following lipid:



or a salt thereof,
wherein X is O or S.

[0142] Other embodiments are within the following claims.

SEQUENCE LISTING

Sequence total quantity: 31
 SEQ ID NO: 1 moltype = AA length = 19
 FEATURE Location/Qualifiers
 source 1..19
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 1
 CYKLVVVGAD GVGKSALTI

-continued

SEQ ID NO: 2	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 2		
CYKLVVVGAV GVGKSALTI		19
SEQ ID NO: 3	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 3		
CYKLVVVGAR GVGKSALTI		19
SEQ ID NO: 4	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 4		
CYKLVVVGAA GVGKSALTI		19
SEQ ID NO: 5	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 5		
CYKLVVVGAS GVGKSALTI		19
SEQ ID NO: 6	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 6		
CYKLVVVGAC GVGKSALTI		19
SEQ ID NO: 7	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 7		
CYKLVVVGAG DVGKSALTI		19
SEQ ID NO: 8	moltype = DNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 8		
tcgtcgtttt gtcgttttgc cgtt		24
SEQ ID NO: 9	moltype = AA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 9		
YKLVVVGAGG VGKSALTI		18
SEQ ID NO: 10	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 10		
tccatgacgt tcctgacgtt		20
SEQ ID NO: 11	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	

-continued

	mol_type = other DNA organism = synthetic construct	
SEQUENCE: 11 ggtccatgac gttcctgacg tt		22
SEQ ID NO: 12 FEATURE source	moltype = AA length = 18 Location/Qualifiers 1..18 mol_type = protein organism = synthetic construct	
SEQUENCE: 12 YKLVVVGADG VGKSALTI		18
SEQ ID NO: 13 FEATURE source	moltype = AA length = 17 Location/Qualifiers 1..17 mol_type = protein organism = synthetic construct	
SEQUENCE: 13 KLVVVGADGV GKSALTI		17
SEQ ID NO: 14 FEATURE source	moltype = AA length = 10 Location/Qualifiers 1..10 mol_type = protein organism = synthetic construct	
SEQUENCE: 14 KLVVVGADGV		10
SEQ ID NO: 15 FEATURE source	moltype = AA length = 9 Location/Qualifiers 1..9 mol_type = protein organism = synthetic construct	
SEQUENCE: 15 LVVVGADGV		9
SEQ ID NO: 16 FEATURE source	moltype = AA length = 9 Location/Qualifiers 1..9 mol_type = protein organism = synthetic construct	
SEQUENCE: 16 VVVGADGVG		9
SEQ ID NO: 17 FEATURE source	moltype = AA length = 9 Location/Qualifiers 1..9 mol_type = protein organism = synthetic construct	
SEQUENCE: 17 VVGADGVGK		9
SEQ ID NO: 18 FEATURE source	moltype = AA length = 9 Location/Qualifiers 1..9 mol_type = protein organism = synthetic construct	
SEQUENCE: 18 VGADGVGKS		9
SEQ ID NO: 19 FEATURE source	moltype = AA length = 9 Location/Qualifiers 1..9 mol_type = protein organism = synthetic construct	
SEQUENCE: 19 GADGVGKSA		9
SEQ ID NO: 20 FEATURE source	moltype = AA length = 9 Location/Qualifiers 1..9 mol_type = protein organism = synthetic construct	
SEQUENCE: 20 ADGVGKSAL		9

-continued

SEQ ID NO: 21	moltype = AA length = 9	
FEATURE	Location/Qualifiers	
source	1..9	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 21		
DGVGKSALT		9
SEQ ID NO: 22	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 22		
CYKLVVVGAT GVGKSALTI		19
SEQ ID NO: 23	moltype = AA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 23		
YKLVVVGADG VGKSALTI		18
SEQ ID NO: 24	moltype = AA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 24		
YKLVVVGAVG VGKSALTI		18
SEQ ID NO: 25	moltype = AA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 25		
YKLVVVGARG VGKSALTI		18
SEQ ID NO: 26	moltype = AA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 26		
YKLVVVGAAAG VGKSALTI		18
SEQ ID NO: 27	moltype = AA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 27		
YKLVVVGASG VGKSALTI		18
SEQ ID NO: 28	moltype = AA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 28		
YKLVVVGACG VGKSALTI		18
SEQ ID NO: 29	moltype = AA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 29		
YKLVVVGATG VGKSALTI		18
SEQ ID NO: 30	moltype = AA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	

-continued

	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 30		
YKLVVVGAGD VGKSALTI		18
SEQ ID NO: 31	moltype = AA length = 9	
FEATURE	Location/Qualifiers	
source	1..9	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 31		
KLVVVGADG		9

What is claimed is:


1. A compound comprising a mutant KRAS sequence and a lipid, wherein the mutant KRAS sequence is conjugated to the lipid by a linker, and (i) the linker comprises one or more polyethylene glycol blocks, (ii) the lipid is 1,2-distearoyl-sn-glycero-3-phosphoethanolamine (DSPE), and (iii) the mutant KRAS sequence consists of an amino acid sequence selected from the group consisting of CYKLVVVGADGVGKSALTI (SEQ ID NO:1), CYKLVVVGAVGVGKSALTI (SEQ ID NO:2), CYKLVVVGARGVGKSALTI (SEQ ID NO:3), CYKLVVVGAAGVGKSALTI (SEQ ID NO:4), CYKLVVVGASGVGKSALTI (SEQ ID NO:5), CYKLVVVGACGVGKSALTI (SEQ ID NO:6), CYKLVVVGATGVGKSALTI (SEQ ID NO:22), and CYKLVVVGAGDVGKSALTI (SEQ ID NO:7).

2. The compound of claim 1, wherein the mutant KRAS sequence consists of an amino acid sequence selected from the group consisting of YKLVVVGADGVGKSALTI (SEQ ID NO:23), YKLVVVGAVGVGKSALTI (SEQ ID NO:24), YKLVVVGARGVGKSALTI (SEQ ID NO:25), YKLVVVGAAGVGKSALTI (SEQ ID NO:26), YKLVVVGASGVGKSALTI (SEQ ID NO:27), YKLVVVGACGVGKSALTI (SEQ ID NO:28), YKLVVVGATGVGKSALTI (SEQ ID NO:29), and YKLVVVGAGDVGKSALTI (SEQ ID NO:30).

3. The compound of claim 1 or 2, wherein the mutant KRAS sequence, at its N-terminus, is conjugated to the linker through a cysteine-maleimide linkage.

4. The compound of claim 1 or 2, wherein linker comprises 48 repeat units of polyethylene glycol.

5. The compound of claim 1 or 2, wherein the mutant KRAS sequence, at its N-terminus, is conjugated to the following structure:



6. A composition comprising one or more compounds of claim 1 or 2, and a pharmaceutically acceptable carrier.


7. The composition of claim 6, wherein the composition comprises (1) a compound comprising the amino acid sequence YKLVVVGADGVGKSALTI (SEQ ID NO:23), (2) a compound comprising the amino acid sequence

YKLVVVGAVGVGKSALTI (SEQ ID NO:24), (3) a compound comprising the amino acid sequence YKLVVVGARGVGKSALTI (SEQ ID NO:25), (4) a compound comprising the amino acid sequence YKLVVVGAAGVGKSALTI (SEQ ID NO:26), (5) a compound comprising the amino acid sequence YKLVVVGASGVGKSALTI (SEQ ID NO:27), (6) a compound comprising the amino acid sequence YKLVVVGACGVGKSALTI (SEQ ID NO:28) or a compound comprising the amino acid sequence YKLVVVGATGVGKSALTI (SEQ ID NO:29), and (7) a compound comprising the amino acid sequence YKLVVVGAGDVGKSALTI (SEQ ID NO:30).

8. The composition of claim 6, wherein the composition comprises (1) a compound comprising the amino acid sequence CYKLVVVGADGVGKSALTI (SEQ ID NO:1), (2) a compound comprising the amino acid sequence CYKLVVVGAVGVGKSALTI (SEQ ID NO:2), (3) a compound comprising the amino acid sequence CYKLVVVGARGVGKSALTI (SEQ ID NO:3), (4) a compound comprising the amino acid sequence CYKLVVVGAAGVGKSALTI (SEQ ID NO:4), (5) a compound comprising the amino acid sequence CYKLVVVGASGVGKSALTI (SEQ ID NO:5), (6) a compound comprising the amino acid sequence CYKLVVVGACGVGKSALTI (SEQ ID NO:6) or a compound comprising the amino acid sequence CYKLVVVGATGVGKSALTI (SEQ ID NO:22), and (7) a compound

comprising the amino acid sequence CYKLVVVGAGDVGKSALTI (SEQ ID NO:7).

9. The composition of claim 6, wherein the composition further comprises a compound consisting of the nucleotide sequence 5'-TCGTCGTTTTGTCGTTTTGTCGTT-3' (SEQ ID NO:8), which, at its 5' end, is bonded or linked by a linker to the following lipid:



or a salt thereof,
wherein X is O or S.

10. The composition of claim 6, wherein the composition comprises 700 μ g of each compound.

11. The composition of claim 9, wherein the nucleotide sequence is bonded to the lipid.

12. A method of treating a cancer in a human patient, the method comprising administering the composition of claim 6 the patient.


13. The method of claim 12, wherein the method further comprises administering an adjuvant.

14. The method of claim 13, wherein the adjuvant comprises a CpG nucleotide sequence.

15. The method of claim 14, wherein the CpG nucleotide sequence comprises 5'-TCGTCGTTTGTGCGTTTGTGCGTT-3' (SEQ ID NO: 8).

16. The method of claim 12, wherein 0.1 mg, 0.5 mg, or 2.5 mg of the adjuvant is administered.


17. The method of claim 12, wherein the method further comprises administering to the patient a compound consisting of the nucleotide sequence 5'-TCGTCGTTTGTGCGTTTGTGCGTT-3' (SEQ ID NO:8), which, at its 5' end, is bonded or linked by a linker to the following lipid:



or a salt thereof,
wherein X is O or S.

18. The method of claim 17, wherein the nucleotide sequence is bonded to the lipid.

19. The method of claim 17, wherein 0.1 mg, 0.5 mg, or 2.5 mg of the compound consisting of the nucleotide sequence 5'-TCGTCGTTTGTGCGTTTGTGCGTT-3' (SEQ ID NO:8), which, at its 5' end, is bonded or linked by a linker to the following lipid:




or a salt thereof,
wherein X is O or S,
is administered.

20. The method of claim 12, wherein the cancer is a pancreatic cancer, a lung cancer, or a colorectal cancer.

21. The method of claim 12, wherein all internucleoside groups connecting the nucleosides in 5'-TCGTCGTTTGTGCGTTTGTGCGTT-3' (SEQ ID NO:8) are phosphorothioates.

22. A kit comprising (i) a compound of claim 1 or 2, and (ii) a compound consisting of the nucleotide sequence 5'-TCGTCGTTTGTGCGTTTGTGCGTT-3' (SEQ ID NO:8), which, at its 5' end, is bonded or linked by a linker to the following lipid:



or a salt thereof,
wherein X is O or S.

* * * * *